

Evaluation of protective effect of N-acetyl cysteine on arsenic-induced hepatotoxicity

Pudari Hemalatha,
Alla Gopala Reddy,
Yerradoddi Ramana
Reddy¹,
Pabbathi Shivakumar

Department of Veterinary Pharmacology and Toxicology, ¹Animal Nutrition, College of Veterinary Science, Rajendranagar, Hyderabad, Andhra Pradesh, India

Address for correspondence:

Prof. A. Gopala Reddy, Department of Veterinary Pharmacology and Toxicology, College of Veterinary Science, Rajendranagar, Hyderabad, Andhra Pradesh, India. E-mail: gopalareddy123@rediffmail.com

Abstract

Objective: The present study was aimed to study protective role of N-acetyl cysteine (NAC) was assessed against arsenic (As)-induced hepatotoxicity in rats. **Methods:** Twenty four male Wistar rats were divided into 4 groups of 6 animals each and treated as follows: Group 1: sham control, 2: arsenic control (sodium arsenite @ 10 mg/kg b. wt orally for 4 wks), 3: Pre-treatment with NAC (@ 300 mg/kg orally for 2 wks) followed by sodium arsenite along with NAC (as per above doses) and 4: Sodium arsenite + NAC (as per above doses for 4 wks). **Results:** The concentration of thiobarbituric acid reacting substances (TBARS) and protein carbonyls was significantly ($P<0.05$) increased, while the concentration of reduced glutathione (GSH), and the activity of CYP450, Na⁺ - K⁺ ATPase and Mg²⁺ ATPase in liver were significantly ($P<0.05$) reduced in group 2 as compared to control. Groups 3 and 4 revealed improvement in the parameters in study. **Conclusion:** The study revealed that arsenic induces hepatotoxicity by inducing oxidative stress and supplementation of NAC is beneficial in countering the adverse effects.

Key words: Arsenic, hepatotoxicity, N-acetyl cysteine

INTRODUCTION

Arsenic is a major global health concern due to its wide distribution and adverse health effects on humans, animals, birds, aquatic life and plants through polluted ground water and food chains. It is a known carcinogen that has been associated with cancers of the skin, lung, urinary bladder, and possibly liver, kidney and prostate in humans. Besides the natural sources, arsenic-contaminated herbicides, insecticides and rodenticides are also potential vehicles of arsenic toxicity.^[1] Organic arsenicals are generally considered non-toxic, whereas inorganic forms are toxic. Inorganic arsenic exists predominantly in trivalent (As³⁺) and pentavalent (As⁵⁺) forms, where trivalent compounds are more toxic than pentavalent ones.^[2] Liver is the target of arsenic-induced carcinogenesis.^[3] Arsenic-induced

global DNA hypomethylation was also seen in mouse livers chronically exposed to inorganic arsenic.^[4-7] The present work was undertaken in male *Wistar Kyoto* rats to study the protective effect of NAC on arsenic-induced hepatotoxicity.

MATERIALS AND METHODS

Male albino rats of *Wistar Kyoto* strain weighing about 200-220 g were procured from National Institute of Nutrition (NIN), Hyderabad, India. The animals were housed in solid bottom polypropylene cages. Animals were placed on commercial standard mash feed for rats (NIN, Hyderabad) and provided water *ad libitum*. Experimental protocol was approved by the Institutional Animal Ethics Committee. Rats were divided into four groups of 6 in each. Group 1 was kept as sham control, 2 was arsenic control (sodiumarsenite@10 mg/kg b. wt orally for 4 weeks), 3 was pre-treated with N-Acetyl cysteine (@ 300 mg/kg orally for 2 weeks) followed by sodium arsenite along with N-Acetyl cysteine (as per above doses for 4 weeks) and 4 was given sodium arsenite + N-acetyl cysteine (as per above doses for 4 weeks). The animals were then euthanized on 29th day and livers were immediately excised, rinsed

Access this article online

Quick Response Code:



Website:
www.jnsbm.org

DOI:
10.4103/0976-9668.116986

Table 1: Results of oxidative stress and enzymes in liver tissue

Group	TBARS (nmol MDA/mg protein)	Protein carbonyls (nmol/mg protein)	GSH (μ mol/mg protein)	CYP ₄₅₀ activity (nmol/mg microsomal protein)	Na ⁺ /K ⁺ ATPase activity (μ mol of Pi liberated/mg microsomal protein/30 min)	Mg ²⁺ ATPase activity (μ mol of Pi liberated/mg microsomal protein/30 min)
1	1.47±0.09 ^a	1.19±0.03 ^a	31.84±4.05 ^c	3.55±0.39 ^c	16.64±1.25 ^c	13.96±1.11 ^c
2	3.06±0.16 ^c	4.07±0.25 ^c	14.12±1.47 ^a	1.16±0.19 ^a	5.16±0.28 ^a	4.94±0.42 ^a
3	1.94±0.10 ^b	2.50±0.14 ^b	26.68±3.29 ^{bc}	2.55±0.20 ^b	14.62±0.92 ^{bc}	11.54±0.97 ^b
4	2.26±0.12 ^b	2.30±0.30 ^b	22.45±2.45 ^b	2.37±0.23 ^b	13.51±0.91 ^b	9.41±0.78 ^b

Values are Mean±SE, n=6; Means with different alphabets as superscripts differ significantly ($P < 0.05$), ANOVA (SPSS version 15)

with ice-cold physiological saline and stored at -20°C for further homogenization to estimate the concentration of thiobarbituric acid reacting substances (TBARS), protein carbonyls and reduced glutathione (GSH), and the activity of CYP₄₅₀, Na⁺-K⁺ ATPase and Mg²⁺ ATPase.

The data were subjected to statistical analysis by applying one way ANOVA using SPSS (version 15.0) and the means were compared by Duncan's multiple comparison test. Significance was set at $P < 0.05$.

RESULTS AND DISCUSSION

The concentration of TBARS (n mol MDA/mg protein) and protein carbonyls (n mol/mg protein) in liver showed a significant ($P < 0.05$) rise in Group 2 (3.06 ± 0.16 and 4.07 ± 0.25 , respectively) as compared to group 1 (1.47 ± 0.09 and 1.19 ± 0.03 , respectively). Groups 3 and 4 showed a significant ($P < 0.05$) decrease as compared to Group 2. The concentration of GSH in liver revealed a significant ($P < 0.05$) reduction in Group 2 ($14.12 \pm 1.47 \mu$ mol/mg protein) as compared to the remaining groups [Table 1].

Oxidative stress mediated by reactive oxygen species (ROS) and reactive nitrogen species (RNS) is the cause for arsenic toxicity.^[8] In the present study, concentration of TBARS and protein carbonyls were increased in the liver of arsenic toxic group suggesting an ongoing oxidative stress. Similar results were obtained by Demerdash *et al.*,^[9] Flora *et al.*,^[10] and Sharma *et al.*^[11] Arsenic produces oxidative damage by disturbing the prooxidant-antioxidant balance, because it has very high affinity for sulfhydryl groups in GSH (non-enzymatic antioxidant), which might have implications in the maintenance of thiol-disulfide balance.^[12] Arsenic also induces oxidative tissue damage through interference with GSH utilization.^[13] N-Acetylcysteine (NAC) is a thiol-containing antioxidant that has been used to reduce various conditions of oxidative stress. Its antioxidant action is attributed to GSH synthesis; therefore maintaining intracellular GSH levels^[14,15] and scavenging reactive oxygen species (ROS).^[16] It is also known as potent metal chelator.^[17] NAC has a strong ability to restore the impaired pro-oxidant/antioxidant balance in metal poisoning.

The activity of CYP₄₅₀ (n mol/mg microsomal protein), Na⁺-K⁺ ATPase (μ mol Pi liberated/mg microsomal protein/30 min) and Mg²⁺ ATPase (μ mol Pi liberated/mg microsomal protein/30 min) in liver revealed a significant ($P < 0.05$) reduction in Group 2 (1.16 ± 0.19 , 5.16 ± 0.28 and 4.94 ± 0.42 , respectively) as compared to Group 1 (3.55 ± 0.39 , 16.64 ± 1.25 and 13.96 ± 1.11 , respectively) [Table 1]. Groups 3 and 4 showed a significant ($P < 0.05$) increase in the activity of CYP₄₅₀ and ATPases as compared to Group 2. The altered activity of ATPases may be attributed to the membrane lipid peroxidation resulting in structural derangement. Earlier reports confirm that low levels of ATP in hepatic cells with arsenic and treatment with NAC improved the activity of Na⁺/K⁺ ATPases.^[10] The activity of CYP₄₅₀ was significantly reduced in Group 2 as compared to control in this study, which is in agreement with the findings of Noreault *et al.*^[18]

CONCLUSION

The study revealed that arsenic induces toxic effects in liver due to generation of reactive oxygen species with eventual oxidative damage. Supplementation of NAC is beneficial in preventing these toxic effects to certain extent. Pre-treatment with NAC was found more protective as compared to NAC co-treatment against arsenic-induced toxicity.

REFERENCES

- Gupta R, Flora SJ. Protective role of Aloe vera against arsenic toxicity in rats. *Phytother Res* 2005;19:23-8.
- Duker AA, Carranza EJ, Hale M. Arsenic geo chemistry and health. *Environ Int* 2005;31:631-41.
- Liu J, Waalkes MP. Liver is a target of arsenic carcinogenesis. *Toxicol Sci* 2008;105:24-32.
- Okoji RS, Yu RC, Maronpot RR, Froines JR. Sodium arsenite administration via drinking water increases genome-wide and Ha-ras DNA hypomethylation in methyl-deficient C57BL/6J mice. *Carcinogenesis* 2002;23:777-85.
- Chen H, Li S, Liu J, Diwan BA, Barrett JC, Waalkes MP. Chronic inorganic arsenic exposure induces hepatic global and individual gene hypomethylation: Implications for arsenic hepatocarcinogenesis. *Carcinogenesis* 2004;25:1779-86.
- Xie Y, Trouba KJ, Liu J, Waalkes MP, Germolec DR. Biokinetics and subchronic toxic effects of oral arsenite, arsenate monomethylarsonic

- acid, and dimethylarsinic acid in v-Ha-ras transgenic (Tg.AC) mice. *Environ Health Perspect* 2004;112:1255-63.
7. Xie Y, Liu J, Benbrahim-Tallaa L, Ward JM, Logsdon D, Diwan BA, *et al.* Aberrant DNA methylation and gene expression in livers of newborn mice transplacentally exposed to a hepatocarcinogenic dose of inorganic arsenic. *Toxicology* 2007;236:7-15.
 8. Von Ehrenstein OS, Poddar S, Yuan Y, Mazumder DG, Eskenazi B, Basu A, *et al.* Children's intellectual function in relation to arsenic exposure. *Epidemiology* 2007;18:44-51.
 9. El-Demerdash FM, Yousef MI, Radwan FM. Ameliorating effect of curcumin on sodium arsenite-induced oxidative damage and lipid peroxidation in different rat organs. *Food Chem Toxicol* 2009;47:249-54.
 10. Flora SJ, Mehta A, Gupta R. Prevention of arsenic-induced hepatic apoptosis by concomitant administration of garlic extracts in mice. *Chem Biol Interact* 2009;177:227-33.
 11. Sharma A, Sharma MK, Kumar M. Modulatory role of *Embllica officinalis* fruit extract against arsenic induced oxidative stress in Swiss albino mice. *Chem Biol Interact* 2009;180:20-30.
 12. Yamanaka K, Heseqwa A, Sawamuna R, Okada S. Cellular response to oxidative damage in lung induced by the administration of dimethylarsinic acid, a major metabolite of inorganic arsenics, in mice. *Toxicol Appl Pharmacol* 1991;108:205-13.
 13. Bhadauria S, Flora SJ. Response of arsenic-induced oxidative stress, DNA damage, and metal imbalance to combined administration of DMSA and monoisoamyl-DMSA during chronic arsenic poisoning in rats. *Cell Biol Toxicol* 2007;23:91-104.
 14. Moldeus P, Cotgreave IA. N-acetylcysteine. *Methods Enzymol* 1994;234:482-92.
 15. Gurer H, Ozunes H, Neal R, Spitz DR, Ercal N. Antioxidant effects of N-acetylcysteine and succimer in red blood cells from lead exposed rats. *Toxicology* 1998;120:181-9.
 16. Arouma OL, Halliwell B, Hoey BM, Butler J. The antioxidant action of N-acetyl cysteine in reaction with hydrogen peroxide, hydroxyl radical, superoxide and hypochlorous acid. *Free Radical Biol Med* 1989;6:593-7.
 17. Banner W, Koch M, Capin DM, Hopf SB, Chang S, Tong TG. Experimental chelation therapy in chromium, lead, and boron intoxication with N-acetylcysteine and other compounds. *Toxicol Appl Pharmacol* 1986;83:142-7.
 18. Noreault LT, Kostrubsky VE, Wood SG, Nichols RC, Strom SC, Trask HW, *et al.* Arsenite decreases CYP3A4 and RXR in primary human hepatocytes. *Drug Metab Dispos* 2005;33:993-1003.

How to cite this article: Hemalatha P, Reddy AG, Reddy YR, Shivakumar P. Evaluation of protective effect of N-acetyl cysteine on arsenic-induced hepatotoxicity. *J Nat Sc Biol Med* 2013;4:393-5.

Source of Support: Nil. **Conflict of Interest:** None declared.