



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



## Short Communication

## *In vitro* protective efficacy of Lithium chloride against *Mycoplasma hyopneumoniae* infection



Hassan Z.A. Ishag<sup>a,b</sup>, Yu-zi Wu<sup>a</sup>, Mao-jun Liu<sup>a</sup>, Qi-yan Xiong<sup>a</sup>, Zhi-xin Feng<sup>a</sup>,  
Ruo-song Yang<sup>a</sup>, Guo-qing Shao<sup>a,\*</sup>

<sup>a</sup> Institute of Veterinary Medicine, Jiangsu Academy of Agricultural Sciences, Key Laboratory of Veterinary Biological Engineering and Technology, Ministry of Agriculture, National Research Center for Engineering and Technology of Veterinary Bio-products, Nanjing 210014, China

<sup>b</sup> College of Veterinary Sciences, University of Nyala, Nyala, Sudan

## ARTICLE INFO

## Article history:

Received 15 September 2015

Received in revised form 8 March 2016

Accepted 28 March 2016

## Keywords:

*Mycoplasma hyopneumoniae*

LiCl

Inhibition

Apoptosis

## ABSTRACT

*Mycoplasma hyopneumoniae* (*M. hyopneumoniae*) infection affects the swine industry. Lithium chloride (LiCl), is a drug used to treat bipolar disorder and has also shown activity against bacterial and viral infections. Herein, we evaluated the antibacterial activity of LiCl on PK-15 cells infected with *M. hyopneumoniae*. Incubation of LiCl (40 mM) with cells for 24 h, did not significantly affect the cell viability. The qRT-PCR showed ~80% reduction in *M. hyopneumoniae* genome when LiCl added post-infection. A direct effect of LiCl on bacteria was also observed. However, treatment of cells with LiCl prior infection, does not protect against the infection. Antibacterial activity of LiCl was further confirmed by IFA, which demonstrated a reduction in the bacterial protein. With 40 mM LiCl, the apoptotic cell death, production of nitric oxide and superoxide anion induced by *M. hyopneumoniae*, were prevented by ~80%, 60% and 58% respectively. Moreover, caspase-3 activity was also reduced (82%) in cells treated with 40 mM LiCl. LiCl showed activity against various strains of *M. hyopneumoniae* examined in our study. Collectively, our data showed that LiCl inhibited the infection of *M. hyopneumoniae* through anti-apoptotic mechanism.

© 2016 Elsevier Ltd. All rights reserved.

*Mycoplasma hyopneumoniae* (*M. hyopneumoniae*) causes porcine enzootic pneumonia (PEP) that affects the swine industry (Clark et al., 1991). Treatment of PEP is limited to antibiotics and vaccines that provide partial protection (Haesebrouck et al., 2004). Development of therapeutic agents against the infection is required.

Lithium chloride (LiCl) is a drug used to treat bipolar disorder (Bowden et al., 2005). It modulates several biological functions, including apoptosis, glycogen synthesis and inflammation (Machado-Vieira et al., 2009). LiCl active against viruses including avian coronavirus (Li et al., 2009) and transmissible gastroenteritis virus (TGEV) (Ren et al., 2011) in addition to bacteria such as *Francisella tularensis* (Zhang et al., 2009) and *Pseudomonas aeruginosa* (Chen et al., 2013). However, it un-clear whether LiCl could also inhibit the infection of *M. hyopneumoniae*.

Reactive oxygen species (ROS) (Beere and Green, 2001), nitric oxide (NO) (Brüne, 2003) and caspases, are important apoptosis inducers. LiCl was found to prevent apoptosis by inhibiting production of NO (Wang et al., 2013), ROS (Lumetti and Galli, 2014) and caspases (Ren et al., 2011). *M. hyopneumoniae* infection was reported to induce apoptosis (Bai et al., 2013). Thus, we aimed to investigate the inhibitory properties

of LiCl on PK-15 cells infected with *M. hyopneumoniae* and to underline the possible mechanism.

Swine tracheal epithelial cells (STEC) are the ideal model to study *Mycoplasma in vitro*. However, porcine kidney-15 (PK-15) cells model used in our study was previously established in our laboratory with time and titer above 4 h and  $1 \times 10^5$  CCU/ml (Che et al., 2013; Che et al., 2014; Che et al., 2012; Zhang et al., 2013) and was also described by others (Burnett et al., 2006; Wang et al., 2016).

*M. hyopneumoniae* isolates Xiaoling Wei-1 (XLW-1), Nanjing (NJ) and attenuated strain (168 L) were grown in KM2 medium (Bai et al., 2013). PK-15 cells were cultured in DMEM containing 10% FBS.

Cytotoxicity of LiCl (MW = 42.39, Sigma (U.S.A.)) on PK-15 cells was analyzed by MTT. We observed that LiCl at concentrations and time used in our had no toxic effect on PK-15 cells (data not shown).

Cells ( $1 \times 10^5$  cells/well) in 24-well plates were infected with *M. hyopneumoniae* ( $1 \times 10^6$  CFU/ml) at 37 °C for 10 h, washed and inoculated with LiCl (10–40 mM) for 12 h. After which, the cells were washed, total DNA was extracted and the effect of LiCl was evaluated by qRT-PCR following previous methods established in our laboratory (Wu et al., 2012). Briefly, the genome sequence of *M. hyopneumoniae* (GenBank CP002274.1) was used to PCR amplify 460 bp of p97 gene (GenBank: ADQ90328.1). This fragment was cloned in pMD18-T vector (pMD18-T-P97) to generate a log-linear standard curve, from which the DNA copy numbers was determined by qRT-PCR (based on

\* Corresponding author.

E-mail address: [gqshaojaas@163.com](mailto:gqshaojaas@163.com) (G. Shao).


Ct value). The qRT-PCR was performed on ABI PRISM 7500 Sequence Detection System (Applied Biosystems, USA) with SYBR<sup>®</sup> Premix Ex Taq<sup>™</sup> Kit (Takara). The qRT-PCR mixture (25 µl in total) consisted of 12.5 µl Premix Ex Taq<sup>™</sup> (2×), 0.5 µl ROX Reference Dye (50×), 1 µl Forward primer (10 µM), 1 µl Reverse primer (10 µM), 0.125 µl Probe, 9 µl ddH<sub>2</sub>O and 1 µl DNA while the cycling condition was set at 95 °C for 10 min followed by 40 cycles of 95 °C for 15 s and 60 °C for 60 s. An about 90 bp of p97 gene was detected using p97 primers Forward- 5' CCAGAACCAAATTCCTTCGCTG 3' and Reverse- 5' ACTGGCTGAACCTT ATCTGGGCTA 3' and p97 probe 5' FAM-AGCAGATCTTAGTCAAAGTGCC CGTG-TAMRA 3' which labeled at the 5' with the reporter dye 6-carboxyfluorescein (FAM) and at the 3' with a Black-Hole-Quencher.

We found that LiCl induce a dose-dependent decrease in the amount of bacterial DNA. At 40 mM, LiCl caused approximately 80% reduction in the relative amplification of p97 gene of *M. hyopneumoniae* (Fig. 1A). This was consistent with the inhibitory effect of LiCl on DNA synthesis of herpes simplex virus (Ziaie and Kefalides, 1989). Previous studies shown that *M. hyopneumoniae* introduced into PK-15 has infected the cell after 10 h (Che et al., 2014). Herein, we added LiCl to cells 10 hrs post-infection, indicating that, the inhibition of LiCl occurred during bacterial replication. We further investigated the direct effect of LiCl on cells or bacteria. LiCl-treated cells (1 × 10<sup>5</sup> cells/well) were infected with untreated bacteria (1 × 10<sup>6</sup> CFU/ml), and untreated cells were infected with LiCl-treated bacteria (1 × 10<sup>6</sup> CFU/ml). A direct effect of LiCl on bacteria was also observed but not on cells (Fig. 1B and Fig. C) which is consistent with the effect of LiCl on IBV (Harrison et al., 2007).


We further infected cells (1 × 10<sup>5</sup> cells/well) in 96-well plates with *M. hyopneumoniae* (1 × 10<sup>6</sup> CFU/ml) at 37 °C for 10 h. Cells washed and inoculated LiCl (10–40 mM) for 12 h. Expression level of *M. hyopneumoniae* surface antigen (p46) was evaluated by IFA using p46

monoclonal antibody (our laboratory) and FITC-conjugated goat anti-mouse secondary antibodies (Boster Biological Technology Company, China). Fluorescence was measured immediately at excitation 485 nm and emission of 535 nm using an infinite F200 Tecan Microplate Reader (Huppert et al., 2010), and the data were acquired using i-control software (Tecan Trading AG, Ma'nnedorf, Switzerland). We also observed that LiCl inhibited p46 expression (Fig. 1D) which indicated that LiCl treatment led to substantial inhibition of *M. hyopneumoniae* infection.

It has been reported that, *M. hyopneumoniae* infection could induce apoptosis in PBMC and STEC cells (Bai et al., 2015; Ni et al., 2014). As the infection model of *M. hyopneumoniae* with PK-15 cells was already developed in our laboratory, it's un-clear whether PK-15 cells undergoes apoptosis upon infection with this bacterium. In this study, by using AO/EB staining (Normal/Apoptotic/Necrotic Cell Detection kit, Beyotime Biotech., China), we observed that, the infection of PK-15 cells with this bacterium could also induce apoptosis in PK-15 cells (Fig. 2A). We further investigated the ability of LiCl to prevent *M. hyopneumoniae* induced apoptosis. Briefly, cells (1 × 10<sup>5</sup> cells/well) on 96-wells plate infected with *M. hyopneumoniae* (1 × 10<sup>6</sup> CFU/ml) for 10 h, washed and treated with or without LiCl (10–40 mM) for 12 h. Cells washed again, added 35 µl of AO/EB solution, examined by fluorescence microscopy and the apoptosis was quantified. Number of positive cells were calculated by selection of five view fields. Apoptotic rate was calculated as follows: apoptotic rate (%) = number of positive staining cells/number of total cells × 100%. Viable cells appeared green with intact nuclei. Nonviable cells had bright orange chromatin. Apoptotic cells appeared shrunk with condensation and fragmentation of nuclei. Apoptotic cells were distinguished from necrotic cells because the latter appeared orange with a normal nucleus (Renvoize et al.,



**Fig. 1.** Antibacterial activity of LiCl against *M. hyopneumoniae* (1 × 10<sup>6</sup> CCU/ml) infected PK-cells (1 × 10<sup>5</sup> cells/well). Cells infected with *M. hyopneumoniae*, XLW-1 for 10 h and treated with LiCl for 12 h. (A) DNA extracted and quantified by qRT-PCR (log<sub>10</sub> copies of DNA/well). Data presented as Means ± SD (n = 3). \*p < 0.008 and \*\*p < 0.001 vs. control. (B) *M. hyopneumoniae* treated with LiCl (direct inactivation) before inoculating into cells. (C) Cells treated with LiCl then infected with *M. hyopneumoniae*, XLW-1 (pre-infection). DNA was extracted and quantified by qRT-PCR as above. A direct effect on bacteria was observed but not in cells. Data presented as Means ± SD (n = 3) \*p < 0.05 and p < 0.54 vs. control. (D) Effect of LiCl treatment was evaluated by IFA. Cells washed, fixed and probed with primary (p46 monoclonal antibody) and secondary-FITC conjugated antibodies. FITC-fluorescence was measured at OD 535. \*\*p < 0.001 vs. control.



**Fig. 2.** Identification of anti-apoptotic mechanism of LiCl against PK-15 cells ( $1 \times 10^5$  cells/well) infected with *M. hyopneumoniae* ( $1 \times 10^6$  CCU/ml). PK-15 cells were infected with *M. hyopneumoniae*, XLW-1 for 10 h and treated with or without LiCl for 12 h. (A) Phenotype of apoptosis and anti-apoptotic effect of LiCl were analyzed by AO/EB staining. +Ve denotes for untreated cells while –Ve denotes for neither infected nor treated cells. Most representative fields are shown. (B) Calculated apoptosis rate is provided. Data presented as Means  $\pm$  SD ( $n = 3$ ). \*\* $P < 0.001$  vs. control. (C) LiCl treatment inhibits production of NO and (D) superoxide anion induced by *M. hyopneumoniae* infection. NO production and superoxide anion were evaluated by Griess assay and superoxide anion assay respectively in PK-15 cells infected with *M. hyopneumoniae* and treated with LiCl. Data presented as mean  $\pm$  SD ( $n = 3$ ). \* $P < 0.05$  and \*\*\* $P < 0.001$  vs. control. (E) LiCl inhibited caspase-3 activity. Caspase-3 activity was measured in PK-15 cell lysate infected with  $\sim 1 \times 10^6$  CCU/ml *M. hyopneumoniae* for 10 h prior to LiCl treatment. LiCl inhibited caspase-3 activity in a dose-dependent manner. Data presented as Means  $\pm$  SD ( $n = 3$ ). \*\* $P < 0.01$  vs. control.

1998). As shown in (Fig. 2A), *M. hyopneumoniae* induced apoptosis were significantly decreased in a dose-dependent manner after treatment with LiCl. Counting apoptotic cells at 24 hpi, confirmed this (Fig. 2B) and indicated  $\sim 80\%$  inhibition of apoptosis at 40 mM LiCl. These results imply that LiCl maybe a universal anti-apoptotic agent with limited selectivity to cell types and apoptosis inducers. Although the result is contradictory with studies on *Burkholderia pseudomallei* (Maniam et al., 2013) in which pretreatment of macrophages and A549 epithelial lung cells inhibited the infection, these data on post-infection treatment indicate that whether LiCl executes an anti-apoptotic or a pro-apoptotic effect is largely associated to the cell type and stimulating factor.

Low concentrations of NO inhibits apoptosis, and excessive production of NO triggers apoptosis (Shen et al., 1998). We evaluated whether LiCl inhibits production of NO in PK-15 cells infected with *M. hyopneumoniae* by using NO assay kit (Beyotime Biotech., China). Cells ( $1 \times 10^5$  cells/well) infected with *M. hyopneumoniae* ( $1 \times 10^6$  CFU/ml) and treated with or without LiCl (10–40 mM). Supernatants were mixed with Griess-I and Griess-II and examined at 570 nm. NO concentrations in the supernatants was reduced by 60% at 40 mM LiCl (Fig. 2C) and this might correlate with the cells protection.

Superoxide anion is involved in the early apoptosis. We treated cells ( $1 \times 10^5$  cells/well) with or without LiCl post-infection and the level of ROS was measured by a superoxide assay kit (Beyotime Biotech., China).

Compared to controls, addition of LiCl reduced ROS production by 58% (Fig. 2D). This inhibitory activity of LiCl against NO and ROS was previously observed (Lumetti and Galli, 2014; Wang et al., 2013). Inhibitors of GSK3 $\beta$  increase CREB (cAMP Response Element-Binding Protein) DNA binding activity and IL-10 expression, that reduce the production of inducible nitric oxide synthase (iNOS) and NO cells (Bécherel et al., 1995; Seyler et al., 1997). Different results were reported regarding LiCl treatment. LiCl was found to potentiate LPS-induced iNOS expression in astrocytes, but not in mouse macrophages (Feinstein, 1998), and LiCl alone slightly increased iNOS expression in hepatocytes and *in vivo* (Du et al., 2006). These variable findings may signify stimulus or cell type-selective regulatory effects of GSK3 $\beta$  inhibitors on iNOS expression. However, majority of reports likely indicate attenuation of NO production by GSK inhibitor. Excessive ROS lead to mitochondrial dysfunction and activates caspases and apoptosis (Lu et al., 2011), suggesting that LiCl might act through the mitochondrial pathway.

*M. hyopneumoniae* infection could activate caspase-3 (Bai et al., 2015; 2013). By using caspase-3 activity assay kit (Beyotime Biotech., China), we evaluated whether LiCl blocks *M. hyopneumoniae* activated caspase-3. Cells ( $1 \times 10^5$  cells/well) infected for 10 h as indicated above and treated with or without LiCl (10–40 mM) for 12 h. Cells lysed in lysis buffer ( $\sim 10^8$  cells/ml), centrifuged and the supernatant was quantified by Bradford method (Beyotime Biotech, China). An

about 50  $\mu$ l of cells lysate (100  $\mu$ g) was mixed with 40  $\mu$ l reaction buffer and 10  $\mu$ l caspase-3 substrate (Ac-DEVD-pNA). After incubating at 37 °C for 2 h, caspase-3 activity was measured at 405 nm. As shown in (Fig. 2E), *M. hyopneumoniae* infection significantly increased caspase-3 activity. When PK-15 cells treated with LiCl, caspase-3 activity was significantly reduced by 82%, suggesting that LiCl may act upstream of caspase-3 activation. LiCl induced cells neuro-protection against glutamate was found to be associated with up-regulation of anti-apoptotic protein Bcl-2, down-regulation of pro-apoptotic proteins p53 and Bax, and suppressed release of cytochrome c from mitochondria (Ghribi et al., 2002), thereby preventing caspase-3 activation and apoptosis (Adams and Cory, 1998). Thus, anti-apoptotic mechanism of LiCl may also be available in other cell protection mechanisms, including the prevention of *M. hyopneumoniae*-induced apoptosis.

Various *M. hyopneumoniae* strains (186L, NJ and XLW-1) were tested for their susceptibility to LiCl. Cells ( $1 \times 10^5$  cells/well) were infected with these strains ( $1 \times 10^6$  CFU/ml) and treated with LiCl. At 40 mM LiCl, the apoptosis was significantly inhibited by 80% (data not shown) in cells infected with these strains which indicate that, the inhibition by LiCl may be a general feature of *M. hyopneumoniae* or other Mollicutes.

In conclusion, our data indicated that LiCl inhibited *M. hyopneumoniae* infection on PK-15 cells via antiapoptotic mechanism. These results are potentially relevant to the search for antimicrobials against *M. hyopneumoniae* infections, as well as other Mollicutes.

### Competing interests

The authors declare that they have no competing interests.

### Acknowledgments

This work was supported by Postdoctoral Fund of Jiangsu Academy of Agricultural Science, China (6511318) National Natural Sciences Foundation of China (31100136, 3111339).

### References

- Adams, J.M., Cory, S., 1998. The Bcl-2 protein family: arbiters of cell survival. *Science* 281, 1322–1326.
- Bai, F., Ni, B., Liu, M., Feng, Z., Xiong, Q., Xiao, S., Shao, G., 2013. *Mycoplasma hyopneumoniae*-derived lipid-associated membrane proteins induce apoptosis in porcine alveolar macrophage via increasing nitric oxide production, oxidative stress, and caspase-3 activation. *Vet. Immunol. Immunopathol.* 155, 155–161.
- Bai, F., Ni, B., Liu, M., Feng, Z., Xiong, Q., Shao, G., 2015. *Mycoplasma hyopneumoniae*-derived lipid-associated membrane proteins induce inflammation and apoptosis in porcine peripheral blood mononuclear cells in vitro. *Vet. Microbiol.* 175, 58–67.
- Bécherel, P.-A., Le Goff, L., Ktorza, S., Ouaz, F., Mencia-Huerta, J.-M., Dugas, B., Debré, P., Mossalayi, M.D., Arock, M., 1995. Interleukin-10 inhibits IgE-mediated nitric oxide synthase induction and cytokine synthesis in normal human keratinocytes. *Eur. J. Immunol.* 25, 2992–2995.
- Beere, H.M., Green, D.R., 2001. Stress management—heat shock protein-70 and the regulation of apoptosis. *Trends Cell Biol.* 11, 6–10.
- Bowden, C.L., Grunze, H., Mullen, J., Brecher, M., Paulsson, B., Jones, M., Vagero, M., Svensson, K., 2005. A randomized, double-blind, placebo-controlled efficacy and safety study of quetiapine or lithium as monotherapy for mania in bipolar disorder. *J. Clin. Psychiatry* 66, 111–121.
- Brüne, B., 2003. Nitric oxide: NO apoptosis or turning it ON? *Cell Death Differ.* 10, 864–869.
- Burnett, T.A., Dinkla, K., Rohde, M., Chhatwal, G.S., Uphoff, C., Srivastava, M., Cordwell, S.J., Geary, S., Liao, X., Minion, F.C., 2006. P159 is a proteolytically processed, surface adhesin of *Mycoplasma hyopneumoniae*: defined domains of P159 bind heparin and promote adherence to eukaryote cells. *Mol. Microbiol.* 60, 669–686.
- Che, Qiao-lin, Xiong, Qi-yan, Feng, Zhi-xin, Zhang, Xu, Liu, Mao-jun, Guo-qing, S., 2012. Establishment of an indirect immunofluorescence assay for detection of adherence of *Mycoplasma hyopneumoniae* to host cells. *J. Agr. Sci.* 28, 1069–1073.
- Che, Qiao-lin, Feng, Zhi-xin, Bai, Fang fang, Liu, Mao-jun, Guo-qing, Shao, 2013. Adhesion of different virulent *Mycoplasma hyopneumoniae* to host cells. *Chin. J. Prev. Vet. Med.* 35, 725–728.
- Che, Qiao-lin, Liu, Bei-bei, Liu, Mao-jun, Xiong, Qi-yan, Guo-qing, Shao, 2014. Invasion of host cells by different virulent *Mycoplasma hyopneumoniae*. *Acta Vet. Zootch. Sin.* 45, 981–988.
- Chen, K., Wu, Y., Zhu, M., Deng, Q., Nie, X., Li, M., Wu, M., Huang, X., 2013. Lithium chloride promotes host resistance against *Pseudomonas aeruginosa* keratitis. *Mol. Vis.* 19, 1502–1514.
- Clark, L., Armstrong, C., Freeman, M., Scheidt, A., Sands-Freeman, L., Knox, K., 1991. Investigating the transmission of *Mycoplasma hyopneumoniae* in a swine herd with enzootic pneumonia. *Vet. Med. (USA)* 86, 543–550.
- Du, Q., Park, K.S., Guo, Z., He, P., Nagashima, M., Shao, L., Sahai, R., Geller, D.A., Hussain, S.P., 2006. Regulation of human nitric oxide synthase 2 expression by Wnt  $\beta$ -catenin signaling. *Cancer Res.* 66, 7024–7031.
- Feinstein, D.L., 1998. Potentiation of astroglial nitric oxide synthase type-2 expression by lithium chloride. *J. Neurochem.* 71, 883–886.
- Ghribi, O., Herman, M.M., Spaulding, N.K., Savory, J., 2002. Lithium inhibits aluminum-induced apoptosis in rabbit hippocampus, by preventing cytochrome c translocation, Bcl-2 decrease, Bax elevation and caspase-3 activation. *J. Neurochem.* 82, 137–145.
- Haesebrouck, F., Pasmans, F., Chiers, K., Maes, D., Ducatelle, R., Decostere, A., 2004. Efficacy of vaccines against bacterial diseases in swine: what can we expect? *Vet. Microbiol.* 100, 255–268.
- Harrison, S.M., Tarpey, I., Rothwell, L., Kaiser, P., Hiscox, J.A., 2007. Lithium chloride inhibits the coronavirus infectious bronchitis virus in cell culture. *Avian Pathol.* 36, 109–114.
- Huppert, J., Closhen, D., Croxford, A., White, R., Kulig, P., Pietrowski, E., Bechmann, I., Becher, B., Luhmann, H.J., Waisman, A., 2010. Cellular mechanisms of IL-17-induced blood-brain barrier disruption. *FASEB J.* 24, 1023–1034.
- Li, J., Yin, J., Sui, X., Li, G., Ren, X., 2009. Comparative analysis of the effect of glycyrrhizin diammonium and lithium chloride on infectious bronchitis virus infection in vitro. *Avian Pathol.* 38, 215–221.
- Lu, T.-H., Su, C.-C., Chen, Y.-W., Yang, C.-Y., Wu, C.-C., Hung, D.-Z., Chen, C.-H., Cheng, P.-W., Liu, S.-H., Huang, C.-F., 2011. Arsenic induces pancreatic  $\beta$ -cell apoptosis via the oxidative stress-regulated mitochondria-dependent and endoplasmic reticulum stress-triggered signaling pathways. *Toxicol. Lett.* 201, 15–26.
- Lumetti, S., Galli, C., 2014. Hyperhydrophilic Surfaces Enhance The Effects of Oxidative Stress on Bone Cells and LiCl Can Reverse Them.
- Machado-Vieira, R., Manji, H.K., Zarate Jr., C.A., 2009. The role of lithium in the treatment of bipolar disorder: convergent evidence for neurotrophic effects as a unifying hypothesis. *Bipolar Disord.* 11, 92–109.
- Maniam, P., Shirat, A.F., Sidek, H.M., Ismail, G., Embi, N., 2013. Inhibition of GSK3 attenuates the intracellular multiplication of burkholderia pseudomallei and modulates the inflammatory response in infected macrophages and A549 epithelial lung cells. *Sains Malays.* 42, 725–735.
- Ni, B., Bai, F., Wei, Y., Liu, M., Feng, Z., Xiong, Q., Hua, L., Shao, G., 2014. Apoptosis induced by lipid-associated membrane proteins from *Mycoplasma hyopneumoniae* in a porcine lung epithelial cell line with the involvement of caspase 3 and the MAPK pathway. *Genet. Mol. Res.* 14, 11429–11443.
- Ren, X., Meng, F., Yin, J., Li, G., Li, X., Wang, C., Herliker, G., 2011. Action mechanisms of lithium chloride on cell infection by transmissible gastroenteritis coronavirus. *PLoS One* 6, e18669.
- Renvoize, C., Biola, A., Pallardy, M., Breard, J., 1998. Apoptosis: identification of dying cells. *Cell Biol. Toxicol.* 14, 111–120.
- Seyler, I., Appel, M., Devissaguet, J.-P., Legrand, P., Barratt, G., 1997. Modulation of nitric oxide production in RAW 264.7 cells by transforming growth factor-beta and interleukin-10: differential effects on free and encapsulated immunomodulator. *J. Leukoc. Biol.* 62, 374–380.
- Shen, Y.H., Wang, X.L., Wilcken, D.E., 1998. Nitric oxide induces and inhibits apoptosis through different pathways. *FEBS Lett.* 433, 125–131.
- Wang, H.-M., Zhang, T., Li, Q., Huang, J.-K., Chen, R.-F., Sun, X.-J., 2013. Inhibition of glycogen synthase kinase-3 $\beta$  by lithium chloride suppresses 6-hydroxydopamine-induced inflammatory response in primary cultured astrocytes. *Neurochem. Int.* 63, 345–353.
- Wang, H., Feng, Z., Wu, Y., Wei, Y., Gan, Y., Hua, L., Li, B., Wang, X., Liu, M., Xiong, Q., 2016. The effects of *Mycoplasma hyopneumoniae* on porcine circovirus type 2 replication in vitro PK-15 cells. *Res. Vet. Sci.* 105, 56–61.
- Wu, Yu-zi, Jin, Meng-meng, Bai, Fang-fang, Zhang, Xu, Hua, Li-zhong, Lei, Zhi-hai, Guo-qing, S., 2012. Development of TaqMan -BHQ real-time PCR assay for detection of *Mycoplasma hyopneumoniae* P97. *Chin. Vet. Sci.* 24, 1268–1272.
- Zhang, P., Katz, J., Michalek, S.M., 2009. Glycogen synthase kinase-3 $\beta$  (GSK3 $\beta$ ) inhibition suppresses the inflammatory response to Francisella infection and protects against tularemia in mice. *Mol. Immunol.* 46, 677–687.
- Zhang, Xu, Feng, Zhi-xin, Wu, Yu-zi, Che, Qiao-lin, Liu, Mao-jun, Xiong, Qi-yan, Hua, Li-zhong, Bin, Li, Guo-qing, Shao, 2013. Establishment of *Mycoplasma hyopneumoniae* adhesion on PK-15 model and the adhesion ability in vitro. *Acta Vet. Zootch. Sin.* 44, 772–777.
- Ziaie, Z., Kefalides, N.A., 1989. Lithium chloride restores host protein synthesis in herpes simplex virus-infected endothelial cells. *Biochem. Biophys. Res. Commun.* 160, 1073–1078.