

RESEARCH ARTICLE

Effects of early, combined endurance and resistance training in mechanically ventilated, critically ill patients: A randomised controlled trial

Sabrina Eggmann^{1*}, Martin L. Verra¹, Gere Luder¹, Jukka Takala², Stephan M. Jakob²

1 Department of Physiotherapy, Inselspital, Bern University Hospital, Bern, Switzerland, **2** Department of Intensive Care Medicine, Inselspital, Bern University Hospital, University of Bern, Bern, Switzerland

* sabrina.eggmann@insel.ch



OPEN ACCESS

Citation: Eggmann S, Verra ML, Luder G, Takala J, Jakob SM (2018) Effects of early, combined endurance and resistance training in mechanically ventilated, critically ill patients: A randomised controlled trial. *PLoS ONE* 13(11): e0207428. <https://doi.org/10.1371/journal.pone.0207428>

Editor: Johannes Fleckenstein, University of Bern, SWITZERLAND

Received: June 17, 2018

Accepted: October 30, 2018

Published: November 14, 2018

Copyright: © 2018 Eggmann et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: The minimal dataset has been anonymized and are within the Supporting Information files.

Funding: This study was self-funded and supported by the Departments of Physiotherapy and Intensive Care Medicine at the Inselspital, Bern University Hospital, Bern, Switzerland. None of the authors received funding for this work.

Competing interests: SE, MLV, GL declare that they have no competing interests. JT and SMJ report the following potential conflicts of interest:

Abstract

Introduction

Neuromuscular weakness resulting in severe functional impairment is common in critical care survivors. This study aimed to evaluate effects of an early progressive rehabilitation intervention in mechanically ventilated adults at risk.

Methods

This was a parallel, two-arm, assessor-blinded, randomised controlled trial with 6-months follow-up that was conducted in a mixed ICU of an academic centre in Switzerland. Previously independent, mechanically ventilated, critically ill adults with expected critical care stay ≥ 72 hours ($n = 115$) were randomised to a control group receiving standard physiotherapy including early mobilisation or to an experimental group with early endurance and resistance training combined with mobilisation. Primary endpoints were functional capacity (6-Minute Walk Distance) and functional independence (Functional Independence Measure) at hospital discharge. Secondary endpoints including muscle strength were assessed at critical care discharge. Safety was monitored closely by standard monitoring and predefined adverse events.

Results

Physiotherapy started within 48 hours of critical care admission while 97% of participants were still ventilated and 68% on inotropes. Compared to the control group ($n = 57$), the experimental group ($n = 58$) received significantly more physiotherapy (sessions: 407 vs 377, $p < 0.001$; time/session: 25min vs 18min, $p < 0.001$) and had less days with sedation ($p < 0.001$). Adverse events were rare (0.6%) and without consequences. There were no significant between-group differences in 6-Minute Walk Distance (experimental 123m (IQR 25–280) vs control 100m (IQR 0–300); $p = 0.542$) or functional independence (98 (IQR 66–119) vs 98 (IQR 18–115); $p = 0.308$). Likewise, no differences were found for the secondary

The Department of Intensive Care Medicine has, or has had in the past, research contracts with Orion Corporation, Abbott Nutrition International, B. Braun Medical AG, CSEM SA, Edwards Lifesciences Services GmbH, Kenta Biotech Ltd, Maquet Critical Care AB, Omnicare Clinical Research AG and research & development/consulting contracts with Edwards Lifesciences SA, Maquet Critical Care AB, and Nestlé. The money was paid into a departmental fund; JT and SMJ received no personal financial gain. The Department of Intensive Care Medicine has received unrestricted educational grants from the following organizations for organizing a quarterly postgraduate educational symposium, the Berner Forum for Intensive Care (until 2015): Fresenius Kabi, gsk, MSD, Lilly, Baxter, astellas, AstraZeneca, B | Braun, CSL Behring, Maquet, Novartis, Covidien, Nycomed, Pierre Fabre Pharma AG (formerly known as RobaPharm), Pfizer, Orion Pharma, Bard Medica S.A., Abbott AG, Anandic Medical Systems. The Department of Intensive Care Medicine has received unrestricted educational grants from the following organizations for organizing bi-annual postgraduate courses in the fields of critical care ultrasound, management of ECMO and mechanical ventilation: Pierre Fabre Pharma AG (formerly known as RobaPharm), Pfizer AG, Bard Medica S.A., Abbott AG, Anandic Medical Systems, PanGas AG Healthcare, Orion Pharma, Bracco, Edwards Lifesciences AG, Hamilton Medical AG, Fresenius Kabi (Schweiz) AG, Getinge Group Maquet AG, Dräger Schweiz AG, Teleflex Medical GmbH. No author was employed by any of these companies. None of the companies played a role in this study. Our relationship with the companies does not alter our adherence to PLOS ONE policies on sharing data and materials.

outcomes, except a trend towards improved mental health in the experimental group after 6 months (84 (IQR 68–88) vs 70 (IQR 64–76); $p = 0.023$).

Conclusions

Early endurance and resistance training in mechanically ventilated, intensive care patients does not improve functional capacity or independence at hospital discharge compared to early standard physiotherapy but may improve mental health 6-months after critical care discharge.

Trial registration

German Clinical Trials Register (DRKS): [DRKS00004347](https://www.drks.de/DRKS00004347), registered on 10 September 2012.

Introduction

Intensive care unit acquired weakness (ICUAW) is common in critical illness survivors. It increases health-care costs and impairs quality of life [1]. ICUAW mainly affects skeletal muscles (critical illness myopathy) and peripheral nerves (critical illness polyneuropathy) [2], but its causes are poorly understood [3]. Symptoms of ICUAW develop within the first week of critical illness [4]. Its risk factors include sepsis, multiple organ failure, mechanical ventilation (MV), immobilisation and hyperglycaemia [3]. ICUAW is associated with worse short- and long-term outcomes, including weaning failure [5], prolonged ICU and hospital stays [6], increased mortality [7], and poor functional status with persistent disability in activities of daily living [1, 8–10].

Early rehabilitation and mobilisation may help to reduce the functional disabilities of ICUAW patients. Some promising short-term outcomes have been reported [11, 12], but long-term benefits from randomised controlled trials are warranted [13–15]. Delayed intervention onset (≥ 1 week) [12, 13, 15], low ICUAW-risk in targeted population [11] or unspecific exercise delivery [11, 14] may have contributed to the inconclusive results. Despite these limitations in evidence, many ICUs, including ours, have adopted early mobilisation, also as a consequence of using less sedation. We aimed to investigate a specific rehabilitation intervention that could be implemented early in a highly ill population with a high ICUAW risk.

Resistance training can restore strength after bed-rest in older adults [16]. Combined endurance and resistance training (ERT) enhances exercise capacity and health-related quality of life in patients with chronic obstructive pulmonary disease [17] and prevents cardiovascular and skeletal muscle deconditioning during bedrest [18]. Whether early ERT is feasible, safe and beneficial in critically ill patients of whom many are still mechanically ventilated and treated with opioids or recovering from sedation is unclear. We hypothesised that early ERT combined with mobilisation in the ICU is feasible and might improve function at hospital discharge. This randomised controlled trial compared the effects and safety of early ERT combined with early mobilisation to standard care including early mobilisation in critically ill, mechanically ventilated adults for a period of up to 6-months after hospital discharge.

Methods

Design and setting

This single-centre, parallel, two-arm, assessor-blinded randomised controlled trial with 6-months follow-up was conducted in the tertiary, mixed ICU of the Department of Intensive Care Medicine at the Inselspital, Bern University Hospital, Switzerland. The trial was approved by the local Ethics Committee and prospectively registered in the German Clinical Trials Register (DRKS00004347) on September 10, 2012. It adhered to Good Clinical Practice Guidelines and the CONSORT statement (S1 Checklist). The study protocol has previously been published [19].

Participants

Adults (≥ 18 years) expected to stay on MV for at least 72 hours and who had been independent before the onset of critical illness were eligible for participation. Independence was defined as living without assistance and qualitatively determined by the patient's family or medical-health records. Exclusion criteria were previous muscle weakness, contraindications to cycling, enrolment in another intervention study, palliative care, admission diagnosis that excluded the possibility of walking at hospital discharge, and patients who did not understand German or French [19].

Procedures

Patients were screened daily by a research nurse and a specialist physiotherapist who evaluated cycling eligibility. The expected MV duration was judged by the responsible physician. Subsequently and before study inclusion, an independent medical doctor confirmed eligibility criteria. Physicians and physiotherapists were unaware of the randomisation sequence. A deferred consent was obtained from the next of kin within 72 hours after randomisation and a written informed consent from each patient as early as possible. Unrestricted, computer-generated randomisation (<https://www.random.org/lists/>) was performed by a study nurse partly involved in screening eligible candidates. Participants were allocated at a 1:1 ratio to either group using sequentially numbered opaque sealed envelopes to ensure allocation concealment [20]. Coded data was managed with REDcap [21].

Blinding the responsible ICU staff was impossible, but the physiotherapy assessors were blinded for group allocation and distinct from the therapists providing the intervention. However, logistical reasons prevented therapist separation between the two treatment groups. Discharge decisions were made by the primary care team without the involvement of physiotherapists.

Interventions

Both groups received standard ICU care with protocol-guided sedation, weaning [22] and nutrition [23]. In cases of ICU readmissions, treatment according to initial group allocation was continued.

Standard care (control group). Patients in the control group received European standard physiotherapy [24] including early mobilisation, respiratory therapy and passive or active exercises. These were physiotherapy-initiated and individually tailored, but subject to a medical prescription. Treatments took place once daily on weekdays and on weekends if the responsible therapists deemed an interruption of therapy to be harmful to patient's previous progress (e.g. intensive rehabilitation and weaning period or retained airway secretion in extubated patients).

Study intervention (experimental group). In patients in the experimental group an early, progressive ERT programme combined with early mobilisation (S1 Fig) was initiated and each component implemented as intensive as possible and tolerated in individual patients. To this end, sedation was reduced if medically permitted prior to physiotherapy [19]. Physiotherapists were encouraged to split therapy into two or more sessions to prevent overexertion. Therapy visits occurred from Monday to Friday up to a maximum of three sessions per day and on weekends if deemed helpful to patient's progress.

Endurance training was conducted with a motor-assisted bed-cycle (MOTOmed letto2, Reck-Technik, Betzenweiler, Germany) that allows passive, motor-assisted or active cycling in bed. Patients were comfortably positioned in a supine position with an individual head-of-bed elevation to permit optimal leg movement. Maximal training intensity for unresponsive patients was 20 minutes with a pedalling rate of 20 cycles/min [12]. During each session, patients were prompted verbally and with the MOTOmed ServoCycle feature to actively participate. When patients were able to cycle with motor-assistance for 20 minutes, assistance was gradually decreased. Only then was training duration raised to 30 minutes. Subsequently, resistance and then duration were increased with a defined maximum of 60 minutes with full resistance (S1 Table).

Resistance training included standardised exercises for both upper and lower limbs using weights or manually administered resistance from the therapist. The targeted training intensity was 8–12 repetitions with 2–5 sets (2min rest) on 50–70% of the estimated one-repetition maximum. Once the exercises were performed properly, participants were given an illustrated handout for further training with family members or nurses. In patients unable to perform resistance exercises because of lack of strength or comprehension, physiotherapists used passive movement or tactile facilitation to excite movement.

Early mobilisation was started with in-bed exercises and, if no medical contraindications were present, continued progressively from sitting on the bedside to sitting in a chair to standing and, finally, to walking. The overall aim was patients' active participation in functional activities of daily living in order to promote early independence.

Usual physiotherapy was provided for patients after ICU discharge once to twice a day. Sessions commonly involved functional exercises, cycling, walking, strengthening and breathing techniques.

Outcomes

The two primary outcomes were functional capacity (6-Minute Walking Distance (6MWD) [25]) and the ability to perform activities of daily living (Functional Independence Measure (FIM) [26]) at hospital discharge.

Secondary outcomes were FIM and muscle strength at ICU discharge. Muscle strength was assessed if patients could follow at least 3 out of 5 standardised questions [27]. Assessments included the Medical Research Council (MRC) sum-score for ICUAW diagnosis [2], handgrip strength using the JAMAR dynamometer (Sammons Preston Rolyan, Bolingbrook, IL, USA) [28] and quadriceps muscle strength measured with a handheld dynamometer (Microfet 2, Biometrics, Almere, Netherlands) [29]. Additionally, limitations in range-of-motion were recorded for shoulder flexion, elbow flexion and extension, fist closure, hip flexion, knee flexion and extension and foot dorsiflexion [30]. To assess functional mobility the Timed 'Up & Go' was performed at hospital discharge [31]. Further endpoints were time on MV, ICU and hospital length of stay along with achieved ICU mobility. Quality of life was determined with the Short Form 36 (SF-36) six months after hospital discharge. The SF-36 is a valid and reliable measurement to assess long-term physical and mental health in ICU survivors [32].

To judge training intensity and stability, physiological trends and individually-set limits instead of strict target numbers were used to account for each patient's condition and

variability. Safety was closely monitored using standard ICU monitoring and indirect calorimetry for 30min before, during and 15min after physiotherapy. Physiological data was recorded in a patient data management system that stored 2-minute median values to remove artefacts. Prospectively defined adverse events (AE) included new hemodynamically relevant arrhythmias or otherwise unstable hemodynamics, oxygenation desaturation under 85%, a fall or other injury and any accidental removal of a tube, catheter or similar device. Per definition, AEs occurred during or up to 15min after physiotherapy and persisted despite an intervention or therapy interruption. Other known complications of bedrest, e.g. atelectasis or pressure ulcers, were recorded along with any serious AE which included death, a life-threatening injury or any event that caused an extension of hospital stay.

Statistical analysis

Sample size determination was based on the 6MWD to show a difference of 54m [33] and a mean walking distance of 301m (SD 81) [34]. A statistical power of 80% and an α -level of 0.05 required a sample size of totally 72 patients (36 per group). This was further adjusted by 28 patients, due to two primary outcomes that were expected to highly correlate. Lastly, the overall sample size was increased by 15% to 115 participants because of expected attrition, including mortality. It was decided a priori that refusals to participate (presumed consent not obtained within 72h of randomisation) were compensated for and were not to be included in the intention-to-treat analysis. The number of these additional participants was determined after the inclusion of 115 patients. As a consequence, 130 patients were randomised until the number of 115 participants with presumed consent was reached.

Analysis was by intention-to-treat and per protocol using SPSS (IBM SPSS Statistics Version Premium GradPack 24), and R (Version 3.4.4). In the per protocol analysis, patients who died were allocated the lowest possible outcome for both primary outcomes. The intention-to-treat analysis was conducted with multiple imputation using a linear regression model that imputed all missing values (including death). Sensitivity analyses for both primary outcomes were conducted to test the robustness of the main analysis. Secondary outcomes were analysed as measured. Between group comparisons for normally distributed data included Student's *t*-tests and for non-normally distributed data Mann-Whitney-U tests. Dichotomous data was analysed with Fisher's exact test, repetitive data with repeated measures analysis of variance (ANOVA) and correlations with Spearman correlation coefficients. Continuous data are presented as means with standard deviation (SD) or median with interquartile range (IQR) dependent upon their distribution. Categorical data are given as numbers with percentages. Treatment effects are reported with 95% Confidence intervals (95% CI) as mean differences (MD) for normally distributed data, as estimator for difference in location for non-normal data (Wilcoxon-Mann-Whitney test implemented in R package "coin"), and as incidence rate ratios (IRR) or risk differences (RD) for percentages. IRRs and RDs were calculated with R using Exact Poisson and Proportion test, respectively. Effect sizes were calculated with Cohen's *d* for normally distributed data and with Cohen's *r* for Wilcoxon-Mann-Whitney test statistic in case of non-normal data (using appropriate *z* value as in [35]). The level of significance was set at $p < 0.05$. A preliminary safety analysis has been conducted after the enrolment of 35 participants [36].

Results

Study participants

From October 8, 2012 to April 5, 2016, we randomised 130 patients for whom 15 (12%) patient's proxy declined initial study participation. These were excluded from all further analyses, as requested by the Ethics Committee. Consequently, 115 patients (experimental $n = 58$,

control $n = 57$) were included in the intention-to-treat analysis (Fig 1). Baseline demographics and clinical characteristics were similar between groups (Table 1).

Primary outcomes

No significant differences were found between groups in the intention-to-treat and the per protocol analysis (Table 2). There was a high positive correlation ($p < 0.001$, $r = 0.833$) between the two primary outcomes. Results were robust in sensitivity analyses (S2 Table).

Secondary outcomes

There were no significant differences between groups at ICU and hospital discharge (Table 3). The incidence of ICUAW at ICU discharge, determined by MRC sum-score in cooperative patients, was 58% (23/40) in the experimental and 61% (26/43) in the control group ($n = 83$; $p = 0.826$). Limitations in range-of-motion were observed in 22 (38%) patients of the experimental and in 16 (28%) of the control group ($n = 115$; $p = 0.323$) with a median of 2 (IQR 0–3) affected joints per patient. No joint was particularly vulnerable for contracture. There were no significant differences for ICU ($p = 0.694$) and hospital discharge destinations ($p = 0.555$) between groups. The most frequent discharge destinations for survivors were rehabilitation (40%; experimental $n = 25$, control $n = 21$), local hospital (26.1%; experimental $n = 17$, control $n = 13$) and home (15.7%; experimental $n = 7$, control $n = 11$). Except for mental health ($p = 0.023$), there were no significant differences in quality of life at 6-months after hospital discharge (Table 3). Because of substantial missing data for the SF-36, characteristics for patients with and without the 6-month follow-up were compared. There were no significant differences between patients that completed the SF-36 versus patients that did not return the questionnaire (S3 Table).

Safety

Safety details are given in Table 4. Overall, 25 (3%) of all physiotherapy sessions had to be discontinued (between-groups difference $p = 0.839$). The most frequent reasons for therapy discontinuation were exceedance of individually set limits ($n = 8$, 32%), lack of patient cooperation ($n = 5$, 20%) and fatigue ($n = 3$, 12%). There were 4 (0.6%) AEs, 1 (0.2%) in the experimental group due to an oxygen desaturation while cycling and 3 (0.8%) in the control group during mobilisation (one oxygen desaturation, two unstable haemodynamics). All AEs resolved after therapy discontinuation and did not have further consequences. The incidence of ICU complications did not differ between the two groups. ICU, hospital- and 6-month mortality rates were similar between groups over the study period.

Patients on MV (S1 File) had a significant effect over time (30min before, during and 15min after physiotherapy) for mean arterial pressure (MAP; $p < 0.001$, effect size $\eta^2 = 0.192$), heartrate (HR; $p < 0.001$, $\eta^2 = 0.161$) and oxygen consumption (VO_2 ; $p < 0.001$, $\eta^2 = 0.224$), but not for peripheral oxygen saturation (SpO_2 ; $p = 0.396$, $\eta^2 = 0.017$; repeated measures ANOVA). Post-hoc paired t-test with adjusted levels of significance ($p < 0.017$) found increases from before to during physiotherapy for MAP (mean difference (95% CI): -1.83mmHg (-2.61 to -1.05), $p < 0.001$), HR (-1.21beats/min (-1.76 to -0.65), $p < 0.001$) and for VO_2 (-12.22ml/min (-16.75 to -7.69), $p < 0.001$). Additionally, there was an increase in HR from before to after physiotherapy (-0.83beats/min (-1.47 to -0.18), $p = 0.012$) and a decrease in VO_2 from during to after physiotherapy (6.27ml/min (1.45 to 11.08), $p = 0.011$). Likewise, predominantly spontaneously breathing patients (S1 File) had a significant effect over time for MAP ($p = 0.046$, $\eta^2 = 0.073$) and HR ($p = 0.004$, $\eta^2 = 0.123$), but not for SpO_2 ($p = 0.083$, $\eta^2 = 0.059$). Post-hoc paired t-test with adjusted levels of significance ($p < 0.017$) showed an increase from before to




Fig 1. Study flow diagram (CONSORT).

<https://doi.org/10.1371/journal.pone.0207428.g001>

during physiotherapy for HR (-1.85 beats/min (-3.16 to -0.53), $p = 0.006$). There were no significant differences between groups for any of the vital parameters.

Study interventions

The experimental group trained on 73% of study days (349 daily physiotherapy visits, including 407 individual sessions), whereas the control group received therapy on 65% of study days

Table 1. Baseline demographic and clinical characteristics.

| | Experimental group n = 58 | Control group n = 57 |
|--|------------------------------|-------------------------|
| Age, years, mean (SD) | 65±15 | 63±15 |
| Sex, female, n (%) | 22 (38%) | 16 (28%) |
| Weight, kg, mean (SD) | 82±18 | 80±16 |
| BMI, kg/m ² , mean (SD) | 28±6 | 27±4 |
| APACHE II score, mean (SD) ^a | 22±8 | 23±7 |
| TISS-28 score, median (IQR) | 37 (31–42) | 37 (32–44) |
| TISS-76 score, median (IQR) | 46 (39–56) | 48 (41–56) |
| ICU days until study inclusion, mean (SD) | 1.9±1.2 | 1.8±1.2 |
| SOFA score, median (IQR) ^b | 8 (6–10) | 8 (7–12) |
| ICU diagnosis on ICU admission | | |
| Heart surgery | 13 (11%) | 8 (7%) |
| Neurology / neurosurgery | 5 (4%) | 4 (3%) |
| Other surgery | 5 (4%) | 9 (8%) |
| Gastroenterology | 7 (6%) | 7 (6%) |
| Trauma | 1 (1%) | 3 (3%) |
| Respiratory insufficiency | 13 (11%) | 12 (10%) |
| Hemodynamic insufficiency | 13 (11%) | 13 (11%) |
| Other | 1 (1%) | 1 (1%) |
| Comorbidities on ICU admission (n = 111: experimental n = 54, control n = 57) | | |
| NYHA symptoms (stage 2 to 4) | 22 (41%) | 26 (46%) |
| Dyspnoea symptoms | 15 (28%) | 15 (26%) |
| Hematologic malignancy | 0 (0%) | 5 (9%) |
| Immunosuppression | 4 (7%) | 12 (21%) |
| Liver disease | 9 (17%) | 6 (11%) |
| Chronic dialysis | 0 (0%) | 0 (0%) |
| Restricted in activities of daily living ^c | 8 (15%) | 5 (9%) |
| Laboratory on ICU admission | | |
| CRP, mg/l, median (IQR), n = 106 | 95 (49–276) | 76 (30–147) |
| Creatinine, µmol /l, median (IQR), n = 109 | 97 (70–168) | 111 (78–156) |
| Bilirubin, µmol /l, median (IQR), n = 68 | 15 (8–54) | 18 (8–32) |
| Thrombocytes, 10 ⁹ /L, median (IQR), n = 113 | 117 (102–216) | 108 (67–148) |
| Leucocytes, 10 ⁹ /L, median (IQR), n = 133 | 11 (8–16) | 11 (6–16) |
| Haemoglobin, g/L, median (IQR), n = 115 | 109 (97–119) | 102 (85–115) |

^a at ICU admission

^b at study inclusion

^c inability to walk for longer distances or to participate in social life

Data are presented as median (IQR), mean (SD) or n (%)

Abbreviations: NYHA = New York Heart Association, BMI = Body Mass Index, APACHE = Acute Physiology and Chronic Health Evaluation, TISS = Therapeutic Intervention Scoring System, SOFA = Sequential Organ Failure Assessment, CRP = C-reactive protein

<https://doi.org/10.1371/journal.pone.0207428.t001>

Table 2. Results for the two primary outcomes per protocol and intention-to-treat analysis.

| | Experimental group n = 58 | | Control group n = 57 | | p value | Effect size | Group difference (95% CI) |
|--|------------------------------|------------------|-------------------------|-----------------|-----------|-------------|------------------------------|
| | n | value | n | value | | | |
| Per Protocol analysis | | | | | | | |
| 6 Minute Walking Distance (m) | 54 | 123 (IQR 25–280) | 53 | 100 (IQR 0–300) | p = 0.542 | 0.006 | 0.00 (-35.00 to 65.00) |
| FIM (18–126) | 54 | 98 (IQR 66–119) | 52 | 98 (IQR 18–115) | p = 0.308 | 0.010 | 3.00 (-2.00 to 14.00) |
| Intention-to-treat analysis^a | | | | | | | |
| 6 Minute Walking Distance (m) | 58 | 223±133 | 57 | 246±167 | p = 0.448 | -0.151 | -22.75 (- 81.66 to 36.16) |
| FIM (18–126) | 58 | 101±22 | 57 | 99±24 | p = 0.659 | 0.085 | 1.98 (-6.85 to 10.82) |

^a Multiple imputation from a linear regression model based on age, gender, BMI, weight, APACHE II score, TISS-28, TISS-76, SOFA score, ICU days until randomisation, length of stay in ICU and hospital, duration of mechanical ventilation, FIM score and MRC sum-score at ICU discharge. All missing values (including death before hospital discharge) were imputed (6MWD: n = 33, FIM: n = 32).

<https://doi.org/10.1371/journal.pone.0207428.t002>

(365 visits, 377 sessions; [Table 5](#)). The main reasons for missed physiotherapy visits were week-ends (188, 58%), day of ICU discharge (42, 13%), medical procedures (33, 10%) or cardiorespiratory instability (19, 6%). Patients in the experimental group had a median of 3.5 sessions (minimum 0, maximum 66) and patients in the control group of 3 sessions (minimum 0, maximum 39) ([S2 File](#)). Treatment session duration was slightly longer in the experimental group (25min (IQR 19–27) vs control 18min (IQR 14–21); $p < 0.001$). This was recorded from the beginning to the end of actual therapy delivery and did not include preparation time, cleaning materials and documentation. On average, physiotherapy started within 48 hours (n = 113; median 1.98 days (IQR 1.41–2.86); experimental 47h (IQR 32–68); control 52h (IQR 36–70); $p = 0.468$) after ICU admission and within 4 hours after randomisation. At this time 97% of participants were still ventilated (experimental 57(100%), control 52(93%); $p = 0.057$), 97% sedated (experimental 55(97%), control 55(98%), $p < 0.999$), 68% on vasoactive support (experimental 42(74%), control 35(63%); $p = 0.230$), 19% paralysed (experimental 9(16%), control 12 (21%); $p = 0.477$) and 97% received opioids (experimental 55(97%), control 54(96%); $p < 0.999$). Patients in the experimental group were less likely to receive sedation (incidence risk ratio (IRR) 1.66 (1.26 to 2.18, $p = 0.0002$) and steroids (IRR 1.18 (1.03 to 1.35), $p = 0.016$), but more likely to receive vasopressors (IRR 0.77 (0.64 to 0.92), $p = 0.005$) compared to the control group.

The number of each therapy intervention per group is outlined in [Fig 2](#). Most patients were unable to perform the planned resistance training (sessions experimental 8, control 2; $p = 0.052$) with the exception of some long-stayers ([S4 Table](#)). Type of cycling included passive (141, 76%), mixed (28, 15%) and active training modes (17, 9%). Mean cycling distance was 1183 m (± 721) during an average of 15min (± 5), a pedalling rate of 20r/min (± 7) and a resistance level of 0.2 (± 0.8). Physiotherapy-initiated and overall mobilisations differed between groups; the experimental group had more edge-of-bed mobilisations, while the control group stood more ([S2 File](#)). However, achievement of milestones from time of randomisation to first edge-of-bed were not different (experimental 4d (IQR 2–7), control 5d (IQR 2–7); $p = 0.451$) ([S2 File](#)).

Discussion

This randomised controlled trial did not find an advantage of early ERT combined with mobilisation over standard early mobilisation and rehabilitation. While participants in the experimental group received more therapy and had less days with sedation, control group

Table 3. Results of secondary outcomes.

| | Experimental group n = 58 | | Control group n = 57 | | p value | Effect size | Group difference (95% CI) |
|---------------------------------------|------------------------------|-----------------------|-------------------------|-----------------------|-----------|-------------|------------------------------|
| | n | value | n | value | | | |
| Hospital discharge | | | | | | | |
| Timed "Up & Go" Test (s) | 36 | 19.5 (IQR 11.5–25.0) | 28 | 16.0 (IQR 10.3–29.0) | p = 0.538 | 0.010 | 1.70 (-3.70 to 7.00) |
| Time on mechanical ventilation (days) | 58 | 5.4 (IQR 3.3–12.9) | 57 | 5.0 (IQR 3.6–11.9) | p = 0.830 | -0.002 | -0.17 (-1.65 to 1.49) |
| Length of stay ICU (days) | 58 | 6.1 (IQR 4.0–12.3) | 57 | 6.6 (IQR 4.6–14.7) | p = 0.568 | -0.005 | -0.44 (-2.27 to 1.20) |
| Length of stay hospital (days) | 58 | 25.9 (IQR 14.3–37.2) | 57 | 22.0 (IQR 15.0–39.2) | p = 0.723 | 0.003 | 1.31 (-4.85 to 7.92) |
| ICU discharge | | | | | | | |
| FIM (18–126) | 58 | 28.5 (IQR 21.0–42.0) | 56 | 28.5 (IQR 19.5–41.5) | p = 0.791 | 0.002 | 0.00 (-4.00 to 5.00) |
| MRC sum-score (0–60) | 40 | 42.4±13.1 | 43 | 44.4±11.7 | p = 0.461 | -0.161 | MD -2.02 (-7.45 to 3.41) |
| Handgrip strength (JAMAR) (kg) | 30 | 20.5±12.6 | 40 | 19.6±13.6 | p = 0.780 | 0.069 | MD 0.90 (-5.47 to 7.25) |
| Quadriceps strength (HDD) (kg) | 28 | 7.7±4.0 | 33 | 8.0±3.9 | p = 0.771 | -0.076 | MD -0.30 (-2.32 to 1.73) |
| 6-month follow-up: SF-36 | | | | | | | |
| Physical functioning (0–100) | 36 | 75.0 (IQR 45.0–85.0) | 27 | 75.0 (IQR 50.0–85.0) | p = 0.676 | -0.007 | -1.11 (-15.00 to 10.00) |
| Role physical (0–100) | 36 | 25.0 (IQR 0.0–62.5) | 25 | 33.3 (IQR 25.0–50.0) | p = 0.443 | -0.013 | 0.00 (-25.00 to 0.00) |
| Bodily pain (0–100) | 36 | 80.0 (IQR 51.0–100.0) | 27 | 74 (IQR 41.0–100.0) | p = 0.530 | 0.010 | 0.00 (-9.00 to 20.00) |
| General health (0–100) | 36 | 57.9±18.4 | 25 | 61.1±19.0 | p = 0.513 | -0.162 | MD -3.20 (-12.91 to 6.52) |
| Vitality (0–100) | 36 | 55.4±19.4 | 26 | 48.3±20.1 | p = 0.167 | 0.359 | MD 7.10 (-3.06 to 17.26) |
| Social functioning (0–100) | 36 | 75.0 (IQR 65.3–100.0) | 26 | 68.8 (IQR 50.0–100.0) | p = 0.458 | 0.012 | 0.00 (0.00 to 25.00) |
| Role emotional (0–100) | 35 | 66.7 (IQR 33.3–100.0) | 26 | 50.0 (IQR 0.0–100.0) | p = 0.870 | 0.003 | 0.00 (0.00 to 33.33) |
| Mental health (0–100) | 35 | 84.0 (IQR 68.0–88.0) | 26 | 70.0 (IQR 64.0–76.0) | p = 0.023 | 0.037 | 12.00 (0.00 to 16.00) |
| Physical health (sum-score) | 34 | 40.8±11.1 | 24 | 42.7±10.4 | p = 0.520 | -0.190 | MD -1.87 (-7.65 to 3.91) |
| Mental health (sum-score) | 34 | 49.4±10.3 | 24 | 45.2±11.4 | p = 0.147 | 0.381 | MD 4.23 (-1.53 to 9.99) |

Data are presented as median (IQR), mean (SD) or n (%)

Abbreviations: FIM = Functional Independence Measure (worst score 18: dependent, best score 126: independent), HDD = handheld dynamometer, LOS = length of stay, SF-36 = Short Form 36, version 2 (worst score: 0, best score: 100, sum-score: T-values where the population mean is 50 and the SD is 10; based on US-population 1990), MD = Mean Difference

<https://doi.org/10.1371/journal.pone.0207428.t003>

participants had more out-of-bed mobilisations. Functional capacity and independence did not differ between these two different exercise regimes at hospital discharge. No differences were found in the secondary outcomes, except for a trend towards improved mental health at 6-months after hospital discharge. AEs were rare and a stepwise, individually-supervised training approach starting within 48 hours of ICU admission appears safe. ERT interventions are feasible and highly beneficial for functional improvement in chronic obstructive pulmonary disease [17] and we hypothesised that they may be equally feasible and effective in the critically ill. However, the transfer of these interventions may not be as straightforward as anticipated. Pulmonary rehabilitation is usually delivered over 8 to 12 weeks [17], while our exercise exposure was limited to the time spent in ICU, when patients were recovering from critical illness, mostly still on mechanical ventilation, receiving opioids and sedatives or at least recovering from previously received sedatives. Since the patients in our study spent the greater part of

Table 4. Safety details for the control and experimental groups.

| | Experimental group (n = 58, 478 study days) | Control group (n = 57, 558 study days) | p value | Group difference (95% CI) |
|--|---|--|-----------|---------------------------|
| Safety | | | | |
| PT discontinuation (% of total physiotherapy sessions) | 12 (3%) | 13 (3%) | p = 0.842 | IRR 0.86 (0.36 to 2.03) |
| Adverse event (% per sessions) | 1 (0.2%) | 3 (0.8%) | p = 0.357 | IRR 0.31 (0.01 to 3.85) |
| Mortality | | | | |
| ICU mortality | 9 (16%) | 10 (18%) | p = 0.770 | RD -2% (-16 to 12%) |
| In-hospital mortality | 10 (17%) | 14 (25%) | p = 0.334 | RD -8% (-22 to 7%) |
| Mortality after 180 days | 16 (28%) | 16 (28%) | p = 0.954 | RD 0% (-17% to 16%) |
| ICU complications^a | | | | |
| Delirium-free ICU days | 256 (61%) | 300 (64%) | p = 0.524 | IRR 0.95 (0.80 to 1.12) |
| Atelectasis-free ICU days | 334 (93%) | 402 (91%) | p = 0.739 | IRR 1.03 (0.88 to 1.19) |
| Pneumonia-free ICU days | 293 (94%) | 355 (92%) | p = 0.782 | IRR 1.02 (0.87 to 1.20) |
| Thrombosis-free ICU days | 415 (96%) | 473 (94%) | p = 0.788 | IRR 1.02 (0.90 to 1.17) |
| Decubiti-free ICU days | 411 (96%) | 493 (95%) | p = 0.867 | IRR 1.01 (0.89 to 1.16) |
| Contracture-free ICU days | 448 (96%) | 535 (96%) | p>0.999 | IRR 1.00 (0.88 to 1.13) |

^a Assessed by responsible physician without any prespecified criteria, assessable ICU days only (discretion of physician) (delirium: experimental n = 421, control n = 467, atelectasis: n = 258, n = 442, pneumonia: n = 313, n = 388, thrombosis: n = 431, n = 501, decubiti: n = 427, n = 518, contracture: n = 468, n = 558) Data are presented as n (%)

Definitions: Study days = number of days from study enrolment to ICU discharge including readmissions and weekends where only limited therapy service was available, session = one single physiotherapy treatment

Abbreviations: IRR = Incidence rate ratio, RD = Risk difference, PT = Physiotherapy, ICU = Intensive care unit

<https://doi.org/10.1371/journal.pone.0207428.t004>

their hospital-stay on wards or step-down units (experimental 77%, control 70%) and the main reason for missed therapy visits were weekends (58%), extended protocol-exposure with more resistance training might have enhanced potential group differences. The median ICU-stay of 6 days seemed to have been too short to properly introduce resistance training, which was defined as using weight or resistance. However, assistive exercise (experimental group: 25% of movement exercises) in the very weak patient may affect strength as well. Wright et al [37] reported a higher but still modest percentage of sessions (33%) with this type of strengthening. In their study, similar to ours, sedation and patient fatigue prevented implementation or targeted intensity. Weakness (58% with ICUAW in our study) and ongoing effects of sedative agents despite significant reduction in their use in this group may both have contributed.

The main difference between our study and previous ones reporting benefits from early mobilisation is that the control group was also mobilised early, as per our ICU standard. We started the intervention very early in both groups, while patients were still mechanically ventilated and on vasoactive support, whereas most [12, 13, 15, 37, 38], but not all other intervention groups [11, 39] started substantially later. Nevertheless, our control group received physiotherapy as early as the intervention group in the study of Schweickert et al [11], and more often than treatment groups in other trials [37, 38]. In contrast to our study, Schaller et al [39] found that early mobilisation shortened ICU length of stay and improved functional mobility in surgical ICU patients at hospital discharge. Contribution of treatment of control groups to the different results cannot be evaluated due to insufficient details of care received in Schaller's study. Yet the case mix appears very different: our patients were sicker (mean APACHE II 23 vs 16 in [39]) and had less surgical patients. It is therefore conceivable that the early and intensive physiotherapy in the control group of our study explains the lack of effect

Table 5. Treatment details for the control and experimental groups.

| | Experimental group (n = 58, 478 study days) | Control group (n = 57, 558 study days) | p value | Group difference (95% CI) |
|--|---|--|-----------|-----------------------------|
| Physiotherapy (PT) | | | | |
| Total PT visits (% per study day) | 349 (73%) | 365 (65%) | p = 0.143 | IRR 1.12 (0.96 to 1.30) |
| Total PT sessions (% per study day) | 407 (85%) | 377 (68%) | p<0.001 | IRR 1.26 (1.09 to 1.45) |
| Two or more sessions per day (% of total PT sessions) | 60 (15%) | 15 (4%) | p<0.001 | IRR 3.71 (2.08 to 7.03) |
| Treatment duration per session (min) ^a | 25.0 (IQR 19.5–27.0) | 18.0 (IQR 14.0–21.0) | p<0.001 | 6.00 (4.00 to 8.00) |
| Time from enrolment to first therapy (days) ^b | 0.16 (IQR 0.07–0.20) | 0.16 (IQR 0.07–0.48) | p = 0.387 | -0.02 (-0.07 to 0.03) |
| Time from ICU admission to first therapy (days) ^b | 1.95 (IQR 1.41–2.80) | 2.15 (IQR 1.53–2.89) | p = 0.470 | -0.14 (-0.61 to 0.24) |
| ICU treatment | | | | |
| Length of ICU stay at original hospital (days) | 5.9 (IQR 4.0–10.9) | 6.1 (IQR 4.6–12.7) | p = 0.595 | -0.37 (-1.90 to 1.13) |
| Study days at original hospital | 5.0 (IQR 3.0–10.0) | 5.0 (IQR 4.0–12.0) | p = 0.696 | 0.00 (-2.00 to 1.00) |
| Tracheostomies | 15 (26%) | 15 (26%) | p = 0.956 | RD 0% (-17% to 16%) |
| ICU re-admissions | 8 (14%) | 11 (19%) | p = 0.427 | RD -5% (-19 to 8%) |
| Daily energy intake (kcal) over whole ICU stay ^c | 1225±833 | 1152±867 | p = 0.139 | MD 73.06 (-23.74 to 169.86) |
| ICU medication | | | | |
| Study days without sedation | 132 (28%) | 93 (17%) | p<0.001 | IRR 1.66 (1.26 to 2.18) |
| Study days without opiates | 41 (9%) | 56 (10%) | p = 0.477 | IRR 0.85 (0.56 to 1.30) |
| Study days without insulin infusions | 289 (61%) | 333 (60%) | p = 0.872 | IRR 1.01 (0.86 to 1.19) |
| Study days without NMBAs | 407 (85%) | 468 (84%) | p = 0.839 | IRR 1.02 (0.89 to 1.16) |
| Study days without steroids | 427 (89%) | 422 (76%) | p = 0.016 | IRR 1.18 (1.03 to 1.35) |
| Study days without inotropes and vasopressors | 191 (40%) | 291 (52%) | p = 0.005 | IRR 0.77 (0.64 to 0.92) |

^a Median of all patients' median session duration

^b n = 57 for experimental and n = 56 for control group due to two participants death before receiving the allocated intervention

^c Excluding oral food intake

Data are presented as median (IQR), mean (SD) or n (%)

Definitions: Visits = days with at least one physiotherapy treatment, study days = number of days from study enrolment to ICU discharge including readmissions and weekends where only limited therapy service was available, session = one single physiotherapy treatment

Abbreviations: PT = Physiotherapy, ICU = Intensive care unit, NMBAs = Neuromuscular blocking agents, IRR = Incidence rate ratio, RD = Risk difference, MD = Mean Difference

<https://doi.org/10.1371/journal.pone.0207428.t005>

of additional endurance training. Early beginning of physiotherapy seems crucial due to early onset of ICUAW [4]. Accordingly, Schweickert et al [11] combined early mobilisation (within 1.7d of intubation) with daily sedation-interruption leading to reduced delirium and to significant functional improvement at hospital discharge. Delirium was equally prevalent in our groups, though the incidence was comparable to the intervention group of [11] (39 vs 33%). However, despite the early start, patients in our study achieved the specific mobilisation milestones substantially later [11]. This may also be related to the case mix—our patients were substantially older and sicker. These patients might not always be suitable for early mobilisation [40], but could profit from appropriate exercise interventions over the whole ICU stay [12]. However, recent evidence also highlights the impact of pre-ICU health [41, 42] and post-extubation dysphagia [43] on patients' functional recovery. Our heterogenous sample might therefore conceal responsive subgroups. More research is needed whether sicker patients need longer rehabilitation exposure [44] and for which subgroups recovery will be achievable.

The conducted early endurance training seemed to have unfavourable effects on mobilisation levels leading to less out-of-bed mobilisations in this group. This self-funded, pragmatic study did




Fig 2. Overview of each physiotherapy intervention for the control and experimental groups. * $p < 0.003$ More than one therapy intervention per session (= single physiotherapy treatment) possible, total sessions: 407 (85% of study days) experimental and 377 (68%) control group.

<https://doi.org/10.1371/journal.pone.0207428.g002>

not have additional resources for therapy-delivery. Physiotherapists may therefore have used available resources for cycling instead of early mobilisation. Although more out-of-bed mobilisations did not positively impact function of the control group either. The large AVERT trial even found that early, high-dose out-of-bed mobilisation after stroke enlarges the probability for disability when compared to low-dose, early mobilisation [45]. Dose-response analyses found the greatest benefit for short, frequent mobilisations [46]. This would favour the higher frequency of physiotherapy sessions (11%) in the intervention group, although this percentage might have been insufficient to reach clinical significance. Similarly, the stepwise approach of our study might have been too ambitious for the allocated therapy time (7min extra) and cycling insufficiently dosed (15min versus 30min reported by [47, 48]) to induce musculoskeletal adaptations. However, the small but significant changes in hemodynamic or oxygen-transport variables appeared to have been sufficient to stimulate patients to meet the metabolic demands of exercise by adequate cardiorespiratory adaptations. In contrast, neither passive cycling in deeply sedated patients [49] nor passive chair transfer [50] increased metabolic costs.

Our results are in line with recent studies [13, 15, 37, 38, 51] that failed to show a benefit of early rehabilitation. Heterogenous populations, difficulties in group separation and exercise dose may all have contributed. The discrepancy between planned and actual therapy delivery in these trials implies that the critically ill's ability to participate in functional exercises may be overestimated. More attention should be paid to patient fatigue and on whom to target, as very early muscle preservation might not be feasible in a seriously ill cohort with active catabolic pathways. Resistance training may therefore be more feasible during recovery.

This study had several limitations. First, blinding of participants and physiotherapists was impossible. Second, loss of follow-up at 6 months was substantial and unequally distributed between groups (41% versus 58%) but without any differences in responders and non-responders. Nevertheless, results from the SF-36 are not adequately powered, possibly attributable to type I error and need to be interpreted with caution. Third, we targeted patients at risk of ICUAW who were previously independent. Therefore, data from this trial might not be applicable to patients with previous functional dependency. Fourth, the observed difference of 23m in the 6MWD may be of clinical significance [52], however, to test a difference of 25m (SD 150) would require at least 1128 participants. Similarly, the two primary outcomes should be adjusted for multiplicity to avoid type I error.

This study also has several strengths: patients were included very early after ICU admission and received physiotherapy within 48 hours. Physiotherapy interventions were pragmatic and reflect real-world resources. The novel use of individually set limits instead of strict target numbers allows for more rehabilitation flexibility and appears to be a safe approach with a low incidence of AEs (0.6%). Nevertheless, physiotherapists were experienced in ICU rehabilitation, which is shown in the exceptionally high control group participation. With a total of 1368 edge-of-bed and 1292 out-of-bed mobilisations on 1130 ICU days this trial further underlines the importance of ICU rehabilitation as a team effort. Additionally, the blinded outcome measurements were patient-centred, valid, reliable and relevant to the devastating consequences of ICUAW.

In the light of these data future studies should assure sufficiently large sample sizes, aim to increase exercise-exposure and ensure a targeted, frequent exercise delivery to the right patient. The effect of early exercise on mental health needs further evaluation, considering that one in two survivors might experience psychiatric symptoms [53].

Conclusions

We did not find functional benefits of early exercise training over an active usual care rehabilitation in mechanically ventilated, critically ill adults. Successively-adapted, early exercise training seemed to be safe. Early endurance training might improve long-term mental health, but the most effective rehabilitation intervention to excite neuromuscular adaptations still needs to be determined.

Supporting information

S1 CONSORT checklist.

(DOCX)

S1 Fig. Intervention protocol for the experimental group [19].

(PDF)

S1 Table. Intervention details for the experimental group. Details are based on CONSORT extension for non-pharmacological interventions and the TIDieR Checklist.

(PDF)

S2 Table. Sensitivity analyses for the primary outcomes.
(DOCX)

S3 Table. Patient characteristics for ‘with versus without SF-36’ (mental health component).
(PDF)

S4 Table. Patient characteristics for ‘with versus without resistance training’.
(PDF)

S1 File. Physiological parameters while on mechanical ventilation (S2 Fig) and predominantly spontaneously breathing (S3 Fig) patients for before, during and after physiotherapy.
(PDF)

S2 File. Detailed information on performed study interventions: sessions, mobilisations and achieved mobility-milestones (S4 to S7 Fig, S5 Table).
(PDF)

S3 File. Original study protocol and English translation. This protocol has been approved by the local ethics committee prior to first patient recruitment.
(PDF)

S4 File. Original, de-identified primary and secondary outcome variables.
(XLSX)

Acknowledgments

We sincerely thank all physiotherapists and ICU staff for their dedicated care; without them this study would not have been possible. Special thanks go to coordinating physiotherapists (Valentine Stefanicki, Daphne Seyler, Andrea Perren, Thomas Nydegger) and study nurses (Margaret Jong, Laureta Fazlija, Manuela Akaltan, Michael Lensch, Marianne Roth, Sabine Berger). We also thank Lukas Bütikofer and Irina Irincheeva for the statistical advice and for reviewing the manuscript.

Author Contributions

Conceptualization: Sabrina Eggmann, Martin L. Verra, Jukka Takala, Stephan M. Jakob.

Data curation: Sabrina Eggmann.

Formal analysis: Sabrina Eggmann, Gere Luder, Stephan M. Jakob.

Investigation: Sabrina Eggmann, Martin L. Verra, Gere Luder, Jukka Takala, Stephan M. Jakob.

Methodology: Sabrina Eggmann, Martin L. Verra, Gere Luder, Jukka Takala, Stephan M. Jakob.

Project administration: Sabrina Eggmann, Stephan M. Jakob.

Resources: Sabrina Eggmann, Martin L. Verra, Jukka Takala, Stephan M. Jakob.

Software: Sabrina Eggmann.

Supervision: Sabrina Eggmann, Martin L. Verra, Stephan M. Jakob.

Validation: Sabrina Eggmann.

Visualization: Sabrina Eggmann.

Writing – original draft: Sabrina Eggmann.

Writing – review & editing: Sabrina Eggmann, Martin L. Verra, Gere Luder, Jukka Takala, Stephan M. Jakob.

References

1. Herridge MS, Tansey CM, Matte A, Tomlinson G, Diaz-Granados N, Cooper A, et al. Functional disability 5 years after acute respiratory distress syndrome. *The New England journal of medicine*. 2011; 364(14):1293–304. Epub 2011/04/08. <https://doi.org/10.1056/NEJMoa1011802> PMID: 21470008.
2. Stevens RD, Marshall SA, Cornblath DR, Hoke A, Needham DM, de Jonghe B, et al. A framework for diagnosing and classifying intensive care unit-acquired weakness. *Critical care medicine*. 2009; 37(10 Suppl):S299–308. Epub 2010/02/06. <https://doi.org/10.1097/CCM.0b013e3181b6ef67> PMID: 20046114.
3. Friedrich O, Reid MB, Van den Berghe G, Vanhorebeek I, Hermans G, Rich MM, et al. The Sick and the Weak: Neuropathies/Myopathies in the Critically Ill. *Physiological reviews*. 2015; 95(3):1025–109. Epub 2015/07/03. <https://doi.org/10.1152/physrev.00028.2014> PMID: 26133937; PubMed Central PMCID: PMC4491544.
4. Puthuchery ZA, Rawal J, McPhail M, Connolly B, Ratnayake G, Chan P, et al. Acute skeletal muscle wasting in critical illness. *JAMA: the journal of the American Medical Association*. 2013; 310(15):1591–600. Epub 2013/10/11. <https://doi.org/10.1001/jama.2013.278481> PMID: 24108501.
5. De Jonghe B, Bastuji-Garin S, Durand MC, Malissin I, Rodrigues P, Cerf C, et al. Respiratory weakness is associated with limb weakness and delayed weaning in critical illness. *Critical care medicine*. 2007; 35(9):2007–15. Epub 2007/09/15. PMID: 17855814.
6. Ali NA, O'Brien JM Jr., Hoffmann SP, Phillips G, Garland A, Finley JC, et al. Acquired weakness, hand-grip strength, and mortality in critically ill patients. *American journal of respiratory and critical care medicine*. 2008; 178(3):261–8. Epub 2008/05/31. <https://doi.org/10.1164/rccm.200712-1829OC> PMID: 18511703.
7. Hermans G, Van Mechelen H, Clerckx B, Vanhullebusch T, Mesotten D, Wilmer A, et al. Acute outcomes and 1-year mortality of intensive care unit-acquired weakness. A cohort study and propensity-matched analysis. *American journal of respiratory and critical care medicine*. 2014; 190(4):410–20. Epub 2014/05/16. <https://doi.org/10.1164/rccm.201312-2257OC> PMID: 24825371.
8. van der Schaaf M, Beelen A, Dongelmans DA, Vroom MB, Nollet F. Poor functional recovery after a critical illness: a longitudinal study. *Journal of rehabilitation medicine: official journal of the UEMS European Board of Physical and Rehabilitation Medicine*. 2009; 41(13):1041–8. Epub 2009/11/07. <https://doi.org/10.2340/16501977-0443> PMID: 19893999.
9. Pfoh ER, Wozniak AW, Colantuoni E, Dinglas VD, Mendez-Tellez PA, Shanholtz C, et al. Physical declines occurring after hospital discharge in ARDS survivors: a 5-year longitudinal study. *Intensive care medicine*. 2016; 42(10):1557–66. Epub 2016/09/18. <https://doi.org/10.1007/s00134-016-4530-1> PMID: 27637716.
10. Fan E, Dowdy DW, Colantuoni E, Mendez-Tellez PA, Sevransky JE, Shanholtz C, et al. Physical complications in acute lung injury survivors: a two-year longitudinal prospective study. *Critical care medicine*. 2014; 42(4):849–59. Epub 2013/11/20. <https://doi.org/10.1097/CCM.000000000000040> PMID: 24247473; PubMed Central PMCID: PMC3959239.
11. Schweickert WD, Pohlman MC, Pohlman AS, Nigos C, Pawlik AJ, Esbrook CL, et al. Early physical and occupational therapy in mechanically ventilated, critically ill patients: a randomised controlled trial. *Lancet*. 2009; 373(9678):1874–82. Epub 2009/05/19. [https://doi.org/10.1016/S0140-6736\(09\)60658-9](https://doi.org/10.1016/S0140-6736(09)60658-9) PMID: 19446324.
12. Burtin C, Clerckx B, Robbeets C, Ferdinande P, Langer D, Troosters T, et al. Early exercise in critically ill patients enhances short-term functional recovery. *Critical care medicine*. 2009; 37(9):2499–505. Epub 2009/07/23. <https://doi.org/10.1097/CCM.0b013e3181a38937> PMID: 19623052.
13. Denehy L, Skinner EH, Edbrooke L, Haines K, Warrillow S, Hawthorne G, et al. Exercise rehabilitation for patients with critical illness: a randomized controlled trial with 12 months of follow-up. *Critical care (London, England)*. 2013; 17(4):R156. Epub 2013/07/26. <https://doi.org/10.1186/cc12835> PMID: 23883525.
14. Kayambu G, Boots R, Paratz J. Early physical rehabilitation in intensive care patients with sepsis syndromes: a pilot randomised controlled trial. *Intensive care medicine*. 2015; 41(5):865–74. Epub 2015/04/09. <https://doi.org/10.1007/s00134-015-3763-8> PMID: 25851383.

15. Moss M, Nordon-Craft A, Malone D, Van Pelt D, Frankel SK, Warner ML, et al. A Randomized Trial of an Intensive Physical Therapy Program for Patients with Acute Respiratory Failure. *American journal of respiratory and critical care medicine*. 2016; 193(10):1101–10. Epub 2015/12/15. <https://doi.org/10.1164/rccm.201505-1039OC> PMID: 26651376; PubMed Central PMCID: PMCPMC4872662.
16. Tanner RE, Bruncker LB, Agergaard J, Barrows KM, Briggs RA, Kwon OS, et al. Age-related differences in lean mass, protein synthesis and skeletal muscle markers of proteolysis after bed rest and exercise rehabilitation. *The Journal of physiology*. 2015; 593(18):4259–73. Epub 2015/07/15. <https://doi.org/10.1113/JP270699> PMID: 26173027; PubMed Central PMCID: PMCPMC4594296.
17. McCarthy B, Casey D, Devane D, Murphy K, Murphy E, Lacasse Y. Pulmonary rehabilitation for chronic obstructive pulmonary disease. *The Cochrane database of systematic reviews*. 2015; 2:Cd003793. Epub 2015/02/24. <https://doi.org/10.1002/14651858.CD003793.pub3> PMID: 25705944.
18. Ploutz-Snyder LL, Downs M, Ryder J, Hackney K, Scott J, Buxton R, et al. Integrated resistance and aerobic exercise protects fitness during bed rest. *Medicine and science in sports and exercise*. 2014; 46(2):358–68. Epub 2014/01/21. <https://doi.org/10.1249/MSS.0b013e3182a62f85> PMID: 24441215.
19. Eggmann S, Verra ML, Luder G, Takala J, Jakob SM. Effects of early, combined endurance and resistance training in mechanically ventilated, critically ill patients: a study protocol for a randomised controlled trial. *Trials*. 2016; 17:403. Epub 2016/08/17. <https://doi.org/10.1186/s13063-016-1533-8> PMID: 27527501; PubMed Central PMCID: PMCPMC4986184.
20. Doig GS, Simpson F. Randomization and allocation concealment: a practical guide for researchers. *Journal of critical care*. 2005; 20(2):187–91; discussion 91–3. Epub 2005/09/06. <https://doi.org/10.1016/j.jcrc.2005.04.005> PMID: 16139163.
21. Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data capture (REDCap)—a metadata-driven methodology and workflow process for providing translational research informatics support. *Journal of biomedical informatics*. 2009; 42(2):377–81. Epub 2008/10/22. <https://doi.org/10.1016/j.jbi.2008.08.010> PMID: 18929686; PubMed Central PMCID: PMCPMC2700030.
22. Takala J, Dellinger RP, Koskinen K, St Andre A, Read M, Levy M, et al. Development and simultaneous application of multiple care protocols in critical care: a multicenter feasibility study. *Intensive care medicine*. 2008; 34(8):1401–10. Epub 2008/04/04. <https://doi.org/10.1007/s00134-008-1084-x> PMID: 18385977.
23. Jakob SM, Butikofer L, Berger D, Coslovsky M, Takala J. A randomized controlled pilot study to evaluate the effect of an enteral formulation designed to improve gastrointestinal tolerance in the critically ill patient—the SPIRIT trial. *Critical care (London, England)*. 2017; 21(1):140. Epub 2017/06/11. <https://doi.org/10.1186/s13054-017-1730-1> PMID: 28599662; PubMed Central PMCID: PMCPMC5466775.
24. Norrenberg M, Vincent JL. A profile of European intensive care unit physiotherapists. *European Society of Intensive Care Medicine. Intensive care medicine*. 2000; 26(7):988–94. Epub 2000/09/16. PMID: 10990117.
25. ATS statement: guidelines for the six-minute walk test. *American journal of respiratory and critical care medicine*. 2002; 166(1):111–7. Epub 2002/07/02. <https://doi.org/10.1164/ajrccm.166.1.at1102> PMID: 12091180.
26. Cournan M. Use of the functional independence measure for outcomes measurement in acute inpatient rehabilitation. *Rehabilitation nursing: the official journal of the Association of Rehabilitation Nurses*. 2011; 36(3):111–7. Epub 2011/06/17. PMID: 21675396.
27. De Jonghe B, Sharshar T, Lefaucheur JP, Authier FJ, Durand-Zaleski I, Boussarsar M, et al. Paresis acquired in the intensive care unit: a prospective multicenter study. *JAMA: the journal of the American Medical Association*. 2002; 288(22):2859–67. Epub 2002/12/11. PMID: 12472328.
28. Hermans G, Clerckx B, Vanhullebusch T, Segers J, Vanpee G, Robbeets C, et al. Interobserver agreement of Medical Research Council sum-score and handgrip strength in the intensive care unit. *Muscle & nerve*. 2012; 45(1):18–25. Epub 2011/12/23. <https://doi.org/10.1002/mus.22219> PMID: 22190301.
29. Baldwin CE, Paratz JD, Bersten AD. Muscle strength assessment in critically ill patients with handheld dynamometry: an investigation of reliability, minimal detectable change, and time to peak force generation. *Journal of critical care*. 2013; 28(1):77–86. Epub 2012/04/24. <https://doi.org/10.1016/j.jcrc.2012.03.001> PMID: 22520490.
30. Clavet H, Hebert PC, Fergusson D, Doucette S, Trudel G. Joint contracture following prolonged stay in the intensive care unit. *CMAJ: Canadian Medical Association journal = journal de l'Association medicale canadienne*. 2008; 178(6):691–7. Epub 2008/03/12. <https://doi.org/10.1503/cmaj.071056> PMID: 18332384; PubMed Central PMCID: PMCPMC2263098.
31. Podsiadlo D, Richardson S. The timed "Up & Go": a test of basic functional mobility for frail elderly persons. *Journal of the American Geriatrics Society*. 1991; 39(2):142–8. Epub 1991/02/01. PMID: 1991946.

32. Heyland DK, Hopman W, Coo H, Tranmer J, McColl MA. Long-term health-related quality of life in survivors of sepsis. Short Form 36: a valid and reliable measure of health-related quality of life. *Critical care medicine*. 2000; 28(11):3599–605. Epub 2000/12/01. PMID: [11098960](https://pubmed.ncbi.nlm.nih.gov/11098960/).
33. Redelmeier DA, Bayoumi AM, Goldstein RS, Guyatt GH. Interpreting small differences in functional status: the Six Minute Walk test in chronic lung disease patients. *American journal of respiratory and critical care medicine*. 1997; 155(4):1278–82. Epub 1997/04/01. <https://doi.org/10.1164/ajrccm.155.4.9105067> PMID: [9105067](https://pubmed.ncbi.nlm.nih.gov/9105067/).
34. Bellet RN, Adams L, Morris NR. The 6-minute walk test in outpatient cardiac rehabilitation: validity, reliability and responsiveness—a systematic review. *Physiotherapy*. 2012; 98(4):277–86. Epub 2012/11/06. <https://doi.org/10.1016/j.physio.2011.11.003> PMID: [23122432](https://pubmed.ncbi.nlm.nih.gov/23122432/).
35. Fritz CO, Morris PE, Richler JJ. Effect size estimates: current use, calculations, and interpretation. *Journal of experimental psychology General*. 2012; 141(1):2–18. Epub 2011/08/10. <https://doi.org/10.1037/a0024338> PMID: [21823805](https://pubmed.ncbi.nlm.nih.gov/21823805/).
36. Eggmann S, Verra ML, Luder G, Takala J, Jakob SM. Physiological effects and safety of an early, combined endurance and resistance training in mechanically ventilated, critically ill patients. *Physiotherapy*. 2015; 101:e344–e5. <https://doi.org/10.1016/j.physio.2015.03.553>.
37. Wright SE, Thomas K, Watson G, Baker C, Bryant A, Chadwick TJ, et al. Intensive versus standard physical rehabilitation therapy in the critically ill (EPICC): a multicentre, parallel-group, randomised controlled trial. *Thorax*. 2017. Epub 2017/08/07. <https://doi.org/10.1136/thoraxjnl-2016-209858> PMID: [28780504](https://pubmed.ncbi.nlm.nih.gov/28780504/).
38. Morris PE, Berry MJ, Files DC, Thompson JC, Hauser J, Flores L, et al. Standardized Rehabilitation and Hospital Length of Stay Among Patients With Acute Respiratory Failure: A Randomized Clinical Trial. *JAMA: the journal of the American Medical Association*. 2016; 315(24):2694–702. Epub 2016/07/02. <https://doi.org/10.1001/jama.2016.7201> PMID: [27367766](https://pubmed.ncbi.nlm.nih.gov/27367766/).
39. Schaller SJ, Anstey M, Blobner M, Edrich T, Grabitz SD, Gradwohl-Matis I, et al. Early, goal-directed mobilisation in the surgical intensive care unit: a randomised controlled trial. *Lancet*. 2016; 388(10052):1377–88. Epub 2016/10/07. [https://doi.org/10.1016/S0140-6736\(16\)31637-3](https://doi.org/10.1016/S0140-6736(16)31637-3) PMID: [27707496](https://pubmed.ncbi.nlm.nih.gov/27707496/).
40. Hodgson CL, Stiller K, Needham DM, Tipping CJ, Harrold M, Baldwin CE, et al. Expert consensus and recommendations on safety criteria for active mobilization of mechanically ventilated critically ill adults. *Critical care (London, England)*. 2014; 18(6):658. Epub 2014/12/06. <https://doi.org/10.1186/s13054-014-0658-y> PMID: [25475522](https://pubmed.ncbi.nlm.nih.gov/25475522/); PubMed Central PMCID: [PMCPmc4301888](https://pubmed.ncbi.nlm.nih.gov/PMC4301888/).
41. Ferrante LE, Pisani MA, Murphy TE, Gahbauer EA, Leo-Summers LS, Gill TM. Functional trajectories among older persons before and after critical illness. *JAMA internal medicine*. 2015; 175(4):523–9. Epub 2015/02/11. <https://doi.org/10.1001/jamainternmed.2014.7889> PMID: [25665067](https://pubmed.ncbi.nlm.nih.gov/25665067/); PubMed Central PMCID: [PMCPmc4467795](https://pubmed.ncbi.nlm.nih.gov/PMC4467795/).
42. Griffith DM, Salisbury LG, Lee RJ, Lone N, Merriweather JL, Walsh TS. Determinants of Health-Related Quality of Life After ICU: Importance of Patient Demographics, Previous Comorbidity, and Severity of Illness. *Critical care medicine*. 2018; 46(4):594–601. Epub 2018/01/03. <https://doi.org/10.1097/CCM.0000000000002952> PMID: [29293149](https://pubmed.ncbi.nlm.nih.gov/29293149/).
43. Schefold JC, Berger D, Zurcher P, Lensch M, Perren A, Jakob SM, et al. Dysphagia in Mechanically Ventilated ICU Patients (DYnAMICS): A Prospective Observational Trial. *Critical care medicine*. 2017. Epub 2017/10/13. <https://doi.org/10.1097/ccm.0000000000002765> PMID: [29023260](https://pubmed.ncbi.nlm.nih.gov/29023260/).
44. Mehrholz J, Muckel S, Oehmichen F, Pohl M. First results about recovery of walking function in patients with intensive care unit-acquired muscle weakness from the General Weakness Syndrome Therapy (GymNAST) cohort study. *BMJ open*. 2015; 5(12):e008828. Epub 2015/12/25. <https://doi.org/10.1136/bmjopen-2015-008828> PMID: [26700274](https://pubmed.ncbi.nlm.nih.gov/26700274/); PubMed Central PMCID: [PMCPmc4691758](https://pubmed.ncbi.nlm.nih.gov/PMC4691758/).
45. Efficacy and safety of very early mobilisation within 24 h of stroke onset (AVERT): a randomised controlled trial. *Lancet*. 2015; 386(9988):46–55. Epub 2015/04/22. [https://doi.org/10.1016/S0140-6736\(15\)60690-0](https://doi.org/10.1016/S0140-6736(15)60690-0) PMID: [25892679](https://pubmed.ncbi.nlm.nih.gov/25892679/).
46. Bernhardt J, Churilov L, Ellery F, Collier J, Chamberlain J, Langhorne P, et al. Prespecified dose-response analysis for A Very Early Rehabilitation Trial (AVERT). *Neurology*. 2016; 86(23):2138–45. Epub 2016/02/19. <https://doi.org/10.1212/WNL.0000000000002459> PMID: [26888985](https://pubmed.ncbi.nlm.nih.gov/26888985/); PubMed Central PMCID: [PMCPmc4898313](https://pubmed.ncbi.nlm.nih.gov/PMC4898313/).
47. Kho ME, Molloy AJ, Clarke FJ, Ajami D, McCaughan M, Obrovac K, et al. TryCYCLE: A Prospective Study of the Safety and Feasibility of Early In-Bed Cycling in Mechanically Ventilated Patients. *PloS one*. 2016; 11(12):e0167561. Epub 2016/12/29. <https://doi.org/10.1371/journal.pone.0167561> PMID: [28030555](https://pubmed.ncbi.nlm.nih.gov/28030555/); PubMed Central PMCID: [PMCPMC5193383](https://pubmed.ncbi.nlm.nih.gov/PMC5193383/)
48. Hickmann CE, Castanares-Zapatero D, Deldicque L, Van den Bergh P, Caty G, Robert A, et al. Impact of Very Early Physical Therapy During Septic Shock on Skeletal Muscle: A Randomized Controlled

- Trial. *Critical care medicine*. 2018; 46(9):1436–43. Epub 2018/06/30. <https://doi.org/10.1097/CCM.0000000000003263> PMID: 29957714; PubMed Central PMCID: PMC6110624.
49. Camargo Pires-Neto R, Fogaca Kawaguchi YM, Sayuri Hirota A, Fu C, Tanaka C, Caruso P, et al. Very early passive cycling exercise in mechanically ventilated critically ill patients: physiological and safety aspects—a case series. *PLoS one*. 2013; 8(9):e74182. Epub 2013/09/17. <https://doi.org/10.1371/journal.pone.0074182> PMID: 24040200; PubMed Central PMCID: PMC3767643.
 50. Collings N, Cusack R. A repeated measures, randomised cross-over trial, comparing the acute exercise response between passive and active sitting in critically ill patients. *BMC anesthesiology*. 2015; 15:1. Epub 2015/02/12. <https://doi.org/10.1186/1471-2253-15-1> PMID: 25670916; PubMed Central PMCID: PMC4322801.
 51. Fossat G, Baudin F, Courtes L, Bobet S, Dupont A, Bretagnol A, et al. Effect of In-Bed Leg Cycling and Electrical Stimulation of the Quadriceps on Global Muscle Strength in Critically Ill Adults: A Randomized Clinical Trial. *JAMA: the journal of the American Medical Association*. 2018; 320(4):368–78. Epub 2018/07/26. <https://doi.org/10.1001/jama.2018.9592> PMID: 30043066.
 52. Chan KS, Pfoh ER, Denehy L, Elliott D, Holland AE, Dinglas VD, et al. Construct validity and minimal important difference of 6-minute walk distance in survivors of acute respiratory failure. *Chest*. 2015; 147(5):1316–26. Epub 2015/03/06. <https://doi.org/10.1378/chest.14-1808> PMID: 25742048; PubMed Central PMCID: PMC4420183.
 53. Bienvenu OJ, Friedman LA, Colantuoni E, Dinglas VD, Sepulveda KA, Mendez-Tellez P, et al. Psychiatric symptoms after acute respiratory distress syndrome: a 5-year longitudinal study. *Intensive care medicine*. 2018; 44(1):38–47. Epub 2017/12/28. <https://doi.org/10.1007/s00134-017-5009-4> PMID: 29279973; PubMed Central PMCID: PMC6020022.