

Inhibitory activity of *Halobacillus trueperi* S61 and its active extracts on potato dry rot

Shuo Shen^{a,b,c,d,e}, Wei Li^{a,b,c,d,e}, and Jian Wang^{a,b,c,d,e}

^aAcademy of Agriculture and Forestry, Qinghai University, Xining, Qinghai 810016 China; ^bKey Laboratory of Potato Breeding in Qinghai Province, Xining, Qinghai 810016 China; ^cState Key Laboratory of Plateau Ecology and Agriculture, Qinghai University, Xining, Qinghai 810016 China; ^dKey Laboratory of Qinghai Tibet Plateau Biotechnology, Ministry of Education Xining, Qinghai 810016 China; ^eNorthwest Potato Engineering Research Center, Ministry of Education, Xining, Qinghai 810016 China

ABSTRACT


This study investigated the inhibitory activity of *Halobacillus trueperi* S61 and its active extract on potato dry rot pathogens and aimed at contributing to biocontrol agent development during potato storage. Three kinds of pathogens were isolated as target pathogenic fungi from dry rot tubers and determined as *Fusarium acuminatum* (Qing 9A-2), *Fusarium equiseti* (Qing 9A-5-8) and *Fusarium tricinctum* (Qing 9A-1-1) by morphological and molecular identification. The strain *Halobacillus trueperi* S61 and its extract exhibited a higher inhibitory rate on both three pathogens (56.32–65.75 and 1.67–51.11%), notably the best suppression efficiency is presented in *Halobacillus trueperi* S61 and 40 mg/mL ethyl acetate extract. In terms of in vivo effects, both *Halobacillus trueperi* S61 and its ethyl acetate extract effectively reduced the decayed fruit and weight loss rate (0–20% and 7.59–16.56%) and enhanced the defensive enzymatic activities to improve resistance. In addition, strain S61 could be colonized on potato tubers, especially the highest amount of 1.55×10^7 CFU/mL on fifth day for variety Xiazhai 65. Overall, *Halobacillus trueperi* S61 and its ethyl acetate extract could be considered as potential approach for biocontrol potato dry rot.



ARTICLE HISTORY

Received 26 November 2021
Revised 27 December 2021
Accepted 27 December 2021

KEYWORDS

Potato dry rot; *Halophilic bacteria* S61; inhibitory activity; biocontrol



CONTACT Wei Li  lwbabyw@163.com  Academy of Agriculture and Forestry, Qinghai University, Xining, Qinghai 810016, China

Novelty Statement This study investigated the inhibitory activity of *Halobacillus trueperi* S61 and its active extract on potato dry rot pathogens and aimed at contributing to biocontrol agent development during potato storage. The dominant pathogens of potato dry rot were *F. acuminatum*, *F. equiseti* and *F. tricinctum* and strong pathogenicity acting on variety Xiazhai 65. The strain S61 and its extract exhibited a higher inhibitory rate on both three pathogens, notably the best suppression efficiency is presented in *Halobacillus trueperi* S61 in vivo and vitro. Overall, *Halobacillus trueperi* S61 could be considered as potential approach for biocontrol potato dry rot.

© 2022 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

Potato (*Solanum tuberosum* L.) as the fourth largest food crop and the main non-cereal crop, has contributed to food and nutrition security in developing countries [1]. The global potato cultivated area is 19.2 Mha with an annual production of 376.8 MT, of which 90.3 tons is produced in China [2]. However, potato crops are vulnerable to various phytopathogens further threatening quality and yield. Among them, post-harvest diseases induced by fungal pathogens cause significant economic losses worldwide [3,4]. For instance, early or late blight, *Verticillium* wilt, and powdery scab, especially dry rot as one kind of most serious postharvest diseases which causes 6%–60% yield loss annually and mainly infested by single or multi kinds of *Fusarium* [5]. There are reported 13 species associated with *Fusarium* dry rot worldwide [6] and Bojanowski et al. [7] isolated the predominant pathogens of *F. oxysporum* and *F. avenaceum* in China. The pathogen *Fusarium* capable resistance to low temperature and cause a high-level infectivity during potato storage. Apart from destroying tuber tissue, *Fusarium* also releases toxins such as fusaric acid accelerate transpiration and causes potato tubers to wilted, *enniatin* and *naphthazarins* disturb the energy supply, *zearalenone* damage the selective permeability of cell membranes and suppress the enzyme synthesis during plant growth [8–10]. After suffering from *Fusarium* dry rot, the potato surface appears necrotic wrinkles with brown to black, and the internal tissue infected by white, yellow or pink hyphae which cause shriveled, shrank and lesions, finally reduced the edible and economic value of potatoes [11].

Therefore, it is imperative to effectively prevent dry rot by agricultural, chemical and biological approaches [12–14]. Agricultural control is mainly focused on seed tubers selection and pretreatment, tillage approach, transportation or storage process. Chemical control is spraying chemical agents such as thiabendazole, fenpiclonil and thiabendazole, but it is easy to interfere with soil physicochemical properties then harmful to beneficial microbial metabolism and cause pathogens resistant [6,7]. Microbial biocontrol has attracted more and more attention due to environment-friendly and wide distribution, large number and complex diversity of metabolites of microorganisms. There are

mainly through direct use of microbes or extract active ingredients for prevention and biocontrol pathogens. The mechanism of microbial control has been reported to be direct inhibited pathogens by biocontrol of primary and secondary metabolites as well as induce plants resistance [12]. At present, numerous endophytic microbials are effective for biocontrol plant diseases. The main strains for biocontrol potato dry rot are *Bacillus*, *Pseudomonas* and *Actinomycetes* [15]. Some studies have found that the spraying of *P. fluorescens* and *Enterobacter cloacae* could significantly inhibit the *Fusarium sambucinum* pathogen, notably *Bacillus* is outstanding in inhibiting fungi activity and promoting plant growth [16]. Mnif et al. [17] found that the extract of *Bacillus subtilis* could break hyphae and reduce the sporulation rate of *Fusarium*. Although, large number of active *Bacillus* have been found in extreme salt conditions such as *Bacillus rigiliprofundii* 1MBB1T, *Bacillus iranensis* X5BT and *Bacillus shacheensis* HNA-14 T [1], while only limited studies focus on the biocontrol of dry rot.

The *Halobacillus trueperi* isolated from extreme environments and has great potential to develop novel biocontrol agents due to stronger survivability and flexible adaptability. Additionally, dry rot pathogens widely occur in potato planting areas and their types varied with seasons, geographic locations and cultivars [18]. At present, there are few studies on inhibitory effects of moderately halophilic microbes on potato dry rot pathogen. Accordingly, this study identified the pathogens in three kinds of main potato types and detected pathogenicity, further evaluate the inhibitory activity of *Halobacillus trueperi* S61 and its organic solvent extracts (ethyl acetate, n-butanol, chloroform and petroleum ether extraction) on potato dry rot in vitro and vivo. The hypothesis that *Halobacillus trueperi* S61 enables to effectively inhibit *Fusarium* pathogens and aims at providing novel biocontrol approach for potato dry rot prevention.

2. Material and methods

2.1. Experimental materials

The potato varieties of Xiazhai 65, Qingshu 9 and Qingshu 2 were provided by the Institute of

Biotechnology, Academy of Agriculture and Forestry Sciences (Qinghai University, China). The pathogens of Qing 9A-1-1, Qing 9A-5-8 and Qing 9A-2 were isolated from the diseased Qingshu 9 tuber, after purified then stored at 4°C. The strain *Halobacillus trueperi* S61 was collected from the Qrhan Salt Lake mud, then isolated and purified through the cultivation method. The potato dextrose agar (PDA) medium was made of 200 g glucose and potato, added distilled water to 1000 mL and adjusted pH to 7.2–7.4. The American type culture collection (ATCC) 213 medium comprised 10 g MgSO₄ · 7H₂O, 0.2g CaCl₂ · 2H₂O, 5 g KCl, 2.5 g peptone, 10 g yeast extract, 40 g NaCl and 12 g agar powder, then added distilled water to 1000 mL and adjusted pH to 7.2–7.4.

2.2. Pathogenicity determination and identification of potato dry rot pathogens

2.2.1. Morphological and molecular biological identification of pathogens

The pathogens Qing 9A-1-1, Qing 9A-5-8 and Qing 9A-2 were activated and picked up the hyphae then inoculated on PDA medium plate by inoculation needle, further cultivated at 28°C for 7 days in constant temperature incubator (HZQ-F160A). Observe and record the substrate color, mycelial shape, and colony size from naked eyes, as well as observe mycelia and spore morphology under microscope, finally perform morphological identification according to the *Fusarium* identification manual.

For molecular biological identification, the mycelia of activated potato dry rot pathogens Qing 9A-1-1, Qing 9A-5-8 and Qing 9A-2 were scraped from the culture medium plate, then extracted deoxyribonucleic acid (DNA) according to Ezup column fungal genomic DNA extraction kit (Sangon Biotech, Shanghai, China) and performed polymerase chain reaction (PCR) amplification based on primers ITS1 and ITS4 (5'-CTTGGTCATTTAGAGGAAGTAA-3' and 5'-TCCTCCGCTTATTGATATGC-3'). The amplification system was 2× PCR Mix 10 µL, primer 2 µL, DNA template 1 µL, added distilled water to volume 50 µL. The PCR reaction conditions included pre-denaturation at 95°C for 5 min, denaturation at 94°C for 30 s, annealing at 55°C for 30 s, extension at 72°C for 50 s, perform 35 cycles and repair extension at 72°C for 8 min. After the process, tested 5 µL PCR amplified product by agarose gel electrophoresis, then

sent the recovered product to Sangon for sequencing. The obtained ITS sequence of pathogenic fungi was submitted to GenBank to obtain the accession number, then compared in the GenBank database of the United States Biotechnology Information Center. Finally, draw phylogenetic tree according to Neighbor-Joining method in MEGA 7.0 software.

2.2.2. In vivo pathogenicity determination of pathogens

The concentration of the three pathogens suspensions Qing 9A-2, Qing 9A-1-1, and Qing 9A-5-8 was calculated using hemocytometer under microscope and added distilled water to adjusted 1×10^7 CFU/mL by high-purity water machine (GREEN-Q3-20 T), then named as A, B and C, respectively. The single infection was inoculated with A, B and C individually, and the compound infection was inoculated with AB, AC, BC and ABC. Selected each variety of ten potatoes (Qingshu 9, Qingshu 2 and Xiazhai 65) with consistent size, free of disease and mechanical damage, the tubers were soaked in 2% sodium hypochlorite for 15 min and then disinfected with 70% ethanol. After natural drying, punch one hole in each tuber using an aseptic puncher (diameter 15 mm) and inject 100 µL single or compound infection pathogen suspension. After 7 days storage in the incubator at 28°C and 70%–80% relative humidity, the lesion diameter and decay were measured. According to the following formula to calculate the decayed fruit rate and weight loss rate:

- (1) Decayed fruit rate (%) = $\frac{\text{Decayed fruit number}}{\text{Total fruit number}} \times 100\%$
- (2) Weight loss rate (%) = $\frac{(\text{Weight before storage} - \text{Weight after storage})}{\text{Weight before storage}} \times 100\%$

2.3. Inhibitory activity of strain S61 and its extracts on potato dry rot pathogens

2.3.1. Preparation of organic solvent extract from *Halobacillus trueperi* S61

The active strain S61 was inoculated into ATCC213 liquid medium and cultured in a shaker at 28°C 180 rpm for 7 days. Collected the fermentation broth and filtered through a Brinell funnel, then obtained filtrate and mixed with equal volume of ethyl acetate, petroleum ether, chloroform and n-butanol at 1:1, the

mixture was suspended on rotary evaporator (EYELA N-1300) for three times to obtain strain S61 extracts.

2.3.2 *In vitro* inhibitory activity of *Halobacillus trueperi* S61 and its extracts on potato dry rot pathogens

The inhibitory effect of *Halobacillus trueperi* S61 on potato dry rot was determined by plate confrontation method. Cross petri dishes with marker and pour into PDA medium, then inoculated the activated *Halobacillus trueperi* S61 after cultivating in ATCC213 solid medium at 28°C. After 7 days cultivation, the pathogen diameter is measured and the inhibition rate of S61 on pathogens is calculated according to the following formula:

$$(1) \text{ Inhibition rate (\%)} = (\text{Control colony diameter} - \text{Treated colony diameter}) / \text{Control colony diameter} \times 100\%$$

The *Halobacillus trueperi* S61 extract was prepared into 5, 10, 20 and 40 mg/mL, then 100 µL of the extract was dripped into Oxford cup and compared with distilled water with three replicates. According to the Oxford cup method, the culture dish was crossed with marker and poured into PDA medium, the sterilized Oxford cup was placed in the PDA medium at 2.5 cm from the medium center. *Fusarium* cultured for 7 d were cut by puncher (diameter 10 mm), and one part was placed in the center of each treatment plate. After incubation at 28°C for 4 d, the colony diameter of pathogens is measured and the inhibition rate is calculated.

2.4. *In vivo* inhibition effect of strain S61 and its extracts on potato dry rot pathogens

2.4.1. Potato fruit treatment

Select disease-free potato tubers with consistent shape and size and soak 20 min in 5% sodium hypochlorite, after naturally dried then soaked in alcohol for 2 min, then punched one hole (diameter 1 cm) on the surface of the tuber after naturally dried.

2.4.2. Electron microscopic observation of strain S61 ethyl acetate extract on pathogens

Sterilize petri dishes and pour into PDA, inoculated with Qing 9A-2, 9A-5-8 and 9A-1-1, then cultured in constant temperature incubator for 7 days. Fresh pathogen mycelia were picked and fixed in EP tube

containing glutar-aldehyde fixative then sent to Kechuang Biological Company (Qingdao, China) for scanning electron microscope observation.

2.4.3. *In vivo* inhibition of strain S61 and its ethyl acetate extract on potato dry rot pathogens

The pathogens group is inoculated 20 mL pathogen suspension in the hole, while the control 1 and 2 (CK1 and CK2) were inoculated with 20 mL distilled water and S61 suspension, control 3 and 4 (CK3 and CK4) were inoculated with 20 mL distilled water and 40 mg/mL S61 ethyl acetate extract. Cultivated in constant temperature incubator at 28°C and 60% relative humidity for 7, 14, 21 and 28 days, then measured weight loss rate and decayed fruit rate.

2.4.4. Enzymatic activity determination

In the sterile environment, 1.0 g potato tubers were cut and put into the sterilized EP tube with a sterilized steel ball, and oscillated for 2 min in a 16,000-rpm shaker. The obtained frozen potato powder was fixed by adding phosphate buffer solution, then centrifuged at 5000 rpm and filtered 1 mL of supernatant for analysis. The enzyme activity of peroxidase, polyphenol oxidase, superoxide dismutase and catalase was determined by corresponding determination kit which was purchased from Jiancheng Biological Company (Nanjing, China).

2.5. Colonization of strain *Halobacillus trueperi* S61 on potato tubers

2.5.1. Strain *Halobacillus trueperi* S61 cultivation and fermentation broth preparation

The *Halobacillus trueperi* S61 was activated on ATCC213 medium without antibiotics and cultured in a constant temperature incubator at 28°C until a single colony grew. The single colonies were picked up by burning inoculation needles and transferred to ATCC213 liquid medium, then cultured with shaking at 150 rpm and 28°C to obtain the strain S61 liquid culture medium.

2.5.2. Inoculation of strain *Halobacillus trueperi* S61 on potato tubers

The potato tubers were soaked in 2% sodium hypochlorite for 2 min, rinsed with distilled water and dried at room temperature, disinfected with 70% alcohol. The S61 liquid culture medium was collected and

centrifuged at 4000 rpm for 5 min, 1 mL of distilled water was added after absorbing the supernatant, and adjusted S61 to OD600 of 0.1 then sprayed on the pretreated potato tubers.

2.5.3 Colonization of strain *Halobacillus trueperi* S61 on potato tubers

On the 1st, 2nd, 3rd and 4 to 7th day, 1 g potato was cut and placed in a sterile conical flask containing 10 mL 1% NaCl at 150 rpm for 30 min. Then, extraction of 100 μ L diluted 10 times and spread in ATCC213 solid medium, cultivated 24 h in constant temperature incubator then counted the colonies number and calculated the colonization amount.

3. Result and discussion

In order to fully understand the pathogenesis of potato dry rot and develop possible biological control approaches, present study identified the dominant dry rot pathogens from three main varieties of potatoes through morphological and molecular analysis. Then, the *pathogenicity* through decayed fruit and weight loss as well as infection rate was evaluated. Importantly, *in vitro* and *in vivo* evaluated the inhibitory effect of *Halobacillus trueperi* S61 and its organic extracts on the potato dry rot pathogen and aimed at reveal the *Halobacillus trueperi* S61 biocontrol capacity and contribute to microbial biocontrol of dry rot.

3.1. Pathogenicity determination and identification of potato dry rot pathogens

3.1.1. Morphological and biological identification of pathogens

After pathogens cultivated in incubator at 28°C for 7 days, the characteristics of pathogen Qing 9A-1-1 were shown white hyphae with felt shape while non-discoloring of medium surface (Figure 1(a)), as well as the observed morphology under microscope was mostly rod-shaped hyphae with few branches and round spores (Figure 1(b)). Regarding Qing 9A-2 appeared felt shape hyphae with white and reddish color and the medium surface changed to red color (Figure 1(c)), as well as the observed morphology was rod-shaped hyphae with less branched and numerous oval and ovoid micro-conidia (Figure 1(d)). Concerning Qing 9A-5-8 was demonstrated white color hyphae with flocculent shape, and the culture

medium showed yellowish color (Figure 1(e)), as well as the observed morphology was bent hyphae without branches, mostly septate and round or oval spores (Figure 1(f)). Similarly, Tiwari et al. [1] performed the identification of *Fusarium* dry rot based on morphology, and observed oval, elliptical and slender macro-conidia, white and brick orange mycelia, pale violet tinge and pale-yellow media. Bayona et al. [19] observed potato dry rot symptom with oval and kidney shaped micro-conidia, white and light purple aerial mycelium, as well as tan and light purple medium.

For identified the pathogens, ITS sequence of three pathogens were logged into GenBank database for BLAST retrieval, and the sequences with more than 98% similarity were selected from Genbank database for homology comparison, then build phylogenetic tree (Figure 2). The results demonstrated that the Qing 9A-5-8 (MW737419) and *Fusarium equiseti* (KY365574) gathered one branch (Figure 2(a)), Qing 9A-2 (MW737420) and *Fusarium acuminatum* (KT965739.1) clustered one branch (Figure 2(b)), Qing 9A-1-1 (MW741756) and *Fusarium tricinctum* (MT180474.1) gathered one branch (Figure 2(c)). Thus, three dominant pathogens of potato dry rot Qing 9A-5-8, Qing 9A-2 and Qing 9A-1-1 were affiliated to *Fusarium equiseti*, *Fusarium acuminatum* and *Fusarium tricinctum*. Schisler et al. [20] identified 14 kinds of main *Fusarium* causing potato dry rot and varied depending on geographic differences, among which *F. sambucinum*, *F. solani* and *F. avenaceum* are the dominant pathogens.

3.1.2. Identification the pathogenicity of single and compound pathogens infection on potato

The single and compound infection of three pathogens on distinct variety of potatoes were evaluated (Figure 3). There were obvious shriveled or necrotic brown-black spots on the tuber surface of the three varieties, while the control was asymptomatic. The effect of single pathogens infect on Xiazhai 65 was greatest with the highest decayed fruit rate and weight loss rate (18.30–24.10 and 40–80%), while the lowest pathogenicity rate and weight loss rate identified in Qingshu 2 (9.70–18.70 and 0–20%). Notably the pathogenicity of Qing 9A-1-1 was greatest with 20–60% decayed fruit rate and 12.30–17.50% weight loss rate. Regarding compound pathogens infection, Qing 9A-2 and Qing 9A-1-1 were strongest with highest




Figure 1. Morphological identification of Qing 9A-1-1 (a) on medium and (b) microscope (400), Qing 9A-2 (c) on medium and (d) microscope (400), and Qing A-5-8 (e) on medium and (f) microscope (400).

weight loss rate and decayed fruit rate (15.70–19.10 and 20–80%) as compared to other combinations (7.20–18.90 and 0–40%), the highest value appeared

in Xiazhai 65 (9.4–19.1 and 20–80%) and lowest in Qingshu 2 (8.7–15.7 and 0–20%) (Table 1). Therefore, distinct potato varieties have different sensitivity




Figure 2. Phylogenetic tree based on sequences (a) Qing A-5-8, (b) Qing 9A-2 and (c) Qing 9A-1-1.

responses to distinct pathogens, the pathogenicity of compound infection was lower than single infection which inconsistency with Tiwari et al. [1] who recorded the synergistic effect between *Fusarium spp.*

and combined inoculation shows higher invasiveness than single inoculation. Maleki et al. [4] also pointed out that combined treatment was superior than individually in vitro and vivo condition.



Figure 3. Pathogenicity effect of pathogens on potato in vivo. A was infection effect of pathogen on Qingshu 9, B was infection effect of pathogen on Qingshu 2, and C was infection effect of pathogen on Xiazhai 65. From left to right were infected by sterile water, Qing 9A-2, Qing 9A-1-1, Qing 9A-2 combine Qing 9A-1-1.

As for infection rate of single or compound pathogens demonstrated the fastest infection rate of Qing 9A-2 (83 mm/d), followed by Qing 9A-1-1 and Qing 9A-5-8 (58 and 44 mm/d) when infected on Xiazhai 65. For variety Qingshu 9, the fastest infection rate was 9A-2 (46 mm/d), followed by Qing 9A-1-1 and Qing 9A-5-8 (39 and 24 mm/d), while there is almost no infection on Qingshu 2. The strongest compound infection of Qing 9A-2 combined with Qing 9A-1-1 has highest infection rates on Xiazhai 65, Qingshu 9 and Qingshu 2 (83, 72 and 57 mm/d) (Table 2). Heltoft et al. [21] pointed out that *F. sambucinum*, *F. graminearum*, *F. solani* and *F. oxysporum* had severe lesions on most cultivars. Tiwari et al. [1] also proved higher infection of *F. sambucinum* than *F. oxysporum* (with lesion diameter 38–39.5 and

27.5–29 mm). Sana et al. [16] isolated 450 strains from potato and found Ar 10 shown the highest inhibition activity with 18 mm inhibited diameter. Khedher et al. [3] inoculated *F. solani*, *F. graminearum*, *F. sambucinum*, and *F. oxysporum* conidial suspensions on potato, observed clearly lesion diameter reduction (42.8–52.2%) after treated with *B. subtilis* V26. Xue et al. [8] proved the potato trichothecenes varied with distinct cultivars, *Fusarium* types and storage conditions. Overall, both three pathogens have a certain level infection on the three main variety potatoes, and the single infection ability was strongest in Qing A-2, followed by Qing 9A-1-1 and Qing 9A-5-8. Considering the pathogenicity and weight loss rate, the variety Xiazhai 65 was more susceptible than Qingshu 9 and Qingshu 2 to

Table 1. Effect of single and combine infection on three potato varieties.

Variety	Qing 9A-1-1			Qing 9A-2			Qing 9A-5-8			AB			AC			BC			ABC		
	Weight loss rate	Decayed fruit rate	Weight loss rate	Decayed fruit rate	Weight loss rate	Decayed fruit rate	Weight loss rate	Decayed fruit rate	Weight loss rate	Decayed fruit rate	Weight loss rate	Decayed fruit rate	Weight loss rate	Decayed fruit rate	Weight loss rate	Decayed fruit rate	Weight loss rate	Decayed fruit rate	Weight loss rate	Decayed fruit rate	
Qingshu ₂	18.70%	20%	12.40%	20%	9.70%	0%	15.70%	20%	14.40%	20%	10.80%	0%	8.70%	0%	0%	0%	10.80%	20%	8.70%	0%	
Xiazhai ₆₅	24.10%	60%	20.90%	80%	18.30%	40%	19.10%	80%	18.90%	40%	9.40%	20%	9.90%	20%	20%	20%	9.40%	40%	9.90%	20%	
Qingshu ₉	17.50%	40%	16.60%	60%	12.30%	20%	17.50%	60%	14.60%	40%	9.30%	20%	7.20%	20%	20%	20%	9.30%	40%	7.20%	20%	

A is Qing 9A-2, B is Qing 9A-1-1, and C is Qing 9A-5-8.

Table 2. Infection rate of single infection on three potato varieties.

Pathogen	Qingshu 9	Xiazhai 65	Qingshu 2
Qing 9A-2	$Y = 0.46X + 0.11$	$Y = 0.83X - 0.73$	-
Qing 9A-5-8	$Y = 0.24X + 0.45$	$Y = 0.44X + 0.76$	-
Qing 9A-1-1	$Y = 0.39X - 0.58$	$Y = 0.58X + 0.43$	-
AB	$Y = 0.72X + 0.63$	$Y = 0.83X - 0.36$	$Y = 0.57X + 0.12$
AC	$Y = 0.63X + 0.45$	$Y = 0.74X + 0.16$	$Y = 0.38X + 0.62$
BC	$Y = 0.41X - 0.53$	$Y = 0.34X + 0.41$	-
ABC	$Y = 0.32X - 0.27$	$Y = 2.28X - 3.6$	$Y = 0.32X + 0.81$

A is Qing 9A-2, B is Qing 9A-1-1, and C is Qing 9A-5-8.

Fusarium dry rot pathogens, and compound inoculum *Fusarium acuminatum* and *Fusarium tricinctum* have stronger pathogenicity than individual infection.

3.2. Inhibitory activity of *Halobacillus trueperi* S61 and extracts on potato dry rot pathogens

The inhibitory rate of *Halobacillus trueperi* S61 and its extracts (ethyl acetate, n-butanol, chloroform and petroleum ether extracts prepared into 5.0, 10.0, 20.0 and 40.0 mg/mL concentrations) on pathogens were detected (Table 3). Firstly, the pathogen diameter of *Halobacillus trueperi* S61 was significantly smaller (24.7–35.3 mm) and the inhibition rates of Qing 9A-1-1, Qing 9A-2 and Qing 9A-5-8 were 65.75%, 58.93% and 56.32%, which indicated the inhibitory activity of S61 on potato dry rot pathogens (Figure 4). Khedher et al. [3] reported the *Bacillus subtilis* V26 has obviously antifungal activity with reduced *Fusarium* diameter and mycelial (10.8–19.7%), and highest inhibition rate exhibited on *F. sambucinum* and lowest in *F. oxysporum* (85.3 and 54.7%). Kolaei et al. [5] pointed out that metabisulfite salts inhibited all selected fungi and 200 mM capable inhibit dry rot 100%.

Regarding S61 extracts, the inhibition rates of the three pathogens improved with the increased concentration (5.0–40.0 mg/mL). As to Qing 9A-1-1, the inhibition rate was 36.67–51.11% for ethyl acetate, 4.33–12.67% for n-butanol, 6.67–23.33% for chloroform, 4.67–11.00% for petroleum ether extracts. Regarding Qing 9A-2, the inhibition rate was 17.76–40.00% for ethyl acetate, 3.30–9.76% for n-butanol, 3.67–22.21% for chloroform and 6.00–8.00% for petroleum ether extracts. For Qing 9A-5-8, the inhibition rate was 25.65–38.89% for ethyl acetate, 3.00–8.33% for n-butanol, 6.00–11.34% for chloroform and 1.67–6.67% for petroleum ether extracts (Table 3). Notably,

the S61 ethyl acetate extract has the highest inhibitory rate of 51.11%, 40.00% and 38.89% for three pathogens at the concentration of 40.0 mg/mL. As comparison, the S61 chloroform extract has higher inhibitory rate on Qing 9A-1-1 and Qing 9A-2 (23.33% and 22.21%) at the concentration of 40.0mg/mL while less effect on Qing 9A-5-8 (11.34%), while other extracts present lower inhibitory rate (3.00–12.67%) (Table 3). Ren et al. [7] compared distinct Black spruce extraction and proved ethyl acetate extraction has great antimicrobial potential. Elsherbiny et al. [9] perform methanol extraction of pomegranate peel demonstrated remarkable activity on *F. sambucinum* inhibition with 23.7–75.5% from 1.25–20 mg/ml concentration, greatest in 20 mg/ml to prevent potato dry rot. Kolaei et al. [5] identified there was no effect of organic solvents dichloromethane, chloroform, and ethyl acetate on antimicrobial activity while ethyl acetate extract has higher recovery and activity. The current study exhibited that *Halobacillus trueperi* S61 and its active extracts could inhibit the growth of pathogens, the ethyl acetate extract had the highest activity and the strongest inhibitory activity of S61 was acting on Qing 9A-1-1 (*Fusarium tricinctum*).


3.3. In vivo inhibitory effect of strain S61 and its extracts on potato dry rot pathogens

3.3.1. In vivo inhibition of strain S61 and its ethyl acetate extract on pathogens

Three pathogens were inhibited by *Halobacillus trueperi* S61 ethyl acetate extract, the hyphae all collapsed and shriveled at certain level and part of the hyphae was broken, and the growth point malformed and ultimate swelled (Figure 5(a-c)). Importantly, weight loss rate and decayed fruit rate are essential for potato quality during storage, water loss and pathogen infection during storage are vital factors for weight loss rate and strengthen with time, as well as the decayed fruit rate directly reflected the pathogens infection [22,23]. The weight loss rate of potato tubers was significantly reduced after inoculate S61 suspension (7.59%) and S61 ethyl acetate extract (7.63%) than single or compound infection (8.94–13.67% and 12.33–16.56%) (Table 4). After sprayed S61 suspension, the strongest single infection ability of Qing 9A-2 has 20% incidence and 13.6% weight loss rate, which were 60% and 8.3% lower than unsprayed S61 tubers. The compound infection of Qing 9A-2 and Qing 9A-1-1 had 10%

Table 3. Inhibitory activity of strain S61 and its ethyl acetate extract on pathogens.

Strains	Concentration (mg/ml)	Qing 9A-1-1		Qing 9A-2		Qing 9A-5-8	
		Diameter/mm	Inhibitory rate/%	Diameter/mm	Inhibitory rate/%	Diameter/mm	Inhibitory rate/%
16S strain	-	24.7 ± 2.7b	65.75%	32.0 ± 3.5ab	58.93%	35.3 ± 01.6a	56.32%
Ethyl acetate extract	5	21.0 ± 2.08a	30.00%	25.3 ± 1.49a	16.65%	25.6 ± 1.20a	14.44%
	10	19.0 ± 2.30b	36.67%	25.0 ± 1.52a	17.76%	22.3 ± 0.30b	25.65%
	20	17.3 ± 1.52c	42.20%	20.0 ± 0.50b	33.34%	21.3 ± 0.56b	28.98%
	40	14.6 ± 0.88d	51.11%	18.0 ± 1.12 c	40.00%	18.3 ± 0.86 c	38.89%
N-butanol extract	5	28.7 ± 0.23a	4.33%	29.0 ± 0.16a	3.30%	29.1 ± 0.08a	3.00%
	10	27.9 ± 0.41b	7.67%	28.1 ± 0.18b	6.30%	28.0 ± 0.57a	6.67%
	20	27.5 ± 0.25b	8.43%	28.5 ± 0.47b	5.00%	27.8 ± 0.38b	7.30%
	40	26.6 ± 0.21b	12.67%	27.1 ± 0.25 c	9.76%	27.5 ± 0.86b	8.33%
Chloroform extract	5	28.8 ± 1.79a	6.67%	28.92 ± 1.55b	3.67%	28.2 ± 1.96a	6.00%
	10	25.3 ± 0.95b	15.56%	27.4 ± 1.20 c	8.76%	28.7 ± 1.37a	4.33%
	20	24.3 ± 1.45 c	16.77%	26.6 ± 1.16a	11.34%	27.3 ± 0.96b	9.65%
	40	23.0 ± 0.65 c	23.33%	23.3 ± 1.18b	22.21%	26.7 ± 1.72b	11.34%
Petroleum ether extract	5	28.6 ± 0.60a	4.67%	28.2 ± 1.96a	6.00%	29.5 ± 1.49a	1.67%
	10	28.9 ± 1.51a	3.76%	28.1 ± 1.37a	6.33%	29.5 ± 1.62a	1.67%
	20	27.5 ± 1.41b	8.33%	27.6 ± 0.96b	8.00%	29.0 ± 0.70a	3.33%
	40	27.2 ± 1.81 c	11.00%	27.3 ± 1.72b	8.00%	28.8 ± 1.35b	6.67%
CK		30	-	30		30	-

**Figure 4.** Inhibition of strain S61 (a) Qing 9A-2, (b) Qing 9A-1-1 and (c) Qing 9A-5-8 on three pathogenic bacteria in vitro, as well as confrontation result of Qing 9A-2 and S61 (d), Qing 9A-5-8 and S61 (e), Qing 9A-1-1 and S61 (f) on the medium.

incidence and 11.6% weight loss rate, which were 70% and 7.5% lower than unsprayed S61 tubers (Figure 5(d)). After sprayed S61 ethyl acetate extract, the incidence and weight loss rate of inoculated Qing 9A-2 were 20% and 13.5%, which were 60% and 7.4% lower than non-sprayed tubers.

The compound infection of Qing 9A-2 and Qing 9A-1-1 had 20% incidence which was 60% lower than

non-sprayed tubers, and 15.4% weight loss rates which similarly with control (Figure 5(e)). Khedher et al. [3] evaluated the effect of *B. subtilis* V26 on *F. solani*, *F. graminearum*, *F. sambucinum*, and *F. oxysporum*. Sana et al. [16] detected the antagonistic strains Ar10 treated 72 h shown positive effect on potato tubers protection with highest weight loss rate, necrosis and depth area (85 and 100%). Finally, inoculated




Figure 5. Electron microscopic observation of S61 on pathogens (a) Qing 9A-2, (b) Qing 9A-1-1 and (c) Qing A-5-8, as well as the in vivo effect of spraying S61 (d) and its ethyl acetate extract (e) on Qing 9A-2, and colonization of S61 on potato tubers (f).

Table 4. Control effect of spraying s61 suspension and its ethyl acetate extract.

Treatment	Decayed fruit rate	Weight loss rate	Treatment	Decayed fruit rate	Weight loss rate
16S strain			Ethyl acetate extract		
A	20%	13.67%	A	20%	16.56%
B	0%	9.89%	B	0%	10.32%
AB	10%	11.76%	AB	20%	15.47%
ABC	0%	8.49%	ABC	10%	12.33%
CK1	0%	7.23%	CK3	0%	8.89%
CK2	0%	7.59%	CK4	0%	7.63%

A is Qing 9A-2, B is Qing 9A-1-1, and C is Qing 9A-5-8, CK1 is inoculated sterile water, CK2 is inoculated S61 suspension, CK3 is inoculated sterile water, CK4 is inoculated S61 extract.

Halobacillus trueperi S61 and ethyl acetate extract remarkably reduced weight loss rate and decayed fruit rate, three cultivars exhibited distinct efficacy in terms of pathogenicity and resistance activity, and the *Halobacillus trueperi* S61 demonstrated biocontrol capacity of dry rot pathogen infection.

3.3.2. Response of enzymatic activity after sprayed strain S61 and ethyl acetate extract

The biocontrol of soil-borne pathogens is related to antibiotics and hydrolytic enzymes, and the disease resistance of plants was closely associated with enzyme activity of the defense enzyme systems [3]. It is reported that beneficial microorganisms could activate the defense mechanism of plant reproductive system [24–26]. Peroxidase (POD) as one kind of plant defense enzyme systems and improved environmental resistance, and the activity was lower in control

sprayed with S61 and ethyl acetate extraction ($0.6 \text{ U} \cdot \text{min}^{-1} \text{mg}^{-1}$). While the POD activity of sprayed with ethyl acetate extract was higher than S61 (7.3 – 12.2 and 7.1 – $12.1 \text{ U} \cdot \text{min}^{-1} \text{mg}^{-1}$), indicated that the stronger bio-control effect of *Halobacillus trueperi* S61 (Figure 6(a-b)). Regarding to polyphenol oxidase (PPO) and superoxide dismutase (SOD) that enable to delay plant senescence and pathological changes, the activity continuously increased from 1–14 days (0.35 – 0.80 and 2.41 – $2.93 \text{ U} \cdot \text{min}^{-1} \text{mg}^{-1}$), while the control earlier reached the peak value at 7 days (0.71 and $2.62 \text{ U} \cdot \text{min}^{-1} \text{mg}^{-1}$) and then dropped down, which proved the S61 and its ethyl acetate extract strengthen the resistance to dry rot pathogens (Figure 6c-f). Moreover, catalase (CAT) is capable of resisting the poison of hydrogen peroxide during the tuber infection process. The activity increased initially and achieved maximum value on 21




Figure 6. Response of enzymatic activity after sprayed strain S61 and its extract: peroxidase (POD) after spraying S61 (a) and its ethyl acetate extract (b), oxidase (PPO) after spraying S61 (c) and ethyl acetate extract (d), superoxide dismutase (SOD) after spraying S61 (e) and ethyl acetate extract (f), catalase (CAT) after spraying S61 (g) and ethyl acetate extract (h).

d ($100 \text{ U}\cdot\text{min}^{-1}\text{mg}^{-1}$) for S61, 14 d ($120 \text{ U}\cdot\text{min}^{-1}\text{mg}^{-1}$) for S61 ethyl acetate extract, while 7 d for control ($60 \text{ U}\cdot\text{min}^{-1}\text{mg}^{-1}$), which indicated that both S61 and its ethyl acetate extract were able to inhibit pathogens, and spraying S61 suspension was faster (Figure 6(g-h)).

Therefore, the change in defense enzyme activity associated with stress resistance and the increased activity effectively relieved the infection of potato dry rot pathogen. Similar phenomena also found by Al-Mughrabi et al. [22] who detected the increased activity of CAT, SOD and MDA when potato tubers were infected by potato dry rot. As reported, the *Fusarium* mainly produces variety of enzymes to degrade the cell wall, causing the potato tubers to corroded and further water loss and shriveled [27,28]. Khedhe et al. [3] proved that V26 is capable of producing cellulase, protease, pectinase and glucanase which could lyse some pathogenic fungi to promote plant growth and inhibit disease. *Bacillus* is a well-known bio-surfactant producer with broad-spectrum antibiotic activity and demonstrated significant antifungal and antiviral activities [16,29,30]. Khedher et al. [3] reported *Bacillus spp.* inhibited *F. sambucinum* and produced chitinases to protect potato tubers. Overall, the in vivo assay proved the antagonistic ability of *Halobacillus trueperi* S16 and its ethyl acetate extract on three kinds of *Fusarium* pathogens and sprayed S61 may have stronger competitiveness for nutrient and metabolic space which suppressed dry rot pathogens.

3.4. Colonization of *Halobacillus trueperi* S61 on potato tubers

Biocontrol agents based on bacteria and fungi have an ideal niche to control potato dry rot which attracted widespread attention [28,29]. Previous studies have found that *Bacillus amyloliquefaciens* could form good colonization and endogenous in maize, then used for biocontrol of maize leaf spot. The strains that can be colonize and endogenous in wheat are used for wheat scab/ head blight control [17–21]. In this study, *Halobacillus trueperi* S16 showed distinct activities on distinct kinds of *Fusarium* pathogens, which might be due to diverse factors such as biocontrol agent type or strain efficiency, pathogen type or

invasiveness and the host susceptibility [31]. The number of *Halobacillus trueperi* S16 detected in potato showed an upward trend when spraying for 1 to 5 d, and the colonization of strain S61 in potato tuber reached the maximum of 1.55×10^7 CFU/mL at 5 d. Thereafter, the colonization of strain S61 in potato tuber decreased during 5 to 7 days (Figure 5(f)). This indicated that *Halobacillus trueperi* S16 could form a certain amount of endogeneity and well colonized on Xiazhai 65 potato. In conclusion, the observation in vitro and vivo supported that *Halobacillus trueperi* S16 has great potential as a biocontrol agent for *Fusarium* potato dry rot.

4. Conclusion

The dominant pathogens of potato dry rot were *Fusarium acuminatum*, *Fusarium equiseti* and *Fusarium tricinctum*. Both pathogens have a certain level of pathogenicity on the three main variety potatoes, especially combine *Fusarium acuminatum* and *Fusarium tricinctum* has stronger pathogenicity than individual infection. The variety Xiazhai 65 was more susceptible than Qingshu 9 and Qingshu 2 to *Fusarium* dry rot. The *Halobacillus trueperi* S61 and its active extracts inhibited the growth of potato dry rot as well as enhanced the defense enzyme activity, notably the ethyl acetate extract had the highest activity and the strongest in vitro and vivo inhibitory activity of S61 was acting on Qing 9A-1-1. Overall, *Halobacillus trueperi* S61 is a promising novel biocontrol agent for *Fusarium* potato dry rot prevention.

Highlight

- Dry rot pathogens dominant by *F. acuminatum*, *F. equiseti* and *F. tricinctum*.
- Variety Xiazhai 65 was most susceptible to *Fusarium* dry rot than Qingshu 9 and Qingshu 2.
- The strongest pathogen inhibitory activity of S61 was acting on Qing 9A-1-1 in vitro and vivo.
- The *Halobacillus trueperi* S61 and ethyl acetate extracts enhanced the defense enzyme activity.

- *Halobacillus trueperi* S61 was considered as potential approach for biocontrol potato dry rot.

Acknowledgements

The authors are very grateful for the help of the General Project of Natural Science Foundation of Qinghai Science and Technology Department (2019-ZJ-914), National Modern Agricultural Technology System (CARS-10).

Disclosure statement

No potential conflict of interest was reported by the author(s).

Funding

We are very grateful for the help of the General Project of Natural Science Foundation of Qinghai Science and Technology Department (2019-ZJ-914), National Modern Agricultural Technology System (CARS-10).

References

- [1] Tiwari R, Bashyal B, Shanmugam V, et al. Impact of *Fusarium* dry rot on physicochemical attributes of potato tubers during postharvest storage. *Postharvest Biol Tec.* 2021;181:111638.
- [2] Bahmani K, Hasanzadeh N, Harighi B, et al. Isolation and identification of endophytic bacteria from potato tissues and their effects as biological control agents against bacterial wilt. *Physiol Mol Plant P.* 2021;116:101692.
- [3] Khedher S, Mejdoub-Trabelsi B, Tounsi S. Biological potential of *Bacillus subtilis* V26 for the control of *Fusarium* wilt and tuber dry rot on potato caused by *Fusarium* species and the promotion of plant growth. *Biol Control.* 2021;152:104444.
- [4] Hajian-Maleki H, Baghaee-Ravari S, Moghaddam M. Herbal essential oils exert a preservative effect against the potato soft rot disease. *Sci Hort.* 2021;285:110192.
- [5] Kolaei E, Tweddell R, Avis T. Antifungal activity of sulfur-containing salts against the development of carrot cavity spot and potato dry rot. *Postharvest Biol Tec.* 2012;63(1):55–59.
- [6] Elsherbiny E, Amin B, Baka Z. Efficiency of pomegranate (*Punica granatum* L.) peels extract as a high potential natural tool towards *Fusarium* dry rot on potato tubers. *Postharvest Biol Tec.* 2016;111:256–263.
- [7] Bojanowski A, Avis T, Pelletier S, et al. Management of potato dry rot. *Postharvest Biol Tec.* 2013;84:99–109.
- [8] Chehri K, Ghasempour H, Karimi N. Molecular phylogenetic and pathogenetic characterization of *Fusarium solani* species complex (FSSC), the cause of dry rot on potato in Iran. 2014; *Microbia Pathogenesis.* 67–68:14–19.
- [9] Recep K, Fikretin S, Erkol D, et al. Biological control of the potato dry rot caused by *Fusarium* species using PGPR strains. *Biol Control.* 2009;50(2):194–198.
- [10] Ren J, Tong J, Li P, et al. Chitosan is an effective inhibitor against potato dry rot caused by *Fusarium oxysporum*. *Physiol Mol Plant P.* 2021;113:101601.
- [11] Xue H, Bi Y, Tang Y, et al. Effect of cultivars, *Fusarium* strains and storage temperature on trichothecenes production in inoculated potato tubers. *Food Chem.* 2014;151:236–242.
- [12] Shen F, Wu W, Han X, et al. Study on the occurrence law and green control of grape gray mold from the perspective of ecological balance. *Bioengineered.* 2021;12(1):779–790.
- [13] Sleator RD. Under the microscope: from pathogens to probiotics and back. *Bioengineered.* 2015;6(5):275–282.
- [14] Zaritsky A, Dov E, Borovsky D, et al. Transgenic organisms expressing genes from *Bacillus thuringiensis* to combat insect pests. *Bioengineered.* 2010;1(5):341–344.
- [15] Selim H, Goma N, Essa A. Application of endophytic bacteria for the biocontrol of *Rhizoctonia solani* (*Cantharellales: ceratobasidiaceae*) damping-off disease in cotton seedlings. *Biocontrol Sci Technol.* 2017;27(1):81–95.
- [16] Sana A, Ben S, Ines K, et al. Biological control of the soft rot bacterium *Pectobacterium carotovorum* by *Bacillus amylo-liquefaciens* strain Ar10 producing glycolipid-like compounds. *Microbiol Res.* 2018;217:23–33.
- [17] Mnif I, Hammami I, Triki M, et al. Antifungal efficiency of a lipopeptide biosurfactant derived from *Bacillus subtilis* SPB1 versus the phytopathogenic fungus, *Fusarium solani*. *Environ Sci Pollut R Int.* 2015;22(22):18137–18147.
- [18] Tiwari R, Kumar R, Sharma S, et al. Potato dry rot disease: current status, pathogenomics and management. 3 *Biotech.* 2020;10(11):503.
- [19] Bayona L, Grajales A, Cárdenas M, et al. Isolation and characterization of two strains of *Fusarium oxysporum* causing potato dry rot in *Solanum tuberosum* in Colombia. *Rev Iberoam Micol.* 2011;28(4):166–172.
- [20] Schisler D, Slininger P, Olsen N. Appraisal of selected osmoprotectants and carriers for formulating Gram-negative biocontrol agents active against *Fusarium* dry rot on potatoes in storage. *Biol Control.* 2016;98:1–10.
- [21] Heltoft P, Molteberg E, Nærstad R, et al. Effect of maturity level and potato cultivar on development of *Fusarium* dry rot in Norway. *Potato Res.* 2015;58(3):205–219.
- [22] Al-Mughrabi K. Biological control of *Fusarium* dry rot and other potato tuber diseases using *Pseudomonas fluorescens* and *Enterobacter cloacae*. *Biol Control.* 2010;53(3):280–284.
- [23] Borgi M, Boudebouze S, Mkaouar H, et al. *Bacillus* phytases: current status and future prospects. *Bioengineered.* 2015;6(4):233–236.
- [24] Bonanomi G, Lorito M, Vinale F, et al. Organic amendments, beneficial microbes, and soil microbiota: toward a unified framework for disease suppression. *Annu Rev Phytopathol.* 2018;56(1):1–20.

- [25] Yu P, Zhang Y, Gu D. Production optimization of a heat-tolerant alkaline pectinase from *Bacillus subtilis* ZGL14 and its purification and characterization. *Bioengineered*. 2017;8(5):613–623. 5.
- [26] Zhang C, Li Y, Zhang T, et al. Increasing chitosanase production in *Bacillus cereus* by a novel mutagenesis and screen method. *Bioengineered*. 2021;12(1):266–277.
- [27] Boivin M, Bourdeau N, Barnabe S, et al. Black spruce extracts reveal antimicrobial and sprout suppressive potentials to prevent potato (*Solanum tuberosum* L.) losses during storage. *J Agr Food Res*. 2021;5:100187.
- [28] Aftab M, Zafar A, Iqbal I, et al. Optimization of saccharification potential of recombinant xylanase from *Bacillus licheniformis*. *Bioengineered*. 2018;9(1):159–165.
- [29] Ibrahim M, Griko N, Junker M, et al. *Bacillus thuringiensis*. *Bioengineered*. 2010;1(1):31–50.
- [30] O'Mahony M, Henneberger R, Selvin J, et al. Inhibition of the growth of *Bacillus subtilis* DSM10 by a newly discovered antibacterial protein from the soil metagenome. *Bioengineered*. 2015;6(2):89–98.
- [31] Wang Y, Shao Y, Zou X, et al. Synergistic action between extracellular products from white-rot fungus and cellulase significantly improves enzymatic hydrolysis. *Bioengineered*. 2018;9(1):178–185.