

# Chromosome Segregation in Budding Yeast: Sister Chromatid Cohesion and Related Mechanisms

Adele L. Marston

The Wellcome Trust Centre for Cell Biology, School of Biological Sciences, University of Edinburgh, Edinburgh, EH9 3JR, United Kingdom

**ABSTRACT** Studies on budding yeast have exposed the highly conserved mechanisms by which duplicated chromosomes are evenly distributed to daughter cells at the metaphase–anaphase transition. The establishment of proteinaceous bridges between sister chromatids, a function provided by a ring-shaped complex known as cohesin, is central to accurate segregation. It is the destruction of this cohesin that triggers the segregation of chromosomes following their proper attachment to microtubules. Since it is irreversible, this process must be tightly controlled and driven to completion. Furthermore, during meiosis, modifications must be put in place to allow the segregation of maternal and paternal chromosomes in the first division for gamete formation. Here, I review the pioneering work from budding yeast that has led to a molecular understanding of the establishment and destruction of cohesion.

## TABLE OF CONTENTS

Abstract	31
Introduction	32
Building Mitotic Chromosomes	32
<i>Structure and function of the cohesin complex</i>	32
<i>Discovery of cohesion:</i>	32
<i>The cohesin ring:</i>	33
<i>How does cohesin hold chromosomes together?:</i>	34
<i>Loading cohesin onto chromosomes</i>	34
<i>The pattern of cohesin localization on chromosomes:</i>	35
<i>The cohesin loading reaction:</i>	36
<i>Replication–coupled establishment of cohesion</i>	37
<i>Eco1-directed cohesion establishment:</i>	37
<i>Other factors involved in cohesion establishment:</i>	38
<i>DNA damage-induced cohesion</i>	39
<i>Other structural components of chromosomes</i>	39
<i>The condensin complex:</i>	40
<i>Topoisomerase II:</i>	40
Establishment of Biorientation	40
<i>Role of kinetochore geometry and centromere structure</i>	41

*Continued*

Copyright © 2014 by the Genetics Society of America

doi: 10.1534/genetics.112.145144

Manuscript received July 24, 2013; accepted for publication September 18, 2013

Available freely online through the author-supported open access option.

Address for correspondence: School of Biological Sciences, Michael Swann Building, Mayfield Rd., Edinburgh EH9 3JR, United Kingdom. E-mail: adele.marston@ed.ac.uk

## CONTENTS, *continued*

<i>Error correction</i>	42
Destruction of Sister Chromatid Cohesion and Anaphase Onset	43
<i>Cleavage of cohesin by separase</i>	43
<i>The spindle checkpoint</i>	43
<i>Inhibition of APC–Cdc20:</i>	43
<i>Checkpoint silencing:</i>	45
<i>Other factors regulating anaphase onset</i>	45
Anaphase Progression	46
<i>Separase initiates mitotic exit</i>	46
<i>Committing to anaphase:</i>	47
<i>Nuclear position:</i>	47
<i>Completion of chromosome segregation:</i>	47
<i>Maintaining spindle integrity:</i>	48
Chromosome Segregation During Meiosis	48
<i>Establishment of links between homologs during meiosis I</i>	49
<i>Homologous chromosome pairing:</i>	49
<i>Meiotic recombination:</i>	49
<i>Monoorientation of sister chromatids during meiosis I</i>	50
<i>Stepwise loss of cohesion</i>	51
<i>Biorientation of homologs</i>	51
<i>Alteration of cell cycle controls in meiosis</i>	52
<i>Meiotic prophase I to meiosis I transition:</i>	52
<i>Meiosis I to meiosis II transition:</i>	52
Perspectives	53


**D**URING cell division, chromosomes must be replicated exactly and accurately distributed into daughter cells. The regulated sequence of events that leads to cell division is known as the cell cycle. In S phase of the cell cycle, DNA synthesis and the establishment of sister chromatid cohesion generates two identical sister chromatids that are held tightly together by a conserved protein complex, known as cohesin. In mitosis, the cohesin linkages provide resistance and generate tension to facilitate the attachment of sister chromatids to microtubules emanating from opposite poles. Once all the chromosomes have properly attached to microtubules, an enzyme known as separase becomes active and cleaves cohesin, thereby triggering the separation of sister chromatids to opposite poles (Figure 1). This process is modified during meiosis, which produces haploid gametes from a diploid progenitor cell. During meiosis, two rounds of chromosome segregation follow a single S phase. In meiosis I, the maternal and paternal chromosomes, called homologs, are segregated, whereas sister chromatids are segregated during meiosis II, which resembles mitosis (Figure 2). To achieve this, an additional layer of regulation must be introduced. While conserved in eukaryotes, what we know about the molecular biology of chromosome segregation is derived largely from work on the budding yeast *Saccharomyces cerevisiae*. Here I review the discoveries in budding

yeast that led to an understanding of the molecular biology of chromosome segregation together with the exquisite controls that ensure its accuracy and the modifications that take place to generate gametes.

## Building Mitotic Chromosomes

### *Structure and function of the cohesin complex*

**Discovery of cohesion:** Pioneering studies in budding yeast were instrumental in the discovery of the chromosome segregation machinery that is conserved in all eukaryotes. Early on, it was recognized that the two sister chromatids must be held tightly together at metaphase to resist spindle forces, thereby allowing their attachment to microtubules from opposite poles. However, the nature of this cohesion was not known and two general models were put forward. One model postulated that the intertwining of sister DNA molecules, perhaps due to the persistence of catenations after DNA replication, might provide cohesion (Murray and Szostak 1985). Another, not mutually exclusive, model proposed the existence of proteins that generate molecular bridges between sister chromatids. However, testing these models relied on the establishment of an assay for sister chromatid cohesion. This was initially achieved by the development of a fluorescence *in situ* hybridization assay



**Figure 1** Chromosome segregation during mitosis. Schematic diagram showing the key features of chromosome segregation during budding yeast mitosis.


(FISH) in yeast (Koshland and Hartwell 1987). Using this assay, it was shown that minichromosomes are cohesed at metaphase even though they lack catenations, dispelling the idea that DNA catenation was sufficient to provide the “glue” (Koshland and Hartwell 1987; Guacci *et al.* 1994). This prompted the search for proteins that might mediate cohesion. A key technical development was the ingenious development of a method to label a single chromosome by integration of tandem repeats of bacterial *lacO*, to which ectopically produced LacI–GFP binds (Straight *et al.* 1996). A similar system was developed using *tetO* and TetR–GFP (Michaelis *et al.* 1997). The availability of these methods to label single chromosomes enabled the first cohesion proteins to be identified (Guacci *et al.* 1997; Michaelis *et al.* 1997). These elegant studies isolated mutants incapable of maintaining sister chromatid cohesion when arrested in mitosis. Subsequent studies revealed that sister chromatid cohesion genes fall into functional classes (Table 1). One class of genes encodes the proteins that make up the structural component of cohesion, called cohesin. Others are accessory, loading, or establishment factors. Remarkable progress has been made in understanding how these many gene products interact to generate sister chromatid cohesion.

**The cohesin ring:** The core structural component of cohesin forms a ring, composed of two structural maintenance of chromosome (SMC) proteins, *Smc1* and *Smc3*, and a “kleisin” (from the Greek for closure) subunit, *Scc1/Mcd1* (Guacci *et al.* 1997; Michaelis *et al.* 1997; Losada *et al.* 1998) (Figure 3). A meiosis-specific kleisin, *Rec8*, replaces *Scc1* in meiotic cells and plays several roles important for the segregation of homologous chromosomes (see below). SMC proteins are conserved from prokaryotes to eukaryotes and are composed of globular N and C termini, joined by a large coiled-coil domain that is separated by a central “hinge” domain (Nasmyth and Haering 2005). Like bacterial SMC proteins, insect cell-produced yeast *Smc1* and *Smc3* fold back on themselves at the hinge region to form antiparallel intramolecular coiled coils (Melby *et al.* 1998; Haering *et al.* 2002). This arrangement juxtaposes the Walker A-containing N terminus and Walker B-containing C terminus of a single SMC protein to generate an ATP nucleotide binding domain (NBD) of the ABC family (Hopfner *et al.* 2000; Lowe *et al.* 2001). The N terminus of each SMC protein also con-

tains a signature motif that is required for the activity of ABC family ATPases. *Smc1* NBD crystallized as a dimer with ATP sandwiched between the Walker A motif of one monomer and the signature motif on the other. In reality, *Smc1* and *Smc3* heterodimerize at their hinge domains to create a V-shaped structure (Anderson *et al.* 2002; Haering *et al.* 2002). Therefore, the most likely arrangement is that two molecules of ATP are sandwiched between the *Smc1* and *Smc3* NBDs. Consistently, fluorescence resonance energy transfer (FRET) experiments indicated that *Smc1* and *Smc3* NBD domains are in close proximity (Mc Intyre *et al.* 2007).

The kleisin subunit, *Scc1*, forms a bridge between the NBDs of the *Smc1*–*Smc3* heterodimer, making contacts with *Smc3* at its N terminus and *Smc1* at its C terminus (Haering *et al.* 2002). A crystal structure revealed that the *Scc1* C terminus forms a winged helix domain that contacts the *Smc1* NBD and mutations in this interface demonstrated that this interaction is essential (Haering *et al.* 2004). Interestingly, prior binding of the *Scc1* C terminus to the *Smc1* NBD is required for the *Scc1* N terminus to bind the *Smc3* NBD (Arumugam *et al.* 2003; Haering *et al.* 2004). This may ensure that a single molecule of *Scc1* binds to the *Smc1*–*Smc3* heterodimer. Although ATP binding to *Smc1*’s NBD is required for binding to the *Scc1* C terminus, the interaction of *Scc1*’s N terminus with *Smc3* does not require ATP (Arumugam *et al.* 2003; Gruber *et al.* 2003). An explanation for this observation is offered by the arrangement of a bacterial *Smc*–kleisin complex. While the C terminus of a bacterial kleisin contacts the ATPase head of one *Smc* protein, its N terminus associates with the coil-coiled domain of the other *Smc* subunit (Bürmann *et al.* 2013). It seems likely that *Scc1* adopts a similar asymmetric arrangement in eukaryotic cohesin; however, confirmation will await structural analysis of the *Smc3*–*Scc1* interaction.

The *Scc3* subunit binds to the central domain of *Scc1* and completes cohesin (Haering *et al.* 2002). *Scc3* is essential for the establishment, though not the maintenance, of cohesion (Toth *et al.* 1999; Kulemzina *et al.* 2012). Similarly, *Pds5* protein is also associated with cohesin and important for cohesion establishment (Hartman *et al.* 2000; Panizza *et al.* 2000; Kulemzina *et al.* 2012). Live cell imaging measurements of fluorescently tagged proteins suggest that *Pds5*, *Smc3*, and *Scc3* exist on chromosomes in a 1:1:1 ratio (Chan



**Figure 2** Chromosome segregation during meiosis. Schematic diagram showing the key features of chromosome segregation during budding yeast meiosis.

*et al.* 2012). However, the structures of Pds5 and Scc3 and their molecular function in cohesin establishment are not yet known.

**How does cohesin hold chromosomes together?:** The realization that the Smc1, Smc3, and Scc1 cohesin subunits form a ring-like structure *in vitro* led to the “embrace” model for cohesin (Haering *et al.* 2002). This model proposes that cohesin is the result of topologically embracing the two sister DNA molecules within a cohesin ring and that opening of the ring, due to cleavage of its Scc1 subunit by separase, liberates the sister chromatids, thereby destroying cohesin. Although this model has an attractive simplicity, others have argued that the ring structure of cohesin may not be the relevant cohesive form on chromatin and alternative models have been suggested (Haering *et al.* 2002; Milutinovich and Koshland 2003; Huang *et al.* 2005). Rather than interacting topologically with the DNA, these models suggest that cohesin binds to the DNA of one sister chromatid and then oligomerizes with one or more cohesin molecules bound to the other sister chromatid. Variations of these models include the “snap” model and “bracelet” models which postulate that cohesin oligomerization occurs through the coiled coil or hinge domain, respectively (Milutinovich and Koshland 2003; Huang *et al.* 2005).

Support for the idea that chromosome-bound cohesin is a ring came with the finding that cohesin subunits remain associated with each other, but not with the chromosomes, after cleavage within the coiled-coil domain of Smc3 or at the separase recognition sites in Scc1 (Gruber *et al.* 2003). Evidence that cohesin interacts topologically with the DNA came from experiments showing that cohesin is released from purified and cohesed circular minichromosomes after cleavage of either the DNA or cohesin (Ivanov and Nasmyth 2005, 2007). A more rigorous demonstration that cohesin interacts topologically with DNA came from experiments where all three interfaces in the cohesin ring were covalently

sealed either by use of fusion proteins or the introduction of side chains that allowed specific chemical cross-linking of cohesin subunits (Haering *et al.* 2008). After protein denaturation, this chemically closed cohesin ring maintained its association with 2.3-kb or 26-kb circular minichromosomes, but not with a 42-kb linear minichromosome (Haering *et al.* 2008; Farcas *et al.* 2011). This provides further support for the topological embrace model and is consistent with the idea that sliding of cohesin along chromatin fibers is normally prevented by the presence of chromatin-bound proteins.

The fact that 26-kb circular and 42-kb linear minichromosomes, which, unlike 2.3-kb minichromosomes, are catenated *in vivo*, allowed Nasmyth and colleagues to finally test the contribution of DNA catenations to cohesin. Importantly, they found that the persistence of catenanes after S phase is dependent on cohesin (Farcas *et al.* 2011). Therefore, cohesin holds sister chromosomes together by preventing the resolution of catenanes, as well as through a direct topological embrace. Nevertheless, cohesin is sufficient to hold sister chromatids together in the absence of catenations, whereas the reverse is not true (Farcas *et al.* 2011). This argues that the direct topological embrace of sister chromatids by cohesin is its critical physical property.

#### Loading cohesin onto chromosomes

To provide cohesin, cohesin must first be loaded onto chromosomes before S phase. Loading of cohesin onto chromosomes requires a separate “loader” complex composed of Scc2 and Scc4 proteins (Ciosk *et al.* 2000). A DNA replication-coupled process converts loaded cohesin into functional cohesin (Uhlmann and Nasmyth 1998) after which Scc2/Scc4 are no longer required (Ciosk *et al.* 2000). Recently it has become apparent that cohesin loading occurs at preferred chromosomal sites that are recognized by Scc2/Scc4. Analysis of mutants in the Smc subunits of cohesin that disrupt ATP binding or hydrolysis have provided insight into the cohesin loading reaction.

**Table 1 Genes involved in generating cohesion**

Function	Gene	Features
Core cohesin subunit	Smc1	Coiled-coil ATPase
	Smc3	Coiled-coil ATPase
	Scc1/Mcd1/Rad21	Kleisin subunit, cleaved by separase
	Rec8	Meiosis-specific kleisin, replaces Scc1
Cohesin associated	Scc3	Cohesion establishment
	Pds5	Cohesion establishment
	Wpl1	Destabilizes cohesin's association with chromosomes
Cohesin loading	Scc2	Required for cohesin's association with chromosomes
	Scc4	Required for cohesin's association with chromosomes
Cohesion establishment	Eco1	Acetyl transferase, acetylates Smc3

### **The pattern of cohesin localization on chromosomes:**


Genome-wide studies have examined the localization of cohesion and its loader. Cohesin is present along chromosomes; however, it is not uniformly associated with all regions of the genome. Cohesin-associated regions (CARs) typically extend for 1–4 kb, are spaced at 2- to 35-kb intervals, and tend to correlate with intergenic regions between convergent genes (Blat and Kleckner 1999; Hartman *et al.* 2000; Laloraya *et al.* 2000; Glynn *et al.* 2004; Lengronne *et al.* 2004). However, the most notable feature of cohesin binding to chromosomes is its enrichment in a large (20–50 kb) region surrounding the small (~125 bp) centromere (Blat and Kleckner 1999; Megee *et al.* 1999; Tanaka *et al.* 1999; Glynn *et al.* 2004; Weber *et al.* 2004; Kiburz *et al.* 2005). This region of cohesin enrichment surrounding the centromere defines the budding yeast pericentromere, which differs from that in other eukaryotes in that it lacks heterochromatin. The enrichment of cohesin within the pericentromere is functionally important, as its absence leads to increased chromosome loss (Tanaka *et al.* 2000; Eckert *et al.* 2007; Fernius and Marston 2009; Ng *et al.* 2009). One clear role of pericentromeric cohesion is to facilitate the proper biorientation of sister chromatids on the metaphase spindle, perhaps by generating the appropriate geometry for this interaction (Ng *et al.* 2009). Additionally, pericentromeric cohesion is critical for accurate segregation during meiosis (see below).

Genome-wide mapping of *Scc2* and *Scc4* association reported a pattern that was distinct from that of cohesin (Lengronne *et al.* 2004). These sites of *Scc2/Scc4* association likely represent cohesin-loading sites as it is here that cohesin is first detected upon cell cycle entry (Lengronne *et al.* 2004). Subsequently, cohesin translocates away from its loading sites to generate the pattern observed at metaphase (Lengronne *et al.* 2004). Transcription has been suggested to contribute to cohesin translocation and the chromosomal locations of cohesin are indeed altered by transcription, though it is unclear if this occurs as a result of cohesin sliding along the chromatin fiber or some other method of translocation (Lengronne *et al.* 2004; Bausch *et al.* 2007; Ocampo-Hafalla *et al.* 2007).

The chromosomal features that are recognized by *Scc2/Scc4* and therefore define the sites of cohesin loading are not well understood. However, the best-studied site for

cohesin loading is at the centromere. Initially, centromere sequences were found to promote cohesin recruitment to minichromosomes (Megee and Koshland 1999; Megee *et al.* 1999). Moreover, relocation of a centromere to a chromosomal arm site set up a domain of enriched cohesin surrounding the ectopic centromere, while eliminating cohesin enrichment at the endogenous pericentromere (Tanaka *et al.* 1999; Weber *et al.* 2004). Consistent with the idea that the centromere is a cohesin-loading site, it shows robust association of *Scc2/Scc4* (Lengronne *et al.* 2004; Kogut *et al.* 2009; Hu *et al.* 2011). However, the extent of the *Scc2/Scc4*-associated domain in centromeric regions has been debated. Although one report suggested a similar profile of *Scc2* and cohesin throughout the pericentromere (Kogut *et al.* 2009), others have found that *Scc2/Scc4* is localized predominantly within the core (~125 bp) centromere, in a much narrower domain than cohesin (Lengronne *et al.* 2004; Hu *et al.* 2011). While the former report implies that cohesin loading occurs throughout the pericentromere, the latter suggests that cohesin loaded at the core centromere translocates into the pericentromere. In live cells, GFP-tagged cohesin forms a pericentromeric barrel between clustered sister kinetochores (Yeh *et al.* 2008). In contrast, *Scc2*-GFP forms foci that colocalize with kinetochores, consistent with distinct localizations of cohesin and its loader (Hu *et al.* 2011). Furthermore, cohesin appears at the core centromere earlier in the cell cycle than at the pericentromere (Fernius *et al.* 2013). The available evidence for cohesin enrichment at the pericentromere is therefore most consistent with the loading and translocation model, though mechanistic details are still lacking.

The factors that attract *Scc2/Scc4* to specific sites on chromosomes are poorly defined. Regions of high transcriptional activity by PolII (tRNA genes), PolIII, and PolIII (rDNA) are correlated with *Scc2/Scc4* localization and specific induction of gene expression leads to *Scc2/Scc4* recruitment at that site (D'Ambrosio *et al.* 2008b). However, the low level of *Scc2/Scc4* at these sites has hampered analysis and the factors involved in its recruitment are not known. Recently, however, factors required for *Scc2/Scc4* association with centromeres have been identified. The centromere directs the assembly of the kinetochore, a large multisubunit complex that mediates the binding of chromosomes to




**Figure 3** Cohesin and condensin structure. Models for the relative arrangements of subunits of cohesin and condensin are shown. Both complexes form a tripartite ring made up of two SMC proteins and a kleisin subunit, which is thought to make asymmetric contacts with the SMC proteins. Both complexes have additional subunits associated with the kleisin subunit, though their exact arrangement is not well defined.

microtubules (Westermann and Schleiffer 2013). Mutants lacking components of the *Ctf19* kinetochore subcomplex (many of which are nonessential) fail to enrich cohesin within the pericentromere, though chromosome arm sites are not affected (Eckert *et al.* 2007; Fernius and Marston 2009; Ng *et al.* 2009; Fernius *et al.* 2013). Consistently, *Ctf19* complex mutants show cohesion defects that are specific to centromere-proximal sites (Fernius and Marston 2009; Ng *et al.* 2009). *Scc2/Scc4* fails to associate with the centromere in *Ctf19* complex mutants (Ng *et al.* 2009; Fernius *et al.* 2013). However, artificially tethering *Scc4* to the centromere rescues cohesin recruitment and the cohesion defects of *Ctf19* complex mutants (Fernius *et al.* 2013). This demonstrates that the *Ctf19* complex directs cohesin enrichment within the pericentromere by attracting *Scc2/Scc4* to centromeres. Recently, the *Ctf19* complex was shown to enable the association of the *Dbf4*-dependent kinase (DDK) with kinetochores from telophase to G1 phase (Natsume *et al.* 2013). Furthermore, kinetochore-associated DDK is required to attract *Scc2/Scc4* to centromeres in late G1 to ensure proper cohesin enrichment in the pericentromere (Natsume *et al.* 2013). Interestingly, kinetochore-associated DDK is also independently important to promote the early replication of centromeres (Natsume *et al.* 2013). Centromeric DDK therefore couples early replication of the pericentromere with enhanced cohesin loading. However, it remains unclear how DDK attracts *Scc2/Scc4* and cohesin loading to centromeres or whether DDK recruitment is the only critical role of the *Ctf19* complex in cohesion establishment.

**The cohesin loading reaction:** How cohesin comes to topologically embrace DNA is still very mysterious but some critical steps in the loading reaction are beginning to emerge (Figure 4). The first step in cohesin loading is preassembly of the loading complex composed of cohesin and *Scc2/Scc4*. Although it was initially assumed that the *Scc2/Scc4* loader complex was prebound to DNA, recently the association of *Scc2* with centromeres, at least, was shown to require cohesin itself (Fernius *et al.* 2013). The complete tripartite cohesin ring, with ATP bound to both SMC heads (engaged state)

together with *Scc3*, is required for cohesin loading (Arumugam *et al.* 2006; Hu *et al.* 2011). Cohesin ring assembly allows its interaction with *Scc2/Scc4*, which in turn enables the entire cohesin–*Scc2/Scc4* complex to associate with centromeres and probably other loading sites too (Fernius *et al.* 2013). This explains why cohesin is not associated with chromosomes in early G1 or anaphase cells. Although *Smc1*, *Smc3*, and *Scc2/Scc4* are all present in early G1 cells, *Scc1* is produced only in late G1 and is cleaved in anaphase (Guacci *et al.* 1997; Michaelis *et al.* 1997; Ciosk *et al.* 2000; Uhlmann *et al.* 2000; Kogut *et al.* 2009). Therefore, *Scc1* production upon cell cycle entry is the trigger for cohesin loading. The second step in cohesin loading is the transition from the state where the cohesin–*Scc2/Scc4* complex has docked at its loading site to one where cohesin is encircling DNA and can translocate along it. This transition is blocked by mutations in the *Smc1* and *Smc3* ATPase heads that block ATP hydrolysis, demonstrating that ATP hydrolysis is important for this step (Hu *et al.* 2011). The notion that cohesin topologically embraces chromosomes predicts that the ring must be opened for its loading onto DNA. ATP hydrolysis is stimulated by *Scc1* binding and this could facilitate cohesin ring opening (Arumugam *et al.* 2003). Interestingly, evidence suggests that the interfaces between the *Smc1* and *Smc3* NBD heads and *Scc1* need not be opened for cohesin’s association with DNA (Gruber *et al.* 2006). Instead, the most likely scenario is that cohesin opens at the interface between the hinge domains of *Smc1* and *Smc3* to allow DNA entry (Gruber *et al.* 2006). Given the separation of the site of ATP hydrolysis with the hinge domain by the long SMC coiled coils, this poses a conundrum: How could ATP hydrolysis influence opening of the hinge? To accommodate this idea, it has been suggested that the coiled-coil domains could fold back on themselves to bring the hinge in proximity of the site of ATP hydrolysis (Gruber *et al.* 2006; Hu *et al.* 2011). Intriguingly, insertional mutations in the loop structure of *Smc1*, situated close to the NBDs, do not affect cohesin’s overall association with chromosomes but prevent cohesin enrichment in the pericentromere and CARs, in common with insertional hinge mutants (Milutinovich *et al.* 2007). This implicates the loop region in cohesin loading. Another possibility is that the NBD-proximal motifs in *Smc1* are important for the interaction with *Scc2/Scc4*, which in turn enable ATP hydrolysis to drive a conformational change that leads to opening of the hinge (Figure 4). The hinge likely plays additional roles in cohesion establishment after loading since the crystal structure of the hinge revealed a positively charged channel in which neutralizing mutations caused loss of cohesion, though they did not affect either dimerization or cohesin loading (Kurze *et al.* 2011). Finally, fluorescence recovery after photobleaching (FRAP) experiments have shown that *Scc2* turns over extremely rapidly at the centromere (Hu *et al.* 2011) and *Scc2/Scc4* does not stably copurify with kinetochores (Fernius *et al.* 2013), suggesting that *Scc2/Scc4* dissociates from the cohesin complex after the loading reaction is complete.



**Figure 4** Cohesin cycle. Current model for cohesin loading and establishment. Cohesin loading involves opening of the Smc1 and Smc3 hinge and requires the Scc2/Scc4 protein and ATP binding to the SMC heads. ATP hydrolysis ensues and chromosome entrapment occurs. Cohesin is unstable on chromosomes due to the activity of Wpl1/Rad61, which opens the Smc3-Scc1 interface. Note that the contact point of Scc2/Scc4 on cohesin shown here is speculative. Eco1-dependent acetylation of Smc3 makes cohesin refractory to Wpl1, effectively locking the rings shut. Cohesin is destroyed during anaphase by cleavage of Scc1 by separase. Hos1 deacetylase recycles Smc1-Smc3 for use in the next cell cycle by removing the acetyl mark on Smc3.

### Replication-coupled establishment of cohesin

**Eco1-directed cohesin establishment:** The Eco1 acetyltransferase (also called Ctf7) is not required for cohesin's association with chromosomes, but is needed during DNA replication for cohesion generation (Skibbens *et al.* 1999; Toth *et al.* 1999; Ivanov *et al.* 2002). A clue as to Eco1 function came from the observation that the lethality caused by deletion of the fission yeast *Schizosaccharomyces pombe* ortholog of ECO1, called *eso1*, is suppressed by loss of Pds5 function (which is nonessential in *S. pombe*) (Tanaka *et al.* 2001). This suggested a negative effect of Pds5 on cohesion establishment that is counteracted by Eco1. This idea was upheld in budding yeast with the identification of missense mutations in SMC3, PDS5, and SCC3 as well as null alleles of WPL1/RAD61, all of which can suppress *eco1* loss-of-function mutations (Ben-Shahar *et al.* 2008; Unal *et al.* 2008; Zhang *et al.* 2008; Sutani *et al.* 2009; Rowland *et al.* 2009). The critical substrate of Eco1 is Smc3, which is acetylated on two residues K112 and K113 in its NBD (Ben-Shahar *et al.* 2008; Unal *et al.* 2008; Zhang *et al.* 2008; Rowland *et al.* 2009). Mutations in Smc3 K113 as well

as nearby residues were among those found to suppress the loss of Eco1 function. This indicated that Smc3 is the only critical substrate of Eco1. The other three proteins in which mutations were found to suppress loss of Eco1 function, Pds5, Scc3, and Wpl1, form a complex that is loosely associated with cohesin. Based on the knowledge that Wpl1 is the ortholog of human WAPL1, which was known to promote cohesin's dissociation from chromosomes during prophase (Gandhi *et al.* 2006; Kueng *et al.* 2006), it was proposed that suppressor mutations in Pds5, Scc3, and Wpl1 abolish an "antiestablishment" activity of these proteins (Rowland *et al.* 2009). This model proposed that Pds5 and Scc3, while essential for cohesin establishment, would also have an additional cohesin-destabilizing effect by allowing access of Wpl1 to cohesin. Conversely, the Smc3 suppressor mutations are thought to make cohesin resistant to this destabilizing activity. This leads to the idea that acetylation of Smc3 by Eco1 counteracts the destabilizing activity of Wpl1 on cohesin, ensuring its maintenance on chromosomes (Figure 4).

Although the discovery of Smc3 as the key Eco1 substrate was a crucial step in understanding its molecular function,

how this contributes to cohesion establishment in S phase was still unexplained. Importantly, it was questionable whether Wpl1 really possessed a cohesin-destabilizing activity in budding yeast. Although loss of WAPL function in mammals leads to an increase in chromosomal cohesin, budding yeast lacking *WPL1* have impaired cohesion and cohesin levels are reduced on chromosomes (Warren *et al.* 2004; Rowland *et al.* 2009; Sutani *et al.* 2009). This contradiction was addressed by measuring the dynamicity of cohesin either using FRAP or its ability to be “anchored” outside the nucleus (Chan *et al.* 2012; Lopez-Serra *et al.* 2013). These studies showed that Wpl1 indeed promotes turnover of cohesin on chromosomes and that this is counteracted by Eco1-dependent acetylation of Smc3. The reduced levels of cohesin on chromosomes of *wpl1Δ* cells seem to be a result of generally decreased cellular levels, though the underlying cause remains unknown (Chan *et al.* 2012). Mutations in *Scc3* and *Pds5* that suppress loss of *Eco1* function also reduce cohesin turnover on chromosomes (Chan *et al.* 2012). Some, but not all, of these mutations affect Wpl1 recruitment to cohesin (Sutani *et al.* 2009; Chan *et al.* 2012). Interestingly, measurements using GFP-tagged proteins suggested that although one molecule of each of *Scc3* and *Pds5* are associated with the cohesin ring, Wpl1 is substoichiometric and highly dynamic (Chan *et al.* 2012). This suggests a catalytic mechanism of cohesin dissociation by Wpl1. Remarkably, fusing *Scc1*'s N terminus to *Smc3* suppresses lethality due to loss of *Eco1* function and causes cohesin to be stable on chromosomes. This indicates that Wpl1 exerts its function by disrupting the interface between *Scc1* and *Smc3*. Notably, this cohesin “exit gate” is distinct from the “entry gate,” the hinge domain, involved in cohesin loading (Gruber *et al.* 2006; Chan *et al.* 2012). Acetylation of *Smc3* by *Eco1* locks the exit gate, thereby making cohesin refractory to the effects of Wpl1.

A key outstanding question was whether Wpl1 acts specifically to prevent the initial chromosome entrapment by cohesin (antiestablishment model) or whether it can also dismantle already established cohesin (“antimaintenance” model). *Eco1* is not required after S phase for cohesion establishment (Skibbens *et al.* 1999; Toth *et al.* 1999) and is thought to travel with the replication fork, being recruited by PCNA (Lengronne *et al.* 2006; Moldovan *et al.* 2006). Moreover, *Smc3* acetylation requires prior loading onto chromosomes and appears in S phase (Ben-Shahar *et al.* 2008; Unal *et al.* 2008; Zhang *et al.* 2008; Rowland *et al.* 2009; Sutani *et al.* 2009). Therefore, ordinarily, *Eco1*-dependent “locking” of cohesin at the exit gate is coupled to DNA replication. Although experiments in human cells had suggested that *Eco1*-dependent cohesin acetylation counteracts a replication fork-slowing activity of Wpl1 (Terret *et al.* 2009), there is no evidence for this in budding yeast (Lopez-Serra *et al.* 2013). Indeed, Wpl1 can cause dissociation of nonacetylated cohesin outside S phase, in G2, consistent with an antimaintenance activity of Wpl1 (Chan *et al.* 2012; Lopez-Serra *et al.* 2013). These experiments argue in

favor of a model whereby competition between *Scc2/Scc4*-dependent cohesin loading and Wpl1-dependent cohesin dissociation creates a state of high cohesin dynamicity on chromosomes. *Eco1*-dependent cohesin acetylation at the replication fork locks cohesin around sister chromatids, rendering it refractory to Wpl1 activity and thereby stably bound to chromosomes.

What is the function of the dynamic nonacetylated pool of cohesin, since it is not participating in cohesion? Analysis of a *wpl1Δ* mutant, in which nonacetylated cohesin loses its dynamicity, revealed that chromosomes were more highly condensed. This suggests that cohesin turnover may be important to modulate the state of chromosome compaction. In contrast, Wpl1-dependent cohesin dynamicity does not contribute in a major way to cohesin's roles in transcription or meiosis (Lopez-Serra *et al.* 2013).

During anaphase, cohesin is deacetylated by the *Hos1* deacetylase (Beckouët *et al.* 2010; Borges *et al.* 2010; Xiong *et al.* 2010). Cohesin release from chromosomes, as a result of its cleavage in anaphase, is essential for cohesin deacetylation, though *Hos1* is present earlier (Beckouët *et al.* 2010; Borges *et al.* 2010). How chromosome-bound acetylated cohesin is “shielded” from *Hos1* activity is unknown but *Scc3* and *Pds5* are likely to be involved in this. Cohesin deacetylation allows *Smc1–Smc3* complexes to be recycled for cohesion establishment in the next S phase. Cohesin that is not deacetylated in anaphase, or a mutant version that mimics this state, loads onto chromosomes in the next cell cycle, but fails to establish cohesion. This indicates that cohesin must be acetylated *de novo* during S phase to lock rings shut.

**Other factors involved in cohesion establishment:** *Eco1*, by counteracting Wpl1, ensures the stability of cohesin on chromosomes, an activity that is likely to be important for the longevity of cohesion. However, in the absence of *Eco1–Wpl1*, budding yeast are viable and establish cohesion, albeit less robustly than that of wild-type cells (Ben-Shahar *et al.* 2008; Unal *et al.* 2008; Zhang *et al.* 2008; Rowland *et al.* 2009; Sutani *et al.* 2009). Furthermore, nonacetylatable fission yeast *Smc3* (called Psm3) does not cause lethality, even in the presence of Wpl1 (Feytout *et al.* 2011). This reveals that cohesion exists without *Eco1–Wpl1*. If the fundamental process by which cohesion is established is independent of *Eco1–Wpl1*, how does this work? A simple explanation is that after loading, cohesin rings encircle a chromatin thread and that the DNA replication machinery passes through this ring to synthesize a sister chromatin thread, which is automatically contained within the ring. An alternative model is that factors associated with the replication machinery facilitate cohesin ring opening upon fork passage, and its reclosure in the wake of the polymerase. Several other factors are known to contribute to cohesion establishment, including *Ctf18*, *Csm3*, *Tof1*, *Mrc1*, *Ctf4*, and *Chl1* but their roles are unknown (Hanna *et al.* 2001; Mayer *et al.* 2001; Mayer *et al.* 2004; Petronczki *et al.* 2004;



Skibbens 2004; Warren *et al.* 2004; Xu *et al.* 2007; Fernius and Marston 2009). Loss of each of these factors reduces cohesin acetylation to variable extents, but it is not known whether they are directly involved in facilitating *Eco1* function or whether they participate in an unrelated establishment process that is a prerequisite for cohesin acetylation (Beckouët *et al.* 2010; Borges *et al.* 2013). *Ctf18*, at least may contribute to cohesin acetylation directly, since it is thought to help load *PCNA*, thereby providing a platform for *Eco1* at the replication fork (Bermudez *et al.* 2003; Bylund and Burgers 2005; Lengronne *et al.* 2006). Other factors also function in replication-associated processes. *Csm3*, *Tof1*, and *Mrc1* form a complex that travels with the replication fork and elicits stalling when barriers are encountered (Katou *et al.* 2003; Calzada *et al.* 2005; Tourriere *et al.* 2005; Bando *et al.* 2009). *Ctf4* is an integral part of the replisome and helps to couple polymerase  $\alpha$ /primase to the Mcm replicative helicase (Gambus *et al.* 2006, 2009). *Chl1* is a likely DNA helicase whose molecular function is unknown (Gerring *et al.* 1990). Uncovering the molecular function of these factors in cohesion establishment will be an important priority for the future.

Recently, SUMOylation of several cohesin subunits has been found to occur in the window between cohesin loading and chromosome entrapment (Almedawar *et al.* 2012). SUMO, like ubiquitin, is a small protein modifier that is attached to lysines of target proteins, which can result in a range of effects including changes in localization, stability, or function (see Cubeñas-Potts and Matunis 2013 for review). A role for SUMOylation in the entrapment process was suggested by the observation that *Scc1* fused to a SUMO-deconjugating enzyme reduced cohesin SUMOylation, which led to cohesion defects and lethality (Almedawar *et al.* 2012). SUMOylation is independent of acetylation and does not appear to counteract *Wpl1*, suggesting that it may function in an *Eco1*-independent pathway of cohesion establishment (Almedawar *et al.* 2012).

### **DNA damage-induced cohesion**

Although cohesion establishment is ordinarily restricted to S phase, it can occur later in the cell cycle under conditions of DNA damage. Yeast cells that fail to establish cohesion in S phase are unable to repair damage caused by  $\gamma$ -irradiation (Sjogren and Nasmyth 2001). However, additional cohesin is also recruited post S phase to an  $\sim$ 100-kb domain surrounding the damage site in a manner dependent on *Scc2/Scc4* and the DNA damage checkpoint, and this is essential for repair (Strom *et al.* 2004; Unal *et al.* 2004). DNA damage not only triggers cohesin loading at the break site, but also, remarkably, genome-wide cohesion establishment in G2/M through *Eco1* (Strom *et al.* 2007; Unal *et al.* 2007). However, the existence of separation-of-function mutations in *ECO1* that abolish damage-induced cohesion, but not S phase-induced cohesion suggested that *Eco1* might work through different mechanisms in these two situations (Unal *et al.* 2007; Zhang *et al.* 2008). Indeed, rather than acetylate

*Smc3* as in S phase-generated cohesion, *Scc1* is the likely target of *Eco1* in G2/M in response to DNA damage (Heidinger-Pauli *et al.* 2008, 2009). The acetylation of *Scc1* is thought to occur in response to its phosphorylation of the checkpoint kinase, *Chk1* and like acetylation of *Smc3*, counteract *Wpl1* activity (Heidinger-Pauli *et al.* 2008, 2009). Normally, *Eco1* is limiting after S phase because *Eco1* overexpression can cause cohesion establishment in G2/M phase (Unal *et al.* 2007). Stepwise phosphorylation of *Eco1* by cyclin-dependent kinase (CDK), *Dbf4*-dependent *Cdc7* kinase (DDK), and the GSK3 kinase, *Mck1*, triggers *Eco1* ubiquitination by the SCF (*Skp1*, Cullin, F box) E3 ligase after S phase, leading to its degradation (Lyons and Morgan 2011; Lyons *et al.* 2013). A failure to degrade *Eco1* increases cohesion at metaphase (Lyons and Morgan 2011). In the case of DNA damage, *Eco1* degradation is prevented because *Dbf4-Cdc7* is inhibited, probably as a result of its phosphorylation by the *Rad53* checkpoint kinase (Lopez-Mosqueda *et al.* 2010; Zegerman and Diffley 2010).

*Scc1* is also sumoylated in response to DNA damage by the SUMO E3 ligases *Siz1*, *Siz2*, and *Nse2/Mms21* (a component of the cohesin-related *Smc5-Smc6* complex) (McAleenan *et al.* 2012). The requirement for *Scc1* SUMOylation in damage-induced cohesion seems to be at the establishment step, similar to its role in S phase cohesion (Almedawar *et al.* 2012; McAleenan *et al.* 2012).

Interestingly, a recent report implied that the cohesion that is induced genome-wide in response to DNA damage may have a different function to that that is built around the break site. The translesion synthesis polymerase, *Pol $\eta$*  (*RAD30* in budding yeast) is required for genome-wide, but not local, damage-induced cohesion, perhaps by facilitating *Scc1* acetylation (Enervald *et al.* 2013). It was observed that genome-wide cohesion generation in G2/M appears to be important not for repair, but for segregation, leading to the proposal that it may be needed to reinforce S phase cohesion (Enervald *et al.* 2013). Consistent with this idea, cohesin, which established cohesion in S phase, has been reported to be removed genome-wide upon DNA damage in G2/M (McAleenan *et al.* 2013). This removal of cohesin seems to be required for the efficient repair of DNA lesions by allowing access of repair proteins. Surprisingly, damage-dependent removal of cohesin in G2/M was reported to depend on cleavage of its *Scc1* subunit by separase (McAleenan *et al.* 2013). This is unexpected because global separase activation must be prevented until all chromosomes have achieved biorientation otherwise mis-segregation will occur, resulting in aneuploidy. The prediction is that separase activity or cohesin cleavage must be locally regulated to spare some cohesion from destruction. Future work will be required to fully illuminate the mechanisms underlying the role of cohesion in DNA-damage repair.

### **Other structural components of chromosomes**

Following the duplication of chromosomes and the establishment of linkages between them, sister chromatids must

be prepared for their segregation during mitosis. Chromosomes are organized into rigid structures by condensation and DNA molecules are resolved from each other, while the linkages between sister chromatids are maintained. The key players in this process are the cohesin-related condensin complex and topoisomerase II (Earnshaw *et al.* 1985; Gasser *et al.* 1986; Hirano and Mitchison 1994; Strunnikov *et al.* 1995). Although dramatic structural changes in chromosome organization cannot be observed directly in the small yeast nucleus, both condensin and topoisomerase II are essential for chromosome segregation, suggesting they perform similar functions in yeast too (Dinardo *et al.* 1984; Holm *et al.* 1985; Strunnikov *et al.* 1995; Bhalla *et al.* 2002).

**The condensin complex:** The condensin complex is related to cohesin but it is much less well understood (Thadani *et al.* 2012). Defective condensin leads to reduced chromosome compaction and a failure of chromosomes to segregate during anaphase (Freeman *et al.* 2000; Bhalla *et al.* 2002; Lavoie *et al.* 2004). However, the molecular function of condensin in chromosome segregation remains unclear.

In mammals, two distinct condensin complexes (condensin I and condensin II) exist. Condensin II mediates the premitosis condensation of chromosomes, whereas condensin I assembles resolved metaphase chromosomes after nuclear envelope breakdown at the start of mitosis (Hirano 2005). Budding yeast possesses only condensin I, composed of two Smc subunits (*Smc2* and *Smc4*), a kleisin (*Brn1*), and two HEAT (huntingtin–elongation factor 3–protein phosphatase 2A–TOR1) repeat-containing subunits (*Ycs4* and *Ycg1/Ycs5*) (Figure 3) (reviewed in Hirano 2012).

Defective condensin causes severe chromosome segregation defects characterized by chromosome bridges during anaphase (Saka *et al.* 1994; Strunnikov *et al.* 1995; Bhat *et al.* 1996; Hudson *et al.* 2003; Ono *et al.* 2004; Oliveira *et al.* 2005; Gerlich *et al.* 2006). A likely activity of condensin that allows it to perform its function is the bringing together of distant DNA sequences of the same molecule. Thus, unlike cohesin, which forms intersister linkages, condensin is thought to build intrasister linkages and stabilize chromatin loops (Cuylen *et al.* 2011; Cuylen and Haering 2011; Thadani *et al.* 2012). Condensin forms a ring-like structure, similar to cohesin, although the hinge dimerization interface may adopt a distinct geometry (Anderson *et al.* 2002) (Figure 3). Furthermore, like cohesin, condensin can topologically embrace a minichromosome (Cuylen *et al.* 2011). Condensin can also bind DNA and introduce positive supercoils (Kimura *et al.* 1999; St-Pierre *et al.* 2009). This has led to the proposal that condensin may act enzymatically, rather than structurally, and drive chromosome compaction through the introduction of positive supercoils (Baxter and Aragón 2012). This idea has allowed an extension of the loop stabilization model that can explain why condensin specifically links regions of the same DNA molecule (Baxter and Aragón 2012).


Condensin is highly enriched at centromeres, pericentromeres, telomeres, and the rDNA (D'Ambrosio *et al.* 2008b).

Condensin has also been reported to associate with genes transcribed by RNA PolII such as tRNAs and 5S rDNA, and the association with tRNAs was found to be partially dependent on the *Scs2/Scs4* cohesin loader complex (D'Ambrosio *et al.* 2008b). However, overall chromosomal condensin levels are not grossly affected by *Scs2* inactivation (Ciosk *et al.* 2000), suggesting that the relationship between *Scs2/Scs4* and condensin may not be direct. The recruitment of condensin to the rDNA depends on the replication fork block protein, *Fob1*, as well as the monopolin proteins, *Lrs4* and *Csm1*, which have roles also at the kinetochore during meiosis (see below) (Johzuka and Horiuchi 2009). Indeed, the fission yeast monopolin proteins, *Mde4* and *Pcs1*, are important for condensin association with centromeric regions (Tada *et al.* 2011); however, this is not the case in budding yeast (Brito *et al.* 2010), and the factors responsible to centromeric/pericentromeric condensin association in budding yeast remain unknown. Notably, while condensin is associated with the rDNA throughout the cell cycle, it begins to colocalize with kinetochores from around the time of S phase but is absent at anaphase onset (Bachelier-Bassi *et al.* 2008).

**Topoisomerase II:** Topoisomerase II catalyzes the ATP-dependent transport of one DNA double helix through another to relieve both negative and positive supercoils (Wang 2002). In mammalian cells, topoisomerase II is required for the individualization of chromosomes prior to mitosis (Giménez-Abián *et al.* 2000) and evidence that budding yeast *Top2* helps condense chromosomes was obtained using a *lacO*–*LacI*–GFP reporter system (Vas *et al.* 2007). The activity of *Top2* is required prior to mitosis to remove catenates generated as a result of two converging replication forks colliding (Holm *et al.* 1985). A failure to remove these catenates is manifest during anaphase where chromosome bridges are observed (Holm *et al.* 1985). Interestingly, proper cohesion at centromeres depends on SUMOylation and *Top2* appears to be an important target (Bachant *et al.* 2002). This implies that modulating chromosome topology through *Top2* is also important for proper cohesion.

## Establishment of Biorientation

Having prepared a pair of duplicated chromosomes, the next step in segregation is their attachment to the mitotic spindle. In budding yeast, each kinetochore has a binding site for a single microtubule (Winey *et al.* 1995). This means that the erroneous situation where a kinetochore attaches to microtubules from the same pole (merotelic attachment) is impossible in budding yeast. However, it is still possible that sister kinetochores attach to microtubules from the same pole (syntelic attachment) and this must be avoided. The equal segregation of sister chromatids to daughter cells will occur only when sister kinetochores are attached to microtubules from opposite poles (amphitelic attachment or biorientation) (Figure 5).



**Figure 5** Establishment of biorientation. The possible modes of kinetochore attachment en route to biorientation in mitosis are shown schematically. For details see text.

The steps leading to kinetochore capture by microtubules have been reviewed recently and will be summarized only briefly here (Tanaka 2010). In budding yeast, kinetochores remain bound to microtubules throughout the cell cycle, detaching transiently only for a short window as DNA is replicated and kinetochores reassemble (Winey and O’Toole 2001; Kitamura *et al.* 2007). The majority of initial kinetochore–microtubule interactions are syntelic, and *Ipl1* is required early in mitosis to sever these connections and provide an opportunity for amphitelic attachment to be established (He *et al.* 2000; Biggins and Murray 2001; Tanaka *et al.* 2002). Unattached kinetochores are captured as follows (Tanaka 2010). First, kinetochores attach to the lateral side of microtubules, either directly or via a microtubule nucleated from the kinetochore, a process that involves the XMAP215/ch-TOG protein, *Stu2* (Kitamura *et al.* 2010; Gandhi *et al.* 2011). Second, the captured kinetochore is transported along the microtubule toward the spindle pole body (SPB) by the kinesin-14 protein, *Kar3* (Tanaka *et al.* 2005, 2007). The lateral attachment of the kinetochore is then converted into an end-on attachment to a SPB-derived microtubule, which requires a conserved loop on the *Ndc80* kinetochore protein (Hsu and Toda 2011; Maure *et al.* 2011; Zhang *et al.* 2012). Upon end-on attachment, loading of the *Dam1* complex that can couple the kinetochore to a shrinking microtubule occurs (Westermann *et al.* 2006; Tanaka *et al.* 2007). Third, capture of the sister kinetochore occurs. The mechanisms that ensure that sister kinetochores are captured by microtubules from opposite poles to achieve biorientation are summarized below.

#### **Role of kinetochore geometry and centromere structure**

Do sister kinetochores adopt a particular geometry that helps facilitate their attachment to opposite poles? There is

no doubt that the pericentromeric chromatin surrounding the kinetochore has specialized properties that could facilitate biorientation through establishment of a preferred geometry for capture by microtubules. The first indication of a specialized pericentromeric structure was the observation using GFP-labeled centromeres, that microtubule tension at metaphase was sufficient to pull sister centromeres apart prior to separase activity and cohesin cleavage (Goshima and Yanagida 2000; He *et al.* 2000; Tanaka *et al.* 2000). The domain of separation extends for ~10 kb on either side of the centromere and sister chromosomal arm sequences remain associated at metaphase. This observation has posed the question: What happens to cohesin during the tension-dependent separation of the pericentromere? One possibility is that cohesin is removed in this region. Indeed, chromatin immunoprecipitation (ChIP) experiments have suggested that pericentromeric cohesin levels are reduced when sister centromeres are under tension compared to those not under tension (Eckert *et al.* 2007; Ocampo-Hafalla *et al.* 2007). However, a pericentromeric barrel of cohesin is clearly visible by microscopy at metaphase, indicating that a substantial amount of cohesin remains associated with the pericentromere (Yeh *et al.* 2008; Hu *et al.* 2011). This poses a paradox: in this situation, where sister centromeres separate over distances of 2–4  $\mu\text{m}$ , it seems impossible that they are trapped within the same cohesin ring. One proposal that could reconcile these observations is that pericentromeric cohesin forms intramolecular, rather than intermolecular, linkages. This would enable the pericentromere to adopt a cruciform structure with sister kinetochores protruding in opposite directions (Yeh *et al.* 2008). Cohesin and condensin together with pericentromeric chromatin constitute a “mitotic chromosome

spring” that balances spindle forces and allows the generation of tension (Stephens *et al.* 2011, 2013). The spring-like properties of the pericentromere are thought to be key to the detection of tension upon biorientation. A further attraction of the cruciform model is that it could be envisaged to facilitate a “back to back” geometry of sister kinetochores, thereby enabling their efficient capture from microtubules from opposite poles.

But is kinetochore geometry actually important for biorientation? The observation that sister kinetochores are inherently biased to biorient on the mitotic spindle argues for this possibility (Indjeian and Murray 2007). Defective kinetochore geometry could also explain why cells lacking cohesin enrichment within the pericentromere are slow to achieve biorientation and rely on the error correction machinery (Ng *et al.* 2009). However, it is equally possible that enriched pericentromere cohesin promotes biorientation by strengthening intersister cohesion to facilitate the generation of tension.

Although it is likely that kinetochore geometry facilitates biorientation, it is clear that it cannot be the only important factor and tension-sensing-based mechanisms exist. Indeed, tension is sufficient to allow biorientation in the absence of a back-to-back sister kinetochore configuration because dicentric chromosomes with physically separated kinetochores achieve biorientation (Dewar *et al.* 2004). Furthermore, reconstituted kinetochore–microtubule attachments persist longer under force, indicating that tension directly mechanically stabilizes them (Akiyoshi *et al.* 2010). In practice, sister kinetochore geometry is likely to increase the probability that initial attachments are made to opposite poles.

### Error correction

While tension stabilizes kinetochore–microtubule attachments, conversely, a lack of tension leads to the destabilization of kinetochore–microtubule attachments, providing a further opportunity for the correct attachments to be made. Central to this “error correction” process is the Aurora B kinase (Biggins *et al.* 1999; Tanaka *et al.* 2002). Aurora B is the kinase constituent of the chromosomal passenger complex (CPC) that also contains INCENP (*Sli15*), Survivin (*Bir1*), and Borealin (*Nbl1*) (reviewed in Carmena *et al.* 2012). *Ipl1* phosphorylates a number of substrates in the outer kinetochore that are thought to prevent interactions with microtubules (Cheeseman *et al.* 2002; Akiyoshi *et al.* 2009a; Demirel *et al.* 2012). Since *Ipl1* is associated with the inner kinetochore, it has been proposed that tension physically separates *Ipl1* from its substrates, thereby allowing their dephosphorylation and silencing of the error correction machinery (Tanaka *et al.* 2002; Keating *et al.* 2009). Although not directly tested in budding yeast, support for this model has been obtained from work in mammalian cells (Liu *et al.* 2009; Welburn *et al.* 2010). In contradiction of this model, Campbell and Desai (2013) recently described a truncated form of budding yeast *Sli15*, which does not

accumulate at centromeres but rather associates with microtubules and chromatin, yet is proficient for tension-sensing and chromosome biorientation. Presumably, though not properly regulated, sufficient CPC accumulates at centromeres to disrupt incorrect attachments. This demonstrates that tight regulation of CPC localization at centromeres may not be essential under normal circumstances (Campbell and Desai 2013)

The Shugoshin (*Sgo1*) protein, which is associated with the budding yeast pericentromere (in the same region as the enriched cohesin) (Kiburz *et al.* 2005), also contributes to biorientation (Indjeian *et al.* 2005; Indjeian and Murray 2007). *Sgo1* is recruited to the pericentromere through phosphorylation of H2A by *Bub1* kinase (Kawashima *et al.* 2010). This explains a requirement for the *Bub1* kinase domain and residue S121 on H2A as well as several residues on H3 in biorientation (Fernius and Hardwick 2007; Kawashima *et al.* 2010). In fission yeast the Shugoshin paralog, *Sgo2*, similarly promotes biorientation during mitosis, where its role appears to be recruitment of Aurora B (called *Ark1* in *S. pombe*) to centromeric regions (Kawashima *et al.* 2007; Vanoosthuysen *et al.* 2007). Although CPC subunits colocalize with kinetochores in *sgo1Δ* cells in budding yeast (Kiburz *et al.* 2008; Storchová *et al.* 2011), it seems likely that *Sgo1* affects biorientation through *Ipl1* too. Indeed, the ability of truncated *Sli15*, which clusters on microtubules, to rescue the biorientation defects of *sgo1Δ* cells, is consistent with *Sgo1* promoting biorientation through the CPC (Campbell and Desai 2013). One possible scenario is that there are multiple ways by which *Ipl1* can be recruited to centromeres and that *Sgo1* only promotes *Ipl1* association under certain conditions, for example in response to a lack of tension. Indeed, *Ipl1* is essential, presumably due to a need to sever the attachment of kinetochores to SPBs (Tanaka *et al.* 2002), whereas *Sgo1* is not. Consistently, *Bir1* CPC subunit is also recruited to the kinetochore through a direct interaction with the kinetochore protein, *Ndc10* (Yoon and Carbon 1999; Cho and Harrison 2012).

The *Mps1* kinase is also essential for biorientation and triggers checkpoint arrest both in response to unattached kinetochores and a lack of tension (Maure *et al.* 2007; Liu and Winey 2012). This can be explained, at least in part, by a requirement for *Mps1* for *Bub1* association with the kinetochore, allowing, in turn, canonical checkpoint activation and presumably *Sgo1* and *Ipl1* recruitment to centromeres (Fernius and Hardwick 2007; Storchová *et al.* 2011; London *et al.* 2012). However, *Mps1* is likely to play additional, possibly more direct, functions in biorientation and, consistently, *Mps1* substrates in the outer kinetochore have been identified (Shimogawa *et al.* 2006; Kemmler *et al.* 2009).

The destabilization of kinetochore–microtubule attachments that are not under tension not only provides an opportunity for reorienting these attachments but also serves to arrest the cell cycle until all errors are corrected. Neither *Sgo1* nor *Ipl1* are required to arrest the cell cycle in the presence of unattached kinetochores in budding yeast, though both proteins are required for a cell cycle delay in

response to a lack of tension (achieved experimentally by preventing replication or cohesion establishment) (Biggins and Murray 2001; Indjeian *et al.* 2005). This led to the proposal that *Sgo1* and *Ipl1* indirectly elicit a checkpoint response in response to tension defects by generating unattached kinetochores (Pinsky *et al.* 2006). *Ipl1* may additionally potentiate the checkpoint signal in response to unattached kinetochores. Indeed, mutation of sites in *Mad3* that are phosphorylated by *Ipl1* *in vitro* abrogates the checkpoint response to lack of tension, but not to unattached kinetochores (King *et al.* 2007a).

### **Destruction of Sister Chromatid Cohesion and Anaphase Onset**

The cohesion built in S phase holds sister chromatids together until their separation at anaphase onset. Crucially, cohesion provides the resistance to spindle microtubule forces, enabling sister chromatids to attach to opposite poles at metaphase. It is essential that cohesion is not destroyed until all chromosomes are properly attached to microtubules. A surveillance mechanism, known as the spindle checkpoint, senses improperly attached chromosomes and prevents separase activation to ensure that it is the case. Once biorientation is achieved, the spindle checkpoint is satisfied, separase is activated, and anaphase proceeds.

#### **Cleavage of cohesin by separase**


Cohesin destruction requires the activity of an E3 ubiquitin ligase known as the anaphase promoting complex, or cyclosome (APC/C) (Irniger *et al.* 1995). The APC/C is a huge molecular machine that attaches ubiquitin, a small protein modifier, to lysine residues of target proteins (see Peters 2006 for review). Polyubiquitinated substrates are recognized by the 26S proteasome, which mediates their destruction. The critical substrate of the APC/C at anaphase onset is not cohesin, but an anaphase inhibitor, known as securin or *Pds1* (Cohen-Fix *et al.* 1996; Funabiki *et al.* 1996; Ciosk *et al.* 1998). The APC/C also targets mitotic cyclin for degradation at the metaphase–anaphase transition but this is not required for chromosome separation in budding yeast (Surana *et al.* 1993). Securin binds to and inhibits separase (*Esp1*), a cysteine protease that is required for sister chromatid separation (Ciosk *et al.* 1998). It is cleavage of the *Scc1* subunit of cohesin by separase that is the trigger for sister chromatid separation. Mutation of the separase recognition sites in *Scc1* prevents sister chromatid segregation, though securin is still destroyed (Uhlmann *et al.* 1999). Moreover, artificial production of the tobacco etch virus (TEV) protease in cells where the only copy of *Scc1* has TEV-cleavage sites triggers chromatid separation (Uhlmann *et al.* 2000). Therefore, *Scc1* cleavage is both necessary and sufficient for chromosome segregation. Given that cleavage of *Scc1* will result in opening of the ring, it is easy to envisage why this causes the release of cohesin from chromosomes (Gruber *et al.* 2003).

### **The spindle checkpoint**

**Inhibition of APC–Cdc20:** Broadly speaking, there are two elements to the surveillance mechanisms that ensure an accurate anaphase. First, cell cycle progression must be halted until all the proper attachments have been generated. This is the role of a checkpoint, known as the “spindle assembly checkpoint.” Second, erroneous attachments, that is, where both sister kinetochores have attached to microtubules from the same pole (syntelic attachment), must be prevented or corrected. Syntelic attachments fail to generate tension and are destabilized by the error-correction machinery, providing a further opportunity for sister kinetochores to attach to microtubules from opposite poles (ampitelic attachment or biorientation).

The spindle checkpoint targets the APC to prevent anaphase onset in the presence of unattached kinetochores by stabilizing securin, thereby maintaining separase inhibition (reviewed in Lara-Gonzalez *et al.* 2012) (Figure 6). For the APC to be active, it must associate with a so-called “coactivator” that is thought to present specific substrates to the APC for ubiquitylation. In vegetatively growing budding yeast, there are two possible coactivators, *Cdc20* and *Cdh1*. APC–*Cdc20* is responsible for triggering securin degradation and, consequently, *Cdc20* is essential for anaphase onset. *Cdc20* is also the crucial target of the spindle checkpoint (Hwang *et al.* 1998). *Cdh1* is not required for chromosome segregation, but is activated later in the cell cycle where it promotes mitotic exit by targeting cyclins for degradation (Visintin *et al.* 1997). Degron motifs known as D (destruction) boxes and KEN boxes on substrates are bound by recognition sites for these motifs on *Cdc20* or *Cdh1* (Peters 2006).

Genetic screens in budding yeast identified the components of the spindle checkpoint that are conserved throughout eukaryotes. The isolation of mutants that failed to arrest when microtubules were disrupted by drugs led to the identification of the “budding uninhibited by benzimidazole” (*BUB*) and “mitotic arrest deficient” (*MAD*) genes (Hoyt *et al.* 1991; Li and Murray 1991). Together with the *Mps1* kinase (Winey *et al.* 1995; Weiss and Winey 1996) *Mad1*, *Mad2*, *Mad3* (*BubR1*), *Bub1*, and *Bub3* form the core spindle checkpoint components that inhibit the APC–*Cdc20* in response to the presence of unattached kinetochores. The spindle checkpoint may also detect kinetochores that are attached to microtubules, but which lack intersister tension, though this is controversial. Nevertheless, it is clear that these components work together to generate a signal at the kinetochore that culminates in the inhibition of the APC; however, the mechanism is not completely understood. Briefly, and taking into account a great deal of work in other organisms (see Musacchio and Salmon 2007; Lara-Gonzalez *et al.* 2012 for more detailed reviews), the following general principles of spindle checkpoint function have emerged. The Aurora B and *Mps1* kinases are the most upstream kinetochore components. In many organisms, including fission



**Figure 6** The spindle checkpoint. (A) Generation of the mitotic checkpoint complex (MCC) by an unattached kinetochore. (B) Modes of APC-Cdc20 inhibition by the spindle checkpoint. Pseudosubstrate inhibition, Cdc20 displacement, and Cdc20 sequestration are all thought to contribute to APC-Cdc20 inhibition.

yeast, Aurora B kinase enables Mps1 kinase association with kinetochores, which in turn enables recruitment of other checkpoint components (summarized in Heinrich *et al.* 2012). However, in budding yeast, Aurora B (Ipl1) and Mps1 appear to be recruited to kinetochores independently (Maure *et al.* 2007). Mps1 phosphorylates the kinetochore protein Spc105/KNL1/Blinkin on conserved MELT motifs to enable recruitment of a complex of Bub1 and Bub3 (London *et al.* 2012; Shepperd *et al.* 2012; Yamagishi *et al.* 2012). Bub1 is also a conserved kinase, though its kinase activity is not essential for checkpoint function and is rather required for biorientation through recruitment of Sgo1 (Warren *et al.* 2002; Peters 2006; Fernius and Hardwick 2007). The Bub1-Bub3 complex is present at kinetochores from S phase until metaphase of mitosis (Kerscher *et al.* 2003; Gillett *et al.* 2004). In contrast, Mad1 and Mad2 are visualized on kinetochores only under conditions where they are not expected to be attached to microtubules, whereas Mad3 has not been detected at kinetochores (Gillett *et al.* 2004). Importantly, Bub1 and Bub3 are required for the kinetochore association

of Mad1 and Mad2 upon checkpoint activation (Gillett *et al.* 2004). How Bub1-Bub3 influences Mad1-Mad2 remains a mystery, although a Mad1-Bub1-Bub3 complex has been observed upon checkpoint activation and this appears to be functionally important in triggering cell cycle arrest (Brady and Hardwick 2000). Ipl1 is also required for Mad2 association with kinetochores during checkpoint activation (Gillett *et al.* 2004), perhaps due to a requirement for Ipl1 in generating unattached kinetochores to which Mad1-Mad2 can be recruited (Pinsky *et al.* 2006). Though the kinetochore receptor for Mad1 is not yet known, kinetochore-bound Mad1 plays a key role in generating the “wait anaphase” signal at kinetochores through recruitment of Mad2 from the soluble pool (Chen *et al.* 1998, 1999). These studies led to the Mad2 “template” model, which provides an explanation for the role of the Mad1-Mad2 interaction in generating the “mitotic checkpoint complex” (MCC) (De Antoni *et al.* 2005), a potent APC inhibitor, composed of Mad2, Mad3, Bub3, and Cdc20 (Hardwick *et al.* 2000; Brady and Hardwick 2000) (Figure 6A). Binding to Mad1 converts Mad2 from an “open” (O-Mad2) to a “closed” (C-Mad2) conformation. C-Mad2 that is already bound to Mad1 dimerizes with soluble O-Mad2, generating further C-Mad2. Since Cdc20 binds C-Mad2 in a similar way to Mad1, the Mad1 template catalyses Mad2 binding to Cdc20. Mad3 binds to the same interface of C-Mad2 as O-Mad2 bound to Mad1 (Chao *et al.* 2012; Mariani *et al.* 2012) and Mad3 binds to the same surface of Bub3 as Bub1 (Larsen *et al.* 2007). Exactly how these interactions lead to Mad3 and Bub3 incorporation into the MCC is not well understood.

What is the role of the MCC in APC inhibition? This question has been difficult to answer, not least because the exact identity of the downstream effector that inhibits the APC is uncertain. The picture that has emerged is that Bub3-Mad3 and Mad2 synergistically inhibit the APC, though the relative contribution of Mad3 and Mad2 is controversial (Fang *et al.* 1998; Tang *et al.* 2001; Fang 2002; Davenport *et al.* 2006; Burton and Solomon 2007; Kulukian *et al.* 2009; Foster and Morgan 2012; Lara-Gonzalez *et al.* 2012; Lau and Murray 2012). One potential mode of APC inhibition by the MCC, for which evidence is accumulating, is the “pseudosubstrate model.” The possibility that Mad3 could act as a pseudosubstrate, blocking access of APC-Cdc20 to securin and cyclin was suggested following the realization that Mad3 has two KEN boxes that are important for APC inhibition (Burton and Solomon 2007; King *et al.* 2007b; Sczaniecka *et al.* 2008; Malureanu *et al.* 2009). Indeed, in the MCC crystal structure, a Mad3 KEN box is optimally positioned by Mad2 to obscure the recognition sites for the KEN box degron on Cdc20, providing support for the pseudosubstrate model (Chao *et al.* 2012). Additional mechanisms of APC inhibition are also likely. For instance, in a model where the MCC crystal structure was mapped onto an existing EM density map of the APC-MCC complex, Cdc20 was displaced away from contacts on the APC required to constitute its D box receptor (Chao *et al.* 2012).

Furthermore, in budding yeast, *Mad2* and *Cdc20* form a separate complex independent of MCC, suggesting that *Cdc20* sequestration by *Mad2* may also play a role in APC inhibition (Poddar *et al.* 2005). This idea is supported by the finding that *Mad2* prevents *Cdc20* binding to the APC in an *in vitro* assay using purified budding yeast components (Foster and Morgan 2012). In addition, tethering *Mad2* to *Cdc20* is sufficient to inhibit the APC in budding yeast, though the basis of the inhibition is not known (Lau and Murray 2012). Human *Mad2* also interacts with *Cdc20* through the same site it normally uses to bind the APC (Izawa and Pines 2012). Therefore, there is also substantial evidence that in the absence of *BubR1/Mad3*, the *Mad2-Cdc20* complex fails to bind to, and activate, the APC. In summary, pseudosubstrate inhibition by *Mad3* and disruption of key interfaces between *Cdc20* and the APC by *Mad2* are both likely to contribute to the inhibition of the APC by the spindle checkpoint (Figure 6B). Whether APC inhibition occurs solely in the context of the MCC, or if indeed different subcomplexes of checkpoint proteins elicit inhibition through different mechanisms, is a question that should be addressable using recently developed *in vitro* APC assays and structural analysis.

**Checkpoint silencing:** Once all sister kinetochores have bioriented, the inhibitory signals that prevent cohesin cleavage must be silenced to allow anaphase to proceed. Of paramount importance is the silencing of the spindle checkpoint to allow APC activation. Broadly, there are two types of reversals that must occur. First, the phosphorylations that are put in place by the checkpoint and error correction machinery must be removed. The protein phosphatase 1 (PP1) plays a central role in checkpoint silencing (Akiyoshi *et al.* 2009b; Pinsky *et al.* 2009). One role of PP1 is to reverse the *Mps1*-dependent phosphorylation of *Spc105* to release *Bub1* from the kinetochore (London *et al.* 2012), but there are likely to be many more important substrates and possibly other phosphatases that will be important too. Second, the MCC must be disassembled and recent data suggest that *Cdc20* autoubiquitination in the context of the MCC plays a role in this process, allowing for rapid activation of the APC once the checkpoint is satisfied (Mansfeld *et al.* 2011; Chao *et al.* 2012; Foster and Morgan 2012; Uzunova *et al.* 2012). The *Mnd2/Apc15* subunit of the APC is important for *Cdc20* autoubiquitination, though not for securin or cyclin B ubiquitination (Mansfeld *et al.* 2011; Foster and Morgan 2012; Uzunova *et al.* 2012). Interestingly, free *Cdc20* can also be ubiquitinated in an *Mnd2/Apc15*-independent manner (Foster and Morgan 2012; Uzunova *et al.* 2012). This leads to a model whereby ubiquitylation of free *Cdc20* restricts its cellular levels, enabling a checkpoint response to be mounted, whereas ubiquitylation of *Cdc20* in the context of the MCC serves to disable the checkpoint response (Musacchio and Ciliberto 2012). An implication of this model is that for a sustained checkpoint response, MCC must be constantly produced to counterbalance the effect of *Cdc20* autoubiquitination. Interestingly,

the most upstream checkpoint component, *Mps1*, is ubiquitinated by the APC-*Cdc20* at anaphase onset, leading to its degradation (Palframan *et al.* 2006). This helps explain why no more MCC is produced once the checkpoint is satisfied, allowing rapid APC activation and anaphase onset once the last appropriate kinetochore-microtubule interaction is made (Musacchio and Ciliberto 2012).

It is also essential that the sudden loss of tension between sister kinetochores at anaphase does not reengage the error correction machinery or activate the spindle checkpoint. The *Cdc14* phosphatase, which becomes active during anaphase (see below), dephosphorylates the CPC component *Sli15* (INCENP), causing its dissociation from centromeres and relocation at the spindle midzone, and this is important to prevent reengagement of the spindle checkpoint after anaphase onset (Mirchenko and Uhlmann 2010).

### **Other factors regulating anaphase onset**

In addition to being targeted by the spindle checkpoint, in budding yeast, securin/*Pds1* is also targeted by the DNA damage response machinery to prevent anaphase onset (Wang *et al.* 2001). The DNA damage checkpoint kinase, *Chk1*, phosphorylates *Pds1*, making it resistant to APC-*Cdc20*-dependent destruction (Sanchez *et al.* 1999; Wang *et al.* 2001). Therefore the DNA damage response and the spindle checkpoint both prevent anaphase onset by preventing *Pds1* degradation, though the mechanism by which this is achieved is distinct.

Despite the convergence of regulatory networks on *Pds1*, cells lacking *PDS1* are viable and initiate anaphase onset with normal timing (Alexandru *et al.* 2001). At first impression, this is surprising, since separase would be expected to be hyperactive in *pds1Δ* cells, resulting in precocious loss of cohesion. However, this paradox can be explained because securin also plays a positive role in separase activation. Indeed, mice lacking securin are viable but only when separase activity is not also compromised (Wirth *et al.* 2006). In budding yeast, securin not only inhibits separase but also promotes its accumulation within the nucleus, and facilitates its rapid activation upon securin destruction (Agarwal and Cohen-Fix 2002; Hornig *et al.* 2002). Securin can therefore be thought of as an inhibitory chaperone for separase.

What controls the timing of anaphase onset in cells lacking securin? Since cohesin cleavage is sufficient to trigger chromosome segregation, anaphase onset could be additionally regulated by securin-independent separase inhibition or by making cohesin more resistant to cleavage. In the absence of securin, a particular form of the protein phosphatase 2A, containing its *Cdc55* regulatory subunit, becomes essential (Tang and Wang 2006; Chirolì *et al.* 2007; Clift *et al.* 2009). Conditional mutants lacking both *Cdc55* and securin/*Pds1* cleave cohesin prematurely, implicating *Cdc55* as a regulator of cohesin cleavage (Clift *et al.* 2009). Timely cohesin cleavage depends on phosphorylation of its *Scc1* subunit within its separase-dependent cleavage sites by Polo kinase (*Cdc5*), so that regulation of cohesin

phosphorylation is one additional way to regulate anaphase onset (Alexandru *et al.* 2001). Recently, an elegant method to spatially measure separase activation at the single cell level has been developed. Morgan and colleagues engineered a fragment of the *Scc1* cohesin subunit containing a separase cleavage site between LacI and GFP tethered to lacOs at a specific location on the chromosome (Yaakov *et al.* 2012). Cleavage of this “separase biosensor” leads to release of GFP from the tether and loss of specific fluorescence at this site. Additionally tethering *Cdc55* to the biosensor delayed its cleavage in a similar way to blocking the Polo-dependent phosphorylation sites. This suggests that PP2A-*Cdc55* prevents cohesin cleavage by counteracting the Polo-dependent phosphorylation of *Scc1* (Yaakov *et al.* 2012). In addition to this regulation at the level of cohesin it is possible that *Cdc55*, or indeed other factors, prevents anaphase onset by regulating separase. In mammals, separase is additionally inhibited by the phosphorylation-dependent binding of *Cdk1* (Stemmann *et al.* 2001; Gorr *et al.* 2006), though this is not known to occur in budding yeast, where downregulation of CDK activity is not required for anaphase onset (Surana *et al.* 1993). The separase biosensor will be invaluable in uncovering further mechanisms regulating cohesin cleavage.

## Anaphase Progression

Once cohesin destruction is initiated, anaphase ensues and chromosomes begin to move apart. For successful cell division, chromosomes must be completely partitioned before cytokinesis occurs. Once initiated, several mechanisms govern safe passage through anaphase with the result that sister chromatids are fully partitioned to opposite poles of the cell in preparation for exit from mitosis.


### Separase initiates mitotic exit

Cell cycle transitions are governed by fluctuations in CDK activity (reviewed in Morgan 2007). In budding yeast, there is a single CDK, *Cdc28*, which in turn associates with G1 cyclins (*Cln1*, *Cln2*, and *Cln3*) and S phase (*Clb5* and *Clb6*) and M phase (*Clb1*, *Clb2*, *Clb3*, and *Clb4*) B type cyclins. Following cell cycle entry, S phase Clb-CDKs promote DNA replication and subsequently M phase Clb-CDKs trigger entry into mitosis. At the end of mitosis, Clb-CDKs are inactivated, to restore the G1 state. In budding yeast, Clb-CDK inactivation takes place in two waves. Concomitant with securin degradation at the metaphase-anaphase transition, the APC-*Cdc20* targets a pool of Clbs for destruction; however, this is not essential for anaphase onset and Clb-CDK activity persists until the end of anaphase (Surana *et al.* 1993). The essential *Cdc14* phosphatase triggers Clb-CDKs inactivation at the end of mitosis, both by triggering cyclin degradation and allowing the accumulation of the Clb-CDK inhibitor, *Sic1* (Jaspersen *et al.* 1998; Visintin *et al.* 1998; Zachariae *et al.* 1998). First, *Cdc14* dephosphorylates the APC/C activator, *Cdh1*, which in turn enables the

degradation of Clb cyclins. Second, *Cdc14* increases the stability and levels of *Sic1* by dephosphorylating *Sic1* and the transcription factor, *Swi5*. *Cdc14* is controlled by its localization. For most of the cell cycle, *Cdc14* is sequestered in the nucleolus through binding to its inhibitor, *Cfi1/Net1* (Shou *et al.* 1999; Visintin *et al.* 1999). During anaphase, *Cdc14* is released from the nucleolus into the nucleus and cytoplasm where it can dephosphorylate its substrates and trigger exit from mitosis. Two regulatory networks control *Cdc14* localization and activity (reviewed in Weiss 2012). At the end of mitosis, a Ras-like GTPase cascade known as the mitotic exit network (MEN) becomes active and triggers the full activation of *Cdc14* and exit from mitosis (reviewed in Stegmeier and Amon 2004). Migration of the SPB into the bud plays a critical role in MEN activation, enabling spindle orientation to be monitored to ensure equal nuclear division. The MEN is essential and in its absence, cells arrest with high CDK activity and fail to break down their spindles (reviewed in Caydasi and Pereira 2012). In early anaphase, *Cdc14* is under control of the nonessential *Cdc14* early anaphase release (FEAR) network. *Cdc14* released via the FEAR network cannot by itself trigger CDK inactivation and exit from mitosis, but rather governs safe passage through anaphase.

The FEAR network is composed of a group of proteins that includes separase (*Esp1*), the separase-associated protein, *Slk19*, Polo kinase (*Cdc5*), *Spo12*, the nucleolar replication fork block protein, *Fob1*, the protein phosphatase 2A bound to its *Cdc55* regulatory subunit, the PP2A regulators *Zds1* and *Zds2*, and Clb-CDKs (reviewed in Rock and Amon 2009). Although it remains unclear precisely how these proteins work together to regulate *Cdc14* in early anaphase, the overall picture that has emerged is as follows (Figure 7). Ultimately, *Cdc14* activation occurs because phosphorylation destabilizes the *Cdc14-Cfi1/Net1* interaction, leading to *Cdc14* release (Azzam *et al.* 2004). *Clb1-CDK*, *Clb2-CDK*, *Clb5-CDK*, and Polo kinase (*Cdc5*) all contribute to destabilization of the *Cdc14-Cfi1/Net1* interaction (Azzam *et al.* 2004; Manzoni *et al.* 2010). PP2A-*Cdc55* prevents *Cfi1/Net1* phosphorylation until anaphase, maintaining its tight association with *Cdc14* for the rest of the cell cycle (Queralt *et al.* 2006). *Fob1* also contributes to stabilization of the *Cdc14-Cfi1/Net1* interaction (Stegmeier *et al.* 2004). Activation of separase at anaphase onset, together with *Slk19*, downregulates PP2A-*Cdc55* through its *Zds1* and *Zds2* inhibitors (Queralt *et al.* 2006; Queralt and Uhlmann 2008; Rossio and Yoshida 2011; Calabria *et al.* 2012). This allows *Clb1-CDK* and *Clb2-CDK* to phosphorylate *Cfi1-Net1*, disrupting its interaction with *Cdc14* (Queralt *et al.* 2006; Queralt and Uhlmann 2008). Interestingly, the protease activity of separase is not required for its function in the FEAR network (Sullivan and Uhlmann 2003). *Spo12* is also phosphorylated by CDKs during anaphase and this is thought to somehow impair the ability of *Fob1* to stabilize the *Cdc14-Cfi1/Net1* interaction (Tomson *et al.* 2009). While the FEAR network initiates *Cdc14* release, sustained





**Figure 7** FEAR network and anaphase. Regulation and role of the FEAR network in completing chromosome segregation during early anaphase. For details see text.

release and export into the cytoplasm requires the activity of the MEN (Mohl *et al.* 2009).

Separase activation, therefore, not only triggers chromosome separation but also initiates mitotic exit through its role in the FEAR network. This provides a mechanism by which the onset of chromosome segregation is coordinated with mitotic exit. Importantly, assaults that result in securin stabilization, such as DNA damage or improperly attached kinetochores, will prevent both chromosome segregation and mitotic exit through separase inhibition.

Although Cdc14 released via the FEAR network is insufficient for mitotic exit, it plays numerous roles in ensuring safe passage through anaphase. These include: sharpening the anaphase switch, inactivation of mitotic surveillance mechanisms, segregation of the rDNA, nuclear positioning, spindle stabilization, spindle midzone assembly, and MEN activation. Consistently, Cdc14 interacts with numerous potential substrates in mitosis, although the functional importance of the majority of these has not been investigated (Bloom *et al.* 2011). These substrates must be dephosphorylated with carefully regulated timing to ensure ordered progression through anaphase. This order is due to the fact that substrates are predisposed to be dephosphorylated at a specific threshold of kinase/phosphatase ratio as this decreases during anaphase (Bouchoux and Uhlmann 2011). Perhaps this phenomenon could explain why Cdc14 is required for spindle stability in anaphase but, paradoxically, triggers spindle disassembly moments later during exit from mitosis.

**Committing to anaphase:** Once the decision to destroy cohesin has been made, the linkages between chromosomes

are destroyed rapidly. Cdc14 facilitates rapid cohesin loss by removing stabilizing CDK-dependent phosphorylations on securin, accelerating its proteolysis (Holt *et al.* 2008).

**Nuclear position:** Due to the asymmetric nature of budding yeast cell division, correct nuclear position is particularly important for accurate segregation. One set of chromosomes must be partitioned into the bud, while the other set remains in the mother cell. Cdc14 released through the FEAR network affects nuclear position in anaphase by modulating the forces that cytoplasmic microtubules exert on the cell cortex. In the absence of FEAR-dependent Cdc14 activity, the entire nucleus migrates aberrantly into the bud, suggesting that forces at the mother cell cortex are weaker (Ross and Cohen-Fix 2004). How Cdc14 alters cortical spindle forces is, however, not known.

**Completion of chromosome segregation:** Chromosome segregation must also be driven to completion upon anaphase onset. Not all regions of the genome segregate simultaneously. Centromeres are the first to segregate, followed by chromosome arms, telomeres, and finally the rDNA during midanaphase (D'Amours *et al.* 2004; Sullivan *et al.* 2004; Renshaw *et al.* 2010). Cdc14 is required for efficient segregation of telomeres and is essential for removal of cohesin-independent linkages to allow segregation of the rDNA (D'Amours *et al.* 2004; Sullivan *et al.* 2004). Cdc14 shuts down transcription in the rDNA to allow condensin binding to the rDNA, which in turn enables compaction of the rDNA and its efficient segregation (D'Amours *et al.* 2004; Sullivan *et al.* 2004; Machin *et al.* 2006; Tomson

*et al.* 2006; Clemente-Blanco *et al.* 2009). The critical role of condensin in the rDNA may be to facilitate decatenation by topoisomerase II (D'Ambrosio *et al.* 2008a). In support of this, condensin, as well as spindle forces were shown to drive an increase in positive supercoiling that occurs as chromosomes segregate, and this was proposed to promote decatenation by topoisomerase II (Baxter *et al.* 2011). The supercoiling activity of condensin in anaphase is promoted by Polo kinase (Cdc5), which directly phosphorylates multiple condensin subunits (St-Pierre *et al.* 2009). Condensin also localizes to chromosome arms during anaphase and enables recoiling of stretched chromosomes, which promotes the removal of residual cohesin (Renshaw *et al.* 2010). Using a system to artificially open and re-close condensin rings, accurate segregation in anaphase was recently shown to require intact condensin rings. (Cuylen *et al.* 2013). The critical function of condensin in chromosome arm segregation in anaphase, therefore, may be mediated through the topological entrapment of chromosomes.

**Maintaining spindle integrity:** During anaphase, several proteins relocate from kinetochores to the spindle midzone or are newly recruited to the spindle and/or focused to the midzone to promote spindle stability and elongation. *Cdc14* activation is key to this process as it dephosphorylates several substrates to enable their association with the spindle (Pereira and Schiebel 2003; Higuchi and Uhlmann 2005; Khmelinskii *et al.* 2007). The CPC components are so-called because of their conserved relocalization from kinetochores to the spindle midzone during anaphase (Carmena *et al.* 2012). *Cdc14* dephosphorylates the CPC component *Sli15* to enable this transition (Pereira and Schiebel 2003). Another CPC component, *Ipl1*, is also subject to *Cdc14*-dependent removal of CDK-directed phosphorylation, which enables its association with the microtubule plus end-tracking protein, *Bim1* (yeast *EB1* protein), leading to *Ipl1* concentration at the spindle midzone (Nakajima *et al.* 2011; Zimniak *et al.* 2012). SUMOylation of the kinetochore component, *Mcm21*, is also important for CPC relocalization to the midzone (Vizeacoumar *et al.* 2010). The CPC additionally undergoes self-regulation in anaphase by *Ipl1*-dependent phosphorylation of *Sli15*, which directs it away from regions of microtubule dynamics (Nakajima *et al.* 2011).

CPCs are important for midzone assembly, spindle stability, elongation, and disassembly. Although the mechanism by which they achieve these functions is unclear, the CPC is important for recruitment of many downstream effectors. Interestingly, distinct CPC subcomplexes exist that appear to carry out specific functions. Among the proteins recruited to the midzone in anaphase by CPC components are the kinetochore protein *Ndc10* (*Cbf2*), which binds to *Bir1* (Bouck and Bloom 2005; Widlund *et al.* 2006; Thomas and Kaplan 2007; Rozelle *et al.* 2011). *Ndc10*, *Bir1*, and *Sli15* form an alternative CPC that lacks *Ipl1* and regulates spindle elongation (Rozelle *et al.* 2011). *Ndc10* must be SUMOylated for proper spindle stability, suggesting that


SUMOylation might be generally important for spindle midzone assembly (Montpetit *et al.* 2006). A complex of *Slk19–Esp1* additionally relocates to the spindle midzone in a manner dependent on *Sli15*, and this is also required for spindle stability (Khmelinskii *et al.* 2007).

Another key protein at the midzone is the microtubule-bundling protein, *Ase1*. *Ase1* is focused at the midzone by *Cdc14*-dependent dephosphorylation where it recruits downstream components to enable midzone assembly, including the kinesin-5 protein, *Cin8* (Khmelinskii *et al.* 2007, 2009). *Cdc14* further dephosphorylates *Fin1*, a regulatory subunit of the protein phosphatase 1 (PP1) to trigger its association with spindle poles and microtubules and which is also important for spindle stability (Woodbury and Morgan 2007). Overall, *Cdc14* ensures the integrity of the mitotic spindle in anaphase through dephosphorylation of multiple substrates.

Recently, CPC at the spindle midzone has been reported to be important in ensuring that chromosomes are clear of the division plane prior to cytokinesis as part of the “NoCut” pathway (Norden *et al.* 2006; Mendoza *et al.* 2009). However, several situations where chromosomes fail to clear the division plane do not lead to a delay in cytokinesis. Inactivation of topoisomerase II prior to anaphase prevents chromosome segregation, but not cytokinesis, resulting in the “cut” phenotype (abscission of the nucleus by the cleavage plane). (Baxter and Diffley 2008). A failure to resolve rDNA loci after *Cdc14* inactivation results in the presence of a single unsegregated chromosome in the division plane, yet only a short delay in cytokinesis, leading to severing of the chromosome (Quevedo *et al.* 2012). Similarly, though artificial cleavage of condensin prior to anaphase prevents chromosome arm segregation, cytokinesis occurs with normal timing, generating DNA breaks that persist in the next cell cycle (Cuylen *et al.* 2013).

## Chromosome Segregation During Meiosis

Meiosis is a specialized cell division, which produces gametes with half the ploidy of the progenitor cell. Two gametes fuse to restore normal ploidy in the offspring. Diploid budding yeast undergo meiosis to produce four haploid spores. To achieve the halving of ploidy, DNA replication is followed by two consecutive rounds of chromosome segregation. In meiosis I, the maternal and paternal chromosomes, or homologs, are segregated, whereas during meiosis II, which resembles mitosis, the sister chromatids are separated. This pattern of segregation requires several remarkable modifications to the segregation machinery. First, homologous chromosomes must be linked to ensure their accurate segregation in meiosis I. In budding yeast, meiotic recombination generates chiasmata that are important in holding homologs together. Second, sister chromatids must segregate to the same pole, rather than opposite poles during meiosis I. A protein complex known as monopolin ensures that sister kinetochores are mono-oriented during meiosis I. Third, sister chromatids must retain



**Figure 8** Modifications to chromosomes for segregating homologs during meiosis I. Meiotic recombination, monoorientation of kinetochores, and protection of pericentromeric cohesion specialize meiosis I chromosomes to enable homolog segregation.

cohesive linkages between them until their segregation to opposite poles during meiosis II. Therefore, although cohesion on chromosome arms is lost during meiosis I, cohesion around the centromere is protected until meiosis II. Finally, cell cycle controls must be modified so that meiosis I is followed not by DNA replication, but by another chromosome segregation phase, meiosis II. In recent years, our knowledge of the molecular basis for these changes has increased dramatically and much of what we have learned has been from work using budding yeast. These modifications are summarized briefly below (Figure 8). For more detailed discussion see Marston and Amon (2004) and Brar and Amon (2008).

#### **Establishment of links between homologs during meiosis I**

In budding yeast, the establishment of linkages between homologs to mediate their accurate segregation is dependent on meiotic recombination. Meiotic recombination produces a reciprocal exchange between the homologs, called crossovers (COs). COs in their mature state are known as chiasmata and hold homologs together owing to sister chromatid cohesion.

**Homologous chromosome pairing:** As a prerequisite to productive recombination, homologous chromosomes must find each other through a process known as pairing. In early meiotic prophase, centromeres detach from the spindle pole body and telomeres become tethered to the nuclear mem-

brane. SUN proteins in the nuclear membrane link telomeres in the nucleus to motors in the cytoplasm (Hiraoka and Dernburg 2009). Chromosomes undergo rapid telomere-led prophase movements and this facilitates chromosome pairing (Conrad *et al.* 2008; Koszul *et al.* 2008; Lee *et al.* 2012). In budding yeast, pairing is initiated by nonhomologous coupling of the centromeres (Tsubouchi and Roeder 2005). This process requires the *Zip1* protein, a major component of the synaptonemal complex (SC), a proteinaceous structure that forms a scaffold between the homologs (Tsubouchi and Roeder 2005). It is thought that nonhomologous pairing initiates a homology search, culminating in stabilization of homologous chromosome pairs. Consistently, synapsis is initiated at centromeres (Tsubouchi *et al.* 2008). The transition to homologous pairing requires the initiation of recombination, that is the introduction of double strand breaks (DSBs) by the *Spo11* nuclease (Tsubouchi and Roeder 2005). The early telomere-led movements culminate in the clustering of telomeres at a common site, known as the bouquet stage (Scherthan 2001). Although the precise function of the bouquet is debated, it is thought to somehow optimize homolog interactions during meiotic recombination, perhaps by clustering centromeres to enable synapsis initiation (Subramanian and Hochwagen 2011).

**Meiotic recombination:** Meiotic recombination begins with the deliberate introduction of DSBs throughout the genome

by the *Spo11* endonuclease (Keeney *et al.* 1997). Meiotic chromosomes are organized in loops that radiate from a proteinaceous axis (Zickler and Kleckner 1999). The DSBs are made in the loops that become tethered to the axis by *Spo11* accessory proteins (Blat *et al.* 2002; Panizza *et al.* 2011). DSBs are distributed nonrandomly throughout the genome (Gerton *et al.* 2000; Blitzblau *et al.* 2007; Buhler *et al.* 2007; Pan *et al.* 2011). Regions of the genome that are highly proficient for recombination are known as “hotspots,” whereas recombination-poor zones, such as centromere and telomere-proximal regions, are known as “coldspots.” The local structure of chromatin influences its susceptibility to *Spo11*-mediated breakage (Brachet *et al.* 2012). For example, histone H4 lysine 3 trimethylation has been identified as a predisposing mark for breakage (Borde *et al.* 2009).

Following DSB formation, 5' ends are resected through the activity of the *Exo1* exonuclease and bifunctional exo/endonuclease *Mre11*, leaving single strand 3' ends that invade template DNA to form a so-called D-loop (Mimitou and Symington 2008; Zhu *et al.* 2008; Nicolette *et al.* 2010; Garcia *et al.* 2011). This invasion step and homology search is dependent on the RecA-like *Rad51* and *Dmc1* proteins that form nucleoprotein filaments on single stranded DNA (Bishop *et al.* 1992; Shinohara *et al.* 1992). *Dmc1* is meiosis specific and *Rad51*'s strand exchange activity is inhibited by the meiosis-specific protein, *Hed1*, and *Rad51* (Tsubouchi and Roeder 2006). Rather than promote strand exchange directly, *Rad51* acts as an accessory factor for *Dmc1* in strand exchange (Cloud *et al.* 2012). The meiosis-specific employment of *Dmc1* in strand exchange may contribute to ensuring that repair occurs from the homologous chromosome [thereby enabling CO generation], rather than from the sister chromatid, the predominant mode of repair in vegetative cells (Bishop 2012). The organization of meiotic chromosomes into loops secured on an axis is especially important in imposing a bias toward homologous repair as disruption of the axis or sister chromatid cohesion biases repair toward the sister chromatid (Kim *et al.* 2010).

After strand invasion, there are many possible modes of repair resulting in different genetic outcomes (Serrentino and Borde 2012). The only outcome that generates a linkage between the homologs is a CO event, in which exchanges of homologous chromosome arms occur. The number of COs is far fewer than the number of DSBs and this number is maintained even when fewer DSBs are produced (Martini *et al.* 2006). This “CO homeostasis” ensures that sufficient linkages between the homologs are generated. Meiotic recombination is also subject to a phenomenon known as “CO interference,” which prevents additional COs close to sites which have already been designated as COs. The molecular basis for CO interference is unknown but a current hypothesis invokes a stress along the chromatin fiber that prevents further COs in the vicinity (Kleckner *et al.* 2004). A large fraction of DSBs that become COs are processed by a pathway requiring a group of proteins known as the “ZMMs,” through a Holliday junction intermediate (Lynn *et al.* 2007).

A few COs are also produced through a *Mus81*-dependent pathway (Hollingsworth and Brill 2004). The remainder of DSBs are processed as noncrossovers (NCOs) through several different mechanisms (Serrentino and Borde 2012). The excess DSBs, which eventually lead to NCOs, have been suggested to be functionally important in chromosome pairing by increasing interhomolog interactions (Tessé *et al.* 2003).

Surveillance mechanisms, or checkpoints, monitor the progression of meiotic recombination to ensure that meiosis does not progress in the presence of DNA lesions (MacQueen and Hochwagen 2011). Once all DSBs have been repaired, the *Ndt80* transcription factor becomes active and drives the expression of genes required for pachytene exit and the meiotic divisions (Chu and Herskowitz 1998; Sourirajan and Lichten 2008).

### ***Monoorientation of sister chromatids during meiosis I***

A defining feature of meiosis I is the segregation of homologs to opposite poles rather than sister chromatids, which move toward the same pole. That is, sister kinetochores must be uniquely monooriented during meiosis I. Pioneering experiments in grasshopper spermatocytes demonstrated that this is a property of the kinetochore rather than the meiotic spindle (Paliulis and Nicklas 2000). We know little about how this is achieved, except in budding yeast where factors specifically required for kinetochore monoorientation have been identified. Monopolar attachment depends on a meiosis-specific protein, *Mam1*, two nucleolar proteins, *Lrs4* and *Csm1*, that together with the casein kinase, *Hrr25*, form a complex called monopolin (Toth *et al.* 2000; Rabitsch *et al.* 2003; Petronczki *et al.* 2006). Cells lacking monopolin fail to monoorient sister kinetochores and biorient sister kinetochores instead, which leads to a failure to undergo the first meiotic division after arm cohesion is lost due to the persistence of centromere cohesion (Toth *et al.* 2000; Rabitsch *et al.* 2003; Petronczki *et al.* 2006). Polo kinase (*Cdc5*) is also required for monoorientation, in part due to a requirement for *Cdc5* in release of *Lrs4* and *Csm1* from the nucleolus (Clyne *et al.* 2003; Lee and Amon 2003). *Cdc5* additionally collaborates with the meiosis-specific regulator, *Spo13* and *Dbf4*-dependent kinase, *Cdc7* (DDK) to bring about *Lrs4* phosphorylation, enabling monopolin recruitment to kinetochores (Matos *et al.* 2008).

Homologs of *Lrs4* and *Csm1* exist in fission yeast, but rather than bring about monopolar attachment during meiosis I, they prevent merotelic attachment during mitosis (Gregan *et al.* 2007). Instead, *Rec8*-containing cohesin is important for monoorientation in fission yeast (Watanabe and Nurse 1999). Unlike mitotic *Rad21* containing cohesin (the equivalent of *Mcd1/Sccl*-containing cohesin in budding yeast), *Rec8* cohesin is enriched within the core centromere in fission yeast (Yokobayashi and Watanabe 2005). In budding yeast, monopolin is sufficient to link sister kinetochores during meiosis I, even in the absence of cohesin (Monje-Casas

*et al.* 2007). However, condensin contributes to monoorientation (Brito *et al.* 2010). This has led to the proposal that the essential feature of monooriented kinetochores is that sister centromeres are closely linked either by monopolin in the case of budding yeast or by cohesion in the case of fission yeast (Watanabe 2006).

How does monopolin physically link sister kinetochores? The possibility that two sister kinetochores could be fused into a single microtubule-binding unit was suggested by images of insect kinetochores in meiosis I (Goldstein 1981; Suja *et al.* 1991) and because budding yeast kinetochores were observed to bind a single microtubule by electron microscopy in meiosis I (Winey *et al.* 2005). Alternatively one kinetochore might be “silenced” so that only one kinetochore between the two sisters is competent to bind microtubules. In budding yeast, structural studies of monopolin have led to a model that is most consistent with the kinetochore fusion model. A complex of *Lrs4/Csm1* forms a V-shaped structure, the apices of which form contacts with the *Dsn1* kinetochore subunits, suggesting that monopolin could crosslink two sister kinetochores (Corbett *et al.* 2010; Corbett and Harrison 2012). *Mam1* and *Hrr25* associate with *Csm1* close to the site of *Dsn1* interaction and somehow modulate association of monopolin with the kinetochore (Corbett and Harrison 2012). A major challenge for the future will be to understand how monopolin alters the interface between microtubules and the kinetochore.

### **Stepwise loss of cohesion**

The introduction of at least one CO per pair of homologs generates a physical link between homologs due to sister chromatid cohesion distal to the CO. This provides the tension that allows the homologs to align on the meiosis I spindle, ready for their segregation to opposite poles during meiosis I. Release of cohesion on chromosome arms triggers the segregation of homologs to opposite poles. However, cohesion in centromeric regions must be preserved during meiosis I to ensure the accurate segregation of sister chromatids during meiosis II. During meiosis, the *Scc1* kleisin subunit of cohesin is replaced by its meiosis-specific homolog, *Rec8* (Klein *et al.* 1999). *Rec8* on chromosome arms is cleaved by separase during meiosis I, but *Rec8* in centromeric regions is maintained until meiosis II (Klein *et al.* 1999; Buonomo *et al.* 2000). The ability of pericentromeric cohesin to resist separase activity during meiosis I is a unique property of *Rec8*-containing cohesin and cannot be fulfilled by *Scc1* (Toth *et al.* 2000). Similarly, *Rec8* performs functions in meiotic pairing, recombination, and chromosome axis formation that *Scc1* cannot support (Klein *et al.* 1999; Brar *et al.* 2009). For its cleavage during meiosis I, *Rec8* needs to be phosphorylated (Brar *et al.* 2006; Katis *et al.* 2010; Attner *et al.* 2013). The *Dbf4*-dependent *Cdc7* kinase (DDK), casein kinase 1 $\delta/\epsilon$  (*Hrr25*) and Polo kinase, *Cdc5*, all phosphorylate *Rec8* during meiosis I and promote its cleavage to some extent (Brar *et al.* 2006; Katis *et al.* 2010; Attner *et al.* 2013). Protection of pericentromeric

cohesin during meiosis I requires the conserved Shugoshin (*Sgo1*) protein that is localized in the pericentromere (Katis *et al.* 2004a; Kitajima *et al.* 2004; Marston *et al.* 2004; Kiburz *et al.* 2005; Clift and Marston 2011). *Sgo1* recruits a particular form of the protein phosphatase 2A, that contains its *Rts1* regulatory subunit, to the pericentromere (Kitajima *et al.* 2006; Riedel *et al.* 2006; Xu *et al.* 2009). In the absence of the alternative PP2A regulatory subunit, *Cdc55*, excess PP2A–*Rts1* complexes form and are highly elevated on chromosomes, preventing *Rec8* phosphorylation and cleavage (Bizzari and Marston 2011). Altogether, this leads to a model whereby PP2A–*Rts1* recruitment by *Sgo1* renders pericentromeric *Rec8* resistant to separase activity by maintaining it in the unphosphorylated state.

Other factors that are important for the protection of pericentromeric cohesion during meiosis I include the meiosis I-specific protein, *Spo13* (Katis *et al.* 2004b; Lee *et al.* 2004), which is required for proper localization of *Sgo1* at centromeres (Kiburz *et al.* 2005). Exactly how *Spo13* contributes to cohesion protection is unknown but it is interesting to note that *Spo13* binds to Polo kinase *Cdc5* during meiosis I (Matos *et al.* 2008), given that *Cdc5* is also important for the protection of cohesion during meiosis I under certain conditions (Katis *et al.* 2010). The Aurora kinase *Ipl1* additionally contributes to cohesion protection, apparently by maintaining PP2A–*Rts1* at centromeres (Yu and Koshland 2007). How these additional protective factors work together to regulate *Sgo1*–PP2A–*Rts1* is an important question to address in the future. The question of how cohesin is “deprotected” during meiosis II is also a priority for future study.

### **Biorientation of homologs**

During meiosis I, sister chromatids are monooriented and instead it is the homologous chromosomes that must achieve biorientation. The arm cohesion distal to chiasmata provides the tension that enables homologs to biorient on the meiosis I spindle. As in mitosis, the spindle checkpoint together with *Ipl1* and *Mps1* play an important role in this process. As in mitosis, *Ipl1* is important to trigger the release of kinetochores from microtubules early in meiosis (Monje-Casas *et al.* 2007; Meyer *et al.* 2013). This provides an opportunity for chromosomes to pair. Following recombination and prophase exit, homologs initially tend to make incorrect attachments to the meiosis I spindle that are destabilized by *Ipl1* to provide a further opportunity for the correct attachments to be made (Meyer *et al.* 2013). *Mps1* kinase is required for the conversion of lateral kinetochore–microtubule attachments into stable end-on attachments and is therefore also critical for homolog biorientation (Meyer *et al.* 2013). The spindle checkpoint protein, *Mad2*, also contributes to proper homolog biorientation (Shonn *et al.* 2003). Interestingly, chromosomes with COs far from the centromere are particularly reliant on *Mad2* function, suggesting that their alignment presents a particular challenge (Lacefield and Murray 2007). *Mad2* plays a further role, together with *Mad3*, in

delaying the cell cycle in response to kinetochore–microtubule attachment defects during meiosis I to increase the possibility that biorientation will be achieved (Shonn *et al.* 2000, 2003).

How tension at meiosis I kinetochores is sensed is not clear. However, *Sgo1* protein, which responds to a lack of tension between sister kinetochores during mitosis, plays only a minor role in sensing tension between homologous chromosomes during meiosis I, suggesting that a distinct mechanism is at work during meiosis I (Kiburz *et al.* 2008). Intriguingly, *Zip1*, a major component of the SC complex, persists at centromeres after SC disassembly (Gladstone *et al.* 2009; Newnham *et al.* 2010). Centromere-associated *Zip1* may aid homolog biorientation during meiosis I by coupling homologous kinetochores to provide a favorable geometry for their capture by microtubules from opposite poles.

### **Alteration of cell cycle controls in meiosis**

The extended prophase in which recombination takes place and the existence of two consecutive chromosome segregation phases without an intervening S phase are defining features of meiosis that require a specialization of cell cycle controls. As in mitosis, cell cycle progression is controlled by a single CDK, *Cdc28*, in complex with cyclins (reviewed in Marston and Amon 2004). There are six B-type cyclins in budding yeast, *Clb1–6*. The S phase cyclins *Clb5* and *Clb6* drive DNA replication, as in mitosis (Dirick *et al.* 1998; Stuart and Wittenberg 1998), and the initiation of recombination (Henderson *et al.* 2006). *Clb1*, -3, and -4 are important for the meiotic divisions, but the major mitotic cyclin, *Clb2*, is not produced during meiosis (Grandin and Reed 1993). CDK activity during meiosis is tightly controlled through transcriptional, translational, and post-translational mechanisms (Grandin and Reed 1993; Carlile and Amon 2008). *Clb1*–CDK activity is restricted to meiosis I, whereas *Clb3*–CDK activity is restricted to meiosis II (Carlile and Amon 2008). The meiosis-specific CDK-related kinase, *Ime2*, is also important for meiotic entry and progression (Benjamin *et al.* 2003; Irniger 2011). *Ime2* activity peaks during prophase I, declines during meiosis I, and peaks again during meiosis II (Benjamin *et al.* 2003; Irniger 2011; Berchowitz *et al.* 2013). An elegant study showed that this pattern of *Ime2* activity is important to restrict the translation of a subset of mRNA, including *CLB3*, to meiosis II (Berchowitz *et al.* 2013). At the meiosis I to meiosis II transition, *Ime2*-dependent downregulation of the RNA-binding protein, *Rim4*, relieves the repression on translation of these genes, thereby coordinating the meiotic program (Berchowitz *et al.* 2013).

**Meiotic prophase I to meiosis I transition:** As recombination takes place, exit from prophase is prevented because the recombination checkpoint represses the *Ndt80* transcription factor that is required for the expression of M phase regulators, including cyclins and Polo kinase (Chu and Herskowitz 1998; Hochwagen and Amon 2006; Sourirajan and Lichten 2008). The APC/C additionally prevents the accumula-

tion of M phase regulators during meiotic prophase by targeting them for destruction (Okaz *et al.* 2012). Budding yeast carry a meiosis-specific APC activator, *Ama1*, that appears in S phase (Cooper *et al.* 2000) and targets M phase regulators for destruction to prevent exit from prophase (Okaz *et al.* 2012). However, remarkably, securin (*Pds1*) and *Sgo1* are spared from APC–*Ama1*-dependent destruction in prophase by the *Mnd2/Apc15* APC subunit, which behaves as a substrate-specific inhibitor of the APC in this context (Oelschlaegel *et al.* 2005; Penkner *et al.* 2005). Note that, in contrast, *Mnd2/Apc15* was found to stimulate *Cdc20* autoubiquitin (see above; Foster and Morgan 2012), suggesting that regulation of the APC by *Mnd2/Apc15* might be complex. The spindle checkpoint protein, *Mad2*, additionally prevents premature APC–*Ama1* activity to ensure proper chromosome segregation in meiosis I, probably indirectly through APC–*Cdc20* inhibition (Tsuchiya *et al.* 2011). Simultaneous activation and inhibition of APC–*Ama1* toward distinct substrates is therefore critical for faithful chromosome segregation at meiosis I, yet how this is achieved is so far unknown.

The importance of restricting CDK activity until after prophase I exit was demonstrated by finding that production of the meiotic cyclins *Clb1* and *Clb3* during premeiotic S phase and prophase interferes with the program of meiosis I chromosome segregation (Carlile and Amon 2008; Miller *et al.* 2012). In these cells, kinetochore microtubule attachments are established prematurely so that monoorientation of kinetochores and the protection of centromeric cohesion is precluded (Miller *et al.* 2012). This demonstrates that kinetochore–microtubule interactions must be prevented during prophase I for the key features of meiosis I chromosomes to be established.

**Meiosis I to meiosis II transition:** After chromosomes segregate during mitosis, CDKs are inactivated, which allows for spindle disassembly and return to G1. This state of low CDK activity upon exit from mitosis is permissive for the resetting of origins of DNA replication, in preparation for the next S phase. However, homolog segregation during meiosis I is followed not by S phase, but by another “M” phase, meiosis II. This means that at meiosis I exit, spindles must disassemble but replication origins must not be reset. This predicts that specialized controls regulate the meiosis I to meiosis II transition.

As during exit from mitosis, the *Cdc14* phosphatase is critical for the meiosis I to meiosis II transition (Buonomo *et al.* 2003; Marston *et al.* 2003). *Cdc14* is released from sequestration in the nucleolus at meiosis I exit due to the activity of the FEAR network, however the MEN functions only in meiosis II (Buonomo *et al.* 2003; Marston *et al.* 2003; Kamieniecki *et al.* 2005; Attner and Amon 2012). Cells with impaired *Cdc14* activity undergo only a single meiotic division in which some chromosomes segregate in a meiosis I-like manner, whereas others segregate in a meiosis II-like manner (Buonomo *et al.* 2003; Marston *et al.* 2003). The

important function of *Cdc14* in meiosis appears to be to allow reduplication of the spindle to ensure that meiosis I and meiosis II segregation occur on consecutively built spindles (Marston *et al.* 2003; Bizzari and Marston 2011). As in mitosis, PP2A, together with its *Cdc55* regulatory subunit, plays a critical role in keeping *Cdc14* sequestered in the nucleolus. In the absence of *Cdc55*, *Cdc14* is released prematurely and this prevents spindle assembly during meiosis I (Bizzari and Marston 2011; Kerr *et al.* 2011). How *Cdc14* impinges on spindle assembly and duplication is unclear. It is thought that *Cdc14* effects must be restricted during meiosis I exit to prevent events such as the resetting of replication origins occurring. Interestingly, *Ime2*-dependent phosphorylation events appear to be resistant to *Cdc14* activity, providing a potential mechanism to limit its activity toward certain substrates (Holt *et al.* 2007). Indeed, the *Mcm2–7* replicative helicase is excluded from the nucleus, which contributes to the prevention of replication origin relicensing, from premeiotic S phase onwards, due to *Ime2* and CDK-dependent phosphorylation (Holt *et al.* 2007).

## Perspectives

Remarkable progress in understanding chromosome segregation mechanisms in eukaryotes has been gained from studies in budding yeast. However, many features of this process and its regulation remain elusive. Although the central players are all known, understanding how they function mechanistically and cooperate with each other in the context of the cell cycle are key challenges. Structural and biochemical studies as well as systems level analysis, all combined with the powerful genetics that the yeast system offers, will be pivotal in driving forward the next era of chromosome segregation research.

## Acknowledgments

I apologize to colleagues whose work is not cited directly. I am grateful to Angelika Amon, Tomo Tanaka, Frank Uhlmann, Kevin Hardwick, and Eris Duro for helpful comments on the manuscript and members of my laboratory for useful discussions. Work in my lab is funded by the Wellcome Trust (090903).

## Literature Cited

- Agarwal, R., and O. Cohen-Fix, 2002 Phosphorylation of the mitotic regulator Pds1/securin by Cdc28 is required for efficient nuclear localization of Esp1/separase. *Genes Dev.* 16: 1371–1382.
- Akiyoshi, B., C. R. Nelson, J. A. Ranish, and S. Biggins, 2009a Analysis of Ipl1-mediated phosphorylation of the Ndc80 kinetochore protein in *Saccharomyces cerevisiae*. *Genetics* 183: 1591–1595.
- Akiyoshi, B., C. R. Nelson, J. A. Ranish, and S. Biggins, 2009b Quantitative proteomic analysis of purified yeast kinetochores identifies a PP1 regulatory subunit. *Genes Dev.* 23: 2887–2899.
- Akiyoshi, B., K. K. Sarangapani, A. F. Powers, C. R. Nelson, S. L. Reichow *et al.*, 2010 Tension directly stabilizes reconstituted kinetochore-microtubule attachments. *Nature* 468: 576–579.
- Alexandru, G., F. Uhlmann, K. Mechtler, M. A. Poupart, and K. Nasmyth, 2001 Phosphorylation of the cohesin subunit Scc1 by Polo/Cdc5 kinase regulates sister chromatid separation in yeast. *Cell* 105: 459–472.
- Almedawar, S., N. Colomina, M. Bermúdez-López, I. Pociño-Merino, and J. Torres-Rosell, 2012 A SUMO-dependent step during establishment of sister chromatid cohesion. *Curr. Biol.* 22: 1576–1581.
- Anderson, D. E., A. Losada, H. P. Erickson, and T. Hirano, 2002 Condensin and cohesin display different arm conformations with characteristic hinge angles. *J. Cell Biol.* 156: 419–424.
- Arumugam, P., S. Gruber, K. Tanaka, C. H. Haering, K. Mechtler *et al.*, 2003 ATP hydrolysis is required for cohesin's association with chromosomes. *Curr. Biol.* 13: 1941–1953.
- Arumugam, P., T. Nishino, C. H. Haering, S. Gruber, and K. Nasmyth, 2006 Cohesin's ATPase activity is stimulated by the C-terminal Winged-Helix domain of its kleisin subunit. *Curr. Biol.* 16: 1998–2008.
- Attner, M. A., and A. Amon, 2012 Control of the mitotic exit network during meiosis. *Mol. Biol. Cell* 23: 3122–3132.
- Attner, M. A., M. P. Miller, L.-S. Ee, S. K. Elkin, and A. Amon, 2013 Polo kinase Cdc5 is a central regulator of meiosis I. *Proc. Natl. Acad. Sci. USA* 110: 14278–14283.
- Azzam, R., S. L. Chen, W. Shou, A. S. Mah, G. Alexandru *et al.*, 2004 Phosphorylation by cyclin B-Cdk underlies release of mitotic exit activator Cdc14 from the nucleolus. *Science* 305: 516–519.
- Bachant, J., A. Alcasabas, Y. Blat, N. Kleckner, and S. J. Elledge, 2002 The SUMO-1 isopeptidase Smt4 is linked to centromeric cohesion through SUMO-1 modification of DNA topoisomerase II. *Mol. Cell* 9: 1169–1182.
- Bachelier-Bassi, S., O. Gadal, G. Bourout, and U. Nehrass, 2008 Cell cycle-dependent kinetochore localization of condensin complex in *Saccharomyces cerevisiae*. *J. Struct. Biol.* 162: 248–259.
- Bando, M., Y. Katou, M. Komata, H. Tanaka, T. Itoh *et al.*, 2009 Csm3, Tof1, and Mrc1 form a heterotrimeric mediator complex that associates with DNA replication forks. *J. Biol. Chem.* 284: 34355–34365.
- Bausch, C., S. Noone, J. M. Henry, K. Gaudenz, B. Sanderson *et al.*, 2007 Transcription alters chromosomal locations of cohesin in *Saccharomyces cerevisiae*. *Mol. Cell Biol.* 27: 8522–8532.
- Baxter, J., and J. F. X. Diffley, 2008 Topoisomerase II inactivation prevents the completion of DNA replication in budding yeast. *Mol. Cell* 30: 790–802.
- Baxter, J., and L. Aragón, 2012 A model for chromosome condensation based on the interplay between condensin and topoisomerase II. *Trends Genet.* 28: 110–117.
- Baxter, J., N. Sen, V. L. Martínez, M. E. M. De Carandini, J. B. Schwartzman *et al.*, 2011 Positive supercoiling of mitotic DNA drives decatenation by topoisomerase II in eukaryotes. *Science* 331: 1328–1332.
- Beckouët, F., B. Hu, M. B. Roig, T. Sutani, M. Komata *et al.*, 2010 An Smc3 acetylation cycle is essential for establishment of sister chromatid cohesion. *Mol. Cell* 39: 689–699.
- Ben-Shahar, T. R., S. Heeger, C. Lehane, P. East, H. Flynn *et al.*, 2008 Eco1-dependent cohesin acetylation during establishment of sister chromatid cohesion. *Science* 321: 563–566.
- Benjamin, K. R., C. Zhang, K. M. Shokat, and I. Herskowitz, 2003 Control of landmark events in meiosis by the CDK Cdc28 and the meiosis-specific kinase Ime2. *Genes Dev.* 17: 1524–1539.
- Berchowitz, L. E., A. S. Gajadhar, F. J. van Werven, A. A. De Rosa, M. L. Samoylova *et al.*, 2013 A developmentally regulated

- translational control pathway establishes the meiotic chromosome segregation pattern. *Genes Dev.* 27: 2147–2163.
- Bermudez, V. P., Y. Maniwa, I. Tappin, K. Ozato, K. Yokomori *et al.*, 2003 The alternative Ctf18-Dcc1-Ctf8-replication factor C complex required for sister chromatid cohesion loads proliferating cell nuclear antigen onto DNA. *Proc. Natl. Acad. Sci. USA* 100: 10237–10242.
- Bhalla, N., S. Biggins, and A. W. Murray, 2002 Mutation of YCS4, a budding yeast condensin subunit, affects mitotic and nonmitotic chromosome behavior. *Mol. Biol. Cell* 13: 632–645.
- Bhat, M. A., A. V. Philp, D. M. Glover, and H. J. Bellen, 1996 Chromatid segregation at anaphase requires the barren product, a novel chromosome-associated protein that interacts with Topoisomerase II. *Cell* 87: 1103–1114.
- Biggins, S., and A. W. Murray, 2001 The budding yeast protein kinase Ipl1/Aurora allows the absence of tension to activate the spindle checkpoint. *Genes Dev.* 15: 3118–3129.
- Biggins, S., F. F. Severin, N. Bhalla, I. Sassoon, A. A. Hyman *et al.*, 1999 The conserved protein kinase Ipl1 regulates microtubule binding to kinetochores in budding yeast. *Genes Dev.* 13: 532–544.
- Bishop, D. K., 2012 Rad51, the lead in mitotic recombinational DNA repair, plays a supporting role in budding yeast meiosis. *Cell Cycle* 11: 4105–4106.
- Bishop, D. K., D. Park, L. Xu, and N. Kleckner, 1992 DMCl1: a meiosis-specific yeast homolog of *E. coli* recA required for recombination, synaptonemal complex formation, and cell cycle progression. *Cell* 69: 439–456.
- Bizzari, F., and A. L. Marston, 2011 Cdc55 coordinates spindle assembly and chromosome disjunction during meiosis. *J. Cell Biol.* 193: 1213–1228.
- Blat, Y., and N. Kleckner, 1999 Cohesins bind to preferential sites along yeast chromosome III, with differential regulation along arms vs. the centric region. *Cell* 98: 249–259.
- Blat, Y., R. U. Protacio, N. Hunter, and N. Kleckner, 2002 Physical and functional interactions among basic chromosome organizational features govern early steps of meiotic chiasma formation. *Cell* 111: 791–802.
- Blitzblau, H. G., G. W. Bell, J. Rodriguez, S. P. Bell, and A. Hochwagen, 2007 Mapping of meiotic single-stranded DNA reveals double-stranded-break hotspots near centromeres and telomeres. *Curr. Biol.* 17: 2003–2012.
- Bloom, J., I. M. Cristea, A. L. Procko, V. Lubkov, B. T. Chait *et al.*, 2011 Global analysis of Cdc14 phosphatase reveals diverse roles in mitotic processes. *J. Biol. Chem.* 286: 5434–5445.
- Borde, V., N. Robine, W. Lin, S. Bonfils, V. Géli *et al.*, 2009 Histone H3 lysine 4 trimethylation marks meiotic recombination initiation sites. *EMBO J.* 28: 99–111.
- Borges, V., C. Lehane, L. Lopez-Serra, H. Flynn, M. Skehel *et al.*, 2010 Hos1 deacetylates Smc3 to close the cohesin acetylation cycle. *Mol. Cell* 39: 677–688.
- Borges, V., D. J. Smith, I. Whitehouse, and F. Uhlmann, 2013 An Eco1-independent sister chromatid cohesion establishment pathway in *S. cerevisiae*. *Chromosoma* 122: 121–134.
- Bouchoux, C., and F. Uhlmann, 2011 A quantitative model for ordered Cdk substrate dephosphorylation during mitotic exit. *Cell* 147: 803–814.
- Bouck, D. C., and K. S. Bloom, 2005 The kinetochore protein Ndc10p is required for spindle stability and cytokinesis in yeast. *Proc. Natl. Acad. Sci. USA* 102: 5408–5413.
- Brachet, E., V. Sommermeyer, and V. Borde, 2012 Interplay between modifications of chromatin and meiotic recombination hotspots. *Biol. Cell* 104: 51–69.
- Brady, D. M., and K. G. Hardwick, 2000 Complex formation between Mad1p, Bub1p and Bub3p is crucial for spindle checkpoint function. *Curr. Biol.* 10: 675–678.
- Brar, G. A., and A. Amon, 2008 Emerging roles for centromeres in meiosis I chromosome segregation. *Nat. Rev.* 9: 899–910.
- Brar, G. A., B. M. Kiburz, Y. Zhang, J. E. Kim, F. White *et al.*, 2006 Rec8 phosphorylation and recombination promote the step-wise loss of cohesins in meiosis. *Nature* 441: 532–536.
- Brar, G. A., A. Hochwagen, L.-S. S. Ee, and A. Amon, 2009 The multiple roles of cohesin in meiotic chromosome morphogenesis and pairing. *Mol. Biol. Cell* 20: 1030–1047.
- Brito, I. L., H.-G. Yu, and A. Amon, 2010 Condensins promote coorientation of sister chromatids during meiosis I in budding yeast. *Genetics* 185: 55–64.
- Buhler, C., V. Borde, and M. Lichten, 2007 Mapping meiotic single-strand DNA reveals a new landscape of DNA double-strand breaks in *Saccharomyces cerevisiae*. *PLoS Biol.* 5: e324.
- Buonomo, S. B., R. K. Clyne, J. Fuchs, J. Loidl, F. Uhlmann *et al.*, 2000 Disjunction of homologous chromosomes in meiosis I depends on proteolytic cleavage of the meiotic cohesin Rec8 by separin. *Cell* 103: 387–398.
- Buonomo, S. B., K. P. Rabitsch, J. Fuchs, S. Gruber, M. Sullivan *et al.*, 2003 Division of the nucleolus and its release of CDC14 during anaphase of meiosis I depends on separase, SPO12, and SLK19. *Dev. Cell* 4: 727–739.
- Burton, J. L., and M. J. Solomon, 2007 Mad3p, a pseudosubstrate inhibitor of APCCdc20 in the spindle assembly checkpoint. *Genes Dev.* 21: 655–667.
- Bürmann, F., H.-C. Shin, J. Basquin, Y.-M. Soh, V. Giménez-Oya *et al.*, 2013 An asymmetric SMC-kleisin bridge in prokaryotic condensin. *Nat. Struct. Mol. Biol.* 20: 371–379.
- Bylund, G. O., and P. M. J. Burgers, 2005 Replication protein A-directed unloading of PCNA by the Ctf18 cohesion establishment complex. *Mol. Cell Biol.* 25: 5445–5455.
- Calabria, I., B. Baro, J.-A. Rodriguez-Rodriguez, N. Russiñol, and E. Queralt, 2012 Zds1 regulates PP2A(Cdc55) activity and Cdc14 activation during mitotic exit through its Zds\_C motif. *J. Cell Sci.* 125: 2875–2884.
- Calzada, A., B. Hodgson, M. Kanemaki, A. Bueno, and K. Labib, 2005 Molecular anatomy and regulation of a stable replisome at a paused eukaryotic DNA replication fork. *Genes Dev.* 19: 1905–1919.
- Campbell, C. S., and A. Desai, 2013 Tension sensing by Aurora B kinase is independent of survivin-based centromere localization. *Nature* 497: 118–121.
- Carlile, T. M., and A. Amon, 2008 Meiosis I is established through division-specific translational control of a cyclin. *Cell* 133: 280–291.
- Carmena, M., M. Wheelock, H. Funabiki, and W. C. Earnshaw, 2012 The chromosomal passenger complex (CPC): from easy rider to the godfather of mitosis. *Nat. Rev. Mol. Cell Biol.* 13: 789–803.
- Caydasi, A. K., and G. Pereira, 2012 SPOC alert—when chromosomes get the wrong direction. *Exp. Cell Res.* 318: 1421–1427.
- Chan, K.-L., M. B. Roig, B. Hu, F. Beckouët, J. Metson *et al.*, 2012 Cohesin's DNA Exit Gate Is Distinct from Its Entrance Gate and Is Regulated by Acetylation. *Cell* 150: 961–974.
- Chao, W. C. H., K. Kulkarni, Z. Zhang, E. H. Kong, and D. Barford, 2012 Structure of the mitotic checkpoint complex. *Nature* 484: 208–213.
- Cheeseman, I. M., S. Anderson, M. Jwa, E. M. Green, and J. Kang *et al.*, 2002 Phospho-regulation of kinetochore-microtubule attachments by the Aurora kinase Ipl1p. *Cell* 111: 163–172.
- Chen, R. H., A. Shevchenko, M. Mann, and A. W. Murray, 1998 Spindle checkpoint protein Xmad1 recruits Xmad2 to unattached kinetochores. *J. Cell Biol.* 143: 283–295.
- Chen, R. H., D. M. Brady, D. Smith, A. W. Murray, and K. G. Hardwick, 1999 The spindle checkpoint of budding yeast depends on a tight complex between the Mad1 and Mad2 proteins. *Mol. Biol. Cell* 10: 2607–2618.
- Chiroli, E., V. Rossio, G. Lucchini, and S. Piatti, 2007 The budding yeast PP2ACdc55 protein phosphatase prevents the onset of



- anaphase in response to morphogenetic defects. *J. Cell Biol.* 177: 599–611.
- Cho, U. S., and S. C. Harrison, 2012 Ndc10 is a platform for inner kinetochore assembly in budding yeast. *Nat. Struct. Mol. Biol.* 19: 48–55.
- Chu, S., and I. Herskowitz, 1998 Gametogenesis in yeast is regulated by a transcriptional cascade dependent on Ndt80. *Mol. Cell* 1: 685–696.
- Ciosk, R., W. Zachariae, C. Michaelis, A. Shevchenko, M. Mann *et al.*, 1998 An ESP1/PDS1 complex regulates loss of sister chromatid cohesion at the metaphase to anaphase transition in yeast. *Cell* 93: 1067–1076.
- Ciosk, R., M. Shirayama, A. Shevchenko, T. Tanaka, A. Toth *et al.*, 2000 Cohesin's binding to chromosomes depends on a separate complex consisting of Scc2 and Scc4 proteins. *Mol. Cell* 5: 243–254.
- Clemente-Blanco, A., M. Mayan-Santos, D. A. Schneider, F. Machín, A. Jarmuz *et al.*, 2009 Cdc14 inhibits transcription by RNA polymerase I during anaphase. *Nature* 458: 219–222.
- Clift, D., and A. L. Marston, 2011 The role of Shugoshin in meiotic chromosome segregation. *Cytogenet. Genome Res.* 133: 234–242.
- Clift, D., F. Bizzari, and A. L. Marston, 2009 Shugoshin prevents cohesin cleavage by PP2A(Cdc55)-dependent inhibition of separase. *Genes Dev.* 23: 766–780.
- Cloud, V., Y.-L. Chan, J. Grubb, B. Budke, and D. K. Bishop, 2012 Rad51 is an accessory factor for Dmc1-mediated joint molecule formation during meiosis. *Science* 337: 1222–1225.
- Clyne, R. K., V. L. Katis, L. Jessop, K. R. Benjamin, I. Herskowitz *et al.*, 2003 Polo-like kinase Cdc5 promotes chiasmata formation and cosegregation of sister centromeres at meiosis I. *Nat. Cell Biol.* 5: 480–485.
- Cohen-Fix, O., J. M. Peters, M. W. Kirschner, and D. Koshland, 1996 Anaphase initiation in *Saccharomyces cerevisiae* is controlled by the APC-dependent degradation of the anaphase inhibitor Pds1p. *Genes Dev.* 10: 3081–3093.
- Conrad, M. N., C.-Y. Lee, G. Chao, M. Shinohara, H. Kosaka *et al.*, 2008 Rapid telomere movement in meiotic prophase is promoted by NDJ1, MPS3, and CSM4 and is modulated by recombination. *Cell* 133: 1175–1187.
- Cooper, K. F., M. J. Mallory, D. B. Egeland, M. Jarnik, and R. Strich, 2000 Ama1p is a meiosis-specific regulator of the anaphase promoting complex/cyclosome in yeast. *Proc. Natl. Acad. Sci. USA* 97: 14548–14553.
- Corbett, K. D., and S. C. Harrison, 2012 Molecular architecture of the yeast monopolin complex. *Cell Rep* 1: 583–589.
- Corbett, K. D., C. K. Yip, L. S. Ee, T. Walz, A. Amon *et al.*, 2010 The monopolin complex crosslinks kinetochore components to regulate chromosome-microtubule attachments. *Cell* 142: 556–567.
- Cubeñas-Potts, C., and M. J. Matunis, 2013 SUMO: a multifaceted modifier of chromatin structure and function. *Dev. Cell* 24: 1–12.
- Cuylen, S., and C. H. Haering, 2011 Deciphering condensin action during chromosome segregation. *Trends Cell Biol.* 21: 552–559.
- Cuylen, S., J. Metz, and C. H. Haering, 2011 Condensin structures chromosomal DNA through topological links. *Nat. Struct. Mol. Biol.* 18: 894–901.
- Cuylen, S., J. Metz, A. Hruba, and C. H. Haering, 2013 Entrapment of chromosomes by condensin rings prevents their breakage during cytokinesis. *Dev. Cell* DOI: .10.1016/j.devcel.2013.10.018
- D'Ambrosio, C., G. Kelly, K. Shirahige, and F. Uhlmann, 2008a Condensin-dependent rDNA decatenation introduces a temporal pattern to chromosome segregation. *Curr. Biol.* 18: 1084–1089.
- D'Ambrosio, C., C. K. Schmidt, Y. Katou, G. Kelly, T. Itoh *et al.*, 2008b Identification of cis-acting sites for condensin loading onto budding yeast chromosomes. *Genes Dev.* 22: 2215–2227.
- D'Amours, D., F. Stegmeier, and A. Amon, 2004 Cdc14 and condensin control the dissolution of cohesin-independent chromosome linkages at repeated DNA. *Cell* 117: 455–469.
- Davenport, J., L. D. Harris, and R. Goorha, 2006 Spindle checkpoint function requires Mad2-dependent Cdc20 binding to the Mad3 homology domain of BubR1. *Exp. Cell Res.* 312: 1831–1842.
- De Antoni, A., C. G. Pearson, D. Cimini, J. C. Canman, V. Sala *et al.*, 2005 The Mad1/Mad2 complex as a template for Mad2 activation in the spindle assembly checkpoint. *Curr. Biol.* 15: 214–225.
- Demirel, P. B., B. E. Keyes, M. Chatterjee, C. E. Remington, and D. J. Burke, 2012 A redundant function for the N-terminal tail of Ndc80 in kinetochore-microtubule interaction in *Saccharomyces cerevisiae*. *Genetics* 192: 753–756.
- Dewar, H., K. Tanaka, K. Nasmyth, and T. U. Tanaka, 2004 Tension between two kinetochores suffices for their bi-orientation on the mitotic spindle. *Nature* 428: 93–97.
- DiNardo, S., K. Voelkel, and R. Sternglanz, 1984 DNA topoisomerase II mutant of *Saccharomyces cerevisiae*: topoisomerase II is required for segregation of daughter molecules at the termination of DNA replication. *Proc. Natl. Acad. Sci. USA* 81: 2616–2620.
- Dirick, L., L. Goetsch, G. Ammerer, and B. Byers, 1998 Regulation of meiotic S phase by Ime2 and a Clb5,6-associated kinase in *Saccharomyces cerevisiae*. *Science* 281: 1854–1857.
- Earnshaw, W. C., B. Halligan, C. A. Cooke, M. M. Heck, and L. F. Liu, 1985 Topoisomerase II is a structural component of mitotic chromosome scaffolds. *J. Cell Biol.* 100: 1706–1715.
- Eckert, C. A., D. J. Gravidahl, and P. C. Megee, 2007 The enhancement of pericentric cohesin association by conserved kinetochore components promotes high-fidelity chromosome segregation and is sensitive to microtubule-based tension. *Genes Dev.* 21: 278–291.
- Enervald, E., E. Lindgren, Y. Katou, K. Shirahige, and L. Ström, 2013 Importance of pol $\eta$  for damage-induced cohesion reveals differential regulation of cohesion establishment at the break site and genome-wide. *PLoS Genet.* 9: e1003158.
- Fang, G., 2002 Checkpoint protein BubR1 acts synergistically with Mad2 to inhibit anaphase-promoting complex. *Mol. Biol. Cell* 13: 755–766.
- Fang, G., H. Yu, and M. W. Kirschner, 1998 The checkpoint protein MAD2 and the mitotic regulator CDC20 form a ternary complex with the anaphase-promoting complex to control anaphase initiation. *Genes Dev.* 12: 1871–1883.
- Farcas, A.-M., P. Uluocak, W. Helmhart, and K. Nasmyth, 2011 Cohesin's concatenation of sister DNAs maintains their intertwining. *Mol. Cell* 44: 97–107.
- Fernius, J., and K. G. Hardwick, 2007 Bub1 kinase targets Sgo1 to ensure efficient chromosome biorientation in budding yeast mitosis. *PLoS Genet.* 3: e213.
- Fernius, J., and A. L. Marston, 2009 Establishment of cohesion at the pericentromere by the Ctf19 kinetochore subcomplex and the replication fork-associated factor, Csm3. *PLoS Genet.* 5: e1000629.
- Fernius, J., O. O. Nerusheva, S. Galander, F. de L. Alves, J. Rappsilber *et al.*, 2013 Cohesin-dependent association of scc2/4 with the centromere initiates pericentromeric cohesion establishment. *Curr. Biol.* 23: 599–606.
- Feytout, A., S. Vaur, S. Genier, S. Vazquez, and J.-P. Javerzat, 2011 Psm3 acetylation on conserved lysine residues is dispensable for viability in fission yeast but contributes to Eso1-mediated sister chromatid cohesion by antagonizing Wpl1. *Mol. Cell Biol.* 31: 1771–1786.
- Foster, S. A., and D. O. Morgan, 2012 The APC/C subunit Mnd2/Apc15 promotes Cdc20 autoubiquitination and spindle assembly checkpoint inactivation. *Mol. Cell* 47: 921–932.

- Freeman, L., L. Aragon-Alcaide, and A. Strunnikov, 2000 The condensin complex governs chromosome condensation and mitotic transmission of rDNA. *J. Cell Biol.* 149: 811–824.
- Funabiki, H., H. Yamano, K. Kumada, K. Nagao, T. Hunt *et al.*, 1996 Cut2 proteolysis required for sister-chromatid separation in fission yeast. *Nature* 381: 438–441.
- Gambus, A., R. C. Jones, A. Sanchez-Diaz, M. Kanemaki, F. van Deursen *et al.*, 2006 GINS maintains association of Cdc45 with MCM in replisome progression complexes at eukaryotic DNA replication forks. *Nat. Cell Biol.* 8: 358–366.
- Gambus, A., F. van Deursen, D. Polychronopoulos, M. Foltman, R. C. Jones *et al.*, 2009 A key role for Ctf4 in coupling the MCM2–7 helicase to DNA polymerase alpha within the eukaryotic replisome. *EMBO J.* 28: 2992–3004.
- Gandhi, R., P. J. Gillespie, and T. Hirano, 2006 Human Wapl is a cohesin-binding protein that promotes sister-chromatid resolution in mitotic prophase. *Curr. Biol.* 16: 2406–2417.
- Gandhi, S. R., M. Gierliński, A. Mino, K. Tanaka, E. Kitamura *et al.*, 2011 Kinetochore-dependent microtubule rescue ensures their efficient and sustained interactions in early mitosis. *Dev. Cell* 21: 920–933.
- Garcia, V., S. E. L. Phelps, S. Gray, and M. J. Neale, 2011 Bidirectional resection of DNA double-strand breaks by Mre11 and Exo1. *Nature* 479: 241–244.
- Gasser, S. M., T. Laroche, J. Falquet, E. Boy de la Tour, and U. K. Laemmli, 1986 Metaphase chromosome structure. Involvement of topoisomerase II. *J. Mol. Biol.* 188: 613–629.
- Gerlich, D., T. Hirota, B. Koch, J. M. Peters, and J. Ellenberg, 2006 Condensin I stabilizes chromosomes mechanically through a dynamic interaction in live cells. *Curr. Biol.* 16: 333–344.
- Gerring, S. L., F. Spencer, and P. Hieter, 1990 The CHL 1 (CTF 1) gene product of *Saccharomyces cerevisiae* is important for chromosome transmission and normal cell cycle progression in G2/M. *EMBO J.* 9: 4347–4358.
- Gerton, J. L., J. DeRisi, R. Shroff, M. Lichten, P. O. Brown *et al.*, 2000 Global mapping of meiotic recombination hotspots and coldspots in the yeast *Saccharomyces cerevisiae*. *Proc. Natl. Acad. Sci. USA* 97: 11383–11390.
- Gillett, E. S., C. W. Espelin, and P. K. Sorger, 2004 Spindle checkpoint proteins and chromosome-microtubule attachment in budding yeast. *J. Cell Biol.* 164: 535–546.
- Giménez-Abián, J. F., D. J. Clarke, J. Devlin, and M. I. Giménez-Abián, C. de la Torre *et al.*, 2000 Premitotic chromosome individualization in mammalian cells depends on topoisomerase II activity. *Chromosoma* 109: 235–244.
- Gladstone, M. N., D. Obeso, H. Chuong, and D. S. Dawson, 2009 The synaptonemal complex protein Zip1 promotes bi-orientation of centromeres at meiosis I. *PLoS Genet.* 5: e1000771.
- Glynn, E. F., P. C. Megee, H. G. Yu, C. Mistrot, E. Unal *et al.*, 2004 Genome-wide mapping of the cohesin complex in the yeast *Saccharomyces cerevisiae*. *PLoS Biol.* 2: E259.
- Goldstein, L. S., 1981 Kinetochore structure and its role in chromosome orientation during the first meiotic division in male *D. melanogaster*. *Cell* 25: 591–602.
- Gorr, I. H., A. Reis, D. Boos, M. Wuhr, S. Madgwick *et al.*, 2006 Essential CDK1-inhibitory role for separase during meiosis I in vertebrate oocytes. *Nat. Cell Biol.* 8: 1035–1037.
- Goshima, G., and M. Yanagida, 2000 Establishing biorientation occurs with precocious separation of the sister kinetochores, but not the arms, in the early spindle of budding yeast. *Cell* 100: 619–633.
- Grandin, N., and S. I. Reed, 1993 Differential function and expression of *Saccharomyces cerevisiae* B-type cyclins in mitosis and meiosis. *Mol. Cell Biol.* 13: 2113–2125.
- Gregan, J., C. G. Riedel, A. L. Pidoux, Y. Katou, C. Rumpf *et al.*, 2007 The kinetochore proteins pcs1 and mde4 and heterochromatin are required to prevent merotelic orientation. *Curr. Biol.* 17: 1190–1200.
- Gruber, S., C. H. Haering, and K. Nasmyth, 2003 Chromosomal cohesin forms a ring. *Cell* 112: 765–777.
- Gruber, S., P. Arumugam, Y. Katou, D. Kuglitsch, W. Helmhart *et al.*, 2006 Evidence that loading of cohesin onto chromosomes involves opening of its SMC hinge. *Cell* 127: 523–537.
- Guacci, V., E. Hogan, and D. Koshland, 1994 Chromosome condensation and sister chromatid pairing in budding yeast. *J. Cell Biol.* 125: 517–530.
- Guacci, V., D. Koshland, and A. Strunnikov, 1997 A direct link between sister chromatid cohesion and chromosome condensation revealed through the analysis of MCD1 in *S. cerevisiae*. *Cell* 91: 47–57.
- Haering, C. H., J. Lowe, A. Hochwagen, and K. Nasmyth, 2002 Molecular architecture of SMC proteins and the yeast cohesin complex. *Mol. Cell* 9: 773–788.
- Haering, C. H., D. Schöffnegger, T. Nishino, W. Helmhart, K. Nasmyth *et al.*, 2004 Structure and stability of cohesin's Smc1-kleisin interaction. *Mol. Cell* 15: 951–964.
- Haering, C. H., A.-M. Farcas, P. Arumugam, J. Metson, and K. Nasmyth, 2008 The cohesin ring concatenates sister DNA molecules. *Nature* 454: 297–301.
- Hanna, J. S., E. S. Kroll, V. Lundblad, and F. A. Spencer, 2001 *Saccharomyces cerevisiae* CTF18 and CTF4 are required for sister chromatid cohesion. *Mol. Cell Biol.* 21: 3144–3158.
- Hardwick, K. G., R. C. Johnston, D. L. Smith, and A. W. Murray, 2000 MAD3 encodes a novel component of the spindle checkpoint which interacts with Bub3p, Cdc20p, and Mad2p. *J. Cell Biol.* 148: 871–882.
- Hartman, T., K. Stead, D. Koshland, and V. Guacci, 2000 Pds5p is an essential chromosomal protein required for both sister chromatid cohesion and condensation in *Saccharomyces cerevisiae*. *J. Cell Biol.* 151: 613–626.
- He, X., S. Asthana, and P. K. Sorger, 2000 Transient sister chromatid separation and elastic deformation of chromosomes during mitosis in budding yeast. *Cell* 101: 763–775.
- Heidinger-Pauli, J. M., E. Unal, V. Guacci, and D. Koshland, 2008 The kleisin subunit of cohesin dictates damage-induced cohesion. *Mol. Cell* 31: 47–56.
- Heidinger-Pauli, J. M., E. Unal, and D. Koshland, 2009 Distinct targets of the Eco1 acetyltransferase modulate cohesion in S phase and in response to DNA damage. *Mol. Cell* 34: 311–321.
- Heinrich, S. S., H. H. Windecker, N. N. Hustedt, and S. S. Hauf, 2012 Mph1 kinetochore localization is crucial and upstream in the hierarchy of spindle assembly checkpoint protein recruitment to kinetochores. *J. Cell Sci.* 125: 4720–4727.
- Henderson, K. A., K. Kee, S. Maleki, P. A. Santini, and S. Keeney, 2006 Cyclin-dependent kinase directly regulates initiation of meiotic recombination. *Cell* 125: 1321–1332.
- Higuchi, T., and F. Uhlmann, 2005 Stabilization of microtubule dynamics at anaphase onset promotes chromosome segregation. *Nature* 433: 171–176.
- Hirano, T., 2005 Condensins: organizing and segregating the genome. *Curr. Biol.* 15: R265–R275.
- Hirano, T., 2012 Condensins: universal organizers of chromosomes with diverse functions. *Genes Dev.* 26: 1659–1678.
- Hirano, T., and T. J. Mitchison, 1994 A heterodimeric coiled-coil protein required for mitotic chromosome condensation in vitro. *Cell* 79: 449–458.
- Hiraoka, Y., and A. F. Dernburg, 2009 The SUN rises on meiotic chromosome dynamics. *Dev. Cell* 17: 598–605.
- Hochwagen, A., and A. Amon, 2006 Checking your breaks: surveillance mechanisms of meiotic recombination. *Curr. Biol.* 16: R217–R228.

- Hollingsworth, N. M., and S. J. Brill, 2004 The Mus81 solution to resolution: generating meiotic crossovers without Holliday junctions. *Genes Dev.* 18: 117–125.
- Holm, C., T. Goto, J. C. Wang, and D. Botstein, 1985 DNA topoisomerase II is required at the time of mitosis in yeast. *Cell* 41: 553–563.
- Holt, L. J., J. E. Hutt, L. C. Cantley, and D. O. Morgan, 2007 Evolution of Ime2 phosphorylation sites on Cdk1 substrates provides a mechanism to limit the effects of the phosphatase Cdc14 in meiosis. *Mol. Cell* 25: 689–702.
- Holt, L. J., A. N. Krutchinsky, and D. O. Morgan, 2008 Positive feedback sharpens the anaphase switch. *Nature* 454: 353–357.
- Hopfner, K. P., A. Karcher, D. S. Shin, L. Craig, L. M. Arthur *et al.*, 2000 Structural biology of Rad50 ATPase: ATP-driven conformational control in DNA double-strand break repair and the ABC-ATPase superfamily. *Cell* 101: 789–800.
- Hornig, N. C., P. P. Knowles, N. Q. McDonald, and F. Uhlmann, 2002 The dual mechanism of separase regulation by securin. *Curr. Biol.* 12: 973–982.
- Hoyt, M. A., L. Totis, and B. T. Roberts, 1991 *S. cerevisiae* genes required for cell cycle arrest in response to loss of microtubule function. *Cell* 66: 507–517.
- Hsu, K.-S., and T. Toda, 2011 Ndc80 internal loop interacts with Dis1/TOG to ensure proper kinetochore-spindle attachment in fission yeast. *Curr. Biol.* 21: 214–220.
- Hu, B., T. Itoh, A. Mishra, Y. Katoh, K.-L. Chan *et al.*, 2011 ATP hydrolysis is required for relocating cohesin from sites occupied by its Scc2/4 loading complex. *Curr. Biol.* 21: 12–24.
- Huang, C. E., M. Milutinovich, and D. Koshland, 2005 Rings, bracelet or snaps: fashionable alternatives for Smc complexes. *Philos. Trans. R. Soc. Lond.* 360: 537–542.
- Hudson, D. F., P. Vagnarelli, R. Gassmann, and W. C. Earnshaw, 2003 Condensin is required for nonhistone protein assembly and structural integrity of vertebrate mitotic chromosomes. *Dev. Cell* 5: 323–336.
- Hwang, L. H., L. F. Lau, D. L. Smith, C. A. Mistrot, K. G. Hardwick *et al.*, 1998 Budding yeast Cdc20: a target of the spindle checkpoint. *Science* 279: 1041–1044.
- Indjeian, V. B., and A. W. Murray, 2007 Budding yeast mitotic chromosomes have an intrinsic bias to biorient on the spindle. *Curr. Biol.* 17: 1837–1846.
- Indjeian, V. B., B. M. Stern, and A. W. Murray, 2005 The centromeric protein Sgo1 is required to sense lack of tension on mitotic chromosomes. *Science* 307: 130–133.
- Imrger, S., 2011 The Ime2 protein kinase family in fungi: more duties than just meiosis. *Mol. Microbiol.* 80: 1–13.
- Imrger, S., S. Piatti, C. Michaelis, and K. Nasmyth, 1995 Genes involved in sister chromatid separation are needed for B-type cyclin proteolysis in budding yeast. *Cell* 81: 269–278.
- Ivanov, D., and K. Nasmyth, 2005 A topological interaction between cohesin rings and a circular minichromosome. *Cell* 122: 849–860.
- Ivanov, D., and K. Nasmyth, 2007 A physical assay for sister chromatid cohesion in vitro. *Mol. Cell* 27: 300–310.
- Ivanov, D., A. Schleiffer, F. Eisenhaber, K. Mechtler, C. H. Haering *et al.*, 2002 Eco1 is a novel acetyltransferase that can acetylate proteins involved in cohesion. *Curr. Biol.* 12: 323–328.
- Izawa, D., and J. Pines, 2012 Mad2 and the APC/C compete for the same site on Cdc20 to ensure proper chromosome segregation. *J. Cell Biol.* 199: 27–37.
- Jaspersen, S. L., J. F. Charles, R. L. Tinker-Kulberg, and D. O. Morgan, 1998 A late mitotic regulatory network controlling cyclin destruction in *Saccharomyces cerevisiae*. *Mol. Biol. Cell* 9: 2803–2817.
- Johzuka, K., and T. Horiuchi, 2009 The cis element and factors required for condensin recruitment to chromosomes. *Mol. Cell* 34: 26–35.
- Kamieniecki, R. J., L. Liu, and D. S. Dawson, 2005 FEAR but not MEN genes are required for exit from meiosis I. *Cell Cycle* 4: 1093–1098.
- Katis, V. L., M. Galova, K. P. Rabitsch, J. Gregan, and K. Nasmyth, 2004a Maintenance of cohesin at centromeres after meiosis I in budding yeast requires a kinetochore-associated protein related to MEI-S332. *Curr. Biol.* 14: 560–572.
- Katis, V. L., J. Matos, S. Mori, K. Shirahige, W. Zachariae *et al.*, 2004b Spo13 facilitates monopole recruitment to kinetochores and regulates maintenance of centromeric cohesion during yeast meiosis. *Curr. Biol.* 14: 2183–2196.
- Katis, V. L., J. J. Lipp, R. Imre, A. Bogdanova, E. Okaz *et al.*, 2010 Rec8 phosphorylation by casein kinase 1 and Cdc7-Dbf4 kinase regulates cohesin cleavage by separase during meiosis. *Dev. Cell* 18: 397–409.
- Katou, Y., Y. Kanoh, M. Bando, H. Noguchi, H. Tanaka *et al.*, 2003 S-phase checkpoint proteins Tof1 and Mrc1 form a stable replication-pausing complex. *Nature* 424: 1078–1083.
- Kawashima, S. A., T. Tsukahara, M. Langeegger, S. Hauf, T. S. Kitajima *et al.*, 2007 Shugoshin enables tension-generating attachment of kinetochores by loading Aurora to centromeres. *Genes Dev.* 21: 420–435.
- Kawashima, S. A., Y. Yamagishi, T. Honda, K.-I. Ishiguro, and Y. Watanabe, 2010 Phosphorylation of H2A by Bub1 prevents chromosomal instability through localizing shugoshin. *Science* 327: 172–177.
- Keating, P., N. Rachidi, T. U. Tanaka, and M. J. R. Stark, 2009 Ipl1-dependent phosphorylation of Dam1 is reduced by tension applied on kinetochores. *J. Cell Sci.* 122: 4375–4382.
- Keeney, S., C. N. Giroux, and N. Kleckner, 1997 Meiosis-specific DNA double-strand breaks are catalyzed by Spo11, a member of a widely conserved protein family. *Cell* 88: 375–384.
- Kemmler, S., M. Stach, M. Knapp, J. Ortiz, J. Pfannstiel *et al.*, 2009 Mimicking Ndc80 phosphorylation triggers spindle assembly checkpoint signalling. *EMBO J.* 28: 1099–1110.
- Kerr, G. W., S. Sarkar, K. L. Tibbles, M. Petronczki, J. B. A. Millar *et al.*, 2011 Meiotic nuclear divisions in budding yeast require PP2A(Cdc55)-mediated antagonism of Net1 phosphorylation by Cdk. *J. Cell Biol.* 193: 1157–1166.
- Kerscher, O., L. B. Crotti, and M. A. Basrai, 2003 Recognizing chromosomes in trouble: association of the spindle checkpoint protein Bub3p with altered kinetochores and a unique defective centromere. *Mol. Cell Biol.* 23: 6406–6418.
- Khmelniskii, A., C. Lawrence, J. Roostalu, and E. Schiebel, 2007 Cdc14-regulated midzone assembly controls anaphase B. *J. Cell Biol.* 177: 981–993.
- Khmelniskii, A., J. Roostalu, H. Roque, C. Antony, and E. Schiebel, 2009 Phosphorylation-dependent protein interactions at the spindle midzone mediate cell cycle regulation of spindle elongation. *Dev. Cell* 17: 244–256.
- Kiburz, B. M., D. B. Reynolds, P. C. Megee, A. L. Marston, B. H. Lee *et al.*, 2005 The core centromere and Sgo1 establish a 50-kb cohesin-protected domain around centromeres during meiosis I. *Genes Dev.* 19: 3017–3030.
- Kiburz, B. M., A. Amon, and A. L. Marston, 2008 Shugoshin Promotes Sister Kinetochore Biorientation in *Saccharomyces cerevisiae*. *Mol. Biol. Cell* 19: 1199–1209.
- Kim, K. P., B. M. Weiner, L. Zhang, A. Jordan, J. Dekker *et al.*, 2010 Sister cohesion and structural axis components mediate homolog bias of meiotic recombination. *Cell* 143: 924–937.
- Kimura, K., V. V. Rybenkov, N. J. Crisona, T. Hirano, and N. R. Cozzarelli, 1999 13S condensin actively reconfigures DNA by introducing global positive writhe: implications for chromosome condensation. *Cell* 98: 239–248.
- King, E. M., N. Rachidi, N. Morrice, K. G. Hardwick, and M. J. Stark, 2007a Ipl1p-dependent phosphorylation of Mad3p is required

- for the spindle checkpoint response to lack of tension at kinetochores. *Genes Dev.* 21: 1163–1168.
- King, E. M., S. J. van der Sar, and K. G. Hardwick, 2007b Mad3 KEN Boxes mediate both Cdc20 and Mad3 turnover, and are critical for the spindle checkpoint. *PLoS ONE* 2: e342.
- Kitajima, T. S., S. A. Kawashima, and Y. Watanabe, 2004 The conserved kinetochore protein shugoshin protects centromeric cohesion during meiosis. *Nature* 427: 510–517.
- Kitajima, T. S., T. Sakuno, K. Ishiguro, S. Iemura, T. Natsume *et al.*, 2006 Shugoshin collaborates with protein phosphatase 2A to protect cohesin. *Nature* 441: 46–52.
- Kitamura, E., K. Tanaka, Y. Kitamura, and T. U. Tanaka, 2007 Kinetochore microtubule interaction during S phase in *Saccharomyces cerevisiae*. *Genes Dev.* 21: 3319–3330.
- Kitamura, E., K. Tanaka, S. Komoto, Y. Kitamura, C. Antony *et al.*, 2010 Kinetochores generate microtubules with distal plus ends: their roles and limited lifetime in mitosis. *Dev. Cell* 18: 248–259.
- Kleckner, N., D. Zickler, G. H. Jones, J. Dekker, R. Padmore *et al.*, 2004 A mechanical basis for chromosome function. *Proc. Natl. Acad. Sci. USA* 101: 12592–12597.
- Klein, F., P. Mahr, M. Galova, S. B. Buonomo, C. Michaelis *et al.*, 1999 A central role for cohesins in sister chromatid cohesion, formation of axial elements, and recombination during yeast meiosis. *Cell* 98: 91–103.
- Kogut, I., J. Wang, V. Guacci, R. K. Mistry, and P. C. Megee, 2009 The Scc2/Scc4 cohesin loader determines the distribution of cohesin on budding yeast chromosomes. *Genes Dev.* 23: 2345–2357.
- Koshland, D., and L. H. Hartwell, 1987 The structure of sister minichromosome DNA before anaphase in *Saccharomyces cerevisiae*. *Science* 238: 1713–1716.
- Kozsul, R., K. P. Kim, M. Prentiss, N. Kleckner, and S. Kameoka, 2008 Meiotic chromosomes move by linkage to dynamic actin cables with transduction of force through the nuclear envelope. *Cell* 133: 1188–1201.
- Kueng, S., B. Hegemann, B. H. Peters, J. J. Lipp, A. Schleiffer *et al.*, 2006 Wapl controls the dynamic association of cohesin with chromatin. *Cell* 127: 955–967.
- Kulemzina, I., M. R. Schumacher, V. Verma, J. Reiter, J. Metzler *et al.*, 2012 Cohesin rings devoid of Scc3 and Pds5 maintain their stable association with the DNA. *PLoS Genet.* 8: e1002856.
- Kulukian, A., J. S. Han, and D. W. Cleveland, 2009 Unattached kinetochores catalyze production of an anaphase inhibitor that requires a Mad2 template to prime Cdc20 for BubR1 binding. *Dev. Cell* 16: 105–117.
- Kurze, A., K. A. Michie, S. E. Dixon, A. Mishra, T. Itoh *et al.*, 2011 A positively charged channel within the Smc1/Smc3 hinge required for sister chromatid cohesion. *EMBO J.* 30: 364–378.
- Lacefield, S., and A. W. Murray, 2007 The spindle checkpoint rescues the meiotic segregation of chromosomes whose crossovers are far from the centromere. *Nat. Genet.* 39: 1273–1277.
- Laloraya, S., V. Guacci, and D. Koshland, 2000 Chromosomal addresses of the cohesin component Mcd1p. *J. Cell Biol.* 151: 1047–1056.
- Lara-Gonzalez, P., F. G. Westhorpe, and S. S. Taylor, 2012 The spindle assembly checkpoint. *Curr. Biol.* 22: R966–R980.
- Larsen, N. A., J. Al-Bassam, R. R. Wei, and S. C. Harrison, 2007 Structural analysis of Bub3 interactions in the mitotic spindle checkpoint. *Proc. Natl. Acad. Sci. USA* 104: 1201–1206.
- Lau, D. T. C., and A. W. Murray, 2012 Mad2 and Mad3 cooperate to arrest budding yeast in mitosis. *Curr. Biol.* 22: 180–190.
- Lavoie, B. D., E. Hogan, and D. Koshland, 2004 In vivo requirements for rDNA chromosome condensation reveal two cell-cycle-regulated pathways for mitotic chromosome folding. *Genes Dev.* 18: 76–87.
- Lee, B. H., and A. Amon, 2003 Role of Polo-like kinase CDC5 in programming meiosis I chromosome segregation. *Science* 300: 482–486.
- Lee, B. H., B. M. Kiburz, and A. Amon, 2004 Spo13 maintains centromeric cohesion and kinetochore coorientation during meiosis I. *Curr. Biol.* 14: 2168–2182.
- Lee, C.-Y., M. N. Conrad, and M. E. Dresser, 2012 Meiotic chromosome pairing is promoted by telomere-led chromosome movements independent of bouquet formation. *PLoS Genet.* 8: e1002730.
- Lengronne, A., Y. Katou, S. Mori, S. Yokobayashi, G. P. Kelly *et al.*, 2004 Cohesin relocation from sites of chromosomal loading to places of convergent transcription. *Nature* 430: 573–578.
- Lengronne, A., J. McIntyre, Y. Katou, Y. Kanoh, K. P. Hopfner *et al.*, 2006 Establishment of sister chromatid cohesion at the *S. cerevisiae* replication fork. *Mol. Cell* 23: 787–799.
- Li, R., and A. W. Murray, 1991 Feedback control of mitosis in budding yeast. *Cell* 66: 519–531.
- Liu, D., G. Vader, M. J. M. Vromans, M. A. Lampson, and S. M. A. Lens, 2009 Sensing chromosome bi-orientation by spatial separation of aurora B kinase from kinetochore substrates. *Science* 323: 1350–1353.
- Liu, X., and M. Winey, 2012 The MPS1 family of protein kinases. *Annu. Rev. Biochem.* 81: 561–585.
- London, N., S. Ceto, J. A. Ranish, and S. Biggins, 2012 Phosphoregulation of Spc105 by Mps1 and Pp1 regulates Bub1 localization to kinetochores. *Curr. Biol.* 22: 900–906.
- Lopez-Mosqueda, J., N. L. Maas, Z. O. Jonsson, L. G. Defazio-Eli, J. Wohlschlegel *et al.*, 2010 Damage-induced phosphorylation of Sld3 is important to block late origin firing. *Nature* 467: 479–483.
- Lopez-Serra, L., A. Lengronne, V. Borges, G. Kelly, and F. Uhlmann, 2013 Budding yeast wapl controls sister chromatid cohesion maintenance and chromosome condensation. *Curr. Biol.* 23: 64–69.
- Losada, A., M. Hirano, and T. Hirano, 1998 Identification of Xenopus SMC protein complexes required for sister chromatid cohesion. *Genes Dev.* 12: 1986–1997.
- Lowe, J., S. C. Cordell, and F. van den Ent, 2001 Crystal structure of the SMC head domain: an ABC ATPase with 900 residues antiparallel coiled-coil inserted. *J. Mol. Biol.* 306: 25–35.
- Lynn, A., R. Soucek, and G. V. Börner, 2007 ZMM proteins during meiosis: crossover artists at work. *Chromosome Res.* 15: 591–605.
- Lyons, N. A., and D. O. Morgan, 2011 Cdk1-dependent destruction of Eco1 prevents cohesion establishment after S phase. *Mol. Cell* 42: 378–389.
- Lyons, N. A., B. R. Fonslow, J. K. Diedrich, J. R. Yates, and D. O. Morgan, 2013 Sequential primed kinases create a damage-responsive phosphodegron on Eco1. *Nat. Struct. Mol. Biol.* 20: 194–201.
- Machín, F., J. Torres-Rosell, G. de Piccoli, J. A. Carballo, R. S. Cha *et al.*, 2006 Transcription of ribosomal genes can cause nondisjunction. *J. Cell Biol.* 173: 893–903.
- MacQueen, A. J., and A. Hochwagen, 2011 Checkpoint mechanisms: the puppet masters of meiotic prophase. *Trends Cell Biol.* 21: 393–400.
- Malureanu, L. A., K. B. Jeganathan, M. Hamada, L. Wasilewski, J. Davenport *et al.*, 2009 BubR1 N terminus acts as a soluble inhibitor of cyclin B degradation by APC/C(Cdc20) in interphase. *Dev. Cell* 16: 118–131.
- Mansfeld, J., P. Collin, M. O. Collins, J. S. Choudhary, and J. Pines, 2011 APC15 drives the turnover of MCC-CDC20 to make the spindle assembly checkpoint responsive to kinetochore attachment. *Nat. Cell Biol.* 13: 1234–1243.
- Manzoni, R., F. Montani, C. Visintin, F. Caudron, A. Ciliberto *et al.*, 2010 Oscillations in Cdc14 release and sequestration reveal a circuit underlying mitotic exit. *J. Cell Biol.* 190: 209–222.

- Mariani, L., E. Chirolì, L. Nezi, H. Müller, S. Piatti *et al.*, 2012 Role of the Mad2 dimerization interface in the spindle assembly checkpoint independent of kinetochores. *Curr. Biol.* 22: 1900–1908.
- Marston, A. L., and A. Amon, 2004 Meiosis: cell-cycle controls shuffle and deal. *Nat. Rev. Mol. Cell Biol.* 5: 983–997.
- Marston, A. L., B. H. Lee, and A. Amon, 2003 The Cdc14 phosphatase and the FEAR network control meiotic spindle disassembly and chromosome segregation. *Dev. Cell* 4: 711–726.
- Marston, A. L., W. H. Tham, H. Shah, and A. Amon, 2004 A genome-wide screen identifies genes required for centromeric cohesion. *Science* 303: 1367–1370.
- Martini, E., R. L. Diaz, N. Hunter, and S. Keeney, 2006 Crossover homeostasis in yeast meiosis. *Cell* 126: 285–295.
- Matos, J., J. J. Lipp, A. Bogdanova, S. Guillot, E. Okaz *et al.*, 2008 Dbf4-dependent cdc7 kinase links DNA replication to the segregation of homologous chromosomes in meiosis I. *Cell* 135: 662–678.
- Maure, J.-F., E. Kitamura, and T. U. Tanaka, 2007 Mps1 kinase promotes sister-kinetochore bi-orientation by a tension-dependent mechanism. *Curr. Biol.* 17: 2175–2182.
- Maure, J.-F., S. Komoto, Y. Oku, A. Mino, S. Pasqualato *et al.*, 2011 The Ndc80 loop region facilitates formation of kinetochore attachment to the dynamic microtubule plus end. *Curr. Biol.* 21: 207–213.
- Mayer, M. L., S. P. Gygi, R. Aebersold, and P. Hieter, 2001 Identification of RFC(Ctf18p, Ctf8p, Dcc1p): an alternative RFC complex required for sister chromatid cohesion in *S. cerevisiae*. *Mol. Cell* 7: 959–970.
- Mayer, M. L., I. Pot, M. Chang, H. Xu, V. Aneliunas *et al.*, 2004 Identification of protein complexes required for efficient sister chromatid cohesion. *Mol. Biol. Cell* 15: 1736–1745.
- McIntyre, J., E. G. D. Müller, S. Weitzer, B. E. Snijdsman, T. N. Davis *et al.*, 2007 In vivo analysis of cohesin architecture using FRET in the budding yeast *Saccharomyces cerevisiae*. *EMBO J.* 26: 3783–3793.
- McAleenan, A., A. Clemente-Blanco, V. Cordon-Preciado, N. Sen, M. Esteras *et al.*, 2013 Post-replicative repair involves separate-dependent removal of the kleisin subunit of cohesin. *Nature* 493: 250–254.
- McAleenan, A., V. Cordon-Preciado, A. Clemente-Blanco, I.-C. Liu, N. Sen *et al.*, 2012 SUMOylation of the  $\alpha$ -kleisin subunit of cohesin is required for DNA damage-induced cohesion. *Curr. Biol.* 22: 1564–1575.
- Megee, P. C., and D. Koshland, 1999 A functional assay for centromere-associated sister chromatid cohesion. *Science* 285: 254–257.
- Megee, P. C., C. Mistrot, V. Guacci, and D. Koshland, 1999 The centromeric sister chromatid cohesion site directs Mcd1p binding to adjacent sequences. *Mol. Cell* 4: 445–450.
- Melby, T. E., C. N. Ciampaglio, G. Briscoe, and H. P. Erickson, 1998 The symmetrical structure of structural maintenance of chromosomes (SMC) and MukB proteins: long, antiparallel coiled coils, folded at a flexible hinge. *J. Cell Biol.* 142: 1595–1604.
- Mendoza, M., C. Norden, K. Durrer, H. Rauter, F. Uhlmann *et al.*, 2009 A mechanism for chromosome segregation sensing by the NoCut checkpoint. *Nat. Cell Biol.* 11: 477–483.
- Meyer, R. E., S. Kim, D. Obeso, P. D. Straight, M. Winey *et al.*, 2013 Mps1 and Ipl1/Aurora B act sequentially to correctly orient chromosomes on the meiotic spindle of budding yeast. *Science* 339: 1071–1074.
- Michaelis, C., R. Ciosk, and K. Nasmyth, 1997 Cohesins: chromosomal proteins that prevent premature separation of sister chromatids. *Cell* 91: 35–45.
- Miller, M. P., E. Unal, G. A. Brar, and A. Amon, 2012 Meiosis I chromosome segregation is established through regulation of microtubule-kinetochore interactions. *Elife* 1: e00117.
- Milutinovich, M., and D. E. Koshland, 2003 Molecular biology. SMC complexes—wrapped up in controversy. *Science* 300: 1101–1102.
- Milutinovich, M., E. Unal, C. Ward, R. V. Skibbens, and D. Koshland, 2007 A multi-step pathway for the establishment of sister chromatid cohesion. *PLoS Genet.* 3: e12.
- Mimitou, E. P., and L. S. Symington, 2008 Sae2, Exo1 and Sgs1 collaborate in DNA double-strand break processing. *Nature* 455: 770–774.
- Mirchenko, L., and F. Uhlmann, 2010 Sli15(INCENP) dephosphorylation prevents mitotic checkpoint reengagement due to loss of tension at anaphase onset. *Curr. Biol.* 20: 1396–1401.
- Mohl, D. A., M. J. Huddleston, T. S. Collingwood, R. S. Annan, and R. J. Deshaies, 2009 Dbf2-Mob1 drives relocalization of protein phosphatase Cdc14 to the cytoplasm during exit from mitosis. *J. Cell Biol.* 184: 527–539.
- Moldovan, G. L., B. Pfander, and S. Jentsch, 2006 PCNA controls establishment of sister chromatid cohesion during S phase. *Mol. Cell* 23: 723–732.
- Monje-Casas, F., V. R. Prabhu, B. H. Lee, M. Boselli, and A. Amon, 2007 Kinetochore orientation during meiosis is controlled by Aurora B and the monopolin complex. *Cell* 128: 477–490.
- Montpetit, B., T. R. Hazbun, S. Fields, and P. Hieter, 2006 Sumoylation of the budding yeast kinetochore protein Ndc10 is required for Ndc10 spindle localization and regulation of anaphase spindle elongation. *J. Cell Biol.* 174: 653–663.
- Morgan, D. O., 2007 *The Cell Cycle: Principles of Control*. New Science Press, London.
- Murray, A. W., and J. W. Szostak, 1985 Chromosome segregation in mitosis and meiosis. *Annu. Rev. Cell Biol.* 1: 289–315.
- Musacchio, A., and E. D. Salmon, 2007 The spindle-assembly checkpoint in space and time. *Nat. Rev. Mol. Cell Biol.* 8: 379–393.
- Musacchio, A., and A. Ciliberto, 2012 The spindle-assembly checkpoint and the beauty of self-destruction. *Nat. Struct. Mol. Biol.* 19: 1059–1061.
- Nakajima, Y., A. Cormier, R. G. Tyers, A. Pigula, Y. Peng *et al.*, 2011 Ipl1/Aurora-dependent phosphorylation of Sli15/INCENP regulates CPC-spindle interaction to ensure proper microtubule dynamics. *J. Cell Biol.* 194: 137–153.
- Nasmyth, K., and C. H. Haering, 2005 The structure and function of SMC and kleisin complexes. *Annu. Rev. Biochem.* 74: 595–648.
- Natsume, T., C. A. Müller, Y. Katou, and R. Retkute, M. Gierlinski *et al.*, 2013 Kinetochores coordinate pericentromeric cohesion and early DNA replication by Cdc7-Dbf4 kinase recruitment. *Mol. Cell* 50: 661–674.
- Newnham, L., P. Jordan, B. Rockmill, G. S. Roeder, and E. Hoffmann, 2010 The synaptonemal complex protein, Zip1, promotes the segregation of nonexchange chromosomes at meiosis I. *Proc. Natl. Acad. Sci. USA* 107: 781–785.
- Ng, T. M., W. G. Waples, B. D. Lavoie, and S. Biggins, 2009 Pericentromeric sister chromatid cohesion promotes kinetochore biorientation. *Mol. Biol. Cell* 20: 3818–3827.
- Nicolette, M. L., K. Lee, Z. Guo, M. Rani, J. M. Chow *et al.*, 2010 Mre11-Rad50-Xrs2 and Sae2 promote 5' strand resection of DNA double-strand breaks. *Nat. Struct. Mol. Biol.* 17: 1478–1485.
- Norden, C., M. Mendoza, J. Dobbelaere, C. V. Kotwaliwale, S. Biggins *et al.*, 2006 The NoCut pathway links completion of cytokinesis to spindle midzone function to prevent chromosome breakage. *Cell* 125: 85–98.
- Ocampo-Hafalla, M. T., Y. Katou, K. Shirahige, and F. Uhlmann, 2007 Displacement and re-accumulation of centromeric cohesin during transient pre-anaphase centromere splitting. *Chromosoma* 116: 531–544.
- Oelschlaegel, T., M. Schwickart, J. Matos, A. Bogdanova, A. Camasses *et al.*, 2005 The yeast APC/C subunit Mnd2 prevents premature

- sister chromatid separation triggered by the meiosis-specific APC/C-Ama1. *Cell* 120: 773–788.
- Okaz, E., O. Argüello-Miranda, A. Bogdanova, P. K. Vinod, J. J. Lipp *et al.*, 2012 Meiotic Prophase Requires Proteolysis of M Phase Regulators Mediated by the Meiosis-Specific APC/C(Ama1). *Cell* 151: 603–618.
- Oliveira, R. A., P. A. Coelho, and C. E. Sunkel, 2005 The condensin I subunit Barren/CAP-H is essential for the structural integrity of centromeric heterochromatin during mitosis. *Mol. Cell Biol.* 25: 8971–8984.
- Ono, T., Y. Fang, D. L. Spector, and T. Hirano, 2004 Spatial and temporal regulation of Condensins I and II in mitotic chromosome assembly in human cells. *Mol. Biol. Cell* 15: 3296–3308.
- Palframan, W. J., J. B. Meehl, S. L. Jaspersen, M. Winey, and A. W. Murray, 2006 Anaphase inactivation of the spindle checkpoint. *Science* 313: 680–684.
- Paliulis, L. V., and R. B. Nicklas, 2000 The reduction of chromosome number in meiosis is determined by properties built into the chromosomes. *J. Cell Biol.* 150: 1223–1232.
- Pan, J., M. Sasaki, R. Kniewel, H. Murakami, H. G. Blitzblau *et al.*, 2011 A hierarchical combination of factors shapes the genome-wide topography of yeast meiotic recombination initiation. *Cell* 144: 719–731.
- Panizza, S., T. Tanaka, A. Hochwagen, F. Eisenhaber, and K. Nasmyth, 2000 Pds5 cooperates with cohesin in maintaining sister chromatid cohesion. *Curr. Biol.* 10: 1557–1564.
- Panizza, S., M. A. Mendoza, M. Berlinger, L. Huang, A. Nicolas *et al.*, 2011 Spo11-accessory proteins link double-strand break sites to the chromosome axis in early meiotic recombination. *Cell* 146: 372–383.
- Penkner, A. M., S. Prinz, S. Ferscha, and F. Klein, 2005 Mnd2, an essential antagonist of the anaphase-promoting complex during meiotic prophase. *Cell* 120: 789–801.
- Pereira, G., and E. Schiebel, 2003 Separase regulates INCENP-Aurora B anaphase spindle function through Cdc14. *Science* 302: 2120–2124.
- Peters, J. M., 2006 The anaphase promoting complex/cyclosome: a machine designed to destroy. *Nat. Rev. Mol. Cell Biol.* 7: 644–656.
- Petronczki, M., B. Chwalla, M. F. Siomos, S. Yokobayashi, W. Helmhart *et al.*, 2004 Sister-chromatid cohesion mediated by the alternative RF-Cctf18/Dcc1/Ctf8, the helicase Chl1 and the polymerase-alpha-associated protein Ctf4 is essential for chromatid disjunction during meiosis II. *J. Cell Sci.* 117: 3547–3559.
- Petronczki, M., J. Matos, S. Mori, J. Gregan, A. Bogdanova *et al.*, 2006 Monopolar attachment of sister kinetochores at meiosis I requires casein kinase 1. *Cell* 126: 1049–1064.
- Pinsky, B. A., C. Kung, K. M. Shokat, and S. Biggins, 2006 The Ipl1-Aurora protein kinase activates the spindle checkpoint by creating unattached kinetochores. *Nat. Cell Biol.* 8: 78–83.
- Pinsky, B. A., C. R. Nelson, and S. Biggins, 2009 Protein phosphatase 1 regulates exit from the spindle checkpoint in budding yeast. *Curr. Biol.* 19: 1182–1187.
- Poddar, A., P. T. Stukenberg, and D. J. Burke, 2005 Two complexes of spindle checkpoint proteins containing Cdc20 and Mad2 assemble during mitosis independently of the kinetochore in *Saccharomyces cerevisiae*. *Eukaryot. Cell* 4: 867–878.
- Queralt, E., and F. Uhlmann, 2008 Separase cooperates with Zds1 and Zds2 to activate Cdc14 phosphatase in early anaphase. *J. Cell Biol.* 182: 873–883.
- Queralt, E., C. Lehane, B. Novak, and F. Uhlmann, 2006 Downregulation of PP2A(Cdc55) phosphatase by separase initiates mitotic exit in budding yeast. *Cell* 125: 719–732.
- Quevedo, O., J. Garcia-Luis, E. Matos-Perdomo, L. Aragón, and F. Machín, 2012 Nondisjunction of a single chromosome leads to breakage and activation of DNA damage checkpoint in G2. *PLoS Genet.* 8: e1002509.
- Rabitsch, K. P., M. Petronczki, J. P. Javerzat, S. Genier, B. Chwalla *et al.*, 2003 Kinetochore recruitment of two nucleolar proteins is required for homolog segregation in meiosis I. *Dev. Cell* 4: 535–548.
- Renshaw, M. J., J. J. Ward, M. Kanemaki, K. Natsume, F. J. Nédélec *et al.*, 2010 Condensins promote chromosome recoiling during early anaphase to complete sister chromatid separation. *Dev. Cell* 19: 232–244.
- Riedel, C. G., V. L. Katis, Y. Katou, S. Mori, T. Itoh *et al.*, 2006 Protein phosphatase 2A protects centromeric sister chromatid cohesion during meiosis I. *Nature* 441: 53–61.
- Rock, J. M., and A. Amon, 2009 The FEAR network. *Curr. Biol.* 19: R1063–R1068.
- Ross, K. E., and O. Cohen-Fix, 2004 A role for the FEAR pathway in nuclear positioning during anaphase. *Dev. Cell* 6: 729–735.
- Rossio, V., and S. Yoshida, 2011 Spatial regulation of Cdc55–PP2A by Zds1/Zds2 controls mitotic entry and mitotic exit in budding yeast. *J. Cell Biol.* 193: 445–454.
- Rowland, B. D., M. B. Roig, T. Nishino, A. Kurze, P. Uluocak *et al.*, 2009 Building sister chromatid cohesion: smc3 acetylation counteracts an antiestablishment activity. *Mol. Cell* 33: 763–774.
- Rozelle, D. K., S. D. Hansen, and K. B. Kaplan, 2011 Chromosome passenger complexes control anaphase duration and spindle elongation via a kinesin-5 brake. *J. Cell Biol.* 193: 285–294.
- Saka, Y., T. Sutani, Y. Yamashita, S. Saitoh, M. Takeuchi *et al.*, 1994 Fission yeast cut3 and cut14, members of a ubiquitous protein family, are required for chromosome condensation and segregation in mitosis. *EMBO J.* 13: 4938–4952.
- Sanchez, Y., J. Bachant, H. Wang, F. Hu, D. Liu *et al.*, 1999 Control of the DNA damage checkpoint by chk1 and rad53 protein kinases through distinct mechanisms. *Science* 286: 1166–1171.
- Scherthan, H., 2001 A bouquet makes ends meet. *Nat. Rev. Mol. Cell Biol.* 2: 621–627.
- Szczaniecka, M., A. Feoktistova, K. M. May, J.-S. Chen, J. Blyth *et al.*, 2008 The spindle checkpoint functions of Mad3 and Mad2 depend on a Mad3 KEN box-mediated interaction with Cdc20-anaphase-promoting complex (APC/C). *J. Biol. Chem.* 283: 23039–23047.
- Serrentino, M.-E., and V. Borde, 2012 The spatial regulation of meiotic recombination hotspots: are all DSB hotspots crossover hotspots? *Exp. Cell Res.* 318: 1347–1352.
- Shepherd, L. A., J. C. Meadows, A. M. Sochaj, T. C. Lancaster, J. Zou *et al.*, 2012 Phosphodependent recruitment of Bub1 and Bub3 to Spc7/KNL1 by Mph1 kinase maintains the spindle checkpoint. *Curr. Biol.* 22: 891–899.
- Shimogawa, M. M., B. Graczyk, M. K. Gardner, S. E. Francis, and E. A. White *et al.*, 2006 Mps1 phosphorylation of Dam1 couples kinetochores to microtubule plus ends at metaphase. *Curr. Biol.* 16: 1489–1501.
- Shinohara, A., H. Ogawa, and T. Ogawa, 1992 Rad51 protein involved in repair and recombination in *S. cerevisiae* is a RecA-like protein. *Cell* 69: 457–470.
- Shonn, M. A., R. McCarroll, and A. W. Murray, 2000 Requirement of the spindle checkpoint for proper chromosome segregation in budding yeast meiosis. *Science* 289: 300–303.
- Shonn, M. A., A. L. Murray, and A. W. Murray, 2003 Spindle checkpoint component Mad2 contributes to biorientation of homologous chromosomes. *Curr. Biol.* 13: 1979–1984.
- Shou, W., J. H. Seol, A. Shevchenko, C. Baskerville, D. Moazed *et al.*, 1999 Exit from mitosis is triggered by Tem1-dependent release of the protein phosphatase Cdc14 from nucleolar RENT complex. *Cell* 97: 233–244.
- Sjogren, C., and K. Nasmyth, 2001 Sister chromatid cohesion is required for postreplicative double-strand break repair in *Saccharomyces cerevisiae*. *Curr. Biol.* 11: 991–995.

- Skibbens, R. V., 2004 Chl1p, a DNA helicase-like protein in budding yeast, functions in sister-chromatid cohesion. *Genetics* 166: 33–42.
- Skibbens, R. V., L. B. Corson, D. Koshland, and P. Hieter, 1999 Ctf7p is essential for sister chromatid cohesion and links mitotic chromosome structure to the DNA replication machinery. *Genes Dev.* 13: 307–319.
- Sourirajan, A., and M. Lichten, 2008 Polo-like kinase Cdc5 drives exit from pachytene during budding yeast meiosis. *Genes Dev.* 22: 2627–2632.
- St-Pierre, J., M. Douziech, F. Bazile, M. Pascariu, E. Bonneil *et al.*, 2009 Polo kinase regulates mitotic chromosome condensation by hyperactivation of condensin DNA supercoiling activity. *Mol. Cell* 34: 416–426.
- Stegmeier, F., and A. Amon, 2004 Closing mitosis: the functions of the Cdc14 phosphatase and its regulation. *Annu. Rev. Genet.* 38: 203–232.
- Stegmeier, F., J. Huang, R. Rahal, J. Zmolik, D. Moazed *et al.*, 2004 The replication fork block protein Fob1 functions as a negative regulator of the FEAR network. *Curr. Biol.* 14: 467–480.
- Stemmann, O., H. Zou, S. A. Gerber, S. P. Gygi, and M. W. Kirschner, 2001 Dual inhibition of sister chromatid separation at metaphase. *Cell* 107: 715–726.
- Stephens, A. D., J. Haase, L. Vicci, R. M. Taylor, and K. Bloom, 2011 Cohesin, condensin, and the intramolecular centromere loop together generate the mitotic chromatin spring. *J. Cell Biol.* 193: 1167–1180.
- Stephens, A. D., R. A. Haggerty, P. A. Vasquez, L. Vicci, C. E. Snider *et al.*, 2013 Pericentric chromatin loops function as a nonlinear spring in mitotic force balance. *J. Cell Biol.* 200: 757–772.
- Storchová, Z., J. S. Becker, N. Talarek, S. Kögelsberger, and D. Pellman, 2011 Bub1, Sgo1, and Mps1 mediate a distinct pathway for chromosome biorientation in budding yeast. *Mol. Biol. Cell* 22: 1473–1485.
- Straight, A. F., A. S. Belmont, C. C. Robinett, and A. W. Murray, 1996 GFP tagging of budding yeast chromosomes reveals that protein-protein interactions can mediate sister chromatid cohesion. *Curr. Biol.* 6: 1599–1608.
- Strom, L., H. B. Lindroos, K. Shirahige, and C. Sjogren, 2004 Postreplicative recruitment of cohesin to double-strand breaks is required for DNA repair. *Mol. Cell* 16: 1003–1015.
- Strom, L., C. Karlsson, H. B. Lindroos, S. Wedahl, Y. Katou *et al.*, 2007 Postreplicative formation of cohesion is required for repair and induced by a single DNA break. *Science* 317: 242–245.
- Strunnikov, A. V., E. Hogan, and D. Koshland, 1995 SMC2, a *Saccharomyces cerevisiae* gene essential for chromosome segregation and condensation, defines a subgroup within the SMC family. *Genes Dev.* 9: 587–599.
- Stuart, D., and C. Wittenberg, 1998 CLB5 and CLB6 are required for premeiotic DNA replication and activation of the meiotic S/M checkpoint. *Genes Dev.* 12: 2698–2710.
- Subramanian, V. V., and A. Hochwagen, 2011 Centromere clustering: where synapsis begins. *Curr. Biol.* 21: R920–R922.
- Suja, J. A., J. de la Torre, J. F. Giménez-Abián, C. García de la Vega, and J. S. Rufas, 1991 Meiotic chromosome structure. Kinetochores and chromatid cores in standard and B chromosomes of *Arcyptera fusca* (Orthoptera) revealed by silver staining. *Genome* 34: 19–27.
- Sullivan, M., and F. Uhlmann, 2003 A non-proteolytic function of separase links the onset of anaphase to mitotic exit. *Nat. Cell Biol.* 5: 249–254.
- Sullivan, M., T. Higuchi, V. L. Katis, and F. Uhlmann, 2004 Cdc14 phosphatase induces rDNA condensation and resolves cohesin-independent cohesion during budding yeast anaphase. *Cell* 117: 471–482.
- Surana, U., A. Amon, C. Dowzer, J. McGrew, B. Byers *et al.*, 1993 Destruction of the CDC28/CLB mitotic kinase is not required for the metaphase to anaphase transition in budding yeast. *EMBO J.* 12: 1969–1978.
- Sutani, T., T. Kawaguchi, R. Kanno, T. Itoh, and K. Shirahige, 2009 Budding yeast Wpl1(Rad61)-Pds5 complex counteracts sister chromatid cohesion-establishing reaction. *Curr. Biol.* 19: 492–497.
- Tada, K., H. Susumu, T. Sakuno, and Y. Watanabe, 2011 Condensin association with histone H2A shapes mitotic chromosomes. *Nature* 474: 477–483.
- Tanaka, K., Z. Hao, M. Kai, and H. Okayama, 2001 Establishment and maintenance of sister chromatid cohesion in fission yeast by a unique mechanism. *EMBO J.* 20: 5779–5790.
- Tanaka, K., N. Mukae, H. Dewar, M. van Breugel, E. K. James *et al.*, 2005 Molecular mechanisms of kinetochore capture by spindle microtubules. *Nature* 434: 987–994.
- Tanaka, K., E. Kitamura, Y. Kitamura, and T. U. Tanaka, 2007 Molecular mechanisms of microtubule-dependent kinetochore transport toward spindle poles. *J. Cell Biol.* 178: 269–281.
- Tanaka, T. U., 2010 Kinetochore-microtubule interactions: steps towards bi-orientation. *EMBO J.* 29: 4070–4082.
- Tanaka, T., M. P. Cosma, K. Wirth, and K. Nasmyth, 1999 Identification of cohesin association sites at centromeres and along chromosome arms. *Cell* 98: 847–858.
- Tanaka, T., J. Fuchs, J. Loidl, and K. Nasmyth, 2000 Cohesin ensures bipolar attachment of microtubules to sister centromeres and resists their precocious separation. *Nat. Cell Biol.* 2: 492–499.
- Tanaka, T. U., N. Rachidi, C. Janke, G. Pereira, M. Galova *et al.*, 2002 Evidence that the Ipl1-Sli15 (Aurora kinase-INCENP) complex promotes chromosome bi-orientation by altering kinetochore-spindle pole connections. *Cell* 108: 317–329.
- Tang, X., and Y. Wang, 2006 Pds1/Esp1-dependent and -independent sister chromatid separation in mutants defective for protein phosphatase 2A. *Proc. Natl. Acad. Sci. USA* 103: 16290–16295.
- Tang, Z., R. Bharadwaj, B. Li, and H. Yu, 2001 Mad2-Independent inhibition of APC<sup>Cdc20</sup> by the mitotic checkpoint protein BubR1. *Dev. Cell* 1: 227–237.
- Terret, M.-E., R. Sherwood, S. Rahman, J. Qin, and P. V. Jallepalli, 2009 Cohesin acetylation speeds the replication fork. *Nature* 462: 231–234.
- Tessé, S., A. Storlazzi, N. Kleckner, S. Gargano, and D. Zickler, 2003 Localization and roles of Ski8p protein in *Sordaria* meiosis and delineation of three mechanistically distinct steps of meiotic homolog juxtaposition. *Proc. Natl. Acad. Sci. USA* 100: 12865–12870.
- Thadani, R., F. Uhlmann, and S. Heeger, 2012 Condensin, chromatin crossbarring and chromosome condensation. *Curr. Biol.* 22: R1012–R1021.
- Thomas, S., and K. B. Kaplan, 2007 A Bir1p Sli15p kinetochore passenger complex regulates septin organization during anaphase. *Mol. Biol. Cell* 18: 3820–3834.
- Tomson, B. N., D. D'Amours, B. S. Adamson, L. Aragon, and A. Amon, 2006 Ribosomal DNA transcription-dependent processes interfere with chromosome segregation. *Mol. Cell Biol.* 26: 6239–6247.
- Tomson, B. N., R. Rahal, V. Reiser, F. Monje-Casas, K. Mekhail *et al.*, 2009 Regulation of Spo12 phosphorylation and its essential role in the FEAR network. *Curr. Biol.* 19: 449–460.
- Toth, A., R. Ciosk, F. Uhlmann, M. Galova, A. Schleiffer *et al.*, 1999 Yeast cohesin complex requires a conserved protein, Eco1p(Ctf7), to establish cohesion between sister chromatids during DNA replication. *Genes Dev.* 13: 320–333.
- Toth, A., K. P. Rabitsch, M. Galova, A. Schleiffer, S. B. Buonomo *et al.*, 2000 Functional genomics identifies monopolin: a kinetochore protein required for segregation of homologs during meiosis I. *Cell* 103: 1155–1168.

- Tourriere, H., G. Versini, V. Cordon-Preciado, C. Alabert, and P. Pasero, 2005 Mrc1 and Tof1 promote replication fork progression and recovery independently of Rad53. *Mol. Cell* 19: 699–706.
- Tsubouchi, H., and G. S. Roeder, 2006 Budding yeast Hed1 down-regulates the mitotic recombination machinery when meiotic recombination is impaired. *Genes Dev.* 20: 1766–1775.
- Tsubouchi, T., and G. S. Roeder, 2005 A synaptonemal complex protein promotes homology-independent centromere coupling. *Science* 308: 870–873.
- Tsubouchi, T., A. J. MacQueen, and G. S. Roeder, 2008 Initiation of meiotic chromosome synapsis at centromeres in budding yeast. *Genes Dev.* 22: 3217–3226.
- Tsuchiya, D., C. Gonzalez, and S. Lacefield, 2011 The spindle checkpoint protein Mad2 regulates APC/C activity during prometaphase and metaphase of meiosis I in *Saccharomyces cerevisiae*. *Mol. Biol. Cell* 22: 2848–2861.
- Uhlmann, F., and K. Nasmyth, 1998 Cohesion between sister chromatids must be established during DNA replication. *Curr. Biol.* 8: 1095–1101.
- Uhlmann, F., F. Lottspeich, and K. Nasmyth, 1999 Sister-chromatid separation at anaphase onset is promoted by cleavage of the cohesin subunit Scc1. *Nature* 400: 37–42.
- Uhlmann, F., D. Wernic, M. A. Poupard, E. V. Koonin, and K. Nasmyth, 2000 Cleavage of cohesin by the CD clan protease separin triggers anaphase in yeast. *Cell* 103: 375–386.
- Unal, E., A. Arbel-Eden, U. Sattler, R. Shroff, M. Lichten *et al.*, 2004 DNA damage response pathway uses histone modification to assemble a double-strand break-specific cohesin domain. *Mol. Cell* 16: 991–1002.
- Unal, E., J. M. Heidinger-Pauli, and D. Koshland, 2007 DNA double-strand breaks trigger genome-wide sister-chromatid cohesion through Eco1 (Ctf7). *Science* 317: 245–248.
- Unal, E., J. M. Heidinger-Pauli, W. Kim, V. Guacci, I. Onn *et al.*, 2008 A molecular determinant for the establishment of sister chromatid cohesion. *Science* 321: 566–569.
- Uzunova, K., B. T. Dye, H. Schutz, R. Ladurner, G. Petzold *et al.*, 2012 APC15 mediates CDC20 autoubiquitylation by APC/C (MCC) and disassembly of the mitotic checkpoint complex. *Nat. Struct. Mol. Biol.* 19: 1116–1123.
- Vanoosthuysse, V., S. Prykhodzhiy, and K. G. Hardwick, 2007 Shugoshin 2 regulates localization of the chromosomal passenger proteins in fission yeast mitosis. *Mol. Biol. Cell* 18: 1657–1669.
- Vas, A. C. J., C. A. Andrews, K. Kirkland Matesky, and D. J. Clarke, 2007 In vivo analysis of chromosome condensation in *Saccharomyces cerevisiae*. *Mol. Biol. Cell* 18: 557–568.
- Visintin, R., S. Prinz, and A. Amon, 1997 CDC20 and CDH1: a family of substrate-specific activators of APC-dependent proteolysis. *Science* 278: 460–463.
- Visintin, R., K. Craig, E. S. Hwang, S. Prinz, M. Tyers *et al.*, 1998 The phosphatase Cdc14 triggers mitotic exit by reversal of Cdk-dependent phosphorylation. *Mol. Cell* 2: 709–718.
- Visintin, R., E. S. Hwang, and A. Amon, 1999 Cfi1 prevents premature exit from mitosis by anchoring Cdc14 phosphatase in the nucleolus. *Nature* 398: 818–823.
- Vizeacoumar, F. J., and N. van Dyk, F. S. Vizeacoumar, V. Cheung, J. Li *et al.*, 2010 Integrating high-throughput genetic interaction mapping and high-content screening to explore yeast spindle morphogenesis. *J. Cell Biol.* 188: 69–81.
- Wang, H., D. Liu, Y. Wang, J. Qin, and S. J. Elledge, 2001 Pds1 phosphorylation in response to DNA damage is essential for its DNA damage checkpoint function. *Genes Dev.* 15: 1361–1372.
- Wang, J. C., 2002 Cellular roles of DNA topoisomerases: a molecular perspective. *Nat. Rev. Mol. Cell Biol.* 3: 430–440.
- Warren, C. D., D. M. Brady, R. C. Johnston, J. S. Hanna, K. G. Hardwick *et al.*, 2002 Distinct chromosome segregation roles for spindle checkpoint proteins. *Mol. Biol. Cell* 13: 3029–3041.
- Warren, C. D., D. M. Eckley, M. S. Lee, J. S. Hanna, A. Hughes *et al.*, 2004 S-phase checkpoint genes safeguard high-fidelity sister chromatid cohesion. *Mol. Biol. Cell* 15: 1724–1735.
- Watanabe, Y., 2006 A one-sided view of kinetochore attachment in meiosis. *Cell* 126: 1030–1032.
- Watanabe, Y., and P. Nurse, 1999 Cohesin Rec8 is required for reductional chromosome segregation at meiosis. *Nature* 400: 461–464.
- Weber, S. A., J. L. Gerton, J. E. Polancic, J. L. DeRisi, D. Koshland *et al.*, 2004 The kinetochore is an enhancer of pericentric cohesin binding. *PLoS Biol.* 2: E260.
- Weiss, E., and M. Winey, 1996 The *Saccharomyces cerevisiae* spindle pole body duplication gene MPS1 is part of a mitotic checkpoint. *J. Cell Biol.* 132: 111–123.
- Weiss, E. L., 2012 Mitotic exit and separation of mother and daughter cells. *Genetics* 192: 1165–1202.
- Welburn, J. P. L., M. Vleugel, D. Liu, J. R. Yates, M. A. Lampson *et al.*, 2010 Aurora B phosphorylates spatially distinct targets to differentially regulate the kinetochore-microtubule interface. *Mol. Cell* 38: 383–392.
- Westermann, S., and A. Schleiffer, 2013 Family matters: structural and functional conservation of centromere-associated proteins from yeast to humans. *Trends Cell Biol.* 23: 260–269.
- Westermann, S., H.-W. Wang, A. Avila-Sakar, D. G. Drubin, E. Nogales *et al.*, 2006 The Dam1 kinetochore ring complex moves processively on depolymerizing microtubule ends. *Nature* 440: 565–569.
- Widlund, P. O., J. S. Lyssand, S. Anderson, S. Niessen, J. R. Yates *et al.*, 2006 Phosphorylation of the chromosomal passenger protein Bir1 is required for localization of Ndc10 to the spindle during anaphase and full spindle elongation. *Mol. Biol. Cell* 17: 1065–1074.
- Winey, M., and E. T. O’Toole, 2001 The spindle cycle in budding yeast. *Nat. Cell Biol.* 3: E23–E27.
- Winey, M., C. L. Mamay, E. T. O’Toole, D. N. Mastronarde, T. H. J. Giddings *et al.*, 1995 Three-dimensional ultrastructural analysis of the *Saccharomyces cerevisiae* mitotic spindle. *J. Cell Biol.* 129: 1601–1615.
- Winey, M., G. P. Morgan, P. D. Straight, T. H. J. Giddings, and D. N. Mastronarde, 2005 Three-dimensional ultrastructure of *Saccharomyces cerevisiae* meiotic spindles. *Mol. Biol. Cell* 16: 1178–1188.
- Wirth, K. G., G. Wutz, N. R. Kudo, C. Desdouets, A. Zetterberg *et al.*, 2006 Separase: a universal trigger for sister chromatid disjunction but not chromosome cycle progression. *J. Cell Biol.* 172: 847–860.
- Woodbury, E. L., and D. O. Morgan, 2007 Cdk and APC activities limit the spindle-stabilizing function of Fin1 to anaphase. *Nat. Cell Biol.* 9: 106–112.
- Xiong, B., S. Lu, and J. L. Gerton, 2010 Hos1 is a lysine deacetylase for the Smc3 subunit of cohesin. *Curr. Biol.* 20: 1660–1665.
- Xu, H., C. Boone, and G. W. Brown, 2007 Genetic dissection of parallel sister-chromatid cohesion pathways. *Genetics* 176: 1417–1429.
- Xu, Z., B. Cetin, M. Anger, U. S. Cho, W. Helmhart *et al.*, 2009 Structure and function of the PP2A-shugoshin interaction. *Mol. Cell* 35: 426–441.
- Yaakov, G., K. Thorn, and D. O. Morgan, 2012 Separase biosensor reveals that cohesin cleavage timing depends on phosphatase PP2A(Cdc55) regulation. *Dev. Cell* 23: 124–136.
- Yamagishi, Y., C.-H. Yang, Y. Tanno, and Y. Watanabe, 2012 MPS1/Mph1 phosphorylates the kinetochore protein KNL1/Spc7 to recruit SAC components. *Nat. Cell Biol.* 14: 746–752.
- Yeh, E., J. Haase, L. V. Paliulis, A. Joglekar, L. Bond *et al.*, 2008 Pericentric chromatin is organized into an intramolecular loop in mitosis. *Curr. Biol.* 18: 81–90.



- Yokobayashi, S., and Y. Watanabe, 2005 The kinetochore protein Moa1 enables cohesion-mediated monopolar attachment at meiosis I. *Cell* 123: 803–817.
- Yoon, H. J., and J. Carbon, 1999 Participation of Bir1p, a member of the inhibitor of apoptosis family, in yeast chromosome segregation events. *Proc. Natl. Acad. Sci. USA* 96: 13208–13213.
- Yu, H. G., and D. Koshland, 2007 The Aurora kinase Ipl1 maintains the centromeric localization of PP2A to protect cohesin during meiosis. *J. Cell Biol.* 176: 911–918.
- Zachariae, W., M. Schwab, K. Nasmyth, and W. Seufert, 1998 Control of cyclin ubiquitination by CDK-regulated binding of Hct1 to the anaphase promoting complex. *Science* 282: 1721–1724.
- Zegerman, P., and J. F. X. Diffley, 2010 Checkpoint-dependent inhibition of DNA replication initiation by Sld3 and Dbf4 phosphorylation. *Nature* 467: 474–478.
- Zhang, G., C. D. Kelstrup, X.-W. Hu, M. J. Kaas Hansen, M. R. Singleton *et al.*, 2012 The Ndc80 internal loop is required for recruitment of the Ska complex to establish end-on microtubule attachment to kinetochores. *J. Cell Sci.* 125: 3243–3253.
- Zhang, J., X. Shi, Y. Li, B. J. Kim, J. Jia *et al.*, 2008 Acetylation of Smc3 by Eco1 is required for S phase sister chromatid cohesion in both human and yeast. *Mol. Cell* 31: 143–151.
- Zhu, Z., W.-H. Chung, E. Y. Shim, S. E. Lee, and G. Ira, 2008 Sgs1 helicase and two nucleases Dna2 and Exo1 resect DNA double-strand break ends. *Cell* 134: 981–994.
- Zickler, D., and N. Kleckner, 1999 Meiotic chromosomes: integrating structure and function. *Annu. Rev. Genet.* 33: 603–754.
- Zimniak, T., V. Fitz, H. Zhou, F. Lampert, S. Opravil *et al.*, 2012 Spatiotemporal regulation of Ipl1/Aurora activity by direct Cdk1 phosphorylation. *Curr. Biol.* 22: 787–793.

*Communicating editor: M. Tyers*