

Co-Infestation and Spatial Distribution of *Bactrocera carambolae* and *Anastrepha* spp. (Diptera: Tephritidae) in Common Guava in the Eastern Amazon

E. G. Deus,¹ W. A. C. Godoy,² M. S. M. Sousa,³ G. N. Lopes,⁴ C. R. Jesus-Barros, J. G. Silva,⁶ and R. Adaime^{1,3,5,7}

¹Universidade Federal do Amapá, Programa de Pós-Graduação em Biodiversidade Tropical, Rodovia JK, Km 4, Macapá, Amapá 68902-280, Brazil, ²Universidade de São Paulo, Escola Superior de Agricultura Luiz de Queiroz. Av. Pádua Dias, Piracicaba, São Paulo 13418-900, Brazil, ³Universidade Federal do Amapá, Programa de Pós-Graduação em Desenvolvimento Regional, Rodovia JK, Km 4, Macapá, Amapá 68902-280, Brazil, ⁴Universidade Federal do Piauí, Rodovia Municipal Bom Jesus/Viana, Km 01, Bom Jesus, Piauí 64900-000, Brazil, ⁵Embrapa Amapá, Rodovia JK, Km 5, N° 2600, Macapá, Amapá 68903-419, Brazil, and ⁶Universidade Estadual de Santa Cruz, Rodovia Jorge Amado, Km 16, Ilhéus, Bahia 45662-900, Brazil, and ⁷Corresponding author, e-mail: ricardo.adaime@embrapa.br

Subject Editor: Juan Rull

Received 25 March 2016; Accepted 16 July 2016

Abstract

Field infestation and spatial distribution of introduced *Bactrocera carambolae* Drew and Hancock and native species of *Anastrepha* in common guavas [*Psidium guajava* (L.)] were investigated in the eastern Amazon. Fruit sampling was carried out in the municipalities of Calçoene and Oiapoque in the state of Amapá, Brazil. The frequency distribution of larvae in fruit was fitted to the negative binomial distribution. *Anastrepha striata* was more abundant in both sampled areas in comparison to *Anastrepha fraterculus* (Wiedemann) and *B. carambolae*. The frequency distribution analysis of adults revealed an aggregated pattern for *B. carambolae* as well as for *A. fraterculus* and *Anastrepha striata* Schiner, described by the negative binomial distribution. Although the populations of *Anastrepha* spp. may have suffered some impact due to the presence of *B. carambolae*, the results are still not robust enough to indicate effective reduction in the abundance of *Anastrepha* spp. caused by *B. carambolae* in a general sense. The high degree of aggregation observed for both species suggests interspecific co-occurrence with the simultaneous presence of both species in the analysed fruit. Moreover, a significant fraction of uninfested guavas also indicated absence of competitive displacement.

Key words: invasive species, fruit flies, negative binomial distribution, Brazil

Habitat invasion by exotic species is a global problem (Davis 2009). Although only a small fraction of the introduced species reach high population levels, when they do so, they can cause substantial ecological and economic impact, thus posing serious challenges to such diverse areas as biology, agriculture, transportation, and economy (Pimentel et al. 2000, Mooney and Cleland 2001, Peterson 2003, Liebhold and Tobin 2007). In the last decades, biological invasions have become the focus of a growing number of studies involving ecology, economics, evolution, biogeography, and genetics (Malacrida et al. 2007, Lockwood et al. 2013).

Once established, exotic species can impact native biodiversity by altering habitats, spreading diseases, hybridising with or competing with native species (Davis 2009). Interspecific competition is one of the most important interactions resulting from biological invasions and also one of the main factors responsible for the success of invasive species (Holway and Suarez 1999, Parker et al. 2006,

Romero et al. 2014, Devescovi et al. 2015, Park and Potter 2015). Interspecific competition can either result in the complete displacement of one of the species or the evolution of a stable coexistence between the two species. In fact, it has been demonstrated that there are several mechanisms that are important in the invasion process as they can keep interspecific competition under a tolerable threshold (cf. Duyck et al. 2004, 2006a; Davis 2009; Ekesi et al. 2009; Devescovi et al. 2015).

Most agricultural pests are invasive exotic species (Lockwood et al. 2013). Multiple cases of invasion of new areas by tephritids have been reported, such as *Bactrocera dorsalis* (Hendel) in Hawaii, Australia (as *Bactrocera papayae* Drew and Hancock; subsequently eradicated), and Africa *Bactrocera invadens* Drew, Tsuruta and White; both *B. invadens* and *B. papayae* are now considered synonyms of *B. dorsalis* according to Schutze et al. (2015); *Ceratitits capitata* (Wiedemann), *Ceratitits rosa* (Karsch), and *Bactrocera zonata*

(Saunders) on the island of La Réunion; and *Bactrocera carambolae* Drew and Hancock in Suriname, French Guyana, and northern Brazil (Sauers-Muller 1991, Lux et al. 2003, Duyck et al. 2006a, De Meyer et al. 2010, Lemos et al. 2014).

The impact of invasive fruit flies on native species has been intensively studied on the island of La Réunion, where *Ceratitidis catoirii* Guérin-Méneville (native), *C. capitata*, *C. rosa*, and *B. zonata* (exotic) currently occur. These studies indicate that the pattern of interaction observed suggests hierarchical competition, a mode in which one of the species dominates and influences the abundance of other species (Duyck et al. 2006a,b, 2008). In most cases studied so far, complete exclusion after tephritid invasions has not been observed. However, competitive displacements and a significant decline in the abundance of native species have been reported, particularly in the case of *C. catoirii*, which is nearly extinct in La Réunion (Duyck et al. 2004, 2006a, 2008). Invasion processes, even though many times may result in the displacement of native species, do not always do so instantly as they depend on the action of different competition mechanisms also including the mediation by noncompetitive factors, e.g., lack of an alternative host, differential impact of natural enemies, metapopulation structure, and adaptation to local conditions (Reitz and Trumble 2002).

Niche partitioning and refuge use by native species may result from competition and have been largely used to explain the coexistence of species at a regional scale. At the smallest scale (e.g., host plant), it is possible that the coexistence of invasive and native species is related to the ability of species to exploit different climates and to avoid fruit previously infested by exotic invasive species (Duyck et al. 2004, Devescovi et al. 2015).

Spatial distribution can have a substantial effect in allowing the coexistence of Tephritidae species that share the same resource. In the model proposed by Atkinson and Shorrocks (1981) for insects that feed on discrete and ephemeral habitats (e.g., fruit, dead wood, carrion), aggregation seems to be paramount for coexistence. The authors suggest that two processes can lead to a more prolonged coexistence in a two-species system, namely increased resource division and increased aggregation of the competitors (Atkinson and Shorrocks 1981). If we suppose that two species can use distinct nutritional resources when sharing the same substrate, then intraspecific aggregation could help each species to better exploit the available resources, which in its own turn would allow co-occurrence and potentialize stable coexistence. Also, there are various evidences indicating that larval aggregation is important in Diptera to improve burrowing, reduce parasitism, and increase the efficiency of the feeding process by the simultaneous movement of mouth hooks and secretion of salivar and proteolytic enzymes by multiple individuals (Goodbrod and Goff 1990, Durisko et al. 2014). Studies aimed at ascertaining the spatial distribution of fruit flies using both adult trapping and fruit sampling have showed different levels of aggregation. However, the mechanisms responsible for this pattern within Tephritidae are still not completely understood (cf. Satarkar et al. 2009, Birke and Aluja 2011, Soemargono et al. 2011, Jahnke et al. 2014, Devescovi et al. 2015).

Bactrocera carambolae, the carambola fruit fly, is native to Indonesia, Malaysia, and Thailand. It has invaded Suriname (Paramaribo in 1975) and is now found in Guyana, French Guiana, and northern Brazil (Sauers-Müller 2005, Godoy et al. 2011). This species infests more than 150 fruit species and is considered a quarantine pest insect in the Caribbean region (Malavasi et al. 2000a, Haq et al. 2015).

In Brazil, *B. carambolae* was first reported in 1996 in the municipality of Oiapoque, state of Amapá, where it occurs in sympatry

with native species in the genus *Anastrepha*, even sharing the same host species (Godoy et al. 2011, Lemos et al. 2014). In recent studies in Brazil, *B. carambolae* has been reported infesting guava (*Psidium guajava* L.) at a relatively high abundance when compared with *Anastrepha* species, except for *Anastrepha striata* Schiner (Lemos et al. 2014). However, despite being in the country for 20 years and its significant economic importance, data on *B. carambolae* population dynamics, demography, host preference, and its impact on the community of native tephritids are scarce.

In this study, we evaluated field infestation and spatial distribution of *B. carambolae* and native species of *Anastrepha* in common guava in the eastern Amazon, where the carambola fruit fly was detected for the first time in Brazil. We hypothesized that the spatial distribution pattern of fruit fly species infesting guavas in the studied area is aggregated.

Materials and Methods

Study Area and Sampling

Fruit sampling was carried out in the municipalities of Calçoene (02°33'N, 50°56'W) and Oiapoque (03°50'N, 51°50'W) in the northernmost region of the state of Amapá, in the Brazilian Amazon. The climate is defined as Am (monsoon) with annual rainfall over 3,000 mm (Alvares et al. 2013). The native vegetation in the region is typical of upland forest (IEPA 2002).

Collections were carried out in April, June, July, and December of 2013. In each site, six guava trees ~100 m from each other were randomly selected. A total of 30 fruit equally distributed among trees was collected at each sampling occasion, totalling 90 fruit per municipality. Fruit were collected both from the trees and from the ground among those that had recently fallen. The fruit collected from trees were completely ripe, a criterion that was adopted to reduce the probability of underestimating infestation, as such fruit were about to fall from the tree. Fruit collected from the ground were carefully examined and those that had holes or cuts in the peel were not sampled as the larvae could have already left the fruit. The collected fruit were placed individually in plastic containers with a layer of sterilized sand. Every 3 d, all containers were checked and all puparia obtained were counted and transferred into 30-ml plastic containers with a layer of moist vermiculite at the bottom and covered with voile cloth until adults emerged. The containers were placed in a growth chamber at controlled temperature ($26.5 \pm 0.3^\circ\text{C}$), RH ($70 \pm 5\%$), and a photoperiod of 12:12 (L:D) h and checked daily for emerged adults.

Guava was chosen as substrate to investigate larval distribution per fruit as it is the preferred tephritid host in northern Brazil (Zucchi et al. 2011). Moreover, previous studies carried out in the state of Amapá verified that *B. carambolae* has frequently shared this host with native *Anastrepha* spp. (Lemos et al. 2014). Additionally, guava is widespread in the state of Amapá fruiting from December to June and it is either cultivated without any phytosanitary management in backyards in urban and rural areas or grows along roads near the native vegetation.

Insect Identification

Anastrepha and *Bactrocera* specimens were identified according to Zucchi (2000) and Drew and Hancock (1994), respectively. Voucher specimens were deposited at the Laboratório de Entomologia, Empresa Brasileira de Pesquisa Agropecuária - Embrapa, Amapá, Brazil.

Data Analysis

Fruit Infestation and Fruit Fly Abundance. The average levels of infestation were calculated dividing the number of puparia by the mass (grams) of the fruit. We quantified the percentage of fruit infested by only one species of fruit fly, the percentage of coinfested fruit (fruit infested by more than one species), and the percentage of noninfested fruit. Abundance was considered as the proportion of individuals within the same species relative to the total number of individuals in the sample.

Spatial Distribution of Fruit Flies. In order to quantify spatial distribution, fruit infestation levels and abundance were described considering the collection sites, i.e., compiling data from different trees. Males and females were grouped together in the analysis and a table of frequency distribution considering the fruit as the sampling unit was constructed. A frequency class interval was constructed considering data of the total absence of individuals from zero (i.e., uninfested fruit) to the highest abundance of tephritids per fruit. Frequency distribution graphs were prepared for both *B. carambolae* and *Anastrepha* spp. and the mean and variance of the distribution were estimated considering the number of sampling units and the frequency of each unit. To evaluate aggregation level, the k parameter of the negative binomial distribution was also estimated (Southwood and Henderson 2000). The highest aggregation level occurs when the value of k tends towards zero. The best estimate of k was also determined by the method of maximum likelihood (Ludwig and Reynolds 1988). The fit of the negative binomial distribution was tested by comparing the observed frequencies with the expected frequencies by the Pearson χ^2 test (Crawley 2007). The algorithm was written in the R statistical programming language based on Crawley (2007). The analyses were carried out using R software (R Core Team 2014).

Results

Infestation Levels and Abundance

In Calçoene, a total of 910 puparia was obtained and there was emergence of fruit flies from 58.8% of them and from the remaining 41.2% no emergence was registered. The species reared from the collected fruit were *Anastrepha fraterculus* (Wiedemann) (5, 0.9%), *A. striata* Schiner (397, 74.2%), and *B. carambolae* (133, 24.9%). Fruit infestation in this locality ranged from 0.02 to 1.21 puparia/g of fruit (mean 0.34). In Oiapoque, a total of 1,090 puparia was obtained and fruit flies emerged from 76.1% of them and there was no emergence out of the remaining 23.9%. The species reared from the collected fruit were *A. fraterculus* (3, 0.4%), *A. striata* (650, 78.3%), and *B. carambolae* (177, 21.3%). Fruit infestation in this locality ranged from 0.02 to 2.94 puparia/g of fruit (mean 0.41). The abundance of *A. striata* was higher than that of *A. fraterculus* and *B. carambolae* in both sampling areas. No parasitoid emergence was observed in either collection site. The proportion of noninfested fruit was higher in Calçoene, which had a direct impact on the number of puparia obtained. The pattern of fruit infestation observed in both sites is shown on Table 1.

Considering only the coinfested fruit, there was a change in pattern, in which *B. carambolae* became the more abundant species accounting for >50% of the individuals in both sampling areas (Table 2). In Calçoene, *A. striata* and *B. carambolae* co-occurred in eight fruit and *A. striata* and *A. fraterculus* in three fruit. In Oiapoque, *A. striata* and *B. carambolae* co-occurred in 13 fruit, *A. fraterculus* and *A. striata* in one fruit, *A. fraterculus* and

Table 1. Proportion of infested fruit

Locality	Number of infested fruit				
	Only by <i>A. fraterculus</i>	Only by <i>A. striata</i>	Only by <i>B. carambolae</i>	Coinfested	Uninfested
Calçoene	0 (0%)	48 (53.3%)	4 (4.4%)	11 (12.2%)	27 (30%)
Oiapoque	0 (0%)	59 (65.6%)	4 (4.4%)	16 (17.8%)	11 (12.2%)

Table 2. Abundance of *A. fraterculus*, *A. striata*, and *B. carambolae* in coinfested fruit

	Number of coinfested fruit	Puparia	<i>A. fraterculus</i>	<i>A. striata</i>	<i>B. carambolae</i>
Calçoene	11	185	5 (3.7%)	28 (20.7%)	102 (75.6%)
Oiapoque	16	342	3 (1.3%)	85 (35.7%)	150 (63%)

B. carambolae in one fruit and *A. fraterculus*, *A. striata*, and *B. carambolae* in one fruit.

Spatial Distribution

The observed distribution frequency of *A. fraterculus*, *A. striata* e *B. carambolae* is shown in Figure 1 for both sampling areas. In all cases, the variance was higher than the mean, which suggests an aggregated distribution pattern described by the negative binomial distribution.

The values estimated for the aggregation coefficient (k) are close to zero, which also suggests that the fruit fly community distribution in guava is aggregated in both areas sampled, despite the difference in magnitude between them, since the population of *A. striata* in Oiapoque showed a lower degree of aggregation when compared with the others (Table 3).

In both areas, the chi-square values were lower than the critical value in the table ($\chi^2 = 0$, $df = 34$; $\chi^2 = 0$, $df = 34$ e $\chi^2 = 0$, $df = 25$ in Calçoene and $\chi^2 = 0$, $df = 3$; $\chi^2 = 0$, $df = 30$; $\chi^2 = 0$, $df = 53$ in Oiapoque for *A. fraterculus*, *A. striata*, and *B. carambolae*, respectively), indicating that the observed values are significantly explained by the negative binomial distribution, which suggests an aggregated distribution. The expected and observed values of frequency distributions are shown in Figure 2.

Discussion

This is the first study to evaluate the infestation pattern and spatial distribution of *B. carambolae* and *Anastrepha* spp. in areas where there is host overlap. Previous studies carried out in the state of Amapá revealed that *B. carambolae* infests fruit commonly used by *Anastrepha* species (Lemos et al. 2014).

In this study, we verified that *A. striata* is the most abundant species in both areas, and most fruit were infested only by this species (Table 1). In Brazil, *A. striata* is the most abundant species in guava only in the Amazon region, where it is commonly reported as either the only species infesting guavas or the most abundant when other fruit fly species are also present (cf. Silva et al. 2007a,b; Deus et al. 2009; Pereira et al. 2010; Jesus-Barros et al. 2012; Marsaro-Júnior et al. 2013).




Fig. 1. Frequency of *A. fraterculus*, *A. striata*, and *B. carambolae* in the sampling areas indicating the distribution of emerging adults (x axis) from the fruits. Numbers on the x axis determine the class interval, showing data from the total absence of individuals (i.e., uninfested fruit) to the highest abundance of tephritids per fruit.

Table 3. Frequency distribution of *A. fraterculus*, *A. striata*, and *B. carambolae* in guava

Species and locality	Mean (\bar{x})	Variance (s^2)	K^a	K^b
<i>A. fraterculus</i> in Calçoene	0.05	0.12	0.047	0.039
<i>A. striata</i> in Calçoene	4.4	45.07	0.475	0.396
<i>B. carambolae</i> in Calçoene	1.47	23.01	0.101	0.084
<i>A. fraterculus</i> in Oiapoque	0.06	0.12	0.06	0.06
<i>A. striata</i> in Oiapoque	7.2	60.70	0.968	0.852
<i>B. carambolae</i> in Oiapoque	1.94	47.38	0.083	0.071

^ainitial estimate of $K \left(\frac{\bar{x}}{(s^2 - \bar{x})} \right)$.

^bestimated using the method of maximum likelihood $\left(\log_{10} \left(\frac{N}{N_0} \right) = K \log_{10} \left[1 + \left(\frac{\bar{x}}{K} \right) \right] \right)$, where N is the total number of sampling units and N_0 is the number of sampling units with zero individuals.

A. fraterculus represented <1% of the specimens reared. Even though it is the fruit fly species with the widest geographical distribution in Brazil, its frequency and distribution are significantly reduced northwards. In the Southern region, this species has been reported to have a frequency higher than 95% in guava (Nunes et al. 2012, Jahnke et al. 2014). However, for the Northern region, *A. fraterculus* is a secondary species with marginal distribution (Malavasi et al. 2000b, Silva and Ronchi-Teles, 2000). In the state of Amapá, *A. fraterculus* has eight known hosts; however, it has low frequency and low abundance. The highest populational density was observed in fruit of *Mouriri acutiflora* Naudin (Melastomataceae), a typical species of the Amazon forest (Deus and Adaime 2013). So far, there are no studies that explain the low abundance of *A. fraterculus* and the high abundance of *A. striata* in guava in the Northern region, however, it is likely that behavioral aspects together with environmental factors are responsible for the pattern observed here. Therefore, additional studies are necessary to elucidate this question.

B. carambolae represented 24.9 and 21.3% of all fruit flies reared from fruit collected in Calçoene and Oiapoque, respectively,

with a higher abundance in coinfecting fruit (Table 2). Several studies have demonstrated that species within the *dorsalis* complex have a marked preference for ripe fruit (cf Alyokhin et al. 2000, Cornelius et al. 2000, Rattanapun et al. 2009, Quilici et al. 2014). Even though *B. carambolae* belongs to the *dorsalis* complex, there are no studies on the influence of the degree of fruit ripeness on female oviposition decision in this species. Thus, additional studies are necessary to test this hypothesis.

The high aggregation pattern observed in this study corroborates similar findings in other tephritids (cf. Dimou et al. 2003, Satarkar et al. 2009, Soemargono et al. 2011, Jahnke et al. 2014). The evaluation of the spatial distribution of species requires analytical tools that can estimate characteristic parameters and indicate dispersion patterns, which generally are the expression of individual behavior as evidenced by the magnitude of the values found (Taylor 1984). Additionally, factors such as the environmental heterogeneity, resource availability, and presence of natural enemies can influence spatial distribution (Waters 1959, Satarkar et al. 2009).

A. fraterculus, *A. striata*, and *B. carambolae* use guava as a host in their area of natural distribution. Thus, when there is resource overlap, the response of the native species to the presence of the potential exotic competitor is paramount for the persistence of the former. Likewise, the strategy employed by the invading species is crucial for its establishment in the presence of native species. As postulated by Atkinson and Shorrocks (1981), the increase in resource division and aggregation of competitors may decrease the effect of competitive displacement. When it comes to a specific and ephemeral resource, the high degree of aggregation of *A. fraterculus*, *A. striata*, and *B. carambolae* in the fruit indicates potential for the coexistence of tephritids in guava in the northern region of Brazil. There are probably other factors involved, including abiotic factors, which can have an influence upon the spatial distribution and that should be taken into account in future studies.

The results reported herein suggest that *A. striata* is the most abundant species in both sampling areas, even though other




Fig. 2. Expected and observed frequency distribution of *A. fraterculus*, *A. striata*, and *B. carambolae* in the sampling areas.

Tephritidae species are also present. Our results also indicate that the spatial distribution of *Anastrepha fraterculus*, *A. striata*, and *B. carambolae* in guava follows a negative binomial distribution, i.e., it is aggregated.

Finally, we consider that basic biology studies are needed for the understanding of the mechanisms involved in the coexistence of species. Among these studies, we could include the influence of coloration and degree of fruit ripeness in the preference and larval development of *B. carambolae*, the response of *B. carambolae* to volatiles of different fruit species, its distribution pattern in the presence and absence of competitors, and the influence of abiotic factors in community structuring. Such studies are paramount for the

understanding of the processes driving fruit fly community dynamics on guava in the Amazon region. Moreover, the data generated will be a valuable tool for the development of management strategies.

Acknowledgments

We thank our colleague at Embrapa Amapá Carlos Alberto Moraes, for support during the fruit sampling expedition. We would also like to thank Carter Robert Miller for reviewing an earlier version of the article as well as the anonymous reviewers for their helpful suggestions. Thanks are also due to the Conselho Nacional de Desenvolvimento Científico e Tecnológico—CNPq for the PhD Fellowship granted to E. G. Deus. We also thank the Fundação de

Amparo à Pesquisa do Estado do Amapá—FAPEAP for partially funding the project (250.203/064/2014). W. A. C. Godoy, J. G. Silva, and R. Adaime are fellow researchers of CNPq.

References Cited

- Alvares, C. A., J. L. Stape, P. C. Sentelhas, J. L. M. Gonçalves, and G. Sparovek. 2013. Köppen's climate classification map for Brazil. *Meteorol. Z.* 22: 711–728.
- Alyokhin, A. V., R. H. Messing, and J. J. Duan. 2000. Visual and olfactory stimuli and fruit maturity affect trap captures of oriental fruit flies (Diptera: Tephritidae). *J. Econ. Entomol.* 93: 644–649.
- Atkinson, W. D., and B. Shorrocks. 1981. Competition on a divided and ephemeral resource: a simulation model. *J. Anim. Ecol.* 50: 461–471.
- Birke, A., and M. Aluja. 2011. *Anastrepha ludens* and *Anastrepha serpentina* (Diptera: Tephritidae) do not infest *Psidium guajava* (Myrtaceae), but *Anastrepha obliqua* occasionally shares this resource with *Anastrepha striata* in nature. *J. Econ. Entomol.* 104: 1204–1211.
- Cornelius, M. L., J. J. Duan, and R. H. Messing. 2000. Volatile Host Fruit Odors as Attractants for the Oriental Fruit Fly (Diptera: Tephritidae). *J. Econ. Entomol.* 93: 93–100.
- Crawley, M. J. 2007. *The R Book*. John Wiley & Sons, Chichester, West Sussex England.
- Davis, M. A. 2009. *Invasion Biology*. Oxford University Press Inc., New York.
- De Meyer, M., M. P. Robertson, M. W. Mansell, S. Ekesi, K. Tsuruta, W. Mwaiko, J. F. Vaissières, and A. T. Peterson. 2010. Ecological niche and potential geographic distribution of the invasive fruit fly *Bactrocera invadens* (Diptera, Tephritidae). *Bull. Entomol. Res.* 100: 35–48.
- Deus, E. G., and R. Adaime. 2013. Dez anos de pesquisas sobre moscas-das-frutas (Diptera: Tephritidae) no Estado do Amapá: avanços obtidos e desafios futuros. *Biota Amazônia*. 3: 157–160.
- Deus, E. G., R. A. Silva, D. B. Nascimento, C. F. Marinho, and R. A. Zucchi. 2009. Hospedeiros e parasitoides de espécies de *Anastrepha* (Diptera, Tephritidae) em dois municípios do Estado do Amapá. *Rev. Agric.* 3: 194–203.
- Devescovi, F., M. C. Liendo, G. E. Bachmann, J. P. Bouvet, F. H. Milla, M. T. Vera, J. L. Cladera, and D. F. Segura. 2015. Fruit infestation patterns by *Anastrepha fraterculus* and *Ceratitidis capitata* reveal that cross-recognition does not lead to complete avoidance of interspecific competition in nature. *Agric. Forest Entomol.* 17: 325–335.
- Dimou, I. J., C. Koutsikopoulos, A. Economopoulos, and J. Lykakis. 2003. The distribution of olive fruit fly captures with MacPhail traps within an olive orchard. *Phytoparasitica*. 31: 124–31.
- Drew, R.A.I., and D. L. Hancock. 1994. The *Bactrocera dorsalis* complex of fruit flies (Diptera: Tephritidae: Dacinae) in Asia. *Bull. Entomol. Res., Suppl.* 2: 68.
- Durisko, Z., R. Kemp, R. Mubasher, and R. Dukas. 2014. Dynamics of social behavior in fruit fly larvae. *Plos One*. 9: 1–8.
- Duyck, P. F., P. David, and S. Quilici. 2004. A review of relationships between interspecific competition and invasions in fruit flies (Diptera: Tephritidae). *Ecol. Entomol.* 29: 511–20.
- Duyck, P. F., P. David, G. Junpd, C. Brunel, R. L. Dupont, and S. Quilici. 2006a. Importance of competition mechanisms in successive invasions by polyphagous tephritids in La Réunion. *Ecology*. 87: 1770–1780.
- Duyck, P. F., P. David, and S. Quilici. 2006b. Climatic niche partitioning following successive invasions by fruit flies in La Réunion. *J. Anim. Ecol.* 75: 518–526.
- Duyck, P. F., P. David, S. Pavoine, and S. Quilici. 2008. Can host-range allow niche differentiation of invasive polyphagous fruit flies (Diptera: Tephritidae) in La Réunion? *Ecol. Entomol.* 33: 439–452.
- Ekesi, S., M. K. Billah, P. W. Nderitu, S. A. Lux, and I. Rwomushana. 2009. Evidence for competitive displacement of *Ceratitidis cosyra* by the invasive fruit fly *Bactrocera invadens* (Diptera: Tephritidae) on mango and mechanisms contributing to the displacement. *J. Econ. Entomol.* 102: 981–91.
- Godoy, M.J.S., W.S.P. Pacheco, R. R. Portal, J. M. Pires Filho, and L. M. M. Moraes. 2011. Programa Nacional de Erradicação da Mosca-da-Carambola, pp. 134–158. *In* R. A. Silva, W. P. Lemos, and R. A. Zucchi (eds.), *Moscas-das-frutas na Amazônia Brasileira: Diversidade, Hospedeiros e Inimigos Naturais*. Embrapa Amapá, Macapá.
- Goodbrod, J. R., and M. L. Goff. 1990. Effects of larval population density on rates of development and interactions between two species of *Chrysomya* (Diptera: Calliphoridae) in laboratory culture. *J. Med. Entomol.* 27: 338–343.
- Haq, I.U., M.J.B. Vreysen, C. Cacères, T. E. Shelly, and J. Hendrichs. 2015. Optimizing methyl eugenol aromatherapy to maximize posttreatment effects to enhance mating competitiveness of male *Bactrocera carambolae* (Diptera: Tephritidae). *Insect Sci.* 22: 661–669.
- Holway, D. A., and A. V. Suarez. 1999. Animal behavior: an essential component of invasion biology. *Trends Ecol. Evol.* 14: 328–330.
- (IEPA) Instituto de Pesquisas Científicas e Tecnológicas do Estado do Amapá. 2002. Macrodiagnóstico do Estado do Amapá: primeira aproximação do ZEE. IEPA, Macapá, AP.
- Jahnke, S. M., E. M. Del Ponte, L. R. Redaelli, and D.R.G.P. Rego. 2014. Spatial patterns and associations of *Anastrepha fraterculus* (Diptera: Tephritidae) and its parasitoid *Doryctobracon areolatus* (Hymenoptera: Braconidae) in organic orchards of *Psidium guajava* and *Acca sellowiana*. *Fla. Entomol.* 97: 744–52.
- Jesus-Barros, C. R., R. Adaime, M. N. Oliveira, W. R. Silva, S. V. Costa-Neto, and M. F. Souza-Filho. 2012. *Anastrepha* (Diptera: Tephritidae) species, their hosts and parasitoids (Hymenoptera: Braconidae) in five municipalities of the State of Amapá, Brazil. *Fla. Entomol.* 95: 694–705.
- Lemos, L. N., R. Adaime, C. R. Jesus-Barros, and E. G. Deus. 2014. New hosts of *Bactrocera carambolae* (Diptera: Tephritidae) in Brazil. *Fla. Entomol.* 97: 694–704.
- Liebholt, A. M., and P. C. Tobin. 2007. Population Ecology of Insect Invasions and Their Management. *Annu. Rev. Entomol.* 53: 387–408.
- Lockwood, J. L., M. F. Hoopes, and M. P. Marchetti. 2013. *Invasion Ecology* 2nd edn. John Wiley & Sons, Chichester, West Sussex England.
- Ludwig, J. A., and J. F. Reynolds. 1988. *Statistical ecology: a primer on methods and computing*. John Wiley & Sons, New York.
- Lux, S. A., R. S. Copeland, I. M. White, A. Manrakhan, and M. K. Billah. 2003. A new invasive fruit fly species from the *Bactrocera dorsalis* (Hendel) group detected in East Africa. *Int. J. Trop. Insect Sci.* 23: 355–361.
- Malacrida, A. R., M. Gomulski, S. Bonizzoni, G. Gasperi, and C. R. Guglielmino. 2007. Globalization and fruit fly invasion and expansion: the medfly paradigm. *Genetica*. 131: 1–9.
- Malavasi, A., A. Sauer-Muller, D. Midgarden, V. Kellman, D. Didelot, P. Caplon, and O. Ribeiro. 2000a. Regional programme for the eradication of the carambola fruit fly in South America, pp. 395–399. *In* K. H. Tan (ed.), *Area-Wide Control of Fruit Flies and Other Insect Pests*. Panerbit Universiti, Sains Malaysia.
- Malavasi, A., R. A. Zucchi, and R. L. Sugayama. 2000b. Biogeografia, pp. 93–98. *In* A. Malavasi and R. A. Zucchi (Eds.), *Moscas-das-frutas de Importância Econômica no Brasil: Conhecimento Básico e Aplicado*. São Paulo, Holos
- Marsaro-Junior, A. L., E. G. Deus, B. Ronchi-Teles, R. Adaime, and R. J. Silva-Junior. 2013. Species of *Anastrepha* (Diptera: Tephritidae) captured in a guava orchard (*Psidium guajava* L., Myrtaceae) in Boa Vista, Roraima, Brazil. *Braz. J. Biol. Brazilian Journal Biology*. 73: 879–886.
- Mooney, H. A., and E. E. Cleland. 2001. The evolutionary impact of invasive species. *PNAS*. 98: 5446–5451.
- Nunes, A. D., F. A. Müller, R. S. Gonçalves, M. S. Garcia, V. A. Costa, and D. E. Nava. 2012. Moscas frugívoras e seus parasitoides nos municípios de Pelotas e Capão do Leão, Rio Grande do Sul, Brasil. *Ciência Rural*. 42: 6–12.
- Park, D. S., and D. A. Potter. 2015. A reciprocal test of Darwin's naturalization hypothesis in two mediterranean-climate regions. *Global Ecol. Biogeogr.* 24: 1049–1058.
- Parker, J. D., D. E. Burkepile, and M. E. Hay. 2006. Opposing effects of native and exotic herbivores on plant invasions. *Science* 311: 1459–1461.
- Peterson, A.T. 2003. Predicting the geography of species' invasions via ecological niche modeling. *Q. Rev. Biol.* 78: 419–433.
- Pereira, J. D. B., D. P. Buriti, W. P. Lemos, W. R. Silva, and R. A. Silva. 2010. Espécies de *Anastrepha* Schiner (Diptera: Tephritidae), seus hospedeiros e parasitoides nos estados do Acre e Rondônia, Brasil. *Biota Neotrop.* 3: 441–446.

- Pimentel, D., L. Lach, R. Zuniga, and D. Morrison. 2000. Environmental and economic costs of nonindigenous species in the United States. *BioScience*. 50: 53–65.
- Quilici, S., T. Atiama-Nurbel, T. Brévault., et al. 2014. Plant odors as fruit fly attractants, pp. 119–144. *In* T. Shelly, N. Epsky, and E. B. Jang (eds.), *Trapping and the Detection, Control, and Regulation of Tephritid Fruit Flies*. Springer Dordrecht, Heidelberg, New York, London.
- R Core Team. 2014. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <http://www.R-project.org/>.
- Rattanapun, W., W. Amornsak, and A. R. Clarke. 2009. *Bactrocera dorsalis* preference for and performance on two mango varieties at three stages of ripeness. *Entomol. Exp. Appl.* 131: 243–53.
- Reitz, S. R., and J. T. Trumble. 2002. Competitive displacement among insects and arachnids. *Annu. Rev. Entomol.* 47: 435–65.
- Romero, D., J. C. Báez, F. Ferri-Yáñez, J. J. Bellido, and R. Real. 2014. Modelling favourability for invasive species encroachment to identify areas of native species vulnerability. *Sci. World J.* 2014: 1–9.
- Satarkar, V. R., S. V. Krishnamurthy, J. R. Faleiro, and A. Verghese. 2009. Spatial distribution of major *Bactrocera* fruit flies attracted to methyl eugenol in different ecological zones of Goa, India. *Int. J. Trop. Insect Sci.* 29: 195–201.
- Sauers-Müller, A. V. 1991. An overview of the Carambola fruit fly *Bactrocera* species (Diptera: Tephritidae), found recently in Suriname. *Fla. Entomol.* 74: 432–40.
- Sauers-Müller, A. V. 2005. Host plants of the carambola fruit fly, *Bactrocera carambolae* Drew & Hancock (Diptera: Tephritidae), in Suriname. *South America. Neotrop. Entomol.* 34: 203–14.
- Silva, N. M., and B. Ronchi-Teles. 2000. Amapá, Amazonas, Pará, Rondônia e Roraima pp. 2039. *In* A. Malavasi and R. A. Zucchi (eds.), *Moscas-das-Frutas de Importância Econômica no Brasil: Conhecimento Básico e Aplicado*. Holos, São Paulo.
- Silva, R. A., D. B. Nascimento, E. G. Deus, and G. D. Souza and L. S. P. Oliveira 2007b. Hospedeiros e parasitóides de *Anastrepha* spp. (Diptera: Tephritidae) em Itauba do Pírim. Estado do Amapá. *Ciência Rural.* 2: 557–560.
- Silva, R. A., S. L. O. Xavier, M. F. Souza-Filho, W. R. Silva, D. B. Nascimento, and E. G. Deus. 2007a. Frutíferas hospedeiras e parasitóides (Hym., Braconidae) de *Anastrepha* spp. (Dip., Tephritidae) na Ilha de Santana, Estado do Amapá. *Brasil. Arq. Inst. Biol.* 74: 153–6.
- Schutze, M., K. Mahmood, A. Pavasovic, W. Bo, J. Newman, A. R. Clarke, M. N. Krosch, and S. L. Cameron. 2015. One and the same: integrative taxonomic evidence that *Bactrocera invadens* (Diptera: Tephritidae) is the same species as the oriental fruit fly *Bactrocera dorsalis*. *Syst. Entomol.* 40: 472–86.
- Soemargono, A., M. Muryati, A. Basyim, and M. Istianto. 2011. Spatial distribution pattern of the fruit fly *Bactrocera dorsalis* complex (Diptera: Tephritidae) in a mango orchard. *Agrivita* 33: 207–13.
- Southwood, T. R. E., and P. A. Henderson. 2000. *Ecological Methods*. Blackwell Science. Osney Mead, Oxford.
- Taylor, L. R. 1984. Assessing and interpreting the spatial distribution of insect populations. *Annu. Rev. Entomol.* 29: 231–57.
- Waters, W. E. 1959. A quantitative measure of aggregation in insects. *J. Econ. Entomol.* 52: 1180–4.
- Zucchi, R. A., E. G. Deus, and R. A. Silva. 2011. Espécies de *Anastrepha* e seus hospedeiros na Amazônia brasileira, pp. 51–70. *In* R. A. Silva, W. P. Lemos and R. A. Zucchi (eds.), *Moscas-das-frutas na Amazônia Brasileira: Diversidade, Hospedeiros e Inimigos Naturais*. Embrapa Amapá, Macapá.
- Zucchi, R. A. 2000. Taxonomia, pp. 13–24. *In* A. Malavasi, and R. A. Zucchi (eds.), *Moscas-das-frutas de Importância Econômica no Brasil: Conhecimento Básico e Aplicado*. Holos, São Paulo.