

RESEARCH

Open Access

Conditional overexpression of TGF β 1 promotes pulmonary inflammation, apoptosis and mortality via TGF β R2 in the developing mouse lung

Angara Sureshbabu¹, Mansoor A Syed¹, Chandra Sekhar Boddupalli², Madhav V Dhodapkar², Robert J Homer³, Parviz Minoo⁴ and Vineet Bhandari^{1*}

Abstract

Background: Earlier studies have reported that transforming growth factor beta 1 (TGF β 1) is a critical mediator of hyperoxia-induced acute lung injury (HALI) in developing lungs, leading to impaired alveolarization and a pulmonary phenotype of bronchopulmonary dysplasia (BPD). However, the mechanisms responsible for the TGF β 1-induced inflammatory signals that lead to cell death and abnormal alveolarization are poorly understood. We hypothesized that TGF β 1 signaling via TGF β R2 is necessary for the pathogenesis of the BPD pulmonary phenotype resulting from HALI.

Methods: We utilized lung epithelial cell-specific TGF β 1 overexpressing transgenic and TGF β R2 null mutant mice to evaluate the effects on neonatal mortality as well as pulmonary inflammation and apoptosis in developing lungs. Lung morphometry was performed to determine the impaired alveolarization and multicolor flow cytometry studies were performed to detect inflammatory macrophages and monocytes in lungs. Apoptotic cell death was measured with TUNEL assay, immunohistochemistry and western blotting and protein expression of angiogenic mediators were also analyzed.

Results: Our data reveals that increased TGF β 1 expression in newborn mice lungs leads to increased mortality, macrophage and immature monocyte infiltration, apoptotic cell death specifically in Type II alveolar epithelial cells (AECs), impaired alveolarization, and dysregulated angiogenic molecular markers.

Conclusions: Our study has demonstrated the potential role of inhibition of TGF β 1 signaling via TGF β R2 for improved survival, reduced inflammation and apoptosis that may provide insights for the development of potential therapeutic strategies targeted against HALI and BPD.

Keywords: Transforming growth factor, Oxygen, Inflammation, Cell death, Angiopoietin, Newborn, Pulmonary, Bronchopulmonary dysplasia

Background

Transforming growth factor beta 1 (TGF β 1) is a secretory cytokine that binds to the Type II TGF β receptor (TGF β R2), which then complexes with the type I TGF β receptor (TGF β RI or ALK1 or ALK5). This binding initiates TGF β signaling via smad phosphorylation and nuclear translocation [1]. There is growing evidence that TGF β signaling is involved in the regulation of branching and septation phases of lung development [2,3]. In contrast, TGF β 1 has been documented to

induce inflammation and apoptosis in lung epithelial cells [4,5].

Premature infants require supplemental oxygen and/or mechanical ventilation for prolonged time periods, and may subsequently develop Bronchopulmonary Dysplasia (BPD). BPD, the most common chronic respiratory disease in infants, is characterized by the presence of impaired alveolarization and dysregulated vascularization [6]. Understanding the mechanisms responsible for the development of BPD has focused on the identification of signaling or remodeling molecules that are crucial in lung development and/or response to lung injury. One such signaling pathway that is known to be a central mediator in hyperoxia induced acute lung injury (HALI)

* Correspondence: vineet.bhandari@yale.edu

¹Department of Pediatrics, Yale University School of Medicine, 333 Cedar Street, New Haven, CT 06510, USA

Full list of author information is available at the end of the article

and BPD is TGF β signaling [1,7]. Previous studies have demonstrated increased expression of TGF β , both total and bioactive forms, in bronchoalveolar lavage fluid and tracheal aspirates of infants developing BPD [8,9].

Many studies have demonstrated that TGF β 1 is a critical regulator of HALI [10,11]. Hyperoxia mediated TGF β 1 sets into flow a deluge of inflammation [12,13]. However, the mechanisms responsible for the TGF β 1-induced immunogenicity and the levels of inflammatory signals that lead to cell death are poorly understood. Therefore, this study was aimed to investigate the quantitation of endogenous immune cells regulated by TGF β 1, which result in cell death. In addition, we wanted to assess the role of TGF β 1 in alveolarization and impact on molecular mediators known to be associated with development of HALI and BPD. Furthermore, we wanted to evaluate if all or some of the above-mentioned effects of TGF β 1 signaling were mediated via TGF β 2.

We hypothesized that TGF β 1 signaling via TGF β 2 is necessary for the pathogenesis of HALI and the BPD phenotype resulting from hyperoxia. Therefore, we used lung epithelial cell-specific TGF β 1 transgenic (TG) mice model to demonstrate TGF β 1-induced inflammation and apoptosis in developing lungs. In addition, we also evaluated Angiopoietin (ANGPT) 1 and ANGPT2 protein expression in the above model systems as ANGPT2 has been described to have a critical role in HALI and BPD [14,15]. Furthermore, in order to assess the receptor specificity of these effects, we used lung epithelial cell-specific TGF β 2 null mutant (knockout or KO) mice to evaluate the effects of TGF β 1-induction on mortality as well as pulmonary inflammation and apoptosis in developing lungs.

Methods

Animals

All animal studies were approved by the Institutional Animal Care and Use Committee at the Yale University School of Medicine. The TGF β 1TG mice were generated as previously described [3,16], and express TGF β 1 with maternal exposure to doxycycline (dox) in the drinking water. These triple-TGF β 1 TG mice are expressing the active form of human TGF β 1 and the levels and downstream effects of the same (upon dox activation) over time have been previously reported [16]. Maternal exposure to dox was performed from postnatal (PN) day 1–7 or from PN7 to PN10 [17]. Lung epithelial-cell specific deletion of TGF β 2 (TGF β 2KO) has been recently described and these mice were utilized for our experiments [18].

Bronchoalveolar lavage fluid collection

The mouse lungs were lavaged three times with 300 microliters of PBS using standard techniques and inflation pressure, as previously described [19]. The bronchoalveolar

lavage fluid (BALF) was centrifuged at 400 g for 5 min at 4°C. The cell pellet was re-suspended in 1% FBS for FACS analysis.

Tissue preparation

Mouse pups were sacrificed at PN7 or PN10. They were subjected to a standard protocol for lung inflation (25 cm) [20]. The right lungs were immediately flash frozen using liquid nitrogen and immediately stored at –80°C while the left lungs were fixed in 10% neutral buffered formalin and processed for histological analysis.

Oxygen exposure

For the exposure to hyperoxia (100% O₂), newborn (NB) mice (along with their mothers) were placed in cages in an airtight Plexiglas chamber (55 × 40 × 50 cm), as described previously [21–23]. Exposure to oxygen was initiated on PN1 and continued till PN7 of life. Two lactating dams were used. Mothers were alternated in hyperoxia and room air (RA) every 24 h. The litter size was kept limited to 12 pups to control for the effects of litter size on nutrition and growth. Throughout the experiment, they were given free access to food and water. Oxygen levels were constantly monitored by an oxygen sensor that was connected to a relay switch incorporated into the oxygen supply circuit. The inside of the chamber was kept at atmospheric pressure, and mice were exposed to a 12 h light–dark cycle.

Lung morphometry

H&E stained sections were photographed using an Olympus microscope with an in built digital camera system at 100× magnification (Olympus, Tokyo, Japan). Alveolar size was estimated from the mean chord length of the airspace, as previously described [19]. Alveolar septal wall thickness was estimated using Image J software, adapting the method described previously for bone trabecular thickness, for the lung [19]. 10× magnification images were obtained with Olympus 1X70 Microscope using Cellsens Dimension software. Radial alveolar count (RAC) and secondary septal crests were done as previously described [24,25].

TUNEL assay

For apoptosis quantification, TdT-mediated dUTP nick end labeling (TUNEL) staining was performed according to manufacturer's instructions keeping positive and negative controls (Roche Diagnostics). 20× magnification images were obtained with Olympus 1X70 Microscope using Cellsens Dimension software. Briefly 5 areas were selected (covers almost the entire histologic specimen) from each slide and 200 cells were counted using Cellsens software from Olympus microscope, followed by manually counting of TUNEL positive cells. The TUNEL index was

calculated by randomly selecting 3 of the 5 high-power fields in each slide, counting 200 cells in each area, and expressing the number of TUNEL-positive cells as a percentage [17].

Immunohistochemistry

Whole lungs were isolated from NB PN7 and PN10 mice and immediately fixed in 10% neutral buffered saline. Lung serial sections were stained with antibodies prosurfactant protein C (Catalog #AB3786, Millipore), vWF (DAKO), Myeloperoxidase (Biovision) and cleaved caspase 3 (Catalog #9661, Cell Signaling Technology). Sections were developed with Vectastain ABC kit (Catalog #PK4001, Vector laboratories) followed by hematoxylin staining (Catalog #H3401, Vector laboratories). 10× or 20× magnification images were obtained with Olympus IX70 Microscope using CellSens Dimension software and vWF staining was quantified by image J software.

Myeloperoxidase (MPO) ELISA from NB lung tissue

The snap-frozen lungs were thawed, weighed, and transferred to different tubes on ice containing 1 ml of Tissue Protein Extraction Reagent (T-PER; Pierce Biotechnology). Cells or lung tissues were homogenized at 4°C. Cell or lung homogenates were centrifuged at 10,000 g for 10 min at 4°C. Supernatants were transferred to clean microcentrifuge tubes. Total protein concentrations in the lung tissue homogenates were determined using a Biorad protein assay kit (Catalog #500-0006; Biorad) and MPO levels were evaluated in lung tissue homogenates using a mouse ELISA kit (Catalog #ab155458; Abcam) according to the user's manual.

Western blot analysis

NB mice whole lungs were harvested in lysis buffer and protein concentrations were determined using Bradford assay [19]. Protein extracts (both cellular and tissue homogenates) were loaded on to a Mini-Protean TGX gel electrophoretic system. Proteins were electroblotted onto an immunoblot PVDF membrane (Biorad catalog #162-0177). After transfer, non-specific binding was prevented with 5% skimmed milk TBS-Tween. After blocking, the membrane was incubated with anti-phospho AKT (ser 473) (Catalog #4060), anti-caspase-3 (Catalog #9662), and anti-cleaved caspase 3 (Catalog #9662) obtained from Cell Signaling Technology, USA. Anti-ANGPT1 (Catalog #AB10516), anti-ANGPT2 (Catalog #176002) and anti-Endoglin (Catalog #05-1424) were obtained from Millipore, USA. Chemiluminescent bands were visualized using SuperSignal West Pico or Femto substrate (Thermo Scientific, Rockford, IL). Rabbit polyclonal or monoclonal antibodies reactive to above proteins of interest were purchased from Cell Signaling Technology, Inc. (Danvers, MA). Quantification of western blots was

performed using Image J software version 1.46 (NIH, Bethesda, MD).

Multicolor flow cytometry

After the indicated duration of dox exposure to NB TGFβ1TG and TGFβ1TG X TGFβR2KO mice, they were sacrificed and their lungs were perfused through the right ventricle with 5 ml of ice cold PBS. The lungs were excised and single cell suspensions were prepared using digestion buffer as previously described [26]. Resultant single cell suspensions were stained for respective surface antigens and subsequently acquired on BD-LSRII flow cytometer. FACS analysis was performed with the use of FlowJo software.

Antibodies

The following monoclonal antibodies directed to the respective proteins, conjugated to different fluorochromes were used. Ly6c (clone: HK1.4), F4/80 (clone: BM8), CD45 (clone: 30-F11), CD103 (clone: 2E7) and MHC-II (clone: M5/114.152) were purchased from eBioscience (San Diego, CA). CD11c (clone: HL3) and CD11b (clone: M1/70) were purchased from BD Biosciences (San Jose, CA). CD206 (clone: C068C2) was purchased from BioLegend (San Diego, CA). All the antibodies that were used for FACS were monoclonals and highly specific. These were the concentrations of antibodies: CD11c (1 μg/ml), CD11b (0.05 μg/ml), F4/80 (0.2 μg/ml), CD206 (0.2 μg/ml), Ly6c (0.05 μg/ml), MHC-II (0.05 μg/ml).

Statistical analysis

All statistical analyses were performed using Graph Pad Prism, version 5.0 (GraphPad Software, San Diego CA). The data were expressed as the mean ± SEM of a minimum of 3 independent experiments with a minimum of 4 mice in each group. Groups were compared with the Student's two-tailed unpaired t test, 1-way ANOVA (followed by Tukey's Multiple Comparison post-hoc test) or the Kaplan Meier Survival Analysis, as appropriate, using GraphPad Prism 5.0. A 'p' value of <0.05 was considered statistically significant in all tests.

Results

Effect on survival, in room air, of lung epithelial cell-specific conditionally overexpressing TGFβ1TG mice lacking lung epithelial-cell specific TGFβR2

Since NB TGFβ1TG mice are known to have a high mortality upon dox activation in room air, in the first PN week, as previously reported [3], we evaluated the role of TGFβ1-signaling via TGFβR2 in lung epithelial cells *in vivo* on survival. Hence, we assessed survival in TGFβ1TG mice lacking TGFβR2, upon TGFβ1 activation from PN1 – PN7. TGFβ1TG X TGFβR2KO mice showed

significantly lower mortality as compared to lung targeted overexpressing TGF β 1TG mice (Figure 1).

This suggests that lung-mediated signaling via TGF β R2 has such a significant effect that it can overcome the dramatic mortality that has been observed when increased concentrations of TGF β 1 are present at these critical (saccular/early alveolar) stages of lung development.

Characterization of pulmonary (lung tissue and BALF) inflammatory myeloid compartment, in room air, of lung epithelial cell-specific conditionally overexpressing TGF β 1TG mice lacking lung epithelial-cell specific TGF β R2

It was initially reported that TGF β 1 overexpression in mice lungs leads to mononuclear-rich inflammation [16]. Since our aim in the present manuscript was to understand TGF β 1-mediated immune signaling in hyperoxia-induced cell death in neonates, we decided to focus on evaluating the macrophage/monocyte populations in the lung by flow cytometry. We hypothesized that reduced inflammation in lung tissue myeloid compartment could lead to TGF β 1TG X TGF β R2KO mice survival, and hence we evaluated this response. The mononuclear phagocytic system (monocytes, macrophages and dendritic cells) collectively plays a critical role in the maintenance of tissue integrity [27]. To evaluate the role of the mononuclear phagocytic system, TGF β 1TG, WT and TGF β 1TG X TGF β R2KO mice lungs and BALF cells were phenotypically analyzed for pro-inflammatory macrophages and lung tissue resident macrophages. Monocytes were analyzed for Ly6C^{high} and Ly6C^{low} cells. Lung pro-inflammatory macrophages resemble M1-like macrophages and are CD45⁺ F4/80⁺ CD206⁻ and CD11c⁻, whereas tissue resident macrophages resemble the M2 phenotype, expressing CD45⁺ F4/80⁺ CD206⁺ and CD11c⁺.

We found significant reduction in inflammatory macrophages and Ly6C^{high} monocytes in the lung tissue of the TGF β 1TG X TGF β R2KO mice, compared to TGF β 1TG

mice. No such differences in inflammatory macrophages and Ly6C^{high} monocytes were observed when compared to WT littermate lungs. The numbers of tissue resident macrophages were slightly reduced in the lungs of TGF β 1TG X TGF β R2KO mice when compared to WT littermates. We observed no differences in Ly6C^{low} monocytes among the groups (Figure 2A). Ly6C^{low} monocytes were significantly decreased in the BALF of TGF β 1TG X TGF β R2KO and WT mice, as compared with TGF β 1TG mice (Figure 2B).

Taken together, the above data suggests that signaling via TGF β R2, at least in part, mediates the pulmonary inflammatory response of TGF β 1 in developing lungs.

Effect on lung apoptotic cell death specifically in Type II alveolar epithelial cells (AECs), in room air, of lung epithelial cell-specific conditionally overexpressing TGF β 1TG mice lacking lung epithelial-cell specific TGF β R2

Next, we evaluated cell death in lung targeted overexpressing TGF β 1TG mice and TGF β 1TG X TGF β R2KO mice from PN7 – PN10. We used TUNEL and immunohistochemistry respectively. TGF β 1TG X TGF β R2KO mice showed significantly decreased TUNEL positive index and decreased cleaved caspase 3 staining in Type II AECs (surfactant protein C positive), as compared to TGF β 1TG mice (Figure 3A – C).

Effect on lung morphometry and vessel density, in room air, of lung epithelial cell-specific conditionally overexpressing TGF β 1TG mice lacking lung epithelial-cell specific TGF β R2

Lung architecture was assessed and chord length as well as septal thickness were found to be significantly decreased in TGF β 1TG X TGF β R2KO as compared to TGF β 1TG mice (Figure 4A – C). In concordance, we found the RAC and secondary septal crest numbers as well as vessel density improved in the TGF β 1TG X TGF β R2KO as compared to TGF β 1TG mice (Figure 4D – G). This suggested that absence of TGF β R2 in TGF β 1 signaling




Figure 1 Effect of TGF β 1 exposure on overall survival in maternally exposed doxycycline to NB TGF β 1TG, and TGF β 1TG X TGF β R2KO mice. NB TGF β 1TG, and TGF β 1TG X TGF β R2KO littermate mice were exposed to maternal dox at PN1 - PN7 and then overall survival was assessed. Where NB: newborn; PN: postnatal; TGF β 1TG: TGF β 1 transgenic mice; TGF β 1TG X TGF β R2KO: TGF β 1 transgene mice crossed with TGF β receptor II knockout mice. N = 7 in each group; ***p < 0.01 and ***p < 0.001 TGF β 1TG vs. TGF β 1TG X TGF β R2KO and TGF β R2KO vs. TGF β 1TG respectively.





Figure 2 TGFβ1-induced inflammation in TGFβ1TG and TGFβ1TG X TGFβR2KO mice. Multicolor flow cytometric analysis showed significantly greater staining of inflammatory macrophages and LY6C high monocytes in TGFβ1TG mice lungs as compared to WT controls. TGFβ1TG X TGFβR2KO developing lungs showed significantly decreased staining of inflammatory macrophages and LY6C high monocytes as compared to TGFβ1TG mice. **(B)** Ly6C low monocytes were significantly decreased in the BALF of WT mice, as compared with TGFβ1TG mice. BALF: bronchoalveolar lavage fluid. N = 4 in each group; *p < 0.05, **p < 0.01 and ***p < 0.001.

resulted in decreased apoptotic cell death and preserved alveolar architecture as well as vascular development.

Effect on molecular mediators, in room air, of lung epithelial cell-specific conditionally overexpressing TGFβ1TG mice lacking lung epithelial-cell specific TGFβR2
To further understand the mechanism of TGFβ1-induced cell death, we examined molecular markers

including endoglin (or TGFβR3), ANGPT1, ANGPT2, Phospho-AKT (Ser 473), BAX and cleaved caspase 3. We noted that Phospho-AKT was increased while endoglin, BAX, caspase 3 and cleaved caspase 3 were decreased in TGFβ1TG X TGFβR2KO as compared to TGFβ1TG mice lungs (Figure 5A). Densitometric analysis showed significantly increased ANGPT1 to ANGPT2 ratio in TGFβ1 exposed NB TGFβTG X



TGFβR2KO lungs as compared to TGFβ1TG mice lungs (Figure 5B).

This suggested that the decreased apoptotic cell death response could be due to increased presence of molecular mediators of cell survival in the absence of TGFβR2 in TGFβ1 signaling processes in developing lungs.

Effect on lung architecture, upon hyperoxia exposure, in TGFβR2KO mice

Given the fact that hyperoxia exposure is known to increase TGFβ in developing lungs and lead to abnormal alveolarization [1,11,13,28,29], we finally examined if lack of signaling via its putative receptor TGFβR2, upon




Figure 4 Effect of TGFβ1 on lung morphometry. **(A)** Representative images of lung histology (H&E stain) of NB TGFβ1TG mice and TGFβ1TG X TGFβR2KO mice. Scale bar: 100 μm **(B)** Bar graph showing the mean chord length in TGFβ1TG mice and TGFβ1TG X TGFβR2KO mice. **(C)** Bar graph showing the septal thickness in TGFβ1TG mice and TGFβ1TG X TGFβR2KO mice. **(D)** Bar graph showing the radial alveolar count in TGFβ1TG mice and TGFβ1TG X TGFβR2KO mice. **(E)** Bar graph showing the secondary septal crest numbers in TGFβ1TG mice and TGFβ1TG X TGFβR2KO mice. **(F)** Representative images of lung immunohistochemistry of vessel density (vWF staining) of NB TGFβ1TG mice and TGFβ1TG X TGFβR2KO mice. Scale bar: 100 μm **(G)** Bar graph showing quantification of vessel density in TGFβ1TG mice and TGFβ1TG X TGFβR2KO mice. N = 4, in each group; *p < 0.05, **p < 0.01.





Figure 5 TGFβ1 signaling in NB TGFβ1TG, and TGFβ1TG X TGFβ2KO mice lungs. **(A)** Western blot analysis showed upregulation of phospho AKT, and down regulation of ANGPT2, and pro-apoptotic proteins (BAX; cleaved caspase 3) in NB TGFβ1TG X TGFβ2KO lungs as compared to TGFβ1TG mice. **(B)** Densitometric analysis showed increased ANGPT1 to ANGPT2 ratio in maternal dox exposed NB TGFβ1TG X TGFβ2KO lungs as compared to TGFβ1TG mice. N = 3, in each group; *p < 0.05.

hyperoxia exposure, would have an impact on the pulmonary phenotype. Hence, we assessed the lung architecture in TGFβ2KO mice exposed to hyperoxia from PN1 – PN7 and confirmed our histologic observations by chord length and septal thickness measurements. Chord length and septal thickness were significantly decreased in hyperoxia

exposed TGFβ2KO mice as compared to TGFβ2 flox (control) mice (Figure 6A - C). In addition, we noted a significant decrease in the homing of inflammatory cells (neutrophils), as detected by MPO staining (Figure 6D - E). This protective effect in the hyperoxia-exposed TGFβ2KO mice lungs was further confirmation of our



results of TGFβ1-induced signaling being mediated through this receptor having a significant impact on TGFβ1-mediated effects in developing lungs.


To summarize, we show that the detrimental effects of increased mortality, pulmonary inflammation, cell death and impaired alveolarization of enhanced lung epithelial cell specific TGFβ1 signaling in developing lungs are mediated, in part, via TGFβR2. In addition, specific molecular mediators of cell survival are associated with these responses in developing lungs. A proposed model of these molecular mediators leading to their ultimate effects has been shown in Figure 7.

Discussion

In the present study, lung epithelial cell specific targeted conditional TGFβ1TG and epithelial cell specific TGFβR2 conditional null mutant mice were used to

explore the molecular mechanisms of TGFβ1-induced injury in developing lungs. This context is particularly important for the study of HALI and BPD that arise in the critical developing stages of lungs, with abnormal TGFβ1 signaling *in vivo* [30]. Although the pathogenesis of BPD is not well understood, increased TGFβ (secondary to hyperoxia exposure) signaling in the saccular/early alveolar stages is said to be one of the causative factors for the development of impaired alveolarization secondary to inflammation [1,31,32].

High levels of TGFβ1 (as well as β2 and β3) immunoreactivity have been reported in the developing lung in the bronchial and alveolar epithelium of the NB mouse [13]. Furthermore, hyperoxia exposure increases TGFβ-signaling [13]. A TGFβ-neutralizing antibody (1D11) was effective in improving alveolarization, extracellular cellular matrix assembly and microvascular development in NB mice



exposed to hyperoxia (85% O₂) for 10 PN days [13]. However, 1D11 is a pan-specific TGFβ IgG1 antibody that targets all the 3 isoforms of TGFβ i.e. 1–3 [13]. In our earlier report, exposure of hyperoxia (100% O₂) to the lung epithelial cell targeted TGFβ1TG mice led to increased mortality, which was significantly diminished by using a c-JunNH2-terminal kinase (JNK) pathway inhibitor [17]. In addition, the JNK-pathway inhibitor was able to ameliorate the BPD pulmonary phenotype of impaired alveolarization in the TGFβ1TG mice [17]. Chronic hyperoxia (85% O₂) exposure from PN1 to PN28 in mice led to a 4-fold increase of TGFβR2 (among other TGFβ signaling molecules), along with impaired alveolarization characterized by approximately doubling of the mean linear intercept, in the lung [1].

Our work extends the above observations by providing specificity of the effects of the TGFβ1 isoform as well as its signaling via the specific receptor TGFβR2. In addition, by using lung epithelial cell targeted overexpression and null mutant mouse models, respectively, we are able to localize the effects to be secondary to the production of TGFβ1 and inhibition of its signaling in particular cell types in the developing lung.

As mentioned above, hyperoxia contributes significantly to the development of HALI and BPD in neonates [33]. To this end, we were able to demonstrate the specificity of such a protective response, as evidenced by TGFβ1TG X TGFβR2KO mice having improved survival (up to PN6) as compared to PN1 of TGFβ1TG mice, in room air. The fact that this survival advantage was noted in room air lends credence to our hypothesis that this is a specific

TGFβ1-induced TGFβR2-mediated effect, and not confounded potentially by other mediators that are activated/released upon hyperoxia exposure.

Other investigators have reported that TGFβ1 induced inflammation in adult lung injury model systems [34], but we were unable to locate any such effects reported in NB mice. Adding novel information, we demonstrated diminished pulmonary inflammation in TGFβ1TG mice in which TGFβR2 signaling had been abrogated. In addition, we were able to demonstrate that specifically it was the accumulation of inflammatory macrophages and monocytes in the lungs of TGFβ1TG mice. In contrast, TGFβ1TG X TGFβR2KO NB mice were found to have diminished levels of pro-inflammatory macrophages.

We have reported earlier that hyperoxia exposure increases TGFβ1 expression as well as cell death in A549 cells upon hyperoxia exposure in a dose-dependent manner [17]. Others have shown that exposure to hyperoxia of primary alveolar epithelial Type II cells led to increased sensitivity to TGFβ-induced apoptosis [1]. Furthermore, we provide data that worsening of lung morphometric indices (chord length, septal thickness, RAC and secondary septal crest numbers) and decreased vessel density in the TGFβ1TG mice were dependent, in part, by signaling via TGFβR2. In support of these observations, we provide novel mechanistic *in vivo* data of increased apoptosis via the mitochondrial active BAX dependent pathway in the presence of increased TGFβ1 in developing lungs, which is abrogated by concomitant inhibition of TGFβR2 signaling.

We have previously reported that TGFβ1TG mice lungs have increased expression of apoptotic cell death mediators, specifically Fas-L and caspase-3 [17]. In the present study, we extend those observations by showing increased BAX and cleaved caspase-3.

We found increased ANGPT2, and Phospho-AKT (Ser 473), but no change in ANGPT1. We have previously reported on the role of increased ANGPT2 as a critical contributing factor in HALI in adult [14,35] and neonatal [19,23] mice. In addition, we have also noted a significant association of increased ANGPT2 in hyperoxia-induced human diseases in adults (ALI) [14] as well as neonates (BPD) [14,15]. Since TGFβ1 has been shown to be associated with BPD and overexpression lung models of TGFβ1 mimic the pulmonary phenotype of BPD as reported by us [17] and others [3,36], our data implicates ANGPT2 as a novel downstream molecule amenable to potential therapeutic modulation in HALI in the NB lung. Since ANGPT1 levels did not change and both ANGPT 1 and 2 signal through the Tie2 receptor, our data suggest that targeting the ligand ANGPT2 in HALI would be the preferred option in such a scenario. Endoglin is part of the TGFβR complex and has been shown to be increased in human BPD [37]. Our data would suggest that TGFβ1 signaling increases the expression of endoglin. AKT is a well-known survival factor

reported to have a role in different experimental models of BPD [38–41]; we now implicate AKT as a down streaming signaling molecule in our TGF β 1 overexpressing mouse lung model of BPD. We further confirmed the specificity of the TGF β 1 signaling pathway by noting the reversal of the effects noted above in the TGF β 1TG X TGF β R2KO mice lungs.

Among other models of BPD, cyclical stretch during invasive mechanical ventilation in NB mice in room air and/or hyperoxia (40% O₂) has also been reported to increase TGF β -signaling and apoptotic cell death, including cleaved caspase 3 [42–44]. Activation of AKT has been shown to protect NB rats from HALI and BPD [39].

Investigators have reported additional molecules in NB mice lungs that have been associated with inhibition of TGF β -signaling to improve the pulmonary phenotype relevant to HALI and BPD [11,45–47]. As also mentioned earlier, increased pulmonary levels of TGF β have been reported in association with human BPD [9,48,49]. Interestingly, increased endoglin [37], decreased ANGPT1 [37], and increased ANGPT2 [14,15] in the NB lung have all been associated with human BPD.

To explore the wider importance, we finally subjected TGF β R2KO mice to hyperoxia exposure from PN1 – PN7. Lung architecture (as measured by chord length and septal thickness) was significantly decreased (with equivalency to room air controls), in addition to decreased inflammation, in TGF β R2KO hyperoxia-exposed mice lungs suggesting inhibition of TGF β 1 signaling mediated via its receptor TGF β R2 has a protective role in hyperoxia settings.

In conclusion, our data provides evidence suggesting TGF β R2KO comprehensively prevents the deleterious effects of TGF β 1 signaling by increased ANGPT1/ANGPT2 ratio, phosphorylation of AKT and decreased cleavage caspase 3 expression (including specifically in Type II AECs). Our study has demonstrated the potential role of inhibition of TGF β 1 signaling via TGF β R2 for improved survival, reduced inflammation and apoptosis that may provide insights for the development of novel therapeutic strategies targeted against HALI and BPD.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

Concept and Design: AS, VB. Acquisition of Data: AS, MS, CB. Data Analysis and Interpretation: AS, MS, CB, MD, RH, PM, VB. Drafting and Critical Revision for intellectual content: AS, MS, CB, MD, VB. All authors have approved the version of the submitted manuscript.

Acknowledgements

Supported in part by grants CA106802 (M.V.D.) and CA135110 (M.V.D.) from the NCI of the National Institutes of Health, USA; The Hastings Foundation (P.M.); grants HL107307 (P.M.) and HL85103 (V.B.) from the NHLBI of the National Institutes of Health, USA.

Author details

¹Department of Pediatrics, Yale University School of Medicine, 333 Cedar Street, New Haven, CT 06510, USA. ²Department of Medicine and Yale Cancer Center, Yale University School of Medicine, 333 Cedar Street, New Haven, CT 06510, USA. ³Department of Pathology, Yale University School of Medicine, 333 Cedar Street, New Haven, CT 06520, USA. ⁴Department of Pediatrics, University of Southern California, 1200 North State Street, Los Angeles, CA 90033, USA.

Received: 27 August 2014 Accepted: 19 December 2014

Published online: 16 January 2015

References

- Alejandro-Alcazar MA, Kwapiszewski G, Reiss I, Amarie OV, Marsh LM, Sevilla-Perez J, et al. Hyperoxia modulates TGF-beta/BMP signaling in a mouse model of bronchopulmonary dysplasia. *Am J Physiol Lung Cell Mol Physiol*. 2007;292:L537–49.
- Jankov RP, Keith Tanswell A. Growth factors, postnatal lung growth and bronchopulmonary dysplasia. *Paediatr Respir Rev*. 2004;5 Suppl A:S265–75.
- Vicencio AG, Lee CG, Cho SJ, Eickelberg O, Chuu Y, Haddad GG, et al. Conditional overexpression of bioactive transforming growth factor-beta1 in neonatal mouse lung: a new model for bronchopulmonary dysplasia? *Am J Respir Cell Mol Biol*. 2004;31:650–6.
- Ganeshan K, Johnston LK, Bryce PJ. TGF-beta1 limits the onset of innate lung inflammation by promoting mast cell-derived IL-6. *J Immunol*. 2013;190:5731–8.
- Dabovic B, Chen Y, Choi J, Davis EC, Sakai LY, Todorovic V, et al. Control of lung development by latent TGF-beta binding proteins. *J Cell Physiol*. 2011;226:1499–509.
- Bhandari A, Bhandari V. "New" bronchopulmonary dysplasia - a clinical review. *Clin Pulm Med*. 2011;18:137–43.
- Jugdutt BI. Ventricular remodeling after infarction and the extracellular collagen matrix: when is enough? *Circulation*. 2003;108:1395–403.
- Kotecha S. Cytokines in chronic lung disease of prematurity. *Eur J Pediatr*. 1996;155 Suppl 2:S14–7.
- Lecart C, Cayabyab R, Buckley S, Morrison J, Kwong KY, Warburton D, et al. Bioactive transforming growth factor-beta in the lungs of extremely low birthweight neonates predicts the need for home oxygen supplementation. *Biol Neonate*. 2000;77:217–23.
- Zhao Y, Gilmore BJ, Young SL. Expression of transforming growth factor-beta receptors during hyperoxia-induced lung injury and repair. *Am J Physiol*. 1997;273:L355–62.
- Dasgupta C, Sakurai R, Wang Y, Guo P, Ambalavanan N, Torday JS, et al. Hyperoxia-induced neonatal rat lung injury involves activation of TGF-beta and Wnt signaling and is protected by rosiglitazone. *Am J Physiol Lung Cell Mol Physiol*. 2009;296:L1031–41.
- Speer CP. Pulmonary inflammation and bronchopulmonary dysplasia. *J Perinatol*. 2006;26 Suppl 1:S57–62. discussion S63–54.
- Nakanishi H, Sugiura T, Streisand JB, Lonning SM, Roberts Jr JD. TGF-beta-neutralizing antibodies improve pulmonary alveologenesis and vasculogenesis in the injured newborn lung. *Am J Physiol Lung Cell Mol Physiol*. 2007;293:L151–61.
- Bhandari V, Choo-Wing R, Lee CG, Zhu Z, Nedrelow JH, Chupp GL, et al. Hyperoxia causes angiotensin 2-mediated acute lung injury and necrotic cell death. *Nat Med*. 2006;12:1286–93.
- Aghai ZH, Faqiri S, Saslow JG, Nakhla T, Farhath S, Kumar A, et al. Angiotensin 2 concentrations in infants developing bronchopulmonary dysplasia: attenuation by dexamethasone. *J Perinatol*. 2008;28:149–55.
- Lee CG, Cho SJ, Kang MJ, Chapoval SP, Lee PJ, Noble PW, et al. Early growth response gene 1-mediated apoptosis is essential for transforming growth factor beta1-induced pulmonary fibrosis. *J Exp Med*. 2004;200:377–89.
- Li Z, Choo-Wing R, Sun H, Sureshbabu A, Sakurai R, Rehan VK, et al. A potential role of the JNK pathway in hyperoxia-induced cell death, myofibroblast transdifferentiation and TGF-beta1-mediated injury in the developing murine lung. *BMC Cell Biol*. 2011;12:54.
- Li M, Krishnaveni MS, Li C, Zhou B, Xing Y, Banfalvi A, et al. Epithelium-specific deletion of TGF-beta receptor type II protects mice from bleomycin-induced pulmonary fibrosis. *J Clin Invest*. 2011;121:277–87.
- Sun H, Choo-Wing R, Sureshbabu A, Fan J, Leng L, Yu S, et al. A critical regulatory role for macrophage migration inhibitory factor in hyperoxia-induced injury in the developing murine lung. *PLoS One*. 2013;8:e60560.

20. Choo-Wing R, Syed MA, Harijith A, Bowen B, Pryhuber G, Janer C, et al. Hyperoxia and interferon-gamma-induced injury in developing lungs occur via cyclooxygenase-2 and the endoplasmic reticulum stress-dependent pathway. *Am J Respir Cell Mol Biol*. 2013;48:749–57.
21. Harijith A, Choo-Wing R, Cataltepe S, Yasumatsu R, Aghai ZH, Janer J, et al. A role for matrix metalloproteinase 9 in IFN γ -mediated injury in developing lungs: relevance to bronchopulmonary dysplasia. *Am J Respir Cell Mol Biol*. 2011;44:621–30.
22. Bhandari V, Choo-Wing R, Lee CG, Yusuf K, Nedrelow JH, Ambalavanan N, et al. Developmental regulation of NO-mediated VEGF-induced effects in the lung. *Am J Respir Cell Mol Biol*. 2008;39:420–30.
23. Choo-Wing R, Nedrelow JH, Homer RJ, Elias JA, Bhandari V. Developmental differences in the responses of IL-6 and IL-13 transgenic mice exposed to hyperoxia. *Am J Physiol Lung Cell Mol Physiol*. 2007;293:L142–50.
24. Cooney TP, Thurlbeck WM. The radial alveolar count method of Emery and Mithal: a reappraisal 1–postnatal lung growth. *Thorax*. 1982;37:572–9.
25. Nicola T, Hagood JS, James ML, Macewen MW, Williams TA, Hewitt MM, et al. Loss of Thy-1 inhibits alveolar development in the newborn mouse lung. *Am J Physiol Lung Cell Mol Physiol*. 2009;296:L738–50.
26. Wang Y, Huang Z, Nayak PS, Sanchez-Esteban J: An experimental system to study mechanotransduction in fetal lung cells. *J Vis Exp* 2012. doi:10.3791/3543.
27. Geissmann F, Manz MG, Jung S, Sieweke MH, Merad M, Ley K. Development of monocytes, macrophages, and dendritic cells. *Science*. 2010;327:656–61.
28. Dang H, Yang L, Wang S, Fang F, Xu F. Calcitonin gene-related peptide ameliorates hyperoxia-induced lung injury in neonatal rats. *Tohoku J Exp Med*. 2012;227:129–38.
29. Witsch TJ, Turowski P, Sakkas E, Niess G, Becker S, Herold S, et al. Deregulation of the lysyl hydroxylase matrix cross-linking system in experimental and clinical bronchopulmonary dysplasia. *Am J Physiol Lung Cell Mol Physiol*. 2014;306:L246–59.
30. Warburton D, Schwarz M, Tefft D, Flores-Delgado G, Anderson KD, Cardoso WV. The molecular basis of lung morphogenesis. *Mech Dev*. 2000;92:55–81.
31. Sakurai R, Villarreal P, Husain S, Liu J, Sakurai T, Tou E, et al. Curcumin protects the developing lung against long-term hyperoxic injury. *Am J Physiol Lung Cell Mol Physiol*. 2013;305:L301–11.
32. Bhandari V. Postnatal inflammation in the pathogenesis of bronchopulmonary dysplasia. *Birth Defects Res A Clin Mol Teratol*. 2014;100:189–201.
33. Bhandari V. Hyperoxia-derived lung damage in preterm infants. *Semin Fetal Neonatal Med*. 2010;15:223–9.
34. Murray LA, Chen Q, Kramer MS, Hesson DP, Argentieri RL, Peng X, et al. TGF- β driven lung fibrosis is macrophage dependent and blocked by Serum amyloid P. *Int J Biochem Cell Biol*. 2011;43:154–62.
35. Bhandari V, Choo-Wing R, Harijith A, Sun H, Syed MA, Homer RJ, et al. Increased hyperoxia-induced lung injury in nitric oxide synthase 2 null mice is mediated via angiotensin II. *Am J Respir Cell Mol Biol*. 2012;46:668–76.
36. Gaudie J, Galt T, Bonniaud P, Robbins C, Kelly M, Warburton D. Transfer of the active form of transforming growth factor- β 1 gene to newborn rat lung induces changes consistent with bronchopulmonary dysplasia. *Am J Pathol*. 2003;163:2575–84.
37. De Paepe ME, Patel C, Tsai A, Gundavarapu S, Mao Q. Endoglin (CD105) up-regulation in pulmonary microvasculature of ventilated preterm infants. *Am J Respir Crit Care Med*. 2008;178:180–7.
38. Wu S, Platteau A, Chen S, McNamara G, Whitsett J, Bancalari E. Conditional overexpression of connective tissue growth factor disrupts postnatal lung development. *Am J Respir Cell Mol Biol*. 2010;42:552–63.
39. Alphonse RS, Vadivel A, Coltan L, Eaton F, Barr AJ, Dyck JR, et al. Activation of Akt protects alveoli from neonatal oxygen-induced lung injury. *Am J Respir Cell Mol Biol*. 2011;44:146–54.
40. Park HS, Park JW, Kim HJ, Choi CW, Lee HJ, Kim BI, et al. Sildenafil alleviates bronchopulmonary dysplasia in neonatal rats by activating the hypoxia-inducible factor signaling pathway. *Am J Respir Cell Mol Biol*. 2013;48:105–13.
41. Madurga A, Mizikova I, Ruiz-Camp J, Vadasz I, Herold S, Mayer K, et al. Systemic hydrogen sulfide administration partially restores normal alveolarization in an experimental animal model of bronchopulmonary dysplasia. *Am J Physiol Lung Cell Mol Physiol*. 2014;306:L684–97.
42. Mokres LM, Parai K, Hilgendorff A, Ertsey R, Alvira CM, Rabinovitch M, et al. Prolonged mechanical ventilation with air induces apoptosis and causes failure of alveolar septation and angiogenesis in lungs of newborn mice. *Am J Physiol Lung Cell Mol Physiol*. 2010;298:L23–35.
43. Hilgendorff A, Parai K, Ertsey R, Jain N, Navarro EF, Peterson JL, et al. Inhibiting lung elastase activity enables lung growth in mechanically ventilated newborn mice. *Am J Respir Crit Care Med*. 2011;184:537–46.
44. Hilgendorff A, Parai K, Ertsey R, Juliana Rey-Parra G, Thebaud B, Tamosiuniene R, et al. Neonatal mice genetically modified to express the elastase inhibitor elafin are protected against the adverse effects of mechanical ventilation on lung growth. *Am J Physiol Lung Cell Mol Physiol*. 2012;303:L215–27.
45. Bachiller PR, Nakanishi H, Roberts Jr JD. Transforming growth factor- β modulates the expression of nitric oxide signaling enzymes in the injured developing lung and in vascular smooth muscle cells. *Am J Physiol Lung Cell Mol Physiol*. 2010;298:L324–34.
46. Bozyk PD, Bentley JK, Popova AP, Anyanwu AC, Linn MD, Goldsmith AM, et al. Neonatal periostin knockout mice are protected from hyperoxia-induced alveolar simplification. *PLoS One*. 2012;7:e31336.
47. Vosdoganes P, Lim R, Koulaeva E, Chan ST, Acharya R, Moss TJ, et al. Human amnion epithelial cells modulate hyperoxia-induced neonatal lung injury in mice. *Cytotherapy*. 2013;15:1021–9.
48. Kotecha S, Wangoo A, Silverman M, Shaw RJ. Increase in the concentration of transforming growth factor β -1 in bronchoalveolar lavage fluid before development of chronic lung disease of prematurity. *J Pediatr*. 1996;128:464–9.
49. Bhandari A, Bhandari V. Biomarkers in bronchopulmonary dysplasia. *Paediatr Respir Rev*. 2013;14:173–9.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

