

Two new species of the genus *Chilobrachys* (Araneae, Theraphosidae) from China

Yejie Lin¹, Xunyou Yan¹, Shuqiang Li²

1 Hebei Key Laboratory of Animal Diversity, College of Life Science, Langfang Normal University, Langfang 065000, China **2** Institute of Zoology, Chinese Academy of Sciences, Beijing 100101, China

Corresponding authors: Xunyou Yan (yanxunyou@163.com), Shuqiang Li (lisq@ioz.ac.cn)

Academic editor: Zhiyuan Yao | Received 26 October 2021 | Accepted 15 December 2021 | Published 13 January 2022

<http://zoobank.org/729EAE0B-66E6-4AA4-BDAE-19566B5C176F>

Citation: Lin Y, Yan X, Li S (2022) Two new species of the genus *Chilobrachys* (Araneae, Theraphosidae) from China. ZooKeys 1081: 99–109. <https://doi.org/10.3897/zookeys.1081.77072>

Abstract

Two new species of the genus *Chilobrachys* Karsh, 1892 are described from China: *Chilobrachys dominus* Lin & Li, **sp. nov.** from Yunnan and *C. jinchengi* Lin & Li, **sp. nov.** from Tibet. Photos and a morphological description of the new species are given. The type specimens of the new species are deposited in the Institute of Zoology, Chinese Academy (IZCAS) in Beijing.

Keywords

Asia, diagnosis, taxonomy, type

Introduction

The family Theraphosidae Thorell, 1869 is one of the largest spider families in the world, including 1031 species in 153 genera (WSC 2021). This family is divided into 12 subfamilies (Foley et al. 2019), of which Selenocosmiinae Simon 1889 and Ornithoctoninae Pocock, 1895 are known from Asia.

Selenocosmiinae includes 125 species and subspecies in 12 genera. It can be distinguished from other subfamilies by the maxilla with an oval stridulatory organ on the prolateral surface, originating medially (not ventrally from the oral fringe), the maxilla with opposing modified setae in multiple rows on the retrolateral cheliceral surface,

the posterior sternal sigillae distant from the sternal margins, the more than 60 labial cuspules, and the leg spines only found distally on the metatarsi; mature males lack a tibial apophysis on leg I (Nunn et al. 2016). In the genus *Chilobrachys* Karsh, 1892, the anterior eyes are nearly in a straight line, the stridulating organ consists of short spines on the chelicerae and a single or double row of paddle hairs overlapped by a fringe of hairs on the maxillae, and the palpal organ of the male ending in a long and slender embolus (Raven 1985; Zhu and Zhang 2008).

Four *Chilobrachys* species were hitherto known from China (Fig. 8, Li 2020; Li et al. 2021; Yao et al. 2021): *C. guangxiensis* (Yin & Tan, 2000) (Guangxi, Hainan), *C. hubei* Song & Zhao, 1988 (Hubei, Chongqing), *C. liboensis* Zhu & Zhang, 2008 (Guizhou, Guangxi) and *C. lubricus* Yu et al., 2021 (Yunnan). Here, we describe two new species: *Chilobrachys dominus* sp. nov. from Xishuangbanna, Yunnan and *Chilobrachys jinchengi* sp. nov. from Medog, Tibet.

Materials and methods

All specimens are preserved in 75% ethanol and deposited in the Institute of Zoology, Chinese Academy of Sciences in Beijing (**IZCAS**). Spermathecae were cleared in a trypsin enzyme solution to dissolve non-chitinous tissues. Specimens were examined under a LEICA M205C stereomicroscope. Photomicroscope images were taken with an Olympus C7070 zoom digital camera (7.1 megapixels). Photographs were stacked with Helicon Focus 6.7.1 and processed in Adobe Photoshop CC 2018.

The terminology used in the text and figures follows Bertani (2000). All measurements are in millimetres. Eye sizes were measured as the maximum diameter in either a dorsal or frontal view. Leg measurements are given as follows: total length (femur, patella, tibia, metatarsus, tarsus).

Abbreviations

A	apical keel;	PI	prolateral inferior keel;
ALE	anterior lateral eyes;	PLE	posterior lateral eyes;
AME	anterior median eyes;	PME	posterior median eyes;
MOA	median ocular area;	PS	prolateral superior keel.

Taxonomy

Family Theraphosidae Thorell, 1869

Subfamily Selenocosmiinae Simon, 1889

Genus *Chilobrachys* Karsh, 1892

Type species. *Chilobrachys nitelinus* Karsch, 1892.

Diagnosis. See Raven (1985) and Zhu and Zhang (2008).

***Chilobrachys dominus* Lin & Li, sp. nov.**

<http://zoobank.org/36EC608E-6671-407E-9E1D-10C5497BADCO>

Figs 1A, 2A–H, 3A, B, 4A–C, 8

Type material. *Holotype* ♂ (IZCAS-Ar42676), CHINA: *Yunnan*: Jinghong, Mount Jinuo, Jinuo Road, 22.0556°N, 100.9853°E, elevation 1040 m, 16.XI.2021, Yi Ming leg.



Figure 1. *Chilobrachys* spp. nov., live males **A** *Chilobrachys dominus* sp. nov. **B** *C. jinchengi* sp. nov. Photos by Qianle Lu, Zhengzhong Huang.




Figure 2. *Chilobrachys dominus* sp. nov., holotype male **A** male habitus, dorsal **B** ocular tubercle **C** right palp maxillae **D** sternum **E** stridulatory lyra, lateral view **F** same, ventral view **G** chelicerae, retrolateral view **H** same, prolateral view.



Figure 3. *Chilobrachys dominus* sp. nov., holotype, right palp bulb, rotated horizontally **A** retrolateral view **B** proteral view.

Diagnosis. The male of *Chilobrachys dominus* sp. nov. is similar to that of *C. pococki* (Thorell, 1897) in having a long, strongly curved embolus, but can be distinguished by the embolus bent in at two thirds (vs. halfway in *C. pococki*).

Description. Male (holotype, IZCAS-Ar42676) (Figs 2B–H, 3, 4). Carapace 13.27 long, 11.89 wide, black, with long purple setae. Opisthosoma black, with long



Figure 4. *Chilobrachys dominus* sp. nov., holotype, left palp **A** prolateral view **B** ventral view **C** retrolateral view.

black setae. Eye group 2.08 long, 1.04 wide (Fig. 2F). MOA 0.81 long, anterior width 1.05, posterior width 1.53. Eye sizes and interdistances: ALE 0.52, AME 0.44, PLE 0.37, PME 0.37; ALE–AME 0.17, AME–AME 0.24, PLE–PME 0.02, PME–PME 0.90. Fovea slightly procurved. Chelicerae black, with row of 14 promarginal teeth. Labium wider than long, with 368 cuspules. Sternum yellow brown with 3 pairs of sigilla. Legs with long and short setae. Tarsi I–III with 2 claws, tarsus IV with 3 claws, denticle number: I 4, II 6, III 5, IV 4. Leg measurements: I 43.77 (12.70 + 5.62 + 11.52 + 7.39 + 6.54), II 38.16 (10.54 + 4.88 + 9.92 + 7.63 + 5.19), III 35.95 (9.59 + 4.08 + 7.87 + 8.61 + 5.80), IV 48.11 (12.72 + 4.89 + 11.10 + 12.75 + 6.65). Leg formula: 4123.

Male palpal bulb (Fig. 3A, B; male palp with bulb Fig. 4A–C). Maxillae with lyra setae ventrally. Bulb oval, embolus slender and long, strongly curved at 90°, with A.

Female. unknown.

Etymology. The species is named after the Latin word *dominus*, as this is one of the rarest spiders from China; noun (name) in apposition.

Distribution. Known only from the type locality.




Figure 5. *Chilobrachys jinchengi* sp. nov., holotype (B–H) and paratype (A) males **A** male habitus, dorsal **B** ocular tubercle **C** right palp maxillae **D** sternum **E** stridulatory lyra, lateral view **F** same, ventral view **G** chelicerae, retrolateral view **H** same, prolateral view.

***Chilobrachys jinchengi* Lin & Li, sp. nov.**

<http://zoobank.org/CA76A9EC-61AF-4EBA-92F5-521EF797E229>

Figs 1B, 5A–H, 6A, B, 7A–C, 8

Type material. *Holotype* ♂ (IZCAS-Ar42677), **CHINA: Tibet:** Nyingchi, Medog, from Baibung to Muge, 29.2386°N, 95.1831°E, elevation 1310 m, 21.VIII.2018, Jincheng Liu leg. *Paratype* 1♂ (IZCAS-Ar42678), same data as holotype.




Figure 6. *Chilobrachys jinchengi* sp. nov., holotype, right palp bulb, rotated horizontally **A** retrolateral view **B** proteral view.



Figure 7. *Chilobrachys jinchengi* sp. nov., holotype, left palp **A** prolateral view **B** ventral view **C** retro-lateral view.

Diagnosis. The male of *Chilobrachys jinchengi* sp. nov. is similar to that *C. bicolor* (Pocock, 1895), but can be distinguished by the length ratio of the embolus to the bulb, which is almost 2:1 in *C. jinchengi* sp. nov. and is 1:1 in *C. bicolor*.

Description. Male (holotype, IZCAS-Ar42677) (Figs 5B–H, 6, 7). Carapace 13.12 long, 10.21 wide, brown with long setae. Opisthosoma absent. Eye group 2.32 long, 0.97 wide (Fig. 5F). MOA 0.77 long, anterior width 0.89, posterior width 1.38. Eye sizes and interdistances: ALE 0.51, AME 0.45, PLE 0.42, PME 0.38; ALE–AME 0.34, AME–AME 0.25, PLE–PME 0.12, PME–PME 0.87. Fovea slightly procurved.




Figure 8. Distribution records of *Chilobrachys* species in China 1 *C. dominus* sp. nov. 2 *C. jinchengi* sp. nov. 3 *C. guangxiensis* (Yin & Tan, 2000) 4 *C. hubei* Song & Zhao, 1988 5 *C. liboensis* Zhu & Zhang, 2008 6 *C. lubricus* Yu et al., 2021.

Chelicerae dark brown, with row of 12 promarginal teeth. Labium wider than long, with 367 cuspules. Sternum yellow-brown, with 3 pairs of sigilla. Legs with long and short setae. Tarsi I–III with 2 claws, tarsus IV with 3 claws, denticle number: I 9, II 9, III 9, IV 12. Leg measurements: I 49.58 (14.23 + 7.12 + 11.21 + 9.89 + 7.13), II 45.77 (13.02 + 6.73 + 9.52 + 9.59 + 6.91), III 43.21 (11.51 + 5.48 + 8.24 + 11.51 + 6.47), IV 53.97 (14.32 + 5.87 + 11.22 + 15.96 + 6.60). Leg formula: 4123.

Male palpal bulb (Fig. 6A, B; male palp with bulb Fig. 7A–C). Maxillae with lyra setae ventrally. Bulb oval, embolus bow-shaped, strongly curved at 180°, with A, PI and PS. Distal edge of embolus relatively flat.

Female. unknown.

Etymology. The species is named after Mr. Jincheng Liu, who collected the type material; noun (name) in the genitive case.

Distribution. Known only from the type locality.

Acknowledgements

The manuscript benefited greatly from comments by Zhiyuan Yao, Danni Sherwood, Robert Raven. The English was checked by Sarah Crews. Jincheng Liu, Run Zhou, Zhengzhong Huang, Yi Ming and Xiaoqing Zhang helped in field collections.

References

- Bertani R (2000) Male palpal bulbs and homologous features in Theraphosinae (Araneae, Theraphosidae). *Journal of Arachnology* 28: 29–42. [https://doi.org/10.1636/0161-8202\(2000\)028\[0029:MPBAHF\]2.0.CO;2](https://doi.org/10.1636/0161-8202(2000)028[0029:MPBAHF]2.0.CO;2)
- Foley S, Lüddecke T, Chen DQ, Krehenwinkel H, Künzel S, Longhorn S, Wendt I, Wirth V von, Tänzler R, Vences M, Piel WH (2019) Tarantula phylogenomics: A robust phylogeny of deep theraphosid clades inferred from transcriptome data sheds light on the prickly issue of urticating setae evolution. *Molecular Phylogenetics and Evolution* 140: 106573. <https://doi.org/10.1016/j.ympev.2019.106573>
- Li J, Yan X, Lin Y, Li S, Chen H (2021) Challenging Wallacean and Linnean shortfalls: *Ectosticta* spiders (Araneae, Hypochilidae) from China. *Zoological Research* 42(6): 791–794. <https://doi.org/10.24272/j.issn.2095-8137.2021.212>
- Li S (2020) Spider taxonomy for an advanced China. *Zoological Systematics* 45(2): 73–77. <https://doi.org/10.11865/zs.202011>
- Nunn SC, West RC, Wirth V von (2016) A revision of the selenocosmiine tarantula genus *Phlogiellus* Pocock 1897 (Araneae: Theraphosidae), with description of 4 new species. *International Journal of Zoology* 2016: e9895234. [54 pp.] <https://doi.org/10.1155/2016/9895234>
- Pocock RI (1895) On a new and natural grouping of some of the Oriental genera of Mygalomorphae, with descriptions of new genera and species. *Annals and Magazine of Natural History* (6) 15(86): 165–184. <https://doi.org/10.1080/00222939508677863>
- Raven RJ (1985) The spider infraorder Mygalomorphae (Araneae): cladistics and systematics. *Bulletin of the American Museum of Natural History* 182: 1–180.
- Simon E (1889) Arachnides. In *Voyage de M. E. Simon au Venezuela (décembre 1887–avril 1888)*. 4^e Mémoire. *Annales Société entomologique de France*, Paris (6)9: 169–220.
- WSC (2021) World Spider Catalog, Version 22.5. Natural History Museum Bern. <http://wsc.nmbe.ch> [accessed on 21/10/2021] <https://doi.org/10.24436/2>
- Yao Z, Wang X, Li S (2021) Tip of the iceberg: species diversity of *Pholcus* spiders (Araneae, Pholcidae) in Changbai Mountains, Northeast China. *Zoological Research* 42(3): 267–271. <https://doi.org/10.24272/j.issn.2095-8137.2020.214>
- Zhu MS, Zhang R (2008) Revision of the theraphosid spiders from China (Araneae: Mygalomorphae). *Journal of Arachnology* 36: 425–447. <https://doi.org/10.1636/CA07-94.1>