

POSTER PRESENTATION

Open Access

Prevalance of non-cardiac pathology on cardiovascular magnetic resonance studies

Elisa McAlindon^{1*}, Julian Strange¹, Nathan Manghat¹, Mark CK Hamilton¹, Peter Wilde¹, Peter Drivas², Dudley Pennell², Chiara Bucciarelli-Ducci³

From 2011 SCMR/Euro CMR Joint Scientific Sessions
Nice, France. 3-6 February 2011

Background


Cardiovascular magnetic resonance (CMR) is routinely used in clinical practice. Although the scan focuses on the cardiovascular system, the large field of view used in the initial dataset of images includes the thorax and upper abdomen. The prevalence of incidental non-cardiac findings is not well described. This multicentre retrospective study investigates the prevalence of incidental non-cardiac pathology in clinical patients undergoing CMR.

Methods

We reviewed consecutive CMR clinical reports of two dedicated CMR Units (Royal Brompton Hospital and Bristol Heart Institute) over a period of 3 years, from 2007 to 2010. All the scans were performed in 1.5T Avanto scanner (Siemens, Erlangen, Germany). Non-cardiac pathology was subsequently further classified in a) benign findings or b) findings requiring further characterisation. In addition, all images acquired in both centres during the month of August 2010 were retrospectively reviewed by a trained CMR cardiologist specifically assessing for non-cardiac findings.

Results

A total of 16,518 reports were reviewed and non cardiac pathology identified in 7.5% (n=1,242) patients. The entire image dataset of all the scans performed over a month period in both centres (n=479) were reviewed and revealed 26% (n=123) of patients with at least one non-cardiac finding (21 patients had >1 finding). A total of 144 non-cardiac pathologies were identified, of which 106 were classified as benign findings and 38 as findings



requiring further characterisation and testing (Figure 1). In particular, 10/38 lesions identified (2% of patients) represented possible malignancies (in particular mediastinal and lung lesions).

Conclusions

Whilst the prevalence of incidental findings by reports review (7.5%) is in keeping with the previous small single centre reports, this appears to underestimate the true incidence of non-cardiac pathology. Retrospective image review identified a higher prevalence of non-cardiac pathology (26%) which is similar to previously reported incidence during cardiac computed tomography. In our study, 8% of the patients required further testing following CMR and 2% of the patients had possible malignancies. Follow-up is required to establish the clinical significance of these findings and their impact on patient management.

Author details

¹CMR Unit, Bristol Heart Institute, NIHR Biomedical Research Unit, Bristol, UK. ²CMR Unit, Royal Brompton Hospital, NIHR Biomedical Research Unit, Bristol, UK. ³CMR Unit, Bristol Heart Institute/ CMR Unit, Royal Brompton Hospital, NIHR Biomedical Research Units, Bristol/ London, UK.

Published: 2 February 2011

¹CMR Unit, Bristol Heart Institute, NIHR Biomedical Research Unit, Bristol, UK
Full list of author information is available at the end of the article

doi:10.1186/1532-429X-13-S1-P174

Cite this article as: McAlindon *et al.*: Prevalance of non-cardiac pathology on cardiovascular magnetic resonance studies. *Journal of Cardiovascular Magnetic Resonance* 2011 **13**(Suppl 1):P174.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

