

Research article

Open Access

Investigating the effects of climate variations on bacillary dysentery incidence in northeast China using ridge regression and hierarchical cluster analysis

Desheng Huang^{1,2}, Peng Guan², Junqiao Guo³, Ping Wang⁴ and Baosen Zhou^{*2}

Address: ¹Department of Mathematics, College of Basic Medical Sciences, China Medical University, Shenyang 110001, PR China, ²Department of Epidemiology, School of Public Health, China Medical University, Shenyang 110001, PR China, ³Liaoning Provincial Center for Disease Control and Prevention, Shenyang 110005, PR China and ⁴Division of Infectious Diseases Control, Shenyang Municipal Center for Disease Control and Prevention, Shenyang 110031, PR China

Email: Desheng Huang - dshuang@mail.cmu.edu.cn; Peng Guan - pguan@mail.cmu.edu.cn; Junqiao Guo - qgcdc@hotmail.com; Ping Wang - pwcdc@hotmail.com; Baosen Zhou* - bszhou@mail.cmu.edu.cn

* Corresponding author

Published: 25 September 2008

Received: 22 May 2008

BMC Infectious Diseases 2008, **8**:130 doi:10.1186/1471-2334-8-130

Accepted: 25 September 2008

This article is available from: <http://www.biomedcentral.com/1471-2334/8/130>

© 2008 Huang et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: The effects of climate variations on bacillary dysentery incidence have gained more recent concern. However, the multi-collinearity among meteorological factors affects the accuracy of correlation with bacillary dysentery incidence.

Methods: As a remedy, a modified method to combine ridge regression and hierarchical cluster analysis was proposed for investigating the effects of climate variations on bacillary dysentery incidence in northeast China.

Results: All weather indicators, temperatures, precipitation, evaporation and relative humidity have shown positive correlation with the monthly incidence of bacillary dysentery, while air pressure had a negative correlation with the incidence. Ridge regression and hierarchical cluster analysis showed that during 1987–1996, relative humidity, temperatures and air pressure affected the transmission of the bacillary dysentery. During this period, all meteorological factors were divided into three categories. Relative humidity and precipitation belonged to one class, temperature indexes and evaporation belonged to another class, and air pressure was the third class.

Conclusion: Meteorological factors have affected the transmission of bacillary dysentery in northeast China. Bacillary dysentery prevention and control would benefit from by giving more consideration to local climate variations.

Background

Shigella infections remain an important public health problem in China, especially among children and old people [1,2]. Bacillary dysentery is an infectious disease of

the intestinal tract caused by bacteria of the genus *Shigella* and is spread by contact with patients or carriers and though food or water contaminated by their feces. The increases in temperature have been observed in most



Figure 1
Location of study area (Shenyang) in China.

Compared to Ordinary Least Square (OLS) linear regression, the basic principle of ridge regression is to artificially reduce correlation coefficient r_{ij} of each pair of variables x_i and x_j (including response variable and independent variables) to $r_{ij}/(1+k)$, k is called ridge parameter, and usually $0 < k < 1$. k value was selected when all the regression coefficients were relatively stable and the sign of the coefficients did not change.

Hierarchical cluster analysis

Hierarchical cluster analysis was adopted to group the weather variables. In this study, the agglomerative method with between-group average linkage algorithm was adopted and the measure for similarity was the Pearson correlation.

All the above statistical analyses were performed by Statistical Product and Service Solutions (SPSS 12.0 for windows, SPSS Inc., Chicago, IL, USA).

Ethical review

The present study was reviewed by research institutional review board of China Medical University and found to be

utilization of disease surveillance data and meteorological data not requiring oversight by an ethics committee.

Results

Statistical description of bacillary dysentery in Shenyang city

The distribution of bacillary dysentery incidence in Shenyang between 1950 and 1996 is shown in Figure 2. Figure 2a shows the long-term trend, which is that the incidence of bacillary dysentery in Shenyang rose over periods 1950–1966 and 1970–1981, and declined over periods 1966–1970 and 1981–1996. Figure 2b indicates that there was a seasonal distribution of bacillary dysentery in the study area, with about 85% of cases occurring in summer and autumn. The incidence of bacillary dysentery increased from January to August and decreased from August to December.

Spearman's correlation analysis

The relationship between monthly mean meteorological factors and the monthly incidence of bacillary dysentery in the studied period was examined by Spearman's correlation analysis (Table 1).




Figure 2
Distribution of bacillary dysentery incidence (Shenyang, China, 1950–1996). A) Long-term trend. B) Seasonal distribution.

Due to the fact that the history data covered more than 40 years, four time subgroups were set for analysis based on historical and economic development of China. Table 1 shows that temperatures, precipitation, evaporation and relative humidity were positively related to the monthly incidence of bacillary dysentery, while air pressure was inversely correlated with the incidence. Lagged effects existed in air pressure from 1987 to 1996 and in relative humidity from 1967 to 1976 and from 1987 to 1996.

Multi-collinearity Diagnosis

The diagnosis of multi-collinearity is shown in Table 2 and Table 3. In Table 2, the largest VIF is 164.187, and the average VIF is 54.574, therefore, climatic variables are strongly collinear. In addition, Table 3 shows that the largest condition index is 815.54 which also indicates strong multi-collinearity. With the help of condition index and variance proportion, we found that collinearity existed between the constant and air pressure, the average tem-

perature and maximum temperature, as well as minimum temperature and relative humidity.

Ridge regression analysis

Ridge regression was adopted to examine the relationship between the monthly mean weather variables and the monthly incidence of bacillary dysentery. Ridge trace maps (Figure 3) show that when ridge parameter *k* reached 0.5, ridge regression coefficients were stable. When ridge parameter was set as 0.5, the ridge regression coefficients were then obtained (Table 4). Different patterns for the analysis unit Sub-grouping improved the determinant coefficients of all models. However, determinant coefficients of each unit differ from 0.490 to 0.892. Most models give the following similar results: relative humidity and precipitation have a positive association with incidence from January to July, and have no association from August to December, while evaporation is the opposite. Some results vary different for the four time sub-

Table 1: Spearman correlations between monthly incidence of bacillary dysentery and corresponding monthly meteorological factors (1950–1996, Shenyang, China)

Year	Air pressure	Average air temperature	Maximum temperature	Minimum temperature	Precipitation	Evaporation	Relative humidity
1950–1966	-0.459	0.542	0.521	0.554	0.429	0.419	0.194
1967–1976	-0.715	0.772	0.768	0.790	0.657	0.545	0.647(lag1)
1977–1986	-0.697	0.824	0.827	0.825	0.711	0.641	0.604
1987–1996	-0.750(lag1)	0.861	0.846	0.870	0.791	0.602	0.601(lag1)
1950–1996	-0.493	0.580	0.568	0.593	0.506	0.374	0.332

All correlation coefficients are significant at the 0.01 level (2-tailed)
 Lag1: one month prior

Table 2: Collinearity Statistics for explanatory variables

Variables	Tolerance	VIF
Air pressure	0.235	4.249
Average temperature	0.006	164.187
Maximum temperature	0.009	113.969
Minimum temperature	0.011	87.756
Precipitation	0.388	2.580
Evaporation	0.154	6.505
Relative humidity	0.360	2.775

groups, for example, ridge regression coefficients are not always statistically significant. And also, lagged effects only existed in air pressure from 1987 to 1996 and in relative humidity from 1967 to 1976 and from 1987 to 1996. In order to compare variables effects between models, original variables without lagged values were used for ridge regression analysis directly.

Cluster analysis

Figure 4 shows the effects of climate variations on bacillary dysentery incidence, the regression coefficients were listed above the arrows. Due to the long period, only the model for 1987 to 1996 was listed. During this period, all meteorological factors were divided into three categories. Relative humidity and precipitation belonged to one class, temperature indexes and evaporation belonged to another class, and air pressure was the third class (Figure 4).

Discussion

Although incidence of intestinal infectious diseases has declined considerably in developed countries, they are still major public health problems in developing countries, especially in the economic developing regions[8,9]. The transmission of Shigella is determined by many factors, such as human dietary lifestyle, and its strains' susceptibility to drugs and climate variations. Climate variability that impacts the incubation rate of Shigella is

considered one of the major environmental contributors to Shigella transmission.

Climate change, especially global warming, has already brought and will continue to bring about challenges to public health and communicable disease control and prevention. The effects of climate variations on bacillary dysentery incidence have gained more recent concern. The historical climatic data from 1950 to 1996 demonstrate a significant increase in temperatures in Shenyang city. It has been reported that temperature was the key climatic indicator in the transmission of bacillary dysentery, a potential 1 °C increase in maximum or minimum temperature was associated with up to 12% increase in bacillary dysentery cases in the northern city if other affecting factors remain unchanged[10]. Public health interventions should be developed at this stage to adapt and mitigate the possible consequences of climate change in the future with consideration of local climatic conditions[11]. The model constructed in the present study showed that explanatory climatic variables for the incidence of bacillary dysentery could be divided into three categories, namely relative humidity and precipitation; temperature indexes and evaporation; and air pressure. Among them, only air pressure showed negative correlations, all other variables showed positive correlations. From the point of effect size, temperature indexes and precipitation had greater effect which is of part concordance with the results of other studies in China[12,13]. Precipitation, relative humidity and air pressure may affect the transmission of food-borne diseases by influencing the pathogens and then affect the incidence of bacillary dysentery. The impact of precipitation on diarrhoeal disease is far from conclusion. Different studies obtained inconsistent results [14-16]. The three categories of climatic variables could be used as predictors for the number of bacillary dysentery cases in Shenyang. The results of the present study may be used to assist public health decision making and community health education. Further studies about this issue are necessary.

Table 3: Collinearity Diagnostics between explanatory variables

Condition Index	Variance Proportions							
	Constant	Air pressure	Average temperature	Maximum temperature	Minimum temperature	Precipitation	Evaporation	Relative humidity
1.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
2.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
4.39	0.00	0.00	0.00	0.00	0.00	0.55	0.02	0.00
8.98	0.00	0.00	0.00	0.00	0.01	0.31	0.36	0.02
30.08	0.00	0.00	0.09	0.12	0.39	0.09	0.45	0.43
35.89	0.00	0.00	0.08	0.04	0.56	0.00	0.11	0.54
50.00	0.00	0.00	0.82	0.83	0.04	0.03	0.00	0.00
815.54	1.00	1.00	0.01	0.01	0.00	0.01	0.05	0.01

Table 4: Ridge regression coefficients between monthly climatic variables and logarithmic incidence of bacillary dysentery (with $k = 0.5$, 1950–1996, Shenyang, China)

Duration	R ²	Minimum temperature	Average temperature	Maximum temperature	Air pressure	Relative humidity	Precipitation	Evaporation	Long-term Effect	Season
1950–1966, Jan – Jul	0.718	0.079*	0.072*	0.051*	-0.042	0.053	0.156*	0.032	0.458*	0.110*
1950–1966, Aug – Dec	0.818	0.109*	0.075*	0.067*	-0.093*	-0.047	0.068	0.148*	0.442*	-0.084*
1967–1976, Jan – Jul	0.653	0.138*	0.115*	0.106*	-0.097*	0.149*	0.096	-0.009	0.142*	0.106*
1967–1976, Aug – Dec	0.768	0.101*	0.092*	0.076*	-0.156*	0.103	0.013	0.073*	0.338*	-0.101*
1977–1986, Jan – Jul	0.490	0.072*	0.046	0.063*	-0.033	0.222*	0.119	0.114*	0.111	0.052
1977–1986, Aug – Dec	0.822	0.113*	0.127*	0.153*	-0.118*	-0.041	0.065	0.189*	-0.094*	-0.125*
1987–1996, Jan – Jul	0.822	0.131*	0.115*	0.100*	-0.069*	0.148*	0.207*	-0.017	-0.140*	-0.154*
1987–1996, Aug – Dec	0.892	0.154*	0.144*	0.142*	-0.101*	0.048	0.058	0.125*	-0.132*	-0.135*
1950–1996, Jan – Jul	0.306	0.107*	0.082*	0.080*	-0.030	0.018	0.131*	-0.027	0.045	0.111*
1950–1996, Aug – Dec	0.430	0.134*	0.105*	0.094*	-0.110*	-0.116*	0.062	0.071*	0.016	-0.116*

* $P < 0.05$

Due to the multi-collinearity in independent variables, the regression coefficients are very unstable with excessive standard error. A general approach to treat the existence of variables that are non significant to the effect of response variables is to remove them one by one until all the variables reserved in the equation are significant. In the present study, high multi-collinearity existed between the constant and air pressure, average temperature and maximum temperature, minimum temperature and relative humidity. We proposed a modified method to combine ridge regression and cluster analysis to solve this problem. The adoption of ridge estimated method to construct the regression model improved the effect of multi-collinearity, reduced the standard error of regression coefficients and produced more stable results. The stable regression coefficients then made the estimation of effect more reliable.

Though the meteorological factors are nonmodifiable, we can modify our dietary lifestyle, such as reducing intake of raw vegetables or cleaning vegetables and fruits more thoroughly. Local weather forecasting may have policy implications for bacterial dysentery prevention and control. Strengthening the education of public health will help to eliminate or reduce the epidemic of bacterial dysentery.

One limitation of this study was that we only focused on the relationship between the meteorological factors and the incidence of bacillary dysentery. The socio-economic

status, especially the awareness of health care has been improved in China. So in order to remove the effect of preventive measures, this study used only the historical data before 1996 and four time subgroups that were obtained based on historical and economic development of China. The changes in social and economic conditions need to be addressed in further studies. Bacillary dysentery is a legally mandated notifiable disease in China, as a provincial capital city, Liaoning provincial CDC and Shenyang municipal CDC have kept good records for Shenyang bacillary dysentery incidence. However, there is still another limitation of the present study, which is that under-reporting is possible in the disease surveillance system, especially before the computer and web were widely applied. An investigation of missing reports of notifiable diseases in China in 2005 indicated that there was much to be improved upon in Chinese medical facilities as far as reporting of infectious diseases[17]. So, the correlation between climatic variation and bacillary dysentery may be underestimated. However, there is no evidence to suggest there was any trend in under-reporting during the four time subgroups in Shenyang city. Therefore it is considered that the impact of under-reporting during each time sub group on the results is not significant.

Conclusion

In summary, relative humidity, temperature indexes and atmospheric pressure affected the transmission of the bacillary dysentery in Shenyang, located in the northeast of China. Climate variations have brought and will continue




Figure 3
Ridge trace between climatic variables and logarithmic incidence of bacillary dysentery (1950–1996, Shenyang, China).




Figure 4
Combination of cluster analysis and ridge regression analysis between climatic variables and logarithmic incidence of bacillary dysentery (1987–1996, Shenyang, China). A) 1987–1996, January – July. B) 1987–1996, August – December.

to bring about challenges to communicable disease control and prevention. Bacillary dysentery prevention and control should be taken by giving more consideration to local climate variations and integrated modelling based on data systems is necessary to disease control and prevention.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

DH conceived the study, attracted funding and drafted the manuscript. DH, PG and BZ managed and analyzed the data. PW managed the bacillary dysentery incidence database which was administered and supervised by JG. All authors contributed to the writing of the final version of this paper.

Acknowledgements

This study was financially supported by the National Natural Science Foundation of China, No. 70503028. Guan Peng was supported by the Scientific Research Foundation for the Returned Overseas Chinese Scholars, State Education Ministry, No. [2008]890 and a CMU Development grant. The authors are grateful to all the participants in this study. The authors thank the editor and two reviewers for their insightful comments which led to a significantly improved version of the manuscript. Also, the authors thank Ryan Session (Emory University, USA) for his careful checking of grammar and spelling of the manuscript.

References

- Wang XY, Tao F, Xiao D, Lee H, Deen J, Gong J, Zhao Y, Zhou W, Li W, Shen B, Song Y, Ma J, Li ZM, Wang Z, Su PY, Chang N, Xu JH, Ouyang PY, von Seidlein L, Xu ZY, Clemens JD: **Trend and disease burden of bacillary dysentery in China (1991–2000).** *Bull World Health Organ* 2006, **84**:561-568.
- von Seidlein L, Kim DR, Ali M, Lee H, Wang X, Thiem VD, Canh do G, Chaicumpa W, Agtini MD, Hossain A, Bhutta ZA, Mason C, Sethabutr O, Talukder K, Nair GB, Deen JL, Kotloff K, Clemens J: **A multicentre study of Shigella diarrhoea in six Asian countries: disease burden, clinical manifestations, and microbiology.** *PLoS Med* 2006, **3**:e353.
- Kalkstein LS, Smoyer KE: **The impact of climate change on human health: some international implications.** *Experientia* 1993, **49**:969-979.
- Curriero F, Patz J, Rose J, Lele S: **The association between extreme precipitation and waterborne disease outbreaks in the United States, 1948–1994.** *Am J Public Health* 2001, **91**:1194-1199.
- Lama JR, Seas CR, Leon-Barua R, Gotuzzo E, Sack RB: **Environmental temperature, cholera, and acute diarrhoea in adults in Lima.** *Peru J Health Popul Nutr* 2004, **22**:399-403.
- Yang GH, Stroup DF, Thacker SB: **National public health surveillance in China: implications for public health in China and the United States.** *Biomed Environ Sci* 1997, **10**:1-13.
- Hocking RR: *Methods and Applications of Linear Models* Wiley, New York; 1996.
- Gauci C, Gilles H, O'Brien S, Mamo J, Stabile I, Ruggeri FM, Calleja N, Spiteri G: **Estimating the burden and cost of infectious intestinal disease in the Maltese community.** *Epidemiol Infect* 2007, **135**:1290-1298.
- Lee H, Kotloff K, Chukaserm P, Samosornsuk S, Chompook P, Deen JL, Von Seidlein L, Clemens JD, Wanpen C: **Shigellosis remains an important problem in children less than 5 years of age in Thailand.** *Epidemiol Infect* 2005, **133**:469-474.
- Zhang Y, Bi P, Hiller JE, Sun Y, Ryan P: **Climate variations and bacillary dysentery in northern and southern cities of China.** *J Infect* 2007, **55**:194-200.
- Bi P, Zhang Y, Parton KA: **Weather variables and Japanese encephalitis in the metropolitan area of Jinan city, China.** *J Infect* 2007, **55**:551-556.
- Zhang Y, Bi P, Hiller JE: **Weather and the transmission of bacillary dysentery in Jinan, northern China: a time-series analysis.** *Public Health Rep* 2008, **123**:61-66.

13. Chen KC, Lin CH, Qiao QX, Zen NM, Zhen GK, Chen GL, Xie YJ, Lin YJ, Zhuang SF: **The epidemiology of diarrhoeal diseases in southeastern China.** *J Diarrhoeal Dis Res* 1991, **9**:94-99.
14. Kovats RS, Edwards SJ, Hajat S, Armstrong BG, Ebi KL: **The effects of temperature on food poisoning: a time-series analysis of salmonellosis in ten European countries.** *Epidemiol Infect* 2004, **132**:443-453.
15. D'Souza R, Becker N, Hall G, Moodie K: **Does ambient temperature affect foodborne disease?** *Epidemiology* 2004, **15**:86-92.
16. Singh R, Hales S, Wet N, Raj R, Hearnden M, Weinstein P: **The influence of climate variation and change on diarrheal disease in the Pacific Islands.** *Environ Health Perspect* 2001, **109**:155-159.
17. Shi XM, Ma JQ, Wang LP, Zhang CX, Guo Q, Qi XP, Guo Y, Yang GH: **An investigational analysis of missing reports of notifiable diseases in medical facilities all over China in 2005.** *Disease Surveillance* 2006, **21**:193-496. (In Chinese)

Pre-publication history

The pre-publication history for this paper can be accessed here:

<http://www.biomedcentral.com/1471-2334/8/130/prepub>

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

