

# Association between Ambient Air Pollution and Diabetes Mellitus in Europe and North America: Systematic Review and Meta-Analysis

**Ikenna C. Eze,<sup>1,2</sup> Lars G. Hemkens,<sup>3</sup> Heiner C. Bucher,<sup>3</sup> Barbara Hoffmann,<sup>4,5</sup> Christian Schindler,<sup>1,2</sup> Nino Künzli,<sup>1,2</sup> Tamara Schikowski,<sup>1,2,4</sup> and Nicole M. Probst-Hensch<sup>1,2</sup>**

<sup>1</sup>Swiss Tropical and Public Health Institute, Basel, Switzerland; <sup>2</sup>University of Basel, Basel, Switzerland; <sup>3</sup>Basel Institute for Clinical Epidemiology and Biostatistics, University Hospital Basel, Basel, Switzerland; <sup>4</sup>UF-Leibniz Research Institute for Environmental Medicine, Düsseldorf, Germany; <sup>5</sup>Medical Faculty, Heinrich Heine University of Düsseldorf, Düsseldorf, Germany

**BACKGROUND:** Air pollution is hypothesized to be a risk factor for diabetes. Epidemiological evidence is inconsistent and has not been systematically evaluated.

**OBJECTIVES:** We systematically reviewed epidemiological evidence on the association between air pollution and diabetes, and synthesized results of studies on type 2 diabetes mellitus (T2DM).

**METHODS:** We systematically searched electronic literature databases (last search, 29 April 2014) for studies reporting the association between air pollution (particle concentration or traffic exposure) and diabetes (type 1, type 2, or gestational). We systematically evaluated risk of bias and role of potential confounders in all studies. We synthesized reported associations with T2DM in meta-analyses using random-effects models and conducted various sensitivity analyses.

**RESULTS:** We included 13 studies (8 on T2DM, 2 on type 1, 3 on gestational diabetes), all conducted in Europe or North America. Five studies were longitudinal, 5 cross-sectional, 2 case-control, and 1 ecologic. Risk of bias, air pollution assessment, and confounder control varied across studies. Dose-response effects were not reported. Meta-analyses of 3 studies on PM<sub>2.5</sub> (particulate matter  $\leq 2.5$   $\mu\text{m}$  in diameter) and 4 studies on NO<sub>2</sub> (nitrogen dioxide) showed increased risk of T2DM by 8–10% per 10- $\mu\text{g}/\text{m}^3$  increase in exposure [PM<sub>2.5</sub>: 1.10 (95% CI: 1.02, 1.18); NO<sub>2</sub>: 1.08 (95% CI: 1.00, 1.17)]. Associations were stronger in females. Sensitivity analyses showed similar results.

**CONCLUSION:** Existing evidence indicates a positive association of air pollution and T2DM risk, albeit there is high risk of bias. High-quality studies assessing dose-response effects are needed. Research should be expanded to developing countries where outdoor and indoor air pollution are high.

**CITATION:** Eze IC, Hemkens LG, Bucher HC, Hoffmann B, Schindler C, Künzli N, Schikowski T, Probst-Hensch NM. 2015. Association between ambient air pollution and diabetes mellitus in Europe and North America: systematic review and meta-analysis. *Environ Health Perspect* 123:381–389; <http://dx.doi.org/10.1289/ehp.1307823>

## Introduction

Ambient air pollution ranks high among risk factors for the global burden of disease (Lim et al. 2012), and is linked to several chronic noncommunicable conditions such as cardiovascular diseases (Bauer et al. 2010; Brook et al. 2010; Künzli et al. 2010), asthma (Bui et al. 2013; Jacquemin et al. 2012; Künzli et al. 2009), chronic obstructive pulmonary diseases (COPD) (Andersen et al. 2011; Schikowski et al. 2014; Zanobetti et al. 2008), and cancers including lung (Raaschou-Nielsen et al. 2013a), cervical, and brain cancers (Raaschou-Nielsen et al. 2011). Persons with type 2 diabetes mellitus (T2DM) are at increased risk to develop micro- and macrovascular diseases and reduced lung function (Jones et al. 2014; Kinney et al. 2014). Air pollution has also been shown to be more detrimental to diabetic patients, worsening their clinical outcomes (O'Neill et al. 2005; Raaschou-Nielsen et al. 2013b; Whitsel et al. 2009; Zanobetti and Schwartz 2001).

More recent evidence is supportive of an air pollution effect on diabetes risk. Experimental evidence show that possible

pathways may include endothelial dysfunction, overactivity of the sympathetic nervous system (Rajagopalan and Brook 2012), immune response alterations in visceral adipose tissues; endoplasmic reticulum stress resulting in alterations in insulin transduction (Sun et al. 2009), insulin sensitivity, and glucose metabolism; and alterations in mitochondria and brown adipocytes (Liu et al. 2013; Rajagopalan and Brook 2012).

Papazafropoulou et al. (2011) systematically reviewed the etiologic association between environmental pollution and diabetes, taking into account studies on organic pollutants and secondary effects of air pollution on diabetic patients published up to November 2010. They described a positive association between environmental pollution and prevalent diabetes, as well as increased morbidity and mortality among diabetic patients. A number of pertinent studies have been published since this review, and thus far there is, to the best of our knowledge, no meta-analysis of the available evidence. We therefore systematically identified and reviewed the epidemiological evidence on the association between air pollution and diabetes

mellitus, and synthesized the results of studies on the association with T2DM.

## Methods

**Search strategy.** We systematically searched electronic literature databases [MEDLINE (<http://www.nlm.nih.gov/bsd/pmresources.html>), EMBASE (<https://www.embase.com>), and ISI Web of Science (<http://www.webofknowledge.com>)] for pertinent literature published up to 3 February 2014. Terms used in this search included “air pollution,” “air pollutants,” “particulate matter,” “PM<sub>10</sub>,” “PM<sub>2.5</sub>,” “nitrogen dioxide,” “NO<sub>2</sub>,” “NO<sub>x</sub>,” “ozone,” “soot,” “smog,” “diabetes mellitus,” “diabetes,” “T1DM,” “T2DM,” “type 1 DM,” “type 2 DM,” “IDDM,” “NIDDM,” alone and in combination. We applied no filters for study designs. Reference lists of eligible articles were searched for further pertinent articles. After de-duplication, titles and abstracts were screened for eligibility and potentially relevant articles were retrieved as full texts. Screening was performed independently by two reviewers and any discrepancies were resolved by discussion.

**Inclusion and exclusion criteria.** We included only original research published in English as a full publication in a peer-reviewed journal. We accepted any type of study design. In eligible studies, the definition of air pollution and diabetes mellitus

Address correspondence to N.M. Probst-Hensch, Department of Epidemiology and Public Health, Swiss Tropical and Public Health Institute, Socinstrasse 57, 4002 Basel, Switzerland. Telephone: 41 61 284 83 88. E-mail: Nicole.Probst@unibas.ch  
Supplemental Material is available online (<http://dx.doi.org/10.1289/ehp.1307823>).

The Federal Office for Forest, Environment and Landscape provided support for the salary costs related to this study. SAPALDIA (Swiss study on Air Pollution and Lung Disease in Adults) is supported by the Swiss National Science Foundation; the Federal Office for Forest, Environment and Landscape; the Federal Office of Public Health; the Federal Office of Roads and Transport; the cantonal governments of Aargau, Basel-Stadt, Basel-Land, Geneva, Luzern, Ticino, and Zurich; the Swiss Lung League; and the Lung Leagues of Basel-Stadt/Basel-Landschaft, Geneva, Ticino, and Zurich. L.G.H. and H.C.B. are supported by unrestricted grants from Santésuisse.

The authors declare they have no actual or potential competing financial interests.

Received: 30 October 2013; Accepted: 26 January 2015; Advance Publication: 27 January 2015; Final Publication: 1 May 2015.

had to be clearly stated. Air pollution had to be outdoor (ambient, including traffic-related), and we accepted any type of assessment including particle concentration in the air or indicators of long-term traffic exposure. Diabetes mellitus had to be physician diagnosed or based on the use of antidiabetic medications. We included any type of diabetes mellitus (type 1, type 2, and gestational). Eligible studies had to report quantitative measures of association between air pollution and diabetes mellitus, and their 95% confidence intervals (CIs) (or enough data to allow derivation of this association). We excluded studies that were based on the effect of blood markers, and not clearly defining clinical outcomes. Studies testing only whether diabetes status would modify the association between air pollution and health outcomes were not considered in this review. Animal studies were excluded.

For the meta-analysis, only studies on individual type 2 diabetes risk were included. We included all studies that quantified particle concentrations as “per ...  $\mu\text{g}/\text{m}^3$ ” or “ppb.” If the diabetes type was not clearly stated, we considered diagnoses of diabetes in nonpregnant adults ( $\geq 18$  years age) as diagnoses of T2DM because  $> 90\%$  of new diagnoses of adult diabetes is type 2 diabetes (Alberti and Zimmet 1998).

**Data extraction.** We extracted the following data from the eligible studies: year

of study, study setting, study design, year of publication, population demographics, study definition of diabetes and assessment of air pollution exposure, confounder adjustments, and effect modification assessments. We extracted data on the effect estimates (unadjusted and final model) of the association (and their 95% CIs) between air pollution and diabetes.

Data were extracted independently by two reviewers and disagreements were resolved by discussion.

**Meta-analysis.** We used random-effects models to synthesize the associations between air pollution and T2DM (Lau et al. 1997). Random-effect models give more weight to smaller studies and have typically wider CIs because in addition to the within-study variance, they also consider potential variation between the true effects that all included studies estimate. We used fixed-effects models (which assume that all studies share a common true effect) in a sensitivity analysis.

We used risk ratios as measure of association across all studies. When hazard ratios and incidence risk ratios were reported, we directly considered them as risk ratios. Because diabetes is not very common, we considered reported odds ratios as equivalent to risk ratios. For studies with estimates of association from multiple particle concentration sources, we chose the estimates modelled at participants' residences (land-use


regression, kriging, or satellite-based estimates). We used the effect estimates reported by the study authors as “main model” or “fully adjusted model.” We used estimates of association and their standard errors reported as “per  $10 \mu\text{g}/\text{m}^3$ ” of exposure and we converted other reported quantities or units where necessary.

We described the between-study heterogeneity using the  $I^2$  metric and the between-studies' variance using  $\text{Tau}^2$ . We assessed publication bias using the Egger's test for asymmetry (Egger et al. 1997). We conducted sensitivity analyses including only studies that *a*) measured air pollution exposure before DM diagnosis, *b*) comprised both males and females, and *c*) were longitudinal, and we applied a fixed-effects analysis. All analyses were performed with Stata version 12 (StataCorp, College Station, TX, USA) using the “metan” command. *p*-Values were two-tailed and  $p < 0.05$  was considered nominally statistically significant.

For reporting, we followed the Meta-analysis Of Observational Studies in Epidemiology (Stroup et al. 2000) and the Preferred Reporting Items for Systematic Reviews and Meta-Analysis (Moher et al. 2010) guidelines.

## Results

The database search yielded 636 records after de-duplication, which were screened on title/abstract level for eligibility (Figure 1). Sixteen potentially eligible articles were screened on full-text level, and 3 were excluded (Figure 1). Thirteen studies were included (Table 1). There were 5 longitudinal studies (Andersen et al. 2012; Chen et al. 2013; Coogan et al. 2012; Krämer et al. 2010; Puett et al. 2011), 5 cross-sectional studies (Brook et al. 2008; Dijkema et al. 2011; Fleisch et al. 2014; Malmqvist et al. 2013; van den Hooven et al. 2009), 2 case-control studies (Hathout et al. 2002, 2006), and 1 ecologic study (Pearson et al. 2010). Two studies were on type 1 diabetes (Hathout et al. 2002, 2006); 3 studies on gestational diabetes (GDM) (Fleisch et al. 2014; Malmqvist et al. 2013; van den Hooven et al. 2009), and 8 studies on T2DM (Andersen et al. 2012; Brook et al. 2008; Chen et al. 2013; Coogan et al. 2012; Dijkema et al. 2011; Krämer et al. 2010; Pearson et al. 2010; Puett et al. 2011). Seven non-ecological studies on T2DM were selected for quantitative synthesis (with the exclusion of Pearson et al. 2010). Air pollution estimates from these studies were based on land-use regression (Andersen et al. 2012; Brook et al. 2008; Dijkema et al. 2011; Krämer et al. 2010; Puett et al. 2011), kriging (Coogan et al. 2012), and satellite-derived estimates (Chen et al. 2013). All studies were conducted in Europe or North



**Figure 1.** Results of systematic literature search.

America. Tables 1 and 2 and Supplemental Material, Table S1, provide an overview of the 13 eligible studies. Table 3 summarizes the data reported in the studies synthesized in meta-analyses.

In the Supplemental Material, Table S2 provides an overview of potential sources of bias and how they were assessed by the 13 studies. These are discussed in detail below.

**Bias due to outcome assessment.** As shown in Table 2, some studies relied on self-reported, physician-diagnosed DM (Coogan et al. 2012; Dijkema et al. 2011; Krämer et al. 2010), whereas others linked participants to established databases to identify cases (Andersen et al. 2012; Brook et al. 2008; Chen et al. 2013; Hathout et al. 2002, 2006; Malmqvist et al. 2013). Additional steps were taken by some studies with self-reported outcomes to test the validity of the DM diagnosis. These steps included sending a follow-up questionnaire with the same questions about diabetes (Krämer et al. 2010) and confirmation from medical records provided by physicians (Coogan et al. 2012). Dijkema et al. (2011) further tested participants who did not report physician-diagnosed diabetes, to identify undiagnosed cases.

**Bias due to exposure assessment.** The reviewed studies used different approaches to assess exposure of participants to air pollution, including modeled concentrations of various particulate matters, nitrogen oxides ( $\text{NO}_x$ ), sulfates, ozone, and various proxies to estimate traffic-related pollution, with varying buffer levels. The studies are also heterogeneous with regard to the lag time considered for exposure assessment. Only the Danish cohort (Andersen et al. 2012) assessed the impact of different lag times, albeit with little evidence for substantial differences in effects (see Supplemental Material, Table S1). In the absence of a biological basis for the latency between exposure and diagnosis of diabetes, different lag times should be tested. Overall, the diversity of exposure measurement makes it difficult to compare the reported effect estimates across these studies.

**Bias due to confounder adjustment. Indoor air pollution and smoking.** Beyond adjustment for basic DM risk factors at baseline (see Supplemental Material, Table S2), Krämer et al. (2010) also adjusted for environmental tobacco smoke (ETS), indoor heating with fossil fuels, as well as occupational exposure to dust, fumes and extreme temperatures;

Andersen et al. (2012) also adjusted for ETS. One study done in children considered ETS exposure (Hathout et al. 2006).

**Demographics, physical activity, and dietary factors.** The longitudinal studies uniformly adjusted for age, body mass index (BMI), and sex (when study population includes both sexes). The studies on women did not adjust for dietary factors, and all longitudinal studies but one adjusted for alcohol consumption and physical activity (see Supplemental Material, Table S1). The other studies assessed confounding by age and BMI except the case-control studies, which did not consider the children's BMI in their models. The GDM studies mostly considered maternal alcohol consumption (but not dietary factors) whereas the cross-sectional T2DM studies did not consider either factor (see Supplemental Material, Table S1).

**Socioeconomic status.** There was a uniform adjustment for socioeconomic status in all studies, although on different scales. At the individual level, educational attainment as a socioeconomic determinant was most commonly used across studies, and a few studies additionally considered household income and ethnicity (see Supplemental

**Table 1.** Characteristics of the studies on the relationship between air pollution and diabetes mellitus.

| Source                            | Location                                                                           | Years of study   | Study design and duration of follow-up                                                                             | Population (n) and age (years) of participants                                                                                                                                                   |
|-----------------------------------|------------------------------------------------------------------------------------|------------------|--------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Krämer et al. 2010 <sup>a</sup>   | Ruhrgebiet, Germany                                                                | 1990–2006        | Longitudinal: Study on the Influence of Air Pollution on Lung Inflammation and Aging<br>Follow-up: 16 years        | n = 1,775 Caucasian women without T2DM at baseline, 54–55 years                                                                                                                                  |
| Andersen et al. 2012 <sup>a</sup> | Copenhagen and Aarhus, Denmark                                                     | (1993–1997)–2006 | Longitudinal: Danish Diet, Cancer and Health cohort<br>Follow-up: 9.7 years                                        | n = 51,818 Caucasians without DM at baseline, 50–65 years                                                                                                                                        |
| Puett et al. 2011 <sup>a</sup>    | Metropolitan Statistical Areas (MSA) in north-eastern and midwestern states of USA | 1989–2009        | Longitudinal, with 2 cohorts: Nurses' Health Study and Health Professionals Follow-up Study<br>Follow-up: 20 years | n = 74,412 female nurses 30–55 years and 15,048 male health professionals 40–75 years, without T2DM at baseline                                                                                  |
| Coogan et al. 2012 <sup>a</sup>   | Los Angeles, California, USA                                                       | 1995–2005        | Longitudinal: Black Women's Health Study<br>Follow-up: 10 years                                                    | n = 3,992 African-American women, without DM at baseline and 21–69 years                                                                                                                         |
| Chen et al. 2013 <sup>a</sup>     | Ontario, Canada                                                                    | (1996–2005)–2010 | Longitudinal<br>Follow-up: 8 years                                                                                 | n = 62,012 Canadians without DM, ≥ 35 years                                                                                                                                                      |
| Brook et al. 2008 <sup>a</sup>    | Hamilton and Toronto, Ontario, Canada                                              | 1992–1999        | Cross-sectional                                                                                                    | n = 7,634 patients who attended two respiratory clinics in Hamilton and Toronto, ≥ 40 years                                                                                                      |
| van den Hooven et al. 2009        | Rotterdam, Netherlands                                                             | 2002–2006        | Cross-sectional: Generation R study                                                                                | n = 7,399 pregnant women, who had delivery date in the study period, 21–38 years                                                                                                                 |
| Dijkema et al. 2011               | Westfriesland, Netherlands                                                         | 1998–2000        | Cross-sectional: Hoorn Screening Study for T2DM                                                                    | n = 8,018 Caucasian residents, 50–75 years                                                                                                                                                       |
| Malmqvist et al. 2013             | Scania, Sweden                                                                     | 1999–2005        | Cross-sectional: The Swedish Medical Birth Registry.                                                               | n = 81,110 women who had singleton deliveries during the study period                                                                                                                            |
| Hathout et al. 2006               | California, USA                                                                    | 2002–2003        | Case-control<br>Follow-up: retrospectively from birth until diagnosis of T1DM                                      | n = 402 children (102 with T1DM and 300 age-matched controls), 1–12 years, receiving care at Loma Linda University Pediatric Center                                                              |
| Hathout et al. 2002               | California, USA                                                                    | 2002             | Case-control<br>Follow-up: retrospectively from birth until diagnosis of T1DM                                      | n = 100 children (61 cases: 30 had onset ≤ 5 years and 31 > 5 years) (39 age-matched controls: 19 were ≤ 5 years and 20 were > 5 years) receiving care at Loma Linda University Pediatric Center |
| Fleisch et al. 2014               | Boston, Massachusetts, USA                                                         | 1999–2002        | Cross-sectional: Project Viva Cohort                                                                               | n = 2,093 second-trimester pregnant women without known diabetes                                                                                                                                 |
| Pearson et al. 2010               | USA                                                                                | 2004–2005        | Ecologic                                                                                                           | n = 3,082 counties of USA                                                                                                                                                                        |

Abbreviations: T2DM, type 1 diabetes mellitus; T2DM, type 2 diabetes mellitus.

<sup>a</sup>Included in meta-analysis.

Material, Table S1). Few studies considered spatial socioeconomic confounding in forms of unemployment rate, urban/rural residence, neighborhood income and neighborhood

socioeconomic status score (see Supplemental Material, Table S1). Overall, there was sufficient consideration for individual-level socioeconomic status, but the insufficient control

of area-level socioeconomic status may increase the risk of bias.

**Co-morbidities.** Some co-morbidities associated with diabetes may also be associated

**Table 2.** Exposure and outcome definitions.

| Source                            | Outcome                        | Definition of outcome                                                                           | Exposure                                                                      | Definition of exposure                                                                                                                                                                                                                                                                                                                                                                                                             | Exposure estimates                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
|-----------------------------------|--------------------------------|-------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Krämer et al. 2010 <sup>a</sup>   | Incident T2DM                  | Self-reported, physician-diagnosed T2DM                                                         | PM <sub>10</sub> , PM <sub>2.5</sub> , NO <sub>2</sub> , and traffic exposure | 5-year means of PM <sub>10</sub> and NO <sub>2</sub> in an 8-km grid from monitoring stations, before baseline<br><br>Traffic PM and NO <sub>2</sub> in a 1-km grid, in 1 year, from emission inventory<br>Traffic PM <sub>2.5</sub> and NO <sub>2</sub> <sup>b</sup> from a (1-year measurement) LUR model. Distance from the next major road with > 10,000 cars per day                                                          | Median (25th–75th percentile) Monitoring stations (µg/m <sup>3</sup> ):<br>PM <sub>10</sub> : 46.9 (44–54.1)<br>NO <sub>2</sub> : 41.7 (23.3–48.2)<br><br>Traffic emission inventory (tons/year/km <sup>2</sup> ):<br>PM: 0.54 (0.22–1.09)<br>NO <sub>2</sub> : 12 (5.4–24.4)<br>LUR soot (10 <sup>-5</sup> m): 1.89 (1.67–2.06)<br>NO <sub>2</sub> (µg/m <sup>3</sup> ): 34.5 (23.8–38.8)<br>% participants living < 100 m from busy road: 15.8                                                                                                    |
| Andersen et al. 2012 <sup>a</sup> | Incident DM                    | Confirmed DM cases from the Danish National Diabetes Register                                   | NO <sub>2</sub> , NO <sub>x</sub> , traffic exposure                          | 35 <sup>b</sup> - and 15-year mean levels of NO <sub>2</sub> and NO <sub>x</sub> from the Danish AirGIS model before baseline<br>1-year mean NO <sub>2</sub> and NO <sub>x</sub> at baseline<br><br>1-year mean NO <sub>2</sub> and NO <sub>x</sub> at follow-up<br><br>Major road (with annual traffic density of ≥ 10,000) within 50 m of residence.<br>Traffic load within 100 m of residence (10 <sup>3</sup> vehicles/km/day) | Median (IQR)<br>35-year NO <sub>2</sub> and NO <sub>x</sub> (µg/m <sup>3</sup> ): 14.5 (4.9) and 20.9 (11.4)<br>15-year NO <sub>2</sub> and NO <sub>x</sub> (µg/m <sup>3</sup> ): 15.3 (5.6) and 22.1 (12)<br>1-year NO <sub>2</sub> and NO <sub>x</sub> at baseline (µg/m <sup>3</sup> ): 15.4 (5.6) and 20.3 (10.9)<br>1-year NO <sub>2</sub> and NO <sub>x</sub> at follow-up (µg/m <sup>3</sup> ): 15.2 (5.7) and 21.5 (12)<br><br>% major road within 50 m: 8.1<br><br>Traffic load within 100 m (10 <sup>3</sup> vehicles/km/day): 0.34 (1.3) |
| Puett et al. 2011 <sup>a</sup>    | Incident T2DM                  | DM according to the National Diabetes Data Group Criteria <sup>c</sup>                          | PM <sub>2.5</sub> , PM <sub>10</sub> , PM <sub>10-2.5</sub>                   | Average PM <sub>2.5</sub> <sup>b</sup> , PM <sub>10</sub> , and PM <sub>10-2.5</sub> concentrations, from LUR model, 12 months before diagnosis                                                                                                                                                                                                                                                                                    | Mean ± SD<br>PM <sub>2.5</sub> (µg/m <sup>3</sup> ): 18.3 ± 3.1 for HPFS and 17.5 ± 2.7 for NHS<br>PM <sub>10</sub> (µg/m <sup>3</sup> ): 28.5 ± 5.5 for HPFS and 26.9 ± 4.8 for NHS<br>PM <sub>10-2.5</sub> (µg/m <sup>3</sup> ): 10.3 ± 3.3 for HPFS and 9.4 ± 2.9 for NHS                                                                                                                                                                                                                                                                        |
| Coogan et al. 2012 <sup>a</sup>   | Incident T2DM                  | Self-reported, physician-diagnosed T2DM                                                         | PM <sub>2.5</sub> , NO <sub>x</sub> , traffic exposure                        | 1-year mean PM <sub>2.5</sub> <sup>b</sup> during follow-up, assigned by kriging model<br><br>1-year mean NO <sub>x</sub> the year after follow-up, assigned by LUR model                                                                                                                                                                                                                                                          | Mean ± SD<br>PM <sub>2.5</sub> (µg/m <sup>3</sup> ): 20.7 ± 2.1<br>Median (25th–75th percentile) PM <sub>2.5</sub> (µg/m <sup>3</sup> ): 21.1 (20.3–21.6)<br><br>Mean ± SD<br>NO <sub>x</sub> (ppb): 43.3 ± 11<br>Median (25th–75th percentile) NO <sub>x</sub> (ppb): 41.6 (36.9–49.2)                                                                                                                                                                                                                                                             |
| Chen et al. 2013 <sup>a</sup>     | Incident DM                    | Physician-diagnosed DM from Ontario database                                                    | PM <sub>2.5</sub>                                                             | 6-year mean PM <sub>2.5</sub> <sup>b</sup> during baseline/follow-up, obtained from satellite-based estimates at 10 × 10 km resolution                                                                                                                                                                                                                                                                                             | Mean (range) PM <sub>2.5</sub> (µg/m <sup>3</sup> ): 10.6 (2.6–19.1)                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |
| Brook et al. 2008 <sup>a</sup>    | Prevalent DM                   | Physician-diagnosed DM from Ontario Health Insurance Plan and Ontario Health Discharge Database | NO <sub>2</sub>                                                               | NO <sub>2</sub> <sup>b</sup> assigned by LUR models developed from mean field measurements within 3 years, from Hamilton and Toronto, Ontario, Canada                                                                                                                                                                                                                                                                              | Median (25th–75th percentile) NO <sub>2</sub> (ppb)<br>Males:<br>Hamilton: 15.2 (13.9–17.1); Toronto: 23 (20.8–25)<br>Females:<br>Hamilton: 15.3 (14–17); Toronto: 22.9 (20.8–24.7)                                                                                                                                                                                                                                                                                                                                                                 |
| van den Hooven et al. 2009        | Prevalent gestational DM (GDM) | GDM diagnosed according to the Dutch midwifery and obstetric guidelines                         | Traffic exposure                                                              | Distance-weighted traffic density (DWTD) within a 150-m radius around residence (vehicles/24 hr × m)<br>Proximity to a major road (> 10,000 vehicles/day)                                                                                                                                                                                                                                                                          | Median (P25–P75) DWTD (vehicles/24 hr × m): 5.5 × 10 <sup>5</sup> (1.6 × 10 <sup>5</sup> –1.2 × 10 <sup>6</sup> )<br>Proximity to a major road (m): 143 (74–225)                                                                                                                                                                                                                                                                                                                                                                                    |
| Dijkema et al. 2011               | Prevalent T2DM                 | Self-reported physician-diagnosed T2DM. Laboratory-based diagnosis for undetected cases         | NO <sub>2</sub> , traffic exposure                                            | 1-year mean NO <sub>2</sub> assigned by LUR model<br><br>Distance to the nearest main road (≥ 5,000 vehicles/day)<br>Traffic flow at the nearest main road (vehicles/24 hr)<br>Total traffic per 24 hr on all roads within a 250-m circular buffer around the address                                                                                                                                                              | Median (25th–75th percentile) NO <sub>2</sub> (µg/m <sup>3</sup> ): 15.2 (14.2–16.5)<br>Distance to nearest main road (m): 140 (74–220)<br><br>Traffic flow at the nearest main road (10 <sup>3</sup> vehicles/24 hr): 7.31 (5.87–9.67)<br>Traffic within 250-m buffer (10 <sup>3</sup> vehicles/24 hr): 680 (516–882)                                                                                                                                                                                                                              |
| Malmqvist et al. 2013             | Prevalent GDM                  | GDM as defined in the Swedish Medical Birth Registry                                            | NO <sub>x</sub> , traffic exposure                                            | Monthly and trimester means of NO <sub>x</sub> assigned by dispersion modeling at a spatial resolution of 500 × 500 m over the duration of the pregnancy<br><br>Traffic density within a 200-m radius                                                                                                                                                                                                                              | Quartiles of NO <sub>x</sub> exposure (µg/m <sup>3</sup> ):<br>Q1: 2.5–8.9<br>Q2: 9.0–14.1<br>Q3: 14.2–22.6<br>Q4: > 22.7<br><br>Categories of traffic density within 200 m (vehicles/min):<br>1: no road<br>2: < 2<br>3: 2–5<br>4: 5–10<br>5: > 10                                                                                                                                                                                                                                                                                                 |

Table continued



with air pollution. These co-morbidities may include hypertension, myocardial infarction, stroke, asthma, and chronic obstructive pulmonary disease (Brook et al. 2010; Pelle et al. 2012; Vojtková et al. 2012). The longitudinal studies considered some of these co-morbidities (see Supplemental Material, Table S1). Participants with co-morbidities were not excluded from any T2DM study.

**Effect modification.** Several studies reported stronger effects in women compared with men (Andersen et al. 2012; Brook et al. 2008; Chen et al. 2013; Dijkema et al. 2011). Other subgroups reported with potentially increased susceptibility include subjects with low education (Andersen et al. 2012; Chen et al. 2013; Krämer et al. 2010), COPD

(Andersen et al. 2012; Chen et al. 2013), asthma (Andersen et al. 2012), higher waist-to-hip ratio (Andersen et al. 2012), and higher level of subclinical inflammation (Krämer et al. 2010), nonsmokers (Andersen et al. 2012), and subjects < 50 years or > 65 years of age (Chen et al. 2013) (see Supplemental Material, Table S1). No study assessed interaction between different air pollutants, air pollutants and noise, or interaction between air pollutants and genetic polymorphisms.

**Loss to follow-up.** Losses to follow-up and healthy survivor bias present common problems in epidemiological studies. Puett et al. (2011) reported a loss of < 10% in both studied cohorts over 20 years of follow-up, and Coogan et al. (2012) reported < 20%

loss of cohort over 10 years of follow-up. The other longitudinal studies did not report losses to follow-up. None of the studies included sensitivity analyses to estimate the effect of the healthy survivor bias.

**Publication bias.** Although selective reporting and publication bias cannot be ruled out, considering a high probability that negative findings will not be published, we found no indication for such sources of bias (*p*-value of Egger's test > 0.2). Some studies reported negative findings. However, most studies had several markers of air pollution available, and it remains unclear if some markers have been measured but not reported, so some selective reporting may have occurred.

**Table 2.** Continued.

| Source              | Outcome        | Definition of outcome                                                                | Exposure                                                                                    | Definition of exposure                                                                                                                                                                                                                                          | Exposure estimates                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
|---------------------|----------------|--------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Hathout et al. 2006 | Prevalent T1DM | Physician-diagnosed T1DM from the database of Loma Linda University Pediatric Center | O <sub>3</sub> , NO <sub>2</sub> , SO <sub>2</sub> , SO <sub>4</sub> , and PM <sub>10</sub> | Average monthly pollutant exposure (obtained from the U.S. EPA and California Air Resources Board) from birth until diagnosis for cases and until enrollment for controls, assigned to residential ZIP codes                                                    | Mean (95% CI)<br>For cases:<br>O <sub>3</sub> : 29.4 (28, 30.8) ppb<br>SO <sub>4</sub> : 3.6 (3.4, 3.87) µg/m <sup>3</sup><br>SO <sub>2</sub> : 1.6 (1.41, 1.75) ppb<br>NO <sub>2</sub> : 30.3 (28.4, 32.3) ppb<br>PM <sub>10</sub> : 48.6 (45.9, 51.3) µg/m <sup>3</sup><br>For controls:<br>O <sub>3</sub> : 25.8 (25.2, 26.3) ppb<br>SO <sub>4</sub> : 3.3 (3.2, 3.36) µg/m <sup>3</sup><br>SO <sub>2</sub> : 1.5 (1.42, 1.5) ppb<br>NO <sub>2</sub> : 29.7 (29.1, 30.4) ppb<br>PM <sub>10</sub> : 47.4 (46.3, 48.5) µg/m <sup>3</sup> |
| Hathout et al. 2002 | Prevalent T1DM | Physician-diagnosed T1DM from the database of Loma Linda University Pediatric Center | O <sub>3</sub> , NO <sub>2</sub> , SO <sub>2</sub> , SO <sub>4</sub> , and PM <sub>10</sub> | Average monthly pollutant exposure (obtained from the U.S. EPA and California Air Resources Board) from birth until diagnosis for cases and until enrollment for controls, assigned to residential ZIP codes                                                    | Mean ± SD<br>For cases:<br>O <sub>3</sub> : 32.5 ± 5.22 ppb<br>SO <sub>4</sub> : 5.52 ± 0.75 µg/m <sup>3</sup><br>SO <sub>2</sub> : 0.67 ± 0.55 ppb<br>NO <sub>2</sub> : 23.7 ± 7.91 ppb<br>PM <sub>10</sub> : 59.3 ± 12.9 µg/m <sup>3</sup><br>For controls:<br>O <sub>3</sub> : 26.7 ± 9.6 ppb<br>SO <sub>4</sub> : 5.88 ± 1.04 µg/m <sup>3</sup><br>SO <sub>2</sub> : 1.29 ± 0.92 ppb<br>NO <sub>2</sub> : 24.7 ± 7.26 ppb<br>PM <sub>10</sub> : 49.6 ± 14.7 µg/m <sup>3</sup>                                                         |
| Fleisch et al. 2014 | Prevalent GDM  | Failed GCT <sup>d</sup> with ≥ 2 high values on the OGTT <sup>e</sup>                |                                                                                             | PM <sub>2.5</sub> and black carbon from central sites within 40 km of residence<br><br>PM <sub>2.5</sub> and black carbon from spatiotemporal models<br><br>Neighborhood traffic density [(vehicles/day) × km] within 100 m<br>Home roadway proximity (≤ 200 m) | Mean ± SD<br>From central sites:<br>PM <sub>2.5</sub> : 10.9 ± 1.4 µg/m <sup>3</sup><br>Black carbon: 0.9 ± 0.1 µg/m <sup>3</sup><br>From spatiotemporal models:<br>PM <sub>2.5</sub> : 11.9 ± 1.4 µg/m <sup>3</sup><br>Black carbon: 0.7 ± 0.2 µg/m <sup>3</sup><br>Traffic density: 1,621 ± 2,234 (vehicles/day × km)<br>Roadway proximity: 281 ± 13                                                                                                                                                                                    |
| Pearson et al. 2010 | Prevalent DM   | County-level DM prevalence from the Centers for Disease Control and Prevention       | PM <sub>2.5</sub>                                                                           | County annual mean level of PM <sub>2.5</sub> obtained from the U.S. EPA as 36-km model, 12-km model, and surface monitor data                                                                                                                                  | PM <sub>2.5</sub> (µg/m <sup>3</sup> ):<br>2004:<br>36-km model: Q1 mean = 7.71; Q4 mean = 12.11<br>12-km model: Q1 mean = 7.78; Q4 mean = 11.77<br>Ground data: Q1 mean = 9.43; Q4 mean = 12.69<br>2005:<br>36-km model: Q1 mean = 7.69; Q4 mean = 12.75<br>12-km model: Q1 mean = 8.41; Q4 mean = 12.38<br>Ground data: Q1 mean = 9.51; Q4 mean = 13.65                                                                                                                                                                                 |

Abbreviations: AirGIS, Air geographic information system; DM, diabetes mellitus; DWTD, distance-weighted traffic density; EPA, Environmental Protection Agency; GDM, gestational diabetes mellitus; HPFS, Health Professionals Follow-up Study; LUR, land-use regression; NHS, Nurses' Health Study; NO<sub>2</sub>, nitrogen dioxide; NO<sub>x</sub>, nitrogen oxides; O<sub>3</sub>, ozone; OGTT, oral glucose tolerance test; PM, particulate matter; PM<sub>10</sub>, particulate matter ≤ 10 µm in diameter; PM<sub>10-2.5</sub>, particulate matter between 2.5 and 10 µm in diameter; PM<sub>2.5</sub>, particulate matter ≤ 2.5 µm in diameter; SO<sub>2</sub>, sulfur dioxide; SO<sub>4</sub>, sulfate; T1DM, type 1 diabetes mellitus; T2DM, type 2 diabetes mellitus.

<sup>a</sup>Studies included in meta-analysis. <sup>b</sup>Air pollution estimates pooled in the meta-analysis. <sup>c</sup>Elevated plasma glucose concentration on at least two different occasions, one or more DM symptoms and a single elevated plasma glucose concentration, or treatment with hypoglycemic medication. <sup>d</sup>Glucose challenge test: serum glucose 1 hr after a non-fasting 50-g oral glucose load. <sup>e</sup>Oral glucose tolerance test: serum glucose 3 hr after a fasting 100-g glucose load.

**Meta-analysis of studies reporting the association of air pollution and risk of T2DM.** Results of seven studies reporting on risk of T2DM [three on particulate matter with diameter  $\leq 2.5$   $\mu\text{m}$  ( $\text{PM}_{2.5}$ ) and four on nitrogen dioxide ( $\text{NO}_2$ )] were considered for quantitative synthesis. All studies synthesized for  $\text{PM}_{2.5}$  were longitudinal. For  $\text{NO}_2$ , two were longitudinal and two were cross-sectional.

The pooled relative risks of T2DM per 10- $\mu\text{g}/\text{m}^3$  increase in exposure to  $\text{PM}_{2.5}$  (Figure 2) and  $\text{NO}_2$  (Figure 3) were 1.10 (95% CI: 1.02, 1.18) and 1.08 (95% CI: 1.00, 1.17), respectively. The effect was more pronounced in females than in males [ $\text{NO}_2$ : 1.15 (95% CI: 1.05, 1.27) vs. 0.99 (95% CI: 0.93, 1.07);  $\text{PM}_{2.5}$ : 1.14 (95% CI: 1.03, 1.26) vs. 1.04 (95% CI: 0.93, 1.17), respectively] per 10- $\mu\text{g}/\text{m}^3$  increase in exposure. The relative risks were similar across all sensitivity analyses (Table 4). We observed substantial statistical heterogeneity with  $\text{NO}_2$  studies (Table 4). Egger's test was consistently  $> 0.2$  ( $p$ -value) in all cases.

## Discussion

In this systematic review, we considered 13 studies on different types of diabetes. The identified epidemiological evidence is highly diverse: Levels, timing, and assessment of exposure varied, as did the outcome definitions, measures of association, and degree of confounder control. The studies included persons with different age ranges and settings, and some populations included only women. Although there is a risk of bias, the results of the meta-analyses indicate a positive association between traffic-related air pollution and T2DM.

**Pathophysiologic mechanisms of DM-air pollution association.** There is strong evidence supporting the role of inflammation in T2DM (Donath and Shoelson 2011; Sjöholm and Nyström 2006). Chronic activation of inflammatory mechanisms can contribute to chronic insulin resistance and subsequent T2DM. Air pollution has been shown to be inflammatory (Liu et al. 2013; Rajagopalan and Brook 2012). Its potential mechanisms in mediating T2DM include pulmonary and systemic inflammation, directly releasing cytokines, alterations in glucose homeostasis through defective insulin signaling in tissues, immune cells activation in visceral adipose tissues potentiating inflammation (Sun et al. 2009; Xu et al. 2010; Yan et al. 2011), and endoplasmic reticulum stress in the lung and liver in relation with hepatocyte and alveolar cells (Liu et al. 2013; Rajagopalan and Brook 2012).  $\text{PM}_{2.5}$  also acts as a hypothalamic stressor, inducing peripheral inflammation and abnormalities in glucose metabolism (Liu et al. 2013;

Purkayastha et al. 2011).  $\text{PM}_{2.5}$  was also shown to mediate dysfunctional brown adipose and mitochondrial tissues (Liu et al. 2013; Rajagopalan and Brook 2012), which is one of the systemic pathologies in T2DM (Lowell and Shulman 2005).

Chuang et al. (2010) demonstrated that exposure to air pollution [ $\text{PM} \leq 10$   $\mu\text{m}$  ( $\text{PM}_{10}$ ) and ozone] exposure leads to alteration in blood pressure, blood lipids, and hemoglobin A1c, a marker of blood glucose control. Kelishadi et al. (2009) found positive


associations between exposure to  $\text{PM}_{10}$ ,  $\text{NO}_2$ , and insulin resistance among children in Iran. Thiering et al. (2013) later found a positive association between residential proximity to traffic, particulate matter ( $\text{PM}_{10}$ ),  $\text{NO}_2$ , and risk of insulin resistance [homeostatic model assessment (HOMA-IR)] among children who were part of a birth cohort in Germany. Exposure to traffic-related air pollution is also associated with impaired glucose tolerance in pregnancy (Fleisch et al. 2014). Experimental evidence also exists for the association of

**Table 3.** Data synthesized for meta-analysis.

| Source               | Population | Pollutant         | Assignment of individual exposure | Reported fully adjusted estimate (95% CI) <sup>a</sup>         |
|----------------------|------------|-------------------|-----------------------------------|----------------------------------------------------------------|
| Krämer et al. 2010   | Females    | $\text{NO}_2$     | LUR model                         | 1.42 (1.16, 1.73) per 15 $\mu\text{g}/\text{m}^3$ of exposure  |
| Andersen et al. 2012 | Females    | $\text{NO}_2$     | LUR model                         | 1.07 (1.01, 1.13) per 4.9 $\mu\text{g}/\text{m}^3$ of exposure |
|                      | Males      | $\text{NO}_2$     | LUR model                         | 1.01 (0.97, 1.07) per 4.9 $\mu\text{g}/\text{m}^3$ of exposure |
|                      | Both       | $\text{NO}_2$     | LUR model                         | 1.04 (1.00, 1.08) per 4.9 $\mu\text{g}/\text{m}^3$ of exposure |
| Brook et al. 2008    | Females    | $\text{NO}_2$     | LUR model                         | 1.04 (1.00, 1.08) per 1 ppb of exposure                        |
|                      | Males      | $\text{NO}_2$     | LUR model                         | 0.99 (0.95, 1.03) per 1 ppb of exposure                        |
|                      | Both       | $\text{NO}_2$     | LUR model                         | 1.015 (0.98, 1.049) per 1 ppb of exposure                      |
| Puett et al. 2011    | Females    | $\text{PM}_{2.5}$ | LUR model                         | 1.02 (0.94, 1.09) per 4 $\mu\text{g}/\text{m}^3$ of exposure   |
|                      | Males      | $\text{PM}_{2.5}$ | LUR model                         | 1.07 (0.92, 1.24) per 4 $\mu\text{g}/\text{m}^3$ of exposure   |
|                      | Both       | $\text{PM}_{2.5}$ | LUR model                         | 1.03 (0.96, 1.10) per 4 $\mu\text{g}/\text{m}^3$ of exposure   |
| Chen et al. 2013     | Females    | $\text{PM}_{2.5}$ | Satellite-based estimates         | 1.17 (1.03, 1.32) per 10 $\mu\text{g}/\text{m}^3$ of exposure  |
|                      | Males      | $\text{PM}_{2.5}$ | Satellite-based estimates         | 1.03 (0.91, 1.16) per 10 $\mu\text{g}/\text{m}^3$ of exposure  |
|                      | Both       | $\text{PM}_{2.5}$ | Satellite-based estimates         | 1.11 (1.02, 1.21) per 10 $\mu\text{g}/\text{m}^3$ of exposure  |
| Coogan et al. 2012   | Females    | $\text{PM}_{2.5}$ | Kriging model                     | 1.63 (0.78, 3.44) per 10 $\mu\text{g}/\text{m}^3$ of exposure  |
| Dijkema et al. 2011  | Females    | $\text{NO}_2$     | LUR model                         | 1.03 (0.90, 1.16) per 10 $\mu\text{g}/\text{m}^3$ of exposure  |
|                      | Males      | $\text{NO}_2$     | LUR model                         | 0.97 (0.87, 1.09) per 10 $\mu\text{g}/\text{m}^3$ of exposure  |
|                      | Both       | $\text{NO}_2$     | LUR model                         | 1.00 (0.94, 1.06) per 10 $\mu\text{g}/\text{m}^3$ of exposure  |

Abbreviations: LUR, land-use regression;  $\text{NO}_2$ , nitrogen dioxide;  $\text{PM}_{2.5}$ , particulate matter  $\leq 2.5$   $\mu\text{m}$  in diameter.

<sup>a</sup>All odds ratio, hazard ratio, and incident risk ratio estimates were converted to per 10  $\mu\text{g}/\text{m}^3$  of exposure for meta-analysis. Estimates from Dijkema et al. (2011) were derived from reported nonlinear estimates.



**Figure 2.**  $\text{PM}_{2.5}$  and risk of T2DM. Where  $I^2$  is the variation in effect estimates attributable to heterogeneity, D + L (DerSimonian and Laird) overall is the pooled random effect estimate of all studies. I-V (inverse variance) overall is the pooled fixed effects estimate of all studies. Weights are from random-effects analysis. %Weight (D + L) is the weight assigned to each study, based on the inverse of the within- and between-study variance. The size of the blue boxes around the point estimates reflects the weight assigned to each study. The summarized studies were adjusted for age, sex, BMI, smoking, alcohol consumption, and socioeconomic status.

air pollution and type 1 diabetes (T1DM). Ozone is known to alter T-cell dependent immune response, predisposing to autoimmune diseases (Krishna et al. 1998). It may also damage the beta cells of the pancreas possibly as a result of pulmonary reactive oxidative species production and oxidative stress, leading to reduced insulin secretion (Brenner et al. 1993; Kelishadi et al. 2009). Together with sulfate, ozone may have apoptotic properties on the beta cells (Hathout et al. 2006). The use of antioxidant prophylaxis for T1DM also points to the possibility of oxidative or inflammatory mechanisms in T1DM (Albright and Goldstein 1996).

**Strengths and limitations.** Although we applied a very broad search strategy and accepted any study design, there are few published studies on the association of air pollution with T1DM or GDM. In addition, some studies did not allow distinguishing adult T1DM from T2DM. Only three of the seven synthesized studies explicitly analyzed the T2DM risk (Coogan et al. 2012; Dijkema et al. 2011; Krämer et al. 2010). However, because > 90% of adult diabetes diagnoses are T2DM, this is unlikely to substantially affect the conclusions. Overall, the available data are not sufficient to evaluate associations with these diabetes types.

Our analysis on the association with T2DM was based on results from primary studies with unclear to high risk of bias and high diversity among the included studies. We took this into account by using effect estimates modeled to participants' residences, converting all effect estimates to a comparable unit (per 10  $\mu\text{g}/\text{m}^3$  of exposure), stratifying analyses by sex, including only longitudinal studies, and performing other sensitivity analyses.

The high diversity among the studies was reflected in our observation of substantial heterogeneity in the meta-analysis for  $\text{NO}_2$  (Table 4), which synthesized longitudinal and cross-sectional data. This was not observed for  $\text{PM}_{2.5}$ , for which all studies were longitudinal. However, the number of studies was too small to further analyze this heterogeneity.

**Prospects.** Future studies should report scales of exposure assessment (pollutant quantification and traffic exposure proxies) that allow direct comparisons with existing evidence. It would be important to apply comparable models in assigning exposure to participants. Ideally, traffic distance measures should be replaced by objective particle concentration measures and models of near-road traffic-related pollutants such as ultrafine particles of elemental carbon. Also, it would be important to consider various time lags for exposure.


The studies on T1DM found associations with ozone and sulfates. These pollutants can be included in the future models

for T2DM, because pollutants usually occur together in different proportions. Carbon monoxide, lead, oxidative metals, volatile organic compounds, and polycyclic aromatic hydrocarbons are other traffic-related pollutants that may be more deleterious to health but have been given less consideration.

Adjusting for noise exposure is also essential because air pollution and noise can be correlated (Foraster 2013; Kim et al. 2012; Ross et al. 2011; Tétreault et al. 2013) and share health effects. Sørensen et al. (2013) recently reported a positive association between road-traffic noise and incident diabetes, and another large meta-analysis of 10 epidemiologic studies by Cappuccio et al. (2010) found that both quality and quantity of sleep, which are related to noise, were

significant predictors of the risk of T2DM. Consideration of noise is thus necessary in assessing the health effects of air pollution.

Also, socioeconomic variables should be adjusted on the spatial scale, apart from individual-level adjustment. Consideration for this spatial confounding is necessary when individual differences in health outcome are associated with neighborhood characteristics such as neighborhood socioeconomic status (Sheppard et al. 2012). It is crucial that studies on diabetes risk consider established diabetes risk factors including obesity, physical activity, and nutrition. Active and passive smoking should be considered when assessing the effect of air pollution. Lack of information on these creates a high risk for bias.



**Figure 3.**  $\text{NO}_2$  and risk of T2DM. Where  $I^2$  is the variation in effect estimates attributable to heterogeneity, D + L (DerSimonian and Laird) overall is the pooled random-effects estimate of all studies. I-V (inverse variance) overall is the pooled fixed-effects estimate of all studies. Weights are from random-effects analysis. %Weight (D + L) is the weight assigned to each study, based on the inverse of the within- and between-study variance. The size of the blue boxes around the point estimates reflects the weight assigned to each study. The summarized studies were adjusted for age, sex, BMI, smoking, and socioeconomic status.

**Table 4.** Sensitivity analyses and heterogeneity measures.

| Analyses                                            | Population | $\text{NO}_2$     |  | Heterogeneity measures                |  | $\text{PM}_{2.5}$ |  | Heterogeneity measures                |  |
|-----------------------------------------------------|------------|-------------------|--|---------------------------------------|--|-------------------|--|---------------------------------------|--|
|                                                     |            | OR (95% CI)       |  | $I^2$ (%); $p$ -value; $\text{Tau}^2$ |  | OR (95% CI)       |  | $I^2$ (%); $p$ -value; $\text{Tau}^2$ |  |
| Main model (random effects)                         | Males      | 0.99 (0.93, 1.07) |  | 0; 0.744; 0                           |  | 1.04 (0.93, 1.17) |  | 0; 0.486; 0                           |  |
|                                                     | Females    | 1.15 (1.05, 1.27) |  | 46.1; 0.135; 0.0042                   |  | 1.14 (1.03, 1.26) |  | 0; 0.405; 0                           |  |
|                                                     | Overall    | 1.08 (1.00, 1.17) |  | 58.4; 0.025; 0.0063                   |  | 1.10 (1.02, 1.18) |  | 0; 0.473; 0                           |  |
| Studies assessing air pollution before DM diagnosis | Males      | 1.02 (0.92, 1.13) |  | NA; NA; 0                             |  | 1.04 (0.93, 1.17) |  | 0; 0.486; 0                           |  |
|                                                     | Females    | 1.20 (1.10, 1.30) |  | 12.5; 0.285; 0.0006                   |  | 1.13 (1.02, 1.25) |  | 0; 0.344; 0                           |  |
|                                                     | Overall    | 1.12 (1.05, 1.19) |  | 69.8; 0.036; 0.008                    |  | 1.09 (1.01, 1.18) |  | 0; 0.489; 0                           |  |
| Studies including both men and women                | Males      | 0.99 (0.93, 1.07) |  | 0; 0.744; 0                           |  | 1.04 (0.93, 1.17) |  | 0; 0.486; 0                           |  |
|                                                     | Females    | 1.11 (1.01, 1.23) |  | 30.2; 0.238; 0.0023                   |  | 1.13 (1.02, 1.25) |  | 0; 0.344; 0                           |  |
|                                                     | Overall    | 1.05 (0.98, 1.12) |  | 34.9; 0.175; 0.0024                   |  | 1.09 (1.01, 1.18) |  | 0; 0.489; 0                           |  |
| Only longitudinal studies                           | Males      | 1.02 (0.92, 1.13) |  | NA; NA; 0                             |  | 1.04 (0.93, 1.17) |  | 0; 0.486; 0                           |  |
|                                                     | Females    | 1.20 (1.10, 1.30) |  | 12.5; 0.285; 0.0006                   |  | 1.14 (1.03, 1.26) |  | 0; 0.405; 0                           |  |
|                                                     | Overall    | 1.12 (1.05, 1.19) |  | 69.8; 0.036; 0.008                    |  | 1.10 (1.02, 1.18) |  | 0; 0.473; 0                           |  |
| Meta-analysis using fixed-effects model             | Males      | 1.00 (0.93, 1.07) |  | 0; 0.744                              |  | 1.04 (0.93, 1.17) |  | 0; 0.486                              |  |
|                                                     | Females    | 1.15 (1.07, 1.23) |  | 46.1; 0.135                           |  | 1.14 (1.03, 1.26) |  | 0; 0.405                              |  |
|                                                     | Overall    | 1.07 (1.02, 1.13) |  | 58.4; 0.025                           |  | 1.10 (1.02, 1.18) |  | 0; 0.473                              |  |

NA, not applicable.  $I^2$  is the proportion of total variability explained by heterogeneity.  $\text{Tau}^2$  is a measure of among-study variance.



Other forms of bias such as the healthy survivor effect should be taken into account, especially in longitudinal studies. Raaschou-Nielsen et al. (2013b) demonstrated associations between diabetes mortality and NO<sub>x</sub> exposure; thus, diabetes patients exposed to air pollution could die and no longer participate, resulting in incorrect estimates of association if mortality was not taken into consideration.

No included study on this topic was done in developing countries. For generalizability of evidence, research should be extended to developing countries where air pollution (including indoor) is high. This could also help in understanding effects of different air pollution compositions. Indoor air pollution is also associated with diabetes as well as cardiovascular diseases (Lee et al. 2012) and is highly prevalent in developing nations (Lim et al. 2012).

Considering the ambiguity in dose–response relationship in air pollution studies (Smith and Peel 2010), future studies should assess air pollution diabetes association in a dose–response manner. This will help in identifying the point in the dose spectrum where control will yield the most benefits for health policy (Smith and Peel 2010).

Overall, the existing evidence indicates a positive association of air pollution and T2DM risk, although there is high risk of bias. High-quality longitudinal studies are needed (taking into consideration sources and composition of air pollution as well as biomarkers) to improve our understanding of this association.

## REFERENCES

- Alberti KG, Zimmet PZ. 1998. Definition, diagnosis and classification of diabetes mellitus and its complications. Part 1: diagnosis and classification of diabetes mellitus provisional report of a WHO consultation. *Diabet Med* 15:539–553.
- Albright JF, Goldstein RA. 1996. Airborne pollutants and the immune system. *Otolaryngol Head Neck Surg* 114:232–238.
- Andersen ZJ, Hvidberg M, Jensen SS, Ketzel M, Loft S, Sørensen M, et al. 2011. Chronic obstructive pulmonary disease and long-term exposure to traffic-related air pollution: a cohort study. *Am J Respir Crit Care Med* 183:455–461.
- Andersen ZJ, Raaschou-Nielsen O, Ketzel M, Jensen SS, Hvidberg M, Loft S, et al. 2012. Diabetes incidence and long-term exposure to air pollution: a cohort study. *Diabetes Care* 35:92–98.
- Bauer M, Moebus S, Möhlenkamp S, Dragano N, Nonnemacher M, Fuchsluger M, et al. 2010. Urban particulate matter air pollution is associated with subclinical atherosclerosis: results from the HNR (Heinz Nixdorf Recall) study. *Am J Cardiol* 106:1803–1808.
- Brenner HH, Burkart V, Rothe H, Kolb H. 1993. Oxygen radical production is increased in macrophages from diabetes prone BB rats. *Autoimmunity* 15:93–98.
- Brook RD, Jerrett M, Brook JR, Bard RL, Finkelstein MM. 2008. The relationship between diabetes mellitus and traffic-related air pollution. *J Occup Environ Med* 50:32–38.
- Brook RD, Rajagopalan S, Pope CA III, Brook JR, Bhatnagar A, Diez-Roux AV, et al. 2010. Particulate matter air pollution and cardiovascular disease: an update to the scientific statement from the American Heart Association. *Circulation* 121:2331–2378.
- Bui DS, Burgess JA, Matheson MC, Erbas B, Perret J, Morrison S, et al. 2013. Ambient wood smoke, traffic pollution and adult asthma prevalence and severity. *Respirology* 18:1101–1107.
- Cappuccio FP, D'Elia L, Strazzullo P, Miller MA. 2010. Quantity and quality of sleep and incidence of type 2 diabetes: a systematic review and meta-analysis. *Diabetes Care* 33:414–420.
- Chen H, Burnett RT, Kwong JC, Villeneuve PJ, Goldberg MS, Brook RD, et al. 2013. Risk of incident diabetes in relation to long-term exposure to fine particulate matter in Ontario, Canada. *Environ Health Perspect* 121:804–810; doi:10.1289/ehp.1205958.
- Chuang KJ, Yan YH, Cheng TJ. 2010. Effect of air pollution on blood pressure, blood lipids, and blood sugar: a population-based approach. *J Occup Environ Med* 52:258–262.
- Coogan PF, White LF, Jerrett M, Brook RD, Su JG, Seto E, et al. 2012. Air pollution and incidence of hypertension and diabetes mellitus in black women living in Los Angeles. *Circulation* 125:767–772.
- Dijkema MB, Mallat SF, Gehring U, van den Hurk K, Alsema M, van Strien RT, et al. 2011. Long-term exposure to traffic-related air pollution and type 2 diabetes prevalence in a cross-sectional screening-study in the Netherlands. *Environ Health* 10:76; doi:10.1186/1476-069X-10-76.
- Donath MY, Shoelson SE. 2011. Type 2 diabetes as an inflammatory disease. *Nat Rev Immunol* 11:98–107.
- Egger M, Davey Smith G, Schneider M, Minder C. 1997. Bias in meta-analysis detected by a simple, graphical test. *BMJ* 315:629–634.
- Fleisch AF, Gold DR, Rifas-Shiman SL, Koutrakis P, Schwartz JD, Kloog I, et al. 2014. Air pollution exposure and abnormal glucose tolerance during pregnancy: the Project Viva cohort. *Environ Health Perspect* 122:378–383; doi:10.1289/ehp.1307065.
- Foraster M. 2013. Is it traffic-related air pollution or road traffic noise, or both? Key questions not yet settled! *Int J Public Health* 58:647–648.
- Hathout EH, Beeson WL, Ischander M, Rao R, Mace JW. 2006. Air pollution and type 1 diabetes in children. *Pediatr Diabetes* 7:81–87.
- Hathout EH, Beeson WL, Nahab F, Rabadi A, Thomas W, Mace JW. 2002. Role of exposure to air pollutants in the development of type 1 diabetes before and after 5 yr of age. *Pediatr Diabetes* 3:184–188.
- Jacquemine B, Schikowski T, Carsin AE, Hansell A, Krämer U, Sunyer J, et al. 2012. The role of air pollution in adult-onset asthma: a review of the current evidence. *Semin Respir Crit Care Med* 33:606–619.
- Jones PW, Nadeau G, Small JM, Adamek L. 2014. Characteristics of a COPD population categorised using the gold framework by health status and exacerbations. *Respir Med* 108:129–135.
- Kelishadi R, Mirghaffari N, Poursafa P, Gidding SS. 2009. Lifestyle and environmental factors associated with inflammation, oxidative stress and insulin resistance in children. *Atherosclerosis* 203:311–319.
- Kim KH, Ho DX, Brown RJ, Oh JM, Park CG, Ryu IC. 2012. Some insights into the relationship between urban air pollution and noise levels. *Sci Total Environ* 424:271–279.
- Kinney GL, Black-Shinn JL, Wan ES, Make B, Regan E, Lutz S, et al. 2014. Pulmonary function reduction in diabetes mellitus with and without chronic obstructive pulmonary disease. *Diabetes Care* 37:389–395.
- Krämer U, Herder C, Sugiri D, Strassburger K, Schikowski T, Ranft U, et al. 2010. Traffic-related air pollution and incident type 2 diabetes: results from the SALIA cohort study. *Environ Health Perspect* 118:1273–1279; doi:10.1289/ehp.0901689.
- Krishna MT, Madden J, Teran LM, Biscione GL, Lau LC, Withers NJ, et al. 1998. Effects of 0.2 ppm ozone on biomarkers of inflammation in bronchoalveolar lavage fluid and bronchial mucosa of healthy subjects. *Eur Respir J* 11:1294–1300.
- Künzli N, Bridevaux PO, Liu LJ, Garcia-Esteban R, Schindler C, Gerbase MW, et al. 2009. Traffic-related air pollution correlates with adult-onset asthma among never-smokers. *Thorax* 64:664–670.
- Künzli N, Jerrett M, Garcia-Esteban R, Basagaña X, Beckermann B, Gilliland F, et al. 2010. Ambient air pollution and the progression of atherosclerosis in adults. *PLoS One* 5:e9096; doi:10.1371/journal.pone.0009096.
- Lau J, Ioannidis JP, Schmid CH. 1997. Quantitative synthesis in systematic reviews. *Ann Intern Med* 127:820–826.
- Lee MS, Hang JQ, Zhang FY, Dai HL, Su L, Christiani DC. 2012. In-home solid fuel use and cardiovascular disease: a cross-sectional analysis of the Shanghai Putuo study. *Environ Health* 11:18; doi:10.1186/1476-069X-11-18.
- Lim SS, Vos T, Flaxman AD, Danaei G, Shibuya K, Adair-Rohani H, et al. 2012. A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 380:2224–2260.
- Liu C, Ying Z, Harkema J, Sun Q, Rajagopalan S. 2013. Epidemiological and experimental links between air pollution and type 2 diabetes. *Toxicol Pathol* 41:361–373.
- Lowell BB, Shulman GI. 2005. Mitochondrial dysfunction and type 2 diabetes. *Science* 307:384–387.
- Malmqvist E, Jakobsson K, Tinnerberg H, Rignell-Hydbom A, Rylander L. 2013. Gestational diabetes and preeclampsia in association with air pollution at levels below current air quality guidelines. *Environ Health Perspect* 121:488–493; doi:10.1289/ehp.1205736.
- Moher D, Liberati A, Tetzlaff J, Altman DG, PRISMA Group. 2010. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Int J Surg* 8:336–341.
- O'Neill MS, Veves A, Zanobetti A, Sarnat JA, Gold DR, Economides PA, et al. 2005. Diabetes enhances vulnerability to particulate air pollution-associated impairment in vascular reactivity and endothelial function. *Circulation* 111:2913–2920.
- Papazafirovoulou AK, Kardara MS, Pappas SI. 2011. Environmental pollution and diabetes mellitus. *Recent Pat Biomater* 1:44–48.
- Pearson JF, Bachireddy C, Shyamprasad S, Goldfine AB, Brownstein JS. 2010. Association between fine particulate matter and diabetes prevalence in the U.S. *Diabetes Care* 33:2196–2201.
- Pelle AJ, Loerbroks A, Widdershoven JW, Denollet J. 2012. Heart failure and comorbid diabetes mellitus or chronic obstructive pulmonary disease: effects on mood in outpatients. *Int J Cardiol* 154:216–217.
- Puett RC, Hart JE, Schwartz J, Hu FB, Liese AD, Laden F. 2011. Are particulate matter exposures associated with risk of type 2 diabetes? *Environ Health Perspect* 119:384–389; doi:10.1289/ehp.1002344.
- Purkayastha S, Zhang G, Cai D. 2011. Uncoupling the mechanisms of obesity and hypertension by targeting hypothalamic IKK- $\beta$  and NF- $\kappa$ B. *Nat Med* 17:883–887.
- Raaschou-Nielsen O, Andersen ZJ, Beelen R, Samoli E, Stafoggia M, Weinmayr G, et al. 2013a. Air pollution



- and lung cancer incidence in 17 European cohorts: prospective analyses from the European Study of Cohorts for Air Pollution Effects (ESCAPE). *Lancet Oncol* 14:813–822.
- Raaschou-Nielsen O, Andersen ZJ, Hvidberg M, Jensen SS, Ketzel M, Sørensen M, et al. 2011. Air pollution from traffic and cancer incidence: a Danish cohort study. *Environ Health* 10:67; doi:10.1186/1476-069X-10-67.
- Raaschou-Nielsen O, Sørensen M, Ketzel M, Hertel O, Loft S, Tjønneland A, et al. 2013b. Long-term exposure to traffic-related air pollution and diabetes-associated mortality: a cohort study. *Diabetologia* 56:36–46.
- Rajagopalan S, Brook RD. 2012. Air pollution and type 2 diabetes: mechanistic insights. *Diabetes* 61:3037–3045.
- Ross Z, Kheirbek I, Clougherty JE, Ito K, Matte T, Markowitz S, et al. 2011. Noise, air pollutants and traffic: continuous measurement and correlation at a high-traffic location in New York City. *Environ Res* 111:1054–1063.
- Schikowski T, Mills IC, Anderson HR, Cohen A, Hansell A, Kauffmann F, et al. 2014. Ambient air pollution: a cause for COPD? *Eur Respir J* 43:250–263.
- Sheppard L, Burnett RT, Szpiro AA, Kim SY, Jerrett M, Pope CA III, et al. 2012. Confounding and exposure measurement error in air pollution epidemiology. *Air Qual Atmos Health* 5:203–216.
- Sjöholm A, Nyström T. 2006. Inflammation and the etiology of type 2 diabetes. *Diabetes Metab Res Rev* 22:4–10.
- Smith KR, Peel JL. 2010. Mind the gap. *Environ Health Perspect* 118:1643–1645; doi:10.1289/ehp.1002517.
- Sørensen M, Andersen ZJ, Nordsborg RB, Becker T, Tjønneland A, Overvad K, et al. 2013. Long-term exposure to road traffic noise and incident diabetes: a cohort study. *Environ Health Perspect* 121:217–222; doi:10.1289/ehp.1205503.
- Stroup DF, Berlin JA, Morton SC, Olkin I, Williamson GD, Rennie D, et al. 2000. Meta-analysis of observational studies in epidemiology: a proposal for reporting. Meta-analysis Of Observational Studies in Epidemiology (MOOSE) group. *JAMA* 283:2008–2012.
- Sun Q, Yue P, DeJulius JA, Lumeng CN, Kampfrath T, Mikolaj MB, et al. 2009. Ambient air pollution exaggerates adipose inflammation and insulin resistance in a mouse model of diet-induced obesity. *Circulation* 119:538–546.
- Tétreault LF, Perron S, Smargiassi A. 2013. Cardiovascular health, traffic-related air pollution and noise: are associations mutually confounded? A systematic review. *Int J Public Health* 58:649–666.
- Thiering E, Cyrys J, Kratzsch J, Meisinger C, Hoffmann B, Berdel D, et al. 2013. Long-term exposure to traffic-related air pollution and insulin resistance in children: results from the GINIplus and LISAplus birth cohorts. *Diabetologia* 56:1696–1704.
- van den Hooven EH, Jaddoe VW, de Kluizenaar Y, Hofman A, Mackenbach JP, Steegers EA, et al. 2009. Residential traffic exposure and pregnancy-related outcomes: a prospective birth cohort study. *Environ Health* 8:59; doi:10.1186/1476-069X-8-59.
- Vojtková J1, Ciljaková M, Michnová Z, Turcan T. 2012. Chronic complications of diabetes mellitus related to the respiratory system. *Pediatr Endocrinol Diabetes Metab* 18:112–115.
- Whitsel EA, Quibrera PM, Christ SL, Liao D, Prineas RJ, Anderson GL, et al. 2009. Heart rate variability, ambient particulate matter air pollution, and glucose homeostasis: the environmental epidemiology of arrhythmogenesis in the women's health initiative. *Am J Epidemiol* 169:693–703.
- Xu X, Yavar Z, Verdin M, Ying Z, Mihai G, Kampfrath T, et al. 2010. Effect of early particulate air pollution exposure on obesity in mice: role of p47phox. *Arterioscler Thromb Vasc Biol* 30:2518–2527.
- Yan YH, Chou CCK, Lee CT, Liu JY, Cheng TJ. 2011. Enhanced insulin resistance in diet-induced obese rats exposed to fine particles by instillation. *Inhal Toxicol* 23:507–519.
- Zanobetti A, Bind MA, Schwartz J. 2008. Particulate air pollution and survival in a COPD cohort. *Environ Health* 7:48; doi:10.1186/1476-069X-7-48.
- Zanobetti A, Schwartz J. 2001. Are diabetics more susceptible to the health effects of airborne particles? *Am J Respir Crit Care Med* 164:831–833.