

The root hair defective phenotype of *Arabidopsis thaliana* Pol IV subunit mutant *nrpd1a-3* is associated with a deletion in RHD6

Rakesh David^{1, 2}, R. Daniel Kortschak¹ and Iain Searle^{1§}

¹School of Biological Sciences, The University of Adelaide, The University of Adelaide and Shanghai Jiao Tong University Joint International Centre for Agriculture and Health, Adelaide, South Australia 5005, Australia

²Present address: ARC Centre of Excellence in Plant Energy Biology, The School of Agriculture, Food and Wine, University of Adelaide, Waite Campus, Adelaide, South Australia 5005, Australia

[§]To whom correspondence should be addressed: iain.searle@adelaide.edu.au




Figure 1: Root hair defective phenotype of *Pol IV* allele *nrpd1a-3* and identification of a DNA deletion in *Root Hair Defective 6 (RHD6)*. **A.** Root hair phenotype of Col-0 (wild type) and *NRPD1A* T-DNA mutant *nrpd1a-3*. Seedlings were grown in a vertical orientation for 7 days. Scale bar = 0.5 cm. **B.** Root hair numbers of Col-0, C-24, *nrpd1a-1*, *nrpd1a-2*, *nrpd1a-3* and *nrpd1a-4*. Alleles *nrpd1a-1* and *nrpd1a-2* are in the C-24 accession and alleles *nrpd1a-3* and *nrpd1a-4* are in the Col-0 accession. Between 18-20 seedlings roots were examined for each genotype, with an asterisk * denoting a significant difference between wildtype and mutant ($p < 0.005$). Error bars show standard error. **C.** Top, shows a schematic representation of *RHD6* where the green boxes indicate exons and the intervening line black lines indicate introns. Below,

is a genome browser view showing whole genome Illumina re-sequencing reads that aligned to *RHD6* in either Col-0 or *nrpd1a-3*. Green bars indicate sequence reads aligned to the top strand and red bars indicate reads aligned to the bottom strand. In *nrpd1a-3*, the region with no aligned sequence reads indicates the deleted region. Scale bar = 100 bp D. Schematic representation of *RHD6* showing the 1,310 bp deletion in *nrpd1a-3*. The arrows represent oligo-nucleotide primers that were used for semi-quantitative RT-PCR. E. *RHD6* transcripts was amplified from Col-0, *nrpd1a-3* and *nrpd1a-4* by semi-quantitative RT-PCR and the products were separated on an agarose gel. For each genotype two biological replicates were derived from a pool of 10 seedlings. *PP2AA3* was used as a positive control.

Description

In eukaryotes, RNA and chromatin-based pathways control transposable elements (TE) to minimize the deleterious consequences of genetic invasion, transposition, mutation and chromosome instability (Matzke and Moshier, 2014). In higher plants, the multi-subunit nuclear RNA polymerase IV (Pol IV) specializes in transcribing the 24 nucleotide class of small RNAs that target TE's for DNA cytosine methylation and silencing in the RNA-directed DNA methylation (RdDM) pathway (Herr *et al.*, 2005; Onodera *et al.*, 2005). In *Arabidopsis*, the Pol IV has two alternative subunits encoded by the *NRPD1a* and *NRPD1b* and a common subunit encoded by *NRPD2A* (Herr *et al.*, 2005; Kanno *et al.*, 2005; Onodera *et al.*, 2005; Pontier *et al.*, 2005). The null mutant for *NRPD1a* is defective in the RdDM pathway and also displays a late flowering phenotype under short day conditions (Pontier *et al.*, 2005; Eamens *et al.*, 2008).

Interestingly, we observed a root hair defective phenotype in the *nrpd1a-3* mutant allele which has not been previously reported (Fig. 1A). When grown on vertically oriented agar medium, *nrpd1a-3* seedlings lacked root hairs or defective root hair elongation when compared to the Col-0 wild type control which showed normal root hair distribution and length in the root maturation zone (Fig. 1A). Further investigation of other *NRPD1A* mutant alleles, *nrpd1a-1*, *nrpd1a-2*, in the C-24 accession, and *nrpd1a-4* in the Col-0 accession revealed that these alleles had a similar root hair number to the wild type controls (Fig. 1B). To investigate if the root hair defective phenotype in *nrpd1a-3* was a spurious event in our laboratory's seed stock or an earlier event, we re-ordered the same mutant from the ABRC stock Centre and tested the root hair phenotype. We observed the same root hair defective phenotype in both *nrpd1a-3* seed stocks. We next crossed the *nrpd1a-3* mutant (Col-0) to wild type accession C-24, self-fertilized the F_1 to create a F_2 mapping population and mapped polymorphic DNA markers across the 5 chromosomes revealed the gene for the root hair defective phenotype was located between DNA markers *ciw3* and *F26B6* on chromosome 1 (Berendzen *et al.*, 2005). Next we whole-genome sequenced DNA from both Col-0 and *nrpd1a-3* using Illumina short read technology, and after GATK (McKenna *et al.*, 2010) and DELLY (Rausch *et al.*, 2012) analysis of the annotated gene models in the genetic window, we identified only one nucleotide mutation, a 1,310 bp deletion in *Root Hair Defective 6 (RHD6)*, in *nrpd1a-3* (Fig. 1C). *RHD6* is a bHLH transcription factor that positively regulates root hair initiation (Masucci and Schiefelbein, 1994) and loss of function mutations cause a root hair defective phenotype. Semi-quantitative RT-PCR showed *RHD6* mRNA was undetectable in the *nrpd1a-3* roots when compared to Col-0 wild-type and *nrpd1a-4* suggesting that the loss of *RHD6* was the likely candidate for the root hair defective phenotype observed in the *nrpd1a-3* mutant (Fig. 1D). We confirmed the 1,310 bp deletion that deleted part of exon 2, all of exons 3-5 and part of the 3' UTR by PCR and Sanger sequencing. Together the genetic mapping and the undetectable *RHD6* mRNA transcript strongly suggests that the root hair defective phenotype only observed in the *nrpd1a-3* mutant background is caused by the deletion in *RHD6*. In the *Arabidopsis* research community, sometimes phenotypes caused by an unlinked mutation to a gene of interest have been incorrectly associated in the research field for many years (Enders *et al.*, 2015; Habets and Offringa, 2015), and so our discovery of a deletion in *RHD6* in *nrpd1a-3* will allow the community to not incorrectly associate the defective root hair phenotype with POLIV function.

Reagents

For soil grown *A. thaliana* plants, seeds were germinated in soil and seedlings grown under halogen lights at 21°C under 16-hour light (130 $\mu\text{moles}/\text{m}^2/\text{sec}$) and 8-hour dark conditions as previously described (David *et al.*, 2017). For vertical plate experiments, seeds were chlorine gas sterilized as previously described (Burgess *et al.*, 2015) stratified at 4°C in the darkness and then transferred to grow under halogen lights of 16-hour light, 8-hour night conditions (130 $\mu\text{moles}/\text{m}^2/\text{sec}$). Seedlings were grown in a vertical orientation for 7 days on half-strength Murashige and Skoog (MS) medium containing 1% sucrose. Col-0 or C-24 accessions were used as wild-type controls. The *NRPD1A* two SALK T-DNA insertion lines in the Col-0 accessions for *NRPD1A* were *nrpd1a-3* (SALK_128428) and *nrpd1a-4* (SALK_083051). Mutants *nrpd1a-1* and *nrpd1a-2* are in the C-24 accession (Herr *et al.*, 2005). ImageJ was used to measure the number of root hairs longer than 2 mm from an image. Statistical analyses of the data were made using Student's t-test.

RT-PCR and PCR

Total RNA was isolated from seedling roots using the Tri Reagent (Sigma Aldrich) as described by Wang *et al.*, 2017. Briefly, cDNA for semi-quantitative RT-PCR was synthesized using Superscript III kits as per the manufacturer's recommendation (Invitrogen) from 2 mg of total RNA that was oligo(dT) primed. Genomic DNA for PCR or sequencing

was isolated using the Dellaporta procedure (Dellaporta *et al.*, 1983) with small modifications as previously described from two-week-old seedlings grown on agar medium. RT-PCR and PCR reagents and thermal cycling conditions were previously described in David *et al.*, 2017.

Gene name	Locus identifier	Application	Forward primer sequence	Reverse primer sequence
<i>RHD6</i>	AT1G14920	RT-PCR	CCAATGGCACCAAGGTTGATTT	TTTCCCCCGATATTATTACAACGTA
<i>PP2AA3</i>	AT1G13320	RT-PCR control	GGGCAATGCAGCATATAGTTC	TGGGTCTTCACTTAGCTCCAC
<i>RHD6</i>	AT1G66470	detection of deletion	AGGGCAACAACATGAGCTACGGC	TAAGAACACGTATCCCTAAT

Whole-genome sequencing and Bioinformatic analysis

Illumina sequencing libraries were prepared using NEBNext Ultra DNA library Prep kit as per the manufacturer's recommendation and sequenced on the Illumina Hi-Seq system. Illumina sequence reads were trimmed by using TrimGalore! (Krueger, 2015), sequence quality assessed by using FastQC (Andrews, 2010), detection of nucleotide variation was performed by using DELLY (Rausch *et al.*, 2012) and GATK (McKenna *et al.*, 2010). All bioinformatic analysis were performed using default parameters. Sequence data from this article can be found in the SRA accession ID: PRJNA590836.

Gene sequence data from this article can be found in The Arabidopsis Information Resource (TAIR) under the following accession numbers: *NRPD1A* At1g63020; *RHD6* At1g66470 and *PP2AA3* At1g13320.

The *nrpd1a-3 rhd6-4* strain has been submitted to ABRC (Stock ID CS72356).

Acknowledgments: We thank the ACRF Cancer Genomics (Adelaide, Australia) staff for assistance with the genome sequencing.

References

- Andrews S, 2010. FastQC: a quality control tool for high throughput sequence data. Available online at: <https://www.bioinformatics.babraham.ac.uk/projects/fastqc/>
- Berendzen K, Searle I, Ravenscroft D, Koncz C, Batschauer A, Coupland G, Somssich IE, Ulker B. 2005. A rapid and versatile combined DNA/RNA extraction protocol and its application to the analysis of a novel DNA marker set polymorphic between Arabidopsis thaliana ecotypes Col-0 and Landsberg erecta. Plant Methods 1(1): 4. PMID: 16270938.
- Burgess AL, David R, Searle IR. 2015. Conservation of tRNA and rRNA 5-methylcytosine in the kingdom Plantae. BMC Plant Biol 15: 199. PMID: 26268215.
- David R, Burgess A, Parker B, Li J, Pulsford K, Sibbritt T, Preiss T, Searle IR. 2017. Transcriptome-Wide Mapping of RNA 5-Methylcytosine in Arabidopsis mRNAs and Noncoding RNAs. The Plant Cell 29(3): 445-460. PMID: 28062751.
- Eamens A, Vaistij FE, Jones L. 2008. NRPD1a and NRPD1b are required to maintain post-transcriptional RNA silencing and RNA-directed DNA methylation in Arabidopsis. Plant J 55(4): 596-606. PMID: 18433438.
- Enders TA, Oh S, Yang Z, Montgomery BL, Strader LC. 2015. Genome Sequencing of Arabidopsis abp1-5 Reveals Second-Site Mutations That May Affect Phenotypes. Plant Cell 27(7): 1820-1826. PMID: 26106149.
- Habets ME, Offringa R. 2015. Auxin Binding Protein 1: A Red Herring After All? Mol Plant 8(8): 1131-1134. PMID: 25917757.
- Herr AJ, Jensen MB, Dalmay T, Baulcombe DC. 2005. RNA polymerase IV directs silencing of endogenous DNA. Science 308(5718): 118-120. PMID: 15692015.
- Kanno T, Huettel B, Mette MF, Aufsatz W, Jaligot E, Daxinger L, Kreil DP, Matzke M, Matzke AJ. 2005. Atypical RNA polymerase subunits required for RNA-directed DNA methylation. Nat Genet 37(7): 761-765. PMID: 15924141.
- Krueger F. 2015. Trim Galore!: A wrapper tool around Cutadapt and FastQC to consistently apply quality and adapter trimming to FastQ files, with some extra functionality for MspI-digested RRBS-type (Reduced Representation Bisulfite-Seq) libraries methylation. Available online at: https://www.bioinformatics.babraham.ac.uk/projects/trim_galore/
- Masucci JD, Schiefelbein JW. 1994. The *rhd6* Mutation of Arabidopsis thaliana Alters Root-Hair Initiation through an Auxin- and Ethylene-Associated Process. Plant Physiol 106(4): 1335-1346. PMID: 12232412.

Matzke MA, Mosher RA. 2014. RNA-directed DNA methylation: an epigenetic pathway of increasing complexity. *Nat Rev Genet* 15(6): 394-408. PMID: 24805120.

McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, Garimella K, Altshuler D, Gabriel S, Daly M, et al. 2010. The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Res* 20(9): 1297-1303. PMID: 20644199.

Onodera Y, Haag JR, Ream T, Costa Nunes P, Pontes O, Pikaard CS. 2005. Plant nuclear RNA polymerase IV mediates siRNA and DNA methylation-dependent heterochromatin formation. *Cell* 120(5): 613-622. PMID: 15766525.

Pontier D, Yahubyan G, Vega D, Bulski A, Saez-Vasquez J, Hakimi MA, Lerbs-Mache S, Colot V, Lagrange T. 2005. Reinforcement of silencing at transposons and highly repeated sequences requires the concerted action of two distinct RNA polymerases IV in *Arabidopsis*. *Genes Dev* 19(17): 2030-2040. PMID: 16140984.

Rausch T, Zichner T, Schlattl A, Stutz AM, Benes V, Korbel JO. 2012. DELLY: structural variant discovery by integrated paired-end and split-read analysis. *Bioinformatics* 28(18): i333-i339. PMID: 22962449.

Wang D, Qu, Yang L, Zhang Q, Liu Z, Do T, Adelson D, Wang Z, Searle I, Zhu J. 2017 Transposable elements (TEs) contribute to stress-related long intergenic noncoding RNAs in plants. *The Plant J* 90(1):133-146. PMID: 28106309.

Funding: This research was supported by Australian Research Council grant DP190101303.

Author Contributions: Rakesh David: Investigation, Writing - original draft, Writing - review and editing, Visualization. R. Daniel Kortschak: Investigation, Writing - review and editing. Iain Searle: Investigation, Writing - original draft, Writing - review and editing, Visualization, Conceptualization, Funding acquisition, Project administration, Supervision.

Reviewed By: Anonymous

History: Received November 20, 2019 Accepted December 5, 2019 Published December 12, 2019

Copyright: © 2019 by the authors. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International (CC BY 4.0) License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Citation: David, R; Kortschak, RD; Searle, I (2019). The root hair defective phenotype of *Arabidopsis thaliana* Pol IV subunit mutant *nrpd1a-3* is associated with a deletion in RHD6. *microPublication Biology*. <https://doi.org/10.17912/micropub.biology.000196>