



ORAL PRESENTATION

Open Access

Glutathione-s-transferase is a minor allergen in birch pollen because of restricted release from hydrated pollen grains

Stephan Deifl¹*, Christian Zwicker¹, Eva Vejvar², Claudia Kitzmüller¹, Gabriele Gadermaier², Birgit Nagl¹, Susanne Vrtala¹, Gerhard Zlabinger³, Peter Briza², Fatima Ferreira², Barbara Bohle¹

From 5th International Symposium on Molecular Allergology (ISMA 2013)
Vienna, Austria. 6-7 December 2013

Background

Recently, proteomic profiling of birch pollen detected a protein homologous to glutathione-S-transferases (GST) in prominent amounts. In mites, cockroach and fungi, GST are relevant allergens. This tempted us to investigate the allergenicity of GST from birch (bGST).

Methods

bGST was expressed in *Escherichia coli*, purified and characterized by mass spectrometry. BALB/c mice were immunized with bGST or Bet v 1. Antibody and T cell responses were assessed. 217 sera from birch pollen-allergic patients were tested for IgE-reactivity to bGST by ELISA. The allergenicity of bGST was evaluated with IgE-loaded rat basophil leukaemia cells (RBL) expressing the α -chain of the human receptor Fc ϵ RI. Cross-reactivity of IgE between bGST and GST from house dust mite, Der p 8, was assessed with murine and human sera in ELISA. The release kinetics of bGST and Bet v 1 from birch pollen upon hydration were studied by immunoblotting.

Results

Immunization with bGST induced specific IgE and a Th2-dominated cellular immune response comparably to immunization with Bet v 1. Only 13.4% of birch pollen-allergic patients were sensitized to bGST. In RBL assays bGST induced mediator release. GST from birch and house dust mites did not cross-react. In contrast to Bet v 1, bGST showed a limited and delayed release from hydrated birch pollen grains.

Conclusion

bGST induces specific IgE in mice but is of limited sensitizing capacity for humans. In contrast to Bet v 1, the release of bGST from hydrated pollen is restricted. Thus, the minor allergenicity of bGST may be explained by a limited exposure of patients to this protein.

Authors' details

¹Medical University of Vienna, Department of Pathophysiology and Allergy Research, Vienna, Austria. ²University of Salzburg, Department of Molecular Biology, Salzburg, Austria. ³Medical University of Vienna, Institute of Immunology, Vienna, Austria.

Published: 17 March 2014

doi:10.1186/2045-7022-4-S2-O1

Cite this article as: Deifl et al.: Glutathione-s-transferase is a minor allergen in birch pollen because of restricted release from hydrated pollen grains. *Clinical and Translational Allergy* 2014 **4**(Suppl 2):O1.

Submit your next manuscript to BioMed Central
and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



¹Medical University of Vienna, Department of Pathophysiology and Allergy Research, Vienna, Austria

Full list of author information is available at the end of the article