

RESEARCH ARTICLE

Open Access

# Enhanced susceptibility to lipopolysaccharide-induced arthritis and endotoxin shock in interleukin-32 alpha transgenic mice through induction of tumor necrosis factor alpha

Masanori Nakayama<sup>1</sup>, Yasuo Niki<sup>1\*</sup>, Toshiki Kawasaki<sup>1</sup>, Yuki Takeda<sup>1</sup>, Keisuke Horiuchi<sup>1</sup>, Aya Sasaki<sup>2</sup>, Yasunori Okada<sup>2</sup>, Kazuo Umezawa<sup>3</sup>, Hiroyasu Ikegami<sup>1</sup>, Yoshiaki Toyama<sup>1</sup> and Takeshi Miyamoto<sup>1</sup>

## Abstract

**Introduction:** The present study assessed the potential functions of interleukin (IL)-32 $\alpha$  on inflammatory arthritis and endotoxin shock models using IL-32 $\alpha$  transgenic (Tg) mice. The potential signaling pathway for the IL-32-tumor necrosis factor (TNF) $\alpha$  axis was analyzed *in vitro*.

**Methods:** IL-32 $\alpha$  Tg mice were generated under control of a ubiquitous promoter. Two disease models were used to examine *in vivo* effects of overexpressed IL-32 $\alpha$ : Toll-like receptor (TLR) ligand-induced arthritis developed using a single injection of lipopolysaccharide (LPS) or zymosan into the knee joints; and endotoxin shock induced with intraperitoneal injection of LPS and D-galactosamine. TNF $\alpha$  antagonist etanercept was administered simultaneously with LPS in some mice. Using RAW264.7 cells, *in vitro* effects of exogenous IL-32 $\alpha$  on TNF $\alpha$ , IL-6 or macrophage inflammatory protein 2 (MIP-2) production were assessed with or without inhibitors for nuclear factor kappa B (NF $\kappa$ B) or mitogen-activated protein kinase (MAPK).

**Results:** Single injection of LPS, but not zymosan, resulted in development of severe synovitis with substantial articular cartilage degradation in knees of the Tg mice. The expression of TNF $\alpha$  mRNA in inflamed synovia was highly upregulated in the LPS-injected Tg mice. Moreover, the Tg mice were more susceptible to endotoxin-induced lethality than the wild-type control mice 48 hours after LPS challenge; but blockade of TNF $\alpha$  by etanercept protected from endotoxin lethality. In cultured bone marrow cells derived from the Tg mice, overexpressed IL-32 $\alpha$  accelerated production of TNF $\alpha$  upon stimulation with LPS. Of note, exogenously added IL-32 $\alpha$  alone stimulated RAW264.7 cells to express TNF $\alpha$ , IL-6, and MIP-2 mRNAs. Particularly, IL-32 $\alpha$ -induced TNF $\alpha$ , but not IL-6 or MIP-2, was inhibited by dehydroxymethylepoxyquinomicin (DHMEQ) and U0126, which are specific inhibitors of nuclear factor kappa B (NF $\kappa$ B) and extracellular signal regulated kinase1/2 (ERK1/2), respectively.

**Conclusions:** These results show that IL-32 $\alpha$  contributed to the development of inflammatory arthritis and endotoxin lethality. Stimulation of TLR signaling with LPS appeared indispensable for activating the IL-32 $\alpha$ -TNF $\alpha$  axis *in vivo*. However, IL-32 $\alpha$  alone induced TNF $\alpha$  production in RAW264.7 cells through phosphorylation of inhibitor kappa B (I $\kappa$ B) and ERK1/2 MAPK. Further studies on the potential involvement of IL-32 $\alpha$ -TNF $\alpha$  axis will be beneficial in better understanding the pathology of autoimmune-related arthritis and infectious immunity.

\* Correspondence: y.niki@lib.bekkoame.ne.jp

<sup>1</sup>Department of Orthopaedic Surgery, Keio University, 35 Shinanomachi, Shinjuku, Tokyo 160-8582, Japan

Full list of author information is available at the end of the article

## Introduction

Interleukin-32 (IL-32) was originally identified as natural killer (NK) transcript 4, which is induced by IL-18 in NK cells [1]. NK transcript 4 showed cytokine-like characteristics and played a critical role in inflammation and was therefore renamed IL-32. This cytokine is reportedly produced by NK cells, T cells, epithelial cells, monocytes, and fibroblasts after stimulation by IL-2, IL-12, and IL-18 and interferon-gamma [2]. Initially, four isoforms of IL-32 (IL-32 $\alpha$ ,  $\beta$ ,  $\gamma$ , and  $\delta$ ) derived from alternative splicing of a single gene. Among these, IL-32 $\alpha$  is the shortest transcript, whereas IL-32  $\gamma$  is the longest isoform and has the strongest biological activity [2,3]. Two additional isoforms, IL-32 $\epsilon$  and  $\zeta$ , have recently been identified, but these isoforms are not ubiquitously expressed except in T cells [4].

IL-32 has been shown to exhibit properties typical of a proinflammatory cytokine and to drive the induction of other proinflammatory cytokines and chemokines, such as tumor necrosis factor-alpha (TNF $\alpha$ ) and IL-1, IL-6, and IL-8. Owing to such proinflammatory properties, IL-32 has been considered to play a key role in the development of various inflammatory diseases, including rheumatoid arthritis (RA), inflammatory bowel disease [5], mycobacterial [6,7] or viral [8-10] infection, chronic obstructive pulmonary disease [11], and pancreatic tumor [12,13]. Although no receptor or analog of IL-32 has yet been identified in mice, human IL-32 reportedly exerts proinflammatory effects as an inducer of TNF $\alpha$  and other inflammatory cytokines in mice both *in vitro* and *in vivo* [2,14-16].

During the last decade, TNF $\alpha$  and IL-6 became widely perceived as substantial therapeutic targets in RA given that the use of either anti-TNF $\alpha$  or anti-IL-6 therapy could successfully control chronic inflammation in RA. As IL-32 is capable of inducing TNF $\alpha$  and IL-6, this cytokine is increasingly becoming a focus as a potential therapeutic target in RA and other inflammatory disorders. Mounting evidence regarding upstream signaling regulators for IL-32 production has been accumulating in the literature [12,17-20]. However, signaling pathways that are downstream of IL-32 and that lead to TNF $\alpha$  production have yet to be fully elucidated. Most investigators advocate the position that IL-32 augments Toll-like receptor (TLR) signaling, and TLR-2, -3, and -4 are associated with the effects of IL-32 signaling, although the detailed mechanisms remain to be clarified. Only a few studies to date have reported the implications of mitogen-activated protein kinase (MAPK) or nuclear factor kappa B (NF- $\kappa$ B) pathways in IL-32 signaling [2,21-23].

The present study generated IL-32 $\alpha$  transgenic (Tg) mice that overexpressed human IL-32 $\alpha$  under a control of ubiquitous CAG promoter, and it assessed the *in vivo* effects of IL-32 $\alpha$  on TLR signaling in the induction of

arthritis and endotoxin shock models using the Tg mice. In addition, the potential signaling pathway of the IL-32 $\alpha$ -TNF $\alpha$  axis was analyzed *in vitro*.

## Materials and methods

### Reagents

Lipopolysaccharide (LPS) from *Escherichia coli* 0111:B4 and zymosan A from *Saccharomyces cerevisiae* were purchased from Sigma-Aldrich (St. Louis, MO, USA), and D-galactosamine was purchased from Wako Pure Chemical Industries (Osaka, Japan). Etanercept was obtained from Wyeth (Tokyo, Japan). Recombinant human IL-32 $\alpha$  protein (rIL-32 $\alpha$ ) was obtained from Takara Bio (Shiga, Japan). IL-32 $\alpha$ -specific enzyme-linked immunosorbent assay (ELISA) (Human IL-32 $\alpha$  ELISA MAX Deluxe Set) was purchased from BioLegend (San Diego, CA, USA), and TNF $\alpha$ -specific ELISA (Quantikine Mouse TNF $\alpha$ ) and anti-IL-32 $\alpha$  antibody were purchased from R&D Systems (Minneapolis, MN, USA). All other antibodies were purchased from Cell Signaling Technology Japan (Tokyo, Japan). Dehydroxymethylepoxyquinomicin (DHMEQ) (an inhibitor for NF- $\kappa$ B) was provided as previously described [24,25]. MAPK inhibitors U0126, SB203580, and SP600125 - which are inhibitors for ERK1/2, p38, and JNK, respectively - were purchased from Sigma-Aldrich. DHMEQ and MAPK inhibitors were dissolved in 100% dimethyl sulfoxide (DMSO) at 100 mg/mL and were stored in aliquots at -30°C. Before use in cell culture, they were diluted with the medium to a final DMSO concentration of not more than 0.05%.

### Real-time polymerase chain reaction analysis

Total RNAs were isolated from each sample by RNAiso plus (Takara Bio), and cDNAs were synthesized from total RNAs by using a PrimeScript RT reagent Kit (Takara Bio). Real-time polymerase chain reaction (PCR) was performed by using SYBR Premix ExTaq II (Takara Bio) with a DICE Thermal cycler (Takara Bio) in accordance with the instructions of the manufacturer. Results were normalized to glyceraldehyde-3-phosphate dehydrogenase (GAPDH) as the fold change compared with samples. The primer sequences used in this study are presented in Table 1.

### Immunoblot analysis

Mice organs or cell lysates were prepared by using RIPA buffer: 1% Triton X-100, 1% sodium deoxycholate, 0.1% SDS, 150 mM NaCl, 10 mM Tris-HCl, pH 7.5, 5 mM ethylenediaminetetraacetic acid (EDTA), and a protease inhibitor cocktail. An equal volume of protein from each lysate was separated by SDS-PAGE, and separated proteins were transferred to nitrocellulose membranes. After transfer, the membrane was blocked with 5% non-fat dried skim milk for 1 hour at room temperature and then incubated with the primary antibody against the

**Table 1 Primers used in real-time polymerase chain reaction**

Primer	Sense	Sequence
Mouse GAPDH	Forward	5'-CTTTGTCAAGCTCATTTCCTGG-3'
	Reverse	5'-TCTTGCTCAGTGCCTTGC-3'
Human IL-32 $\alpha$	Forward	5'-CCTTGGCTCCTTGAACCTTTTG-3'
	Reverse	5'-CTGTCCACGTCCTGATTCTG-3'
Mouse TNF $\alpha$	Forward	5'-CTTCTGTCTACTGAACCTCGGG-3'
	Reverse	5'-CAGGCTTGTCACTCGAATTTTG-3'
Mouse IL-6	Forward	5'-CAAAGCCAGAGTCCTTCAGAG-3'
	Reverse	5'-GTCCTTAGCCACTCCTCTG-3'
Mouse MIP-2	Forward	5'-GAAGTCATAGCCACTCTCAAGG-3'
	Reverse	5'-CTTCCGTTGAGGGACAGC-3'

GAPDH, glyceraldehyde-3-phosphate dehydrogenase; IL, interleukin; MIP-2, macrophage inflammatory protein 2; TNF $\alpha$ , tumor necrosis factor-alpha.

target molecule overnight. Next, the membrane was incubated with horseradish peroxidase-conjugated secondary antibody for 1 hour at room temperature. After washing, protein was detected by using an enhanced chemiluminescence (ECL) system. The same membrane was then stripped by Tris-HCL (62.5 mM), 2% SDS, and  $\beta$ -mercaptoethanol and blocked with skim milk and incubated with primary antibody against another molecule, following the same procedure mentioned above.

#### Generation of human IL-32 $\alpha$ transgenic mice

We generated IL-32 $\alpha$  Tg mice, which overexpressed human IL-32 $\alpha$  under a control of a ubiquitous CAG promoter constructed by the first intron of the chicken  $\beta$ -actin gene and a portion of the rabbit  $\beta$ -globin gene [26]. The background of these mice was C57BL/6 Jcl (obtained from CLEA Japan, Tokyo, Japan). Wild-type C57BL/6 Jcl (Wt) mice were also obtained from CLEA Japan. All mice were 18 to 20 weeks old when they were used. All animal experiments were conducted in accordance with institutional and national guidelines. IL-32 $\alpha$  insertion was confirmed by amplification of the genome DNA isolated from mouse organs by using real-time PCR. The reason real-time PCR was employed was that we wanted to simultaneously detect and quantify transgene-derived IL-32 $\alpha$  in multiple organs of Tg mice. The amount of TNF $\alpha$  expression in multiple organs was measured by real-time PCR. IL-32 $\alpha$  protein in multiple organs was detected by immunoblot analysis. Levels of IL-32 $\alpha$  in blood serum, knee, and liver lysate were measured by specific ELISA.

#### Inflammatory arthritis model

Inflammatory arthritis was induced as described previously [27-29]. As in these reports, intra-articular injection of LPS was employed in IL-32 $\alpha$  Tg mice to elucidate the arthritogenic capacity of IL-32 $\alpha$ . Knees of Tg and Wt mice were injected once with LPS (1  $\mu$ g, much less than

the reported amount [27,28]) or zymosan (20  $\mu$ g, much less than the reported amount [29]) without any other material as a booster. Injection of phosphate-buffered saline (PBS) served as a control to the contralateral knee of the same mouse. Two weeks after injection, a histopathological examination was performed, and TNF $\alpha$  mRNA expression in synovia of knees was quantitatively measured by using real-time PCR.

#### Endotoxin shock model

Endotoxin shock was induced by intraperitoneal injections of 5  $\mu$ g of LPS and 10 mg of D-galactosamine [30]. Half of the mice were injected with 100  $\mu$ g of etanercept at the same time. All injected mice were closely monitored every hour for the first 16 hours and every 3 to 6 hours thereafter for 48 hours. In an additional experiment, we sacrificed mice at 1, 3, and 6 hours after intraperitoneal injection and extracted the liver, spleen, and blood serum. TNF $\alpha$  mRNA of liver and spleen lysates was measured by real-time PCR, and serum TNF $\alpha$  was measured by ELISA.

#### Cell culture

Bone marrow (BM) was extracted from the leg bones of Tg or Wt mice and cultured for 72 hours in alpha-minimum essential medium (Sigma-Aldrich) containing 10% fetal calf serum (FCS), 100 U/mL penicillin, 100 mg/mL streptomycin, and 1% Gluta Max (Gibco Invitrogen, Carlsbad, CA, USA) supplemented with macrophage colony-stimulating factor (50 ng/mL; Kyowa Hakko Kirin, Tokyo, Japan). RAW 264.7 cells were cultured in Dulbecco's modified Eagle's medium (high-glucose) (Gibco Invitrogen) containing 10% FCS, 100 U/mL penicillin, and 100  $\mu$ g/mL streptomycin.

#### Stimulation of cells

BM cells from Tg and Wt mice ( $10^6$  cells) were incubated under stimulation of LPS (50 ng/mL) for 1, 3, 6, and 24 hours, and the amount of TNF $\alpha$  was measured by real-time PCR. TNF $\alpha$  concentration in culture media of BM cells ( $10^5$  cells) after 24-hour incubation with LPS (5, 10, or 50 ng/mL) was measured by specific ELISA. RAW 264.7 cells ( $2 \times 10^4$  cells) were cultured with rIL-32 $\alpha$  or LPS for 24 hours, and the TNF $\alpha$  concentration in culture media was measured by ELISA. For the analysis of signaling pathway, RAW 264.7 cells ( $2 \times 10^4$  cells) were stimulated with rIL-32 $\alpha$  in combination with the specific inhibitors of NF- $\kappa$ B and MAPKs, such as DHMEQ (0.4, 2, or 10  $\mu$ g/mL), U0126 (0.2, 1, or 5  $\mu$ M), SB203580 (0.2, 1, or 5  $\mu$ M), and SP600125 (0.4, 2, or 10  $\mu$ M). DMSO (final concentration of 0.05%) served as a control. After 24 hours of culture, the level of TNF $\alpha$  in culture media was measured by ELISA. For immunoblot analysis, RAW 264.7 cells ( $2 \times 10^6$  cells) were cultured with rIL-32 $\alpha$  for

5, 10, 30, 60, 90, 120, or 180 minutes, followed by washing with ice-cold PBS and lysis in RIPA detergent buffer. The resultant cell lysates were then immunoblotted by using affinity-purified antibodies against phospho-I $\kappa$ B, I $\kappa$ B, phospho-ERK1/2, ERK1/2, phospho-p38, p38, phospho-JNK, JNK, and  $\beta$ -actin. mRNA expressions for IL-6 and macrophage inflammatory protein 2 (MIP-2) as well as TNF $\alpha$  in IL-32-stimulated RAW 264.7 cells ( $2 \times 10^4$  cells) were measured by real-time PCR, and the inhibitory effects of specific signaling inhibitors, including DHMEQ, U0126, SB203580, and SP600125, were analyzed.

### Statistical analysis


Results are reported as the mean  $\pm$  standard deviation. Statistical analysis was undertaken by using a two-tailed Student *t* test. Differences were considered statistically

significant at a *P* value of less than 0.05. The endotoxin shock model was graphed in Kaplan-Meier format and analyzed by a log-rank test. All experiments were performed in three or four times.

## Results

### Generation of human IL-32 $\alpha$ transgenic mice

Human IL-32 $\alpha$  Tg mice were designed to overexpress human IL-32 $\alpha$  by using CAG promoter. Of seven F0 mice, two mice expressing sufficient levels of IL-32 $\alpha$  mRNA were used to establish lines. The F0 mice and all offspring exhibited no evident pathological phenotype, had a normal body weight, and developed and bred normally. Real-time PCR analysis of the Tg mouse lines demonstrated high levels of IL-32 $\alpha$  mRNA expression in a variety of organs, prominently in the knee joint and cardiac muscle (Figure 1a). Transgene-derived IL-32 $\alpha$



**Figure 1 Overexpression of transgene-derived interleukin-32 alpha (IL-32 $\alpha$ ) in transgenic (Tg) mice.** Levels of transgene-derived human IL-32 $\alpha$  mRNA in various organs were measured by real-time polymerase chain reaction. IL-32 $\alpha$  mRNA expression was prominent in knee joints and heart (a). Transgene-derived IL-32 $\alpha$  protein was not detected in serum from Tg mice (b) but was detected in multiple organs (c). Levels of tumor necrosis factor-alpha (TNF $\alpha$ ) mRNA constitutively expressed in various organs were analyzed for Tg and wild-type (Wt) mice. The relative quantities of Tg were divided by those of Wt of each organ and that ratio is graphed (d). In particular, TNF $\alpha$  expression levels in colon and knee joint of Tg mice corresponded to six- to seven-fold those of their littermates. The values are expressed as the mean  $\pm$  standard deviation for four different mice. GAPDH, glyceraldehyde-3-phosphate dehydrogenase.




protein could be detected in multiple organs (Figure 1b, c) but not in serum from Tg mice (Figure 1b). This result might be because the IL-32 $\alpha$  isotype has been reported to remain intracellularly [2]. Constitutive expression of TNF $\alpha$  mRNA induced by overexpressed IL-32 $\alpha$  was apparent in most organs, and expression levels in the colon and knee joint from Tg mice reached six to seven times the levels seen in littermates (Figure 1d).

#### Single intra-articular injection of LPS, but not zymosan, induced inflammatory synovitis and cartilage degradation in transgenic mice


Mice were sacrificed 2 weeks after a single injection of LPS or zymosan, followed by a histopathological examination of the knee joints. The results indicated that the single injection of LPS, but not zymosan, resulted in the development of severe synovitis with articular cartilage destruction in the knees of Tg mice. Such LPS-induced arthritis did not occur in the knees of Wt mice or in contralateral knees injected with PBS (Figure 2a). The level of TNF $\alpha$  mRNA expressed in inflamed synovia after LPS injection was significantly higher in Tg mice than in Wt mice (Figure 2b).

#### Transgenic mice exhibited severe endotoxin lethality after LPS challenge

As a single intraperitoneal injection of LPS with D-galactosamine has been perceived to be capable of inducing endotoxin shock in mice, the impacts of constitutive expression of IL-32 $\alpha$  and subsequently produced TNF $\alpha$  on endotoxin lethality were investigated. Mice receiving an intraperitoneal injection of LPS started to die in 5 hours, and the survival rates at 48 hours after injection were 41% for Tg mice and 75% for Wt mice (Figure 3a), showing statistical significance ( $P < 0.05$ ). Importantly, blockade of TNF $\alpha$  by simultaneous administration of etanercept protected from endotoxin shock and markedly increased survival rate in both Tg and Wt mice, suggesting that IL-32-induced TNF $\alpha$  played a key role in developing endotoxin shock. TNF $\alpha$  mRNA expression in liver and spleen peaked at 1 hour after injection and at much higher levels than those of Wt mice ( $P < 0.01$ ) (Figure 3b, c). A similar temporal course was observed for the level of TNF $\alpha$  protein in serum (Figure 3d). At 1 hour after injection, levels of TNF $\alpha$  mRNA in the liver and spleen and TNF $\alpha$  protein in serum were significantly higher in Tg mice than in Wt mice ( $P < 0.05$ ).



**Figure 2 Inflammatory arthritis induced by single intra-articular injection of lipopolysaccharide (LPS) in transgenic (Tg) mice.** One microgram of LPS or 20  $\mu$ g of zymosan was injected into the knee joints of Tg and wild-type (Wt) mice. Histopathologically, severe synovitis with articular cartilage destruction was observed 2 weeks after injection in Tg mice injected with LPS but not in Tg mice injected with zymosan or phosphate-buffered saline or in Wt mice ( $\times 100$ ) (a). The level of tumor necrosis factor-alpha (TNF $\alpha$ ) mRNA expressed in inflamed synovia was significantly higher in Tg mice injected with LPS than in Tg mice injected with zymosan or in Wt mice injected with LPS ( $*P < 0.01$ ) (b). The values are expressed as the mean  $\pm$  standard deviation for four different mice. GAPDH, glyceraldehyde-3-phosphate dehydrogenase.



**Figure 3 Interleukin-32 alpha overexpression in mice increased the lethality of lipopolysaccharide (LPS)-induced endotoxin shock.** Endotoxin shock was induced in both transgenic (Tg) mice ( $n = 24$ ) and wild-type (Wt) mice ( $n = 24$ ) by intraperitoneal injection of LPS with D-galactosamine. Twelve mice of each group were simultaneously administered etanercept. All mice were monitored for survival up to 48 hours. Survival rate was significantly worse in the Tg group than in the Wt group ( $*P < 0.05$ ). Etanercept protected against endotoxin shock in both groups (a). Temporal changes in TNF $\alpha$  levels were measured by real-time polymerase chain reaction for liver and spleen lysate or enzyme-linked immunosorbent assay for blood serum. Levels of tumor necrosis factor- $\alpha$  (TNF $\alpha$ ) mRNA and protein peaked at 1 hour after LPS injection in the liver (b), spleen (c), and serum (d) from Tg and Wt mice, but levels were much higher in Tg mice than in Wt mice ( $*P < 0.05$ ,  $**P < 0.01$ ). The values are expressed as the mean  $\pm$  standard deviation for four different mice. GAPDH, glyceraldehyde-3-phosphate dehydrogenase.


#### Endogenously overexpressed IL-32 $\alpha$ accelerated production of TNF $\alpha$ upon stimulation with LPS

To examine the effects of endogenous IL-32 $\alpha$  on TNF $\alpha$  production *in vitro*, BM macrophages derived from IL-32 $\alpha$  Tg and Wt mice were used. The level of TNF $\alpha$  mRNA expression was significantly higher in Tg mice than in Wt mice after stimulation with LPS ( $P < 0.05$ ). Temporal changes in TNF $\alpha$  mRNA expression revealed that the level of TNF $\alpha$  mRNA peaked at 3 hours after LPS stimulation and gradually decreased with time (Figure 4a). LPS increased TNF $\alpha$  secretion into culture media in a dose-dependent manner, and the amounts of TNF $\alpha$  produced by BM macrophages were universally higher in Tg mice than in Wt mice throughout all doses of LPS tested (Figure 4b).

#### Exogenous IL-32 $\alpha$ enhanced TNF $\alpha$ production in RAW 264.7 cells through NF- $\kappa$ B and ERK1/2 signaling pathways

To elucidate the effects of exogenous IL-32 $\alpha$  on TNF $\alpha$  production *in vitro*, rIL-32 $\alpha$  was added to RAW 264.7

cells in culture. Although RAW 264.7 cells constitutively produced substantial amounts of TNF $\alpha$ , rIL-32 $\alpha$  alone as well as LPS could further stimulate RAW 264.7 cells to produce TNF $\alpha$  (Figure 5a). DHMEQ and U0126, as inhibitors of NF- $\kappa$ B and ERK1/2, respectively, reduced IL-32 $\alpha$ -induced TNF $\alpha$  production in a dose-dependent manner, whereas SB203580 and SP600125, as inhibitors of p38 and JNK, respectively, did not (Figure 5b). Immunoblot analysis revealed that exogenous IL-32 $\alpha$  clearly phosphorylated I $\kappa$ B and ERK1/2, both starting at 30 minutes and peaking at 90 minutes for ERK1/2 and at 120 minutes for I $\kappa$ B, whereas significant phosphorylation was not observed in p38 or JNK (Figure 5c). These results supported the finding that DHMEQ and U0126, but not SB203580 and SP600125, inhibited IL-32 $\alpha$ -induced TNF $\alpha$  production. Consequently, exogenous IL-32 $\alpha$ -induced TNF $\alpha$  production was mediated predominantly through the activation of NF- $\kappa$ B and the MEK-ERK signaling pathway.



### Exogenous IL-32 $\alpha$ stimulated IL-6 and MIP-2 expression in RAW 264.7 cells independently of NF- $\kappa$ B and MAPK signaling pathways


The effects of exogenous IL-32 $\alpha$  on IL-6 and MIP-2 production were examined since these cytokines were reportedly induced by IL-32 [3]. rIL-32 $\alpha$  alone stimulated RAW 264.7 cells to express TNF $\alpha$ , IL-6, and MIP-2 mRNAs to a similar degree. Specific signaling inhibitors, DHMEQ and U0126, suppressed the expression of TNF $\alpha$  mRNAs; however, neither of these two inhibitors affected the expression of IL-6 and MIP-2 mRNAs induced by IL-32 $\alpha$  (Figure 6), suggesting that a signaling pathway other than NF- $\kappa$ B and MAPKs might be involved in IL-6 and MIP-2 mRNA expressions.

### Discussion

To date, the arthritogenic role of IL-32 has been elucidated on the basis of accumulated evidence that overexpression of IL-32 $\beta$  in a mouse model using bone transplantation exacerbated collagen-induced arthritis in mice [14] and that intra-articular injection of IL-32 $\gamma$  in mouse knee joints resulted in severe joint inflammation [15]. In this study, although IL-32 $\alpha$  Tg mice did not spontaneously exhibit any abnormal phenotype, intra-articular injection of low-dose LPS resulted in the development of inflammatory arthritis. However, injection of zymosan was not capable of sufficiently inducing TNF $\alpha$

and subsequent arthritis. As LPS is known as a specific ligand of TLR-4, interaction of IL-32 $\alpha$  with TLR-4 may play a critical role in the development of arthritis, and this was also the case in LPS-triggered endotoxin shock in the Tg mice. This endotoxin shock model provided an excellent means to evaluate the effects of IL-32 $\alpha$  on infectious immunity. In the present study, IL-32 $\alpha$  overproduction in Tg mice was associated with severe endotoxin lethality; this was shown to be mediated through the induction of TNF $\alpha$ , because etanercept significantly attenuated the endotoxin shock.

Although the present study clearly demonstrated that LPS, as a TLR-4 agonist, but not the TLR-2 agonist zymosan, might play a key role in potentiating the proinflammatory activity of IL-32 $\alpha$ , how exactly IL-32 $\alpha$  interacted with the TLR-4 signaling pathway remains unclear. Most recently, Heinhuis and colleagues [31] reported that LPS co-stimulation was mandatory to elicit IL-32 bioactivity in THP-1 cells, and the present study obtained similar findings that TNF $\alpha$  production promoted by IL-32 $\alpha$  required co-stimulation with LPS (Figure 4). In terms of the interaction between IL-32 and TLR-2/NOD2 (nucleotide-binding oligomerization domain-containing protein 2) signaling, IL-32 has been reported to stimulate TNF $\alpha$ , IL-6, and IL-8 production by directly increasing expression of TLR-2 and NOD2 [32]. Conversely, the interaction of IL-32 with TLR-4




**Figure 5 Exogenous 'IL-32α induced' TNFα production by RAW 264.7 cells through activation of NF-κB and ERK1/2 MAPK.** Two different concentrations of rIL-32α or LPS were added to RAW 264.7 cells in culture, followed by incubation for 24 hours. The level of TNFα in culture media was measured by a specific enzyme-linked immunosorbent assay (a). IL-32α alone as well as LPS was capable of inducing TNFα secretion into culture media by RAW 264.7 cells. Values are expressed as ± standard deviation (SD) of four determinations. RAW 264.7 cells were cultured with or without rIL-32α for 24 hours in combination with inhibitors for IκB and MAPKs, including DHMEQ, U0126, SB203580, and SP600125 (b). IL-32α-induced TNFα production was inhibited by DHMEQ or U0126 but not by SB203580 or SP600125 (\*P < 0.05, \*\*P < 0.01). Values are expressed as ± SD of four determinations. Phosphorylations of IκB and MAPKs in RAW 264.7 cells were determined by Western blotting by using anti-phospho-IκB, -ERK1/2, -p38, or -JNK antibodies after treatment with rIL-32α (100 ng/mL) (c). rIL-32α stimulated phosphorylation of IκB and ERK1/2, starting at 30 minutes and peaking at 90 (ERK1/2) or 120 (IκB) minutes, whereas significant phosphorylation was not observed in p38 or JNK. Degradation of IκB was also observed with a peak at 90 minutes. These data represent one of three independent experiments. DHMEQ, dehydroxymethyl epoxyquinomicin; DMSO, dimethyl sulfoxide; IκB, inhibitor kappa B; IL, interleukin; LPS, lipopolysaccharide; MAPK, mitogen-activated protein kinase; NF-κB, nuclear factor kappa B; rIL-32α, recombinant human interleukin-32α protein; TNFα, tumor necrosis factor-alpha.

can be speculated to involve the binding of IL-32 to its putative receptor modulates downstream signaling for TLR-4 or other TLRs, since the proinflammatory activities of IL-32 were present even in macrophages derived from TLR-4<sup>-/-</sup> mice, and stimulation with IL-32 plus TLR ligand elicited only additive effects rather than synergistic effects [33]. Two candidate molecules potentially connecting IL-32α and TLR-4 signaling are considered. One is proteinase-3 and the other is proteinase-activated receptor 2 (PAR2); the former reportedly acts as an IL-32-binding protein and cleaves all isoforms of

IL-32 to generate a more active form [34,35], and the latter has been shown to be associated with late NF-κB activation and subsequent TNFα production predominantly through a myeloid differentiation factor 88 (MyD88)-independent pathway [36].

In contrast to mounting evidence on upstream signaling regulators for IL-32, downstream signaling pathways of IL-32 toward TNFα production have not yet been analyzed in detail, and only a small number of reports have focused on different signals in different cell types. The first report of IL-32 advocated that IL-32α





**Figure 6 Exogenous IL-32 $\alpha$  stimulated mRNA expression of IL-6 and MIP-2 independently of NF- $\kappa$ B and MAPK signaling pathways.** RAW 264.7 cells were stimulated with rIL-32 $\alpha$  in combination with inhibitors for I $\kappa$ B and MAPKs, including DHMEQ, U0126, SB203580, and SP600125, for 6 hours. The amounts of mRNA of TNF $\alpha$  (a), IL-6 (b), and MIP-2 (c) were measured by real-time polymerase chain reaction. rIL-32 $\alpha$  alone stimulated RAW 264.7 cells to express IL-6 and MIP-2 as well as TNF $\alpha$  at the mRNA level. None of the inhibitors canceled IL-6 and MIP-2 mRNA expression induced by IL-32 $\alpha$ , whereas DHMEQ and U0126 suppressed TNF $\alpha$  mRNA expression. Values are expressed as mean  $\pm$  standard deviation of triplicate determinations. Ct, control; DHMEQ, dehydroxymethyllepoxyquinomicin; GAPDH, glyceraldehyde-3-phosphate dehydrogenase; I $\kappa$ B, inhibitor kappa B; IL, interleukin; LPS, lipopolysaccharide; MAPK, mitogen-activated protein kinase; MIP-2, macrophage inflammatory protein 2; NF- $\kappa$ B, nuclear factor kappa B; n.s., not significant; rIL-32 $\alpha$ , recombinant human interleukin-32 $\alpha$  protein; TNF $\alpha$ , tumor necrosis factor-alpha.

stimulated TNF $\alpha$  production through activation of NF- $\kappa$ B and p38 MAPK in mouse RAW 267.4 cells; the two peaks of p38 MAPK phosphorylation at 5 and 45 minutes were considered a characteristic finding for this cytokine [2]. Netea and colleagues [21] subsequently reported that IL-32-induced TNF $\alpha$  production by human peripheral blood mononuclear cells (PBMCs) was similarly regulated through phosphorylation of p38 MAPK. On the other hand, ERK1/2 MAPK was dominant in IL-32-induced osteoclastogenesis for human PBMCs [22] and in IL-32-induced IL-6 and IL-8 production by human fibroblast-like synoviocytes [23]. The present study demonstrated that IL-32 $\alpha$ -induced TNF $\alpha$  production was mediated through phosphorylation of

I $\kappa$ B and ERK1/2 in RAW 267.4 cells. Actually, all three components of MAPKs - p38, JNK, and ERK1/2 - were constitutively phosphorylated in RAW 267.4 cells; however, only ERK1/2 phosphorylation was significantly accelerated in response to IL-32 $\alpha$  stimulation. This observation corroborated the fact that the addition of inhibitors for ERK1/2 and NF- $\kappa$ B suppressed each phosphorylation (data not shown and Figure S1 of Additional file 1) and consequently canceled IL-32 $\alpha$ -induced upregulation of TNF $\alpha$  at the mRNA level (Figures 5b and 6). Given the delayed phosphorylation of I $\kappa$ B and ERK1/2 starting at 30 minutes in this study, IL-32 $\alpha$  in RAW 267.4 cells might not directly activate I $\kappa$ B or ERK1/2. Instead, other molecules might play an important role

in I $\kappa$ B or ERK1/2 activation of IL-32 $\alpha$ , or the IL-32 $\alpha$ -TNF $\alpha$  axis might use the MyD88-independent pathway reportedly associated with the late inflammatory response of TLR-4 [37]. Our study also revealed that IL-32 $\alpha$  induced IL-6 and MIP-2 as well as TNF $\alpha$ , but their induction was not canceled by inhibitors for NF- $\kappa$ B or MAPKs. This observation indicates that a signaling pathway other than NF- $\kappa$ B or MAPKs might be involved in IL-6 and MIP-2 expressions.

IL-32 $\gamma$  Tg mice obtained by using a promoter similar to that of the present study reportedly exhibited no apparent phenotype, but once inflammatory colitis was induced with dextran sodium sulfate (DSS) in the Tg mice, severe colitis occurred within 4 days [38]. Interestingly, at more than 6 days after DSS challenge, the degree of colonic inflammation in the Tg mice was significantly reduced, and recovery was more rapid than that in Wt mice because of increased IL-10 levels in serum. In another study, IL-32 $\beta$  was reported to promote the production of IL-10 in human cell lines [39]. According to our data on RAW 264.7 cells, the level of TNF $\alpha$  in culture media peaked at 12 hours after stimulation with IL-32 $\alpha$  and gradually decreased thereafter, whereas IL-10 levels increased from 24 to 96 hours after stimulation (data not shown and Figure S2 of Additional file 2). IL-32 is thus considered to represent a cytokine possessing contradictory properties according to the different phases of the disease. Such paradoxical effects of IL-32 were not observed in our Tg mice. In fact, a single intra-articular injection of LPS in our Tg mice resulted in a transient flare of inflammatory arthritis, characterized by neutrophil infiltration and synovial proliferation, but such inflammation might cease within 2 weeks, followed by amelioration of synovitis with only mild cartilage erosion remaining. On the other hand, the endotoxin shock model using our Tg mice was suitable for examining short-term effects, but not long-term effects, of IL-32 *in vivo* since most mice died within several hours after LPS challenge, and TNF $\alpha$  induced by IL-32 $\alpha$  and LPS was confirmed as an early mediator of endotoxin lethality [40]. The time-dependent and complicated regulation of IL-32 and the relevant molecules of the IL-32-TNF $\alpha$  axis during the course of autoimmune-related arthritis and infectious immunity should be elucidated in future studies.

## Conclusions

This study revealed that IL-32 $\alpha$  contributed to the development of LPS-induced inflammatory arthritis and endotoxin lethality; therefore, stimulation of LPS appears indispensable for activating the IL-32-TNF $\alpha$  axis *in vivo*. However, IL-32 $\alpha$  alone induced TNF $\alpha$  production in RAW 264.7 cells through phosphorylation of I $\kappa$ B and ERK1/2 MAPK.

## Additional material

**Additional file 1: Figure S1.** Phosphorylation of I $\kappa$ B and MAPKs stimulated with rIL-32 $\alpha$  (100 ng/ml) in RAW 264.7 cells was determined by Western blotting using anti-phospho-I $\kappa$ B, -ERK1/2, -p38, and -JNK antibodies. Phosphorylation of I $\kappa$ B and ERK1/2 were observed and inhibited by their specific inhibitors, while significant phosphorylation of p38 or JNK was not observed. This data represents one of three independent experiments.

**Additional file 2: Figure S2.** Levels of TNF $\alpha$  and IL-10 were expressed as a proportion to that in control culture media without IL-32 $\alpha$  stimulation. Level of TNF $\alpha$  peaked at 12 h after stimulation with IL-32 $\alpha$  and gradually decreased thereafter, while levels IL-10 kept increasing from 24 to 96 h after stimulation. Values are expressed as mean  $\pm$  SD of triplicate determinations.

## Abbreviations

BM: bone marrow; DHMEQ: dehydroxymethylleptoxyquinomicin; DMSO: dimethyl sulfoxide; DSS: dextran sodium sulfate; ELISA: enzyme-linked immunosorbent assay; ERK1/2: extracellular signal regulated kinase1/2; FCS: fetal calf serum; I $\kappa$ B: inhibitor kappa B; IL: interleukin; LPS: lipopolysaccharide; MAPK: mitogen-activated protein kinase; MIP: macrophage inflammatory protein; MyD88: myeloid differentiation factor 88; NF- $\kappa$ B: nuclear factor kappa B; NK: natural killer; NOD: nucleotide-binding oligomerization domain-containing protein; PBMC: peripheral blood mononuclear cell; PBS: phosphate-buffered saline; PCR: polymerase chain reaction; RA: rheumatoid arthritis; rIL-32 $\alpha$ : recombinant human interleukin-32 $\alpha$  protein; Tg: transgenic; TLR: Toll-like receptor; TNF: tumor necrosis factor; Wt: wild-type (C57BL/6 Jcl).

## Acknowledgements

This work was partially supported by Grants-in-aid for Scientific Research (C) (#21591959) from the Japan Society for the Promotion of Science.

## Author details

<sup>1</sup>Department of Orthopaedic Surgery, Keio University, 35 Shinanomachi, Shinjuku, Tokyo 160-8582, Japan. <sup>2</sup>Department of Pathology, Keio University, 35 Shinanomachi, Shinjuku, Tokyo 160-8582, Japan. <sup>3</sup>Department of Applied Chemistry, Faculty of Science and Technology, Keio University, Hiyoshi 3-14-1, Kohoku-ku, Yokohama 223-8522, Japan.

## Authors' contributions

MN carried out the animal and molecular experiments, performed the statistical analysis, and drafted the manuscript. YN conceived and designed the study and edited the manuscript. TK taught the experimental procedure and advised MN on this study. YTa and KH were involved in the conception and design of the study. AS and YO performed the histopathological examinations. KU provided the reagents to MN. HI, YTo, and TM helped to supervise the study design and provide valuable advice to MN. All authors read and approved the final manuscript.

## Competing interests

The authors declare that they have no competing interests.

Received: 19 December 2011 Revised: 1 May 2012

Accepted: 21 May 2012 Published: 21 May 2012

## References

1. Dahl CA, Schall RP, He HL, Cairns JS: Identification of a novel gene expressed in activated natural killer cells and T cells. *J Immunol* 1992, **15**:597-603.
2. Kim SH, Han SY, Azam T, Yoon DY, Dinarello CA: Interleukin-32: a cytokine and inducer of TNF-alpha. *Immunity* 2005, **22**:131-142.
3. Choi JD, Bae SY, Hong JW, Azam T, Dinarello CA, Her E, Choi WS, Kim BK, Lee CK, Yoon DY, Kim SJ, Kim SH: Identification of the most active interleukin-32 isoform. *Immunology* 2009, **126**:535-542.
4. Goda C, Kanaji T, Kanaji S, Tanaka G, Arima K, Ohno S, Izuhara K: Involvement of IL-32 in activation-induced cell death in T cells. *Int Immunol* 2006, **18**:233-240.

5. Shioya M, Nishida A, Yagi Y, Ogawa A, Tsujikawa T, Kim-Mitsuyama S, Takayanagi A, Shimizu N, Fujiyama Y, Andoh A: **Epithelial overexpression of interleukin-32alpha in inflammatory bowel disease.** *Clin Exp Immunol* 2007, **149**:480-486.
6. Netea MG, Azam T, Lewis EC, Joosten LA, Wang M, Langenberg D, Meng X, Chan ED, Yoon DY, Ottenhoff T, Kim SH, Dinarello CA: **Mycobacterium tuberculosis induces interleukin-32 production through a caspase-1/IL-18/interferon-gamma-dependent mechanism.** *PLoS Med* 2006, **3**:e277.
7. Bai X, Kim SH, Azam T, McGibney MT, Huang H, Dinarello CA, Chan ED: **IL-32 is a host protective cytokine against Mycobacterium tuberculosis in differentiated THP-1 human macrophages.** *J Immunol* 2010, **184**:3830-3840.
8. Rasool ST, Tang H, Wu J, Li W, Mukhtar MM, Zhang J, Mu Y, Xing HX, Wu J, Zhu Y: **Increased level of IL-32 during human immunodeficiency virus infection suppresses HIV replication.** *Immunol Lett* 2008, **117**:161-167.
9. Li W, Sun W, Liu L, Yang F, Li Y, Chen Y, Fang J, Zhang W, Wu J, Zhu Y: **IL-32: a host proinflammatory factor against influenza viral replication is upregulated by aberrant epigenetic modifications during influenza A virus infection.** *J Immunol* 2010, **185**:5056-5065.
10. Smith AJ, Toledo CM, Wietgreffe SW, Duan L, Schacker TW, Reilly CS, Haase AT: **The immunosuppressive role of IL-32 in lymphatic tissue during HIV-1 infection.** *J Immunol* 2011, **186**:6576-6584.
11. Calabrese F, Baraldo S, Bazzan E, Lunardi F, Rea F, Maestrelli P, Turato G, Lokar-Oliani K, Papi A, Zuin R, Sfriso P, Balestro E, Dinarello CA, Saetta M: **IL-32, a novel proinflammatory cytokine in chronic obstructive pulmonary disease.** *Am J Respir Crit Care Med* 2008, **178**:894-901.
12. Nishida A, Andoh A, Shioya M, Kim-Mitsuyama S, Takayanagi A, Fujiyama Y: **Phosphatidylinositol 3-kinase/Akt signaling mediates interleukin-32alpha induction in human pancreatic periacinar myofibroblasts.** *Am J Physiol Gastrointest Liver Physiol* 2008, **294**:G831-G838.
13. Nishida A, Andoh A, Inatomi O, Fujiyama Y: **Interleukin-32 expression in the pancreas.** *J Biol Chem* 2009, **284**:17868-17876.
14. Shoda H, Fujio K, Yamaguchi Y, Okamoto A, Sawada T, Kochi Y, Yamamoto K: **Interactions between IL-32 and tumor necrosis factor alpha contribute to the exacerbation of immune-inflammatory diseases.** *Arthritis Res Ther* 2006, **8**:R166.
15. Joosten LA, Netea MG, Kim SH, Yoon DY, Oppers-Walgreen B, Radstake TR, Barrera P, van de Loo FA, Dinarello CA, van den Berg WB: **IL-32, a proinflammatory cytokine in rheumatoid arthritis.** *Proc Natl Acad Sci USA* 2006, **103**:3298-3303.
16. Kobayashi H, Huang J, Ye F, Shyr Y, Blackwell TS, Lin PC: **Interleukin-32beta propagates vascular inflammation and exacerbates sepsis in a mouse model.** *PLoS One* 2010, **5**:e9458.
17. Mun SH, Kim JW, Nah SS, Ko NY, Lee JH, Kim JD, Kim do K, Kim HS, Choi JD, Kim SH, Lee CK, Park SH, Kim BK, Kim HS, Kim YM, Choi WS: **Tumor necrosis factor alpha-induced interleukin-32 is positively regulated via the Syk/protein kinase Cdelta/JNK pathway in rheumatoid synovial fibroblasts.** *Arthritis Rheum* 2009, **60**:678-685.
18. Ko NY, Mun SH, Lee SH, Kim JW, Kim do K, Kim HS, Her E, Kim SH, Won HS, Shin HS, Kim HS, Kim YM, Choi WS: **Interleukin-32alpha production is regulated by MyD88-dependent and independent pathways in IL-1beta-stimulated human alveolar epithelial cells.** *Immunobiology* 2011, **216**:32-40.
19. Alsaleh G, Sparsa L, Chatelus E, Ehlinger M, Gottenberg JE, Wachsmann D, Sibilia J: **Innate immunity triggers IL-32 expression by fibroblast-like synoviocytes in rheumatoid arthritis.** *Arthritis Res Ther* 2010, **12**:R135.
20. Pan X, Cao H, Lu J, Shu X, Xiong X, Hong X, Xu Q, Zhu H, Li G, Shen G: **Interleukin-32 expression induced by hepatitis B virus protein X is mediated through activation of NF-kB.** *Mol Immunol* 2011, **48**:1573-1577.
21. Netea MG, Lewis EC, Azam T, Joosten LA, Jaekal J, Bae SY, Dinarello CA, Kim SH: **Interleukin-32 induces the differentiation of monocytes into macrophage-like cells.** *Proc Natl Acad Sci USA* 2008, **105**:3515-3520.
22. Mabileau G, Sabokbar A: **Interleukin-32 promotes osteoclast differentiation but not osteoclast activation.** *PLoS One* 2009, **4**:e4173.
23. Kim YG, Lee CK, Kim SH, Cho WS, Mun SH, Yoo B: **Interleukin-32gamma enhances the production of IL-6 and IL-8 in fibroblast-like synoviocytes via Erk1/2 activation.** *J Clin Immunol* 2010, **30**:260-267.
24. Ariga A, Namekawa J, Matsumoto N, Inoue J, Umezawa K: **Inhibition of tumor necrosis factor-alpha -induced nuclear translocation and activation of NF-kappa B by dehydroxymethylepoxyquinomicin.** *J Biol Chem* 2002, **277**:24625-24630.
25. Umezawa K, Chaicharoenpong C: **Molecular design and biological activities of NF-kappaB inhibitors.** *Mol Cells* 2002, **14**:163-167.
26. Niki Y, Yamada H, Seki S, Kikuchi T, Takaishi H, Toyama Y, Fujikawa K, Tada N: **Macrophage- and neutrophil-dominant arthritis in human IL-1 alpha transgenic mice.** *J Clin Invest* 2001, **107**:1127-1135.
27. Abu-Ghefreh AA, Masocha W: **Enhancement of antinociception by coadministration of minocycline and a non-steroidal anti-inflammatory drug indomethacin in naive mice and murine models of LPS-induced thermal hyperalgesia and monoarthritis.** *BMC Musculoskelet Disord* 2010, **11**:276.
28. Chen WT, Mahmood U, Weissleder R, Tung CH: **Arthritis imaging using a near-infrared fluorescence folate-targeted probe.** *Arthritis Res Ther* 2005, **7**:R310.
29. Frasnelli ME, Tarussio D, Chobaz-Péclat V, Busso N, So A: **TLR2 modulates inflammation in zymosan-induced arthritis in mice.** *Arthritis Res Ther* 2005, **7**:R370.
30. Horiuchi K, Kimura T, Miyamoto T, Takaishi H, Okada Y, Toyama Y, Blobel CP: **Cutting edge: TNF-alpha-converting enzyme (TACE/ADAM17) inactivation in mouse myeloid cells prevents lethality from endotoxin shock.** *J Immunol* 2007, **179**:2686-2689.
31. Heinhuis B, Koenders MI, van Riel PL, van de Loo FA, Dinarello CA, Netea MG, van den Berg WB, Joosten LA: **Tumour necrosis factor alpha-driven IL-32 expression in rheumatoid arthritis synovial tissue amplifies an inflammatory cascade.** *Ann Rheum Dis* 2011, **70**:660-667.
32. Heinhuis B, Koenders MI, van de Loo FA, van Lent PL, Kim SH, Dinarello CA, Joosten LA, van den Berg WB: **IL-32gamma and Streptococcus pyogenes cell wall fragments synergise for IL-1-dependent destructive arthritis via upregulation of TLR-2 and NOD2.** *Ann Rheum Dis* 2010, **69**:1866-1872.
33. Netea MG, Azam T, Ferwerda G, Girardin SE, Walsh M, Park JS, Abraham E, Kim JM, Yoon DY, Dinarello CA, Kim SH: **IL-32 synergizes with nucleotide oligomerization domain (NOD) 1 and NOD2 ligands for IL-1beta and IL-6 production through a caspase 1-dependent mechanism.** *Proc Natl Acad Sci USA* 2005, **102**:16309-16314.
34. Novick D, Rubinstein M, Azam T, Rabinkov A, Dinarello CA, Kim SH: **Proteinase 3 is an IL-32 binding protein.** *Proc Natl Acad Sci USA* 2006, **103**:3316-3321.
35. Kim S, Lee S, Her E, Bae S, Choi J, Hong J, Jaekal J, Yoon D, Azam T, Dinarello CA, Kim S: **Proteinase 3-processed form of the recombinant IL-32 separate domain.** *BMB Rep* 2008, **41**:814-819.
36. Rallabhandi P, Nhu QM, Toshchakov VY, Piao W, Medvedev AE, Hollenberg MD, Fasano A, Vogel SN: **Analysis of proteinase-activated receptor 2 and TLR4 signal transduction: a novel paradigm for receptor cooperativity.** *J Biol Chem* 2008, **283**:24314-24325.
37. Kawai T, Akira S: **Toll-like receptors and their crosstalk with other innate receptors in infection and immunity.** *Immunity* 2011, **34**:637-650.
38. Choi J, Bae S, Hong J, Ryoo S, Jhun H, Hong K, Yoon D, Lee S, Her E, Choi W, Kim J, Azam T, Dinarello CA, Kim SH: **Paradoxical effects of constitutive human IL-32gamma in transgenic mice during experimental colitis.** *Proc Natl Acad Sci USA* 2010, **107**:21082-21086.
39. Kang JW, Choi SC, Cho MC, Kim HJ, Kim JH, Lim JS, Kim SH, Han JY, Yoon DY: **A proinflammatory cytokine interleukin-32beta promotes the production of an anti-inflammatory cytokine interleukin-10.** *Immunology* 2009, **128**:e532-e540.
40. Wang H, Bloom O, Zhang M, Vishnubhakat JM, Ombrellino M, Che J, Frazier A, Yang H, Ivanova S, Borovikova L, Manogue KR, Faist E, Abraham E, Andersson J, Andersson U, Molina PE, Abumrad NN, Sama A, Tracey KJ: **HMG-1 as a late mediator of endotoxin lethality in mice.** *Science* 1999, **285**:248-251.

doi:10.1186/ar3850

**Cite this article as:** Nakayama *et al.*: Enhanced susceptibility to lipopolysaccharide-induced arthritis and endotoxin shock in interleukin-32 alpha transgenic mice through induction of tumor necrosis factor alpha. *Arthritis Research & Therapy* 2012 **14**:R120.