

Research article

Open Access

Solubility enhancement of aggregation-prone heterologous proteins by fusion expression using stress-responsive *Escherichia coli* protein, RpoS

Jin-Seung Park^{†1}, Kyung-Yeon Han^{†1}, Jong-Ho Lee¹, Jong-Am Song¹, Keum-Young Ahn¹, Hyuk-Seong Seo¹, Sang-Jun Jun Sim², Seung-Wook Kim¹ and Jeewon Lee^{*1}

Address: ¹Department of Chemical and Biological Engineering, Korea University, Anam-Dong 5-1, Sungbuk-Ku, Seoul 136-713, South Korea and ²Department of Chemical Engineering, Sungkyunkwan University, Suwon, South Korea

Email: Jin-Seung Park - anorain@korea.ac.kr; Kyung-Yeon Han - ruddus78@korea.ac.kr; Jong-Ho Lee - jjong0614@korea.ac.kr; Jong-Am Song - uga33@korea.ac.kr; Keum-Young Ahn - anggum@korea.ac.kr; Hyuk-Seong Seo - shs5951@korea.ac.kr; Sang-Jun Jun Sim - simsj@skku.edu; Seung-Wook Kim - kimsuw@korea.ac.kr; Jeewon Lee* - leeju@korea.ac.kr

* Corresponding author †Equal contributors

Published: 19 February 2008

Received: 29 August 2007

BMC Biotechnology 2008, 8:15 doi:10.1186/1472-6750-8-15

Accepted: 19 February 2008

This article is available from: <http://www.biomedcentral.com/1472-6750/8/15>

© 2008 Park et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: The most efficient method for enhancing solubility of recombinant proteins appears to use the fusion expression partners. Although commercial fusion partners including maltose binding protein and glutathione-S-transferase have shown good performance in enhancing the solubility, they cannot be used for the proprietary production of commercially value-added proteins and likely cannot serve as universal helpers to solve all protein solubility and folding issues. Thus, novel fusion partners will continue to be developed through systematic investigations including proteome mining presented in this study.

Results: We analyzed the *Escherichia coli* proteome response to the exogenous stress of guanidine hydrochloride using 2-dimensional gel electrophoresis and found that RpoS (RNA polymerase sigma factor) was significantly stress responsive. While under the stress condition the total number of soluble proteins decreased by about 7 %, but a 6-fold increase in the level of RpoS was observed, indicating that RpoS is a stress-induced protein. As an N-terminus fusion expression partner, RpoS increased significantly the solubility of many aggregation-prone heterologous proteins in *E. coli* cytoplasm, indicating that RpoS is a very effective solubility enhancer for the synthesis of many recombinant proteins. RpoS was also well suited for the production of a biologically active fusion mutant of *Pseudomonas putida* cutinase.

Conclusion: RpoS is highly effective as a strong solubility enhancer for aggregation-prone heterologous proteins when it is used as a fusion expression partner in an *E. coli* expression system. The results of these findings may, therefore, be useful in the production of other biologically active industrial enzymes, as successfully demonstrated by cutinase.

Background

Escherichia coli have been widely used as a host to produce valuable commercial, industrial, and therapeutic proteins. There are several disadvantages with this system, however, especially for the expression of eukaryotic proteins. For example, when randomly selected 2,078 full-length genes of *Caenorhabditis elegans* were expressed in *E. coli* cytoplasm, only 11% of genes yielded significant amounts of soluble material [1]. Several different approaches have been taken to resolve the solubility problem in the past and include (1) truncation of long multi-domain proteins into short and separate domains [2]; (2) co-expression of molecular chaperones or foldases [3]; (3) enabled secretion to the periplasm where disulfide bonds can be properly formed with the help of an oxidative environment and Dsb protein families [4]; (4) co-expression of aminoacyl tRNA cognates to amino acids encoded by rare codons [5]; and more recently (5) use of a fusion expression partner [6,7]. The most efficient method for enhancing solubility and folding efficiencies of recombinant proteins appears to be the latter (fusion expression partners) which includes maltose binding protein (MBP) [8], thioredoxin (Trx) [9], human ferritin heavy chains (hFTN-H) [10], and glutathione-S-transferase (GST) [11]. Although these fusion partners have shown good performance in enhancing the solubility and folding of some recombinant proteins [12], they likely cannot serve as universal helpers to solve all protein solubility and folding issues. Thus, novel fusion partners will continue to be developed through systematic investigations including proteome mining.

In the present study, we found that the level of RpoS significantly increased during the stress caused by guanidine hydrochloride through proteome-wide mining involving the stress response of *E. coli* BL21(DE3). As an N-terminus fusion expression partner, RpoS dramatically increased the solubility of the following heterologous proteins (human minipro-insulin (mp-INS), human epidermal growth factor (EGF), human prepro-ghrelin (ppGRN), human interleukin-2 (hIL-2), human activation induced cytidine deaminase (AID), human glutamate decarboxylase (GAD448-585), *Pseudomonas putida* cutinase (CUT), human ferritin light chain (hFTN-L), human granulocyte colony-stimulating factor (G-CSF), and cold autoinflammatory syndrome 1 (NALP3) NACHT domain (NACHT)). We also have demonstrated an increased yield of a biologically active fusion mutant of heterologous bacterial cutinase that is expected to be of significant biotechnical and commercial interest.

Results and discussion

Guanidine hydrochloride-induced proteome response of *E. coli* and finding of the aggregation-resistant protein, RpoS

Exogenous stresses including heat shock and GdnHCl [13-15] often results in the misfolding and aggregation of proteins within the cell. Therefore, it would seem reasonable to presume that intracellular proteins which exist in their native soluble forms even under stress conditions have an intrinsic ability to remain efficiently folded compared to proteins that unfold or aggregate under the same stresses. GdnHCl stress facilitates the release of periplasmic proteins through increased membrane permeability [16-18] and hence could be regarded as an osmolyte. We investigated changes of the proteome profile of *E. coli* when the stress reagent, guanidine hydrochloride, was added to growing bacterial cultures, by applying 2-dimensional gel electrophoresis (2-DE). Compared to non-stress condition, the total number of intracellular soluble proteins was reduced to 731 from 788, which indicates that many proteins aggregated under the stress condition induced by GdnHCl. What was surprising is that the RpoS expression level (i.e. protein spot intensity estimated through analysis of 2-DE gel images, Fig. 1. and Table 1) increased 6-fold compared to the non-stress situation. In addition to RpoS, we found several other proteins (Table 1), the expression level of which more than 1.9-fold increased in response to the stressor GdnHCl, including HSP60 chaperonin (57 kD), chaperone HtpG (71 kD), transcription elongation protein NusA (55 kD), DNA gyrase subunit A (97 kD), formate acetyltransferase 1 (85 kD), succinate dehydrogenase flavoprotein subunit (64 kD), *etc.* Since the synthesis yield of target protein can be significantly reduced, the molecular mass of proteins to be used as fusion expression partner should not be too high. That is, although the large amount of fusion protein is synthesized, the actual amount of the target fusion-free protein could be very low if the fusion partner is too big. RpoS (38 kD) is a relatively small protein among the stress-responsive proteins we found and was highly effective in enhancing the solubility of target proteins.

RpoS, stress responsive RNA polymerase sigma factor is a well-known universal stress regulator controlling many proteins under various stress conditions, e.g. the onset of the stationary phase [19], and carbon starvation [20]. RpoS was previously reported to be induced by osmotic stress [16] and heat shock [21]. We also observed that the RpoS expression level increased 3- to 5-fold in response to heat shock (data not shown). Muffler *et al.* [21] reported that the duration of stability of RpoS in response to heat shock stress is maintained by the direct- or indirect-binding of DnaK. Bound DnaK appears to assist the effective folding of RpoS and also protect RpoS against the action of ClpP, a protease that degrades RpoS [21]. From this point of view, RpoS may serve as a solubility enhancer in

E. coli cytoplasm, when used as a fusion expression partner upon the expression of aggregation-prone heterologous proteins.

Expression of aggregation-prone heterologous proteins using RpoS as fusion partner

We used RpoS as an N-terminus fusion expression partner for the synthesis of numerous heterologous proteins [mpINS, EGF, ppGRN, hIL-2, AID, GAD₄₄₈₋₅₈₅, CUT, hFTN-L, G-CSF, and NACHT (Fig. 2 and Table 2 for expression system construction)]. Stenström *et al.* [22] reported that the codon following an AUG start triplet (+2 codon) significantly affects gene expression in *E. coli*, and the second codon starting with A is most advantageous to achieve an enhanced expression level. The second codon of RpoS is AGT (encoding Ser) that seems to be favorable second

codon based on the report of Stenström *et al.* [22]. As shown in Figure 3B and Table 3, all heterologous proteins expressed directly without RpoS fusion formed insoluble inclusion bodies resulting in nearly negligible solubility. Compared to these results, it seems surprising that when the same heterologous proteins were expressed with the fusion of RpoS, the solubility of these foreign proteins dramatically increased (Fig. 3A and Table 3), thereby indicating that *E. coli* RpoS is a highly effective solubility enhancer for aggregation-prone heterologous proteins. Table 3 compares the effect of RpoS- and GST fusion on the solubility enhancement for heterologous proteins. In the fusion expression of CUT, GAD₄₄₈₋₅₈₅, and mpINS, the effect of RpoS fusion was much higher, whereas GST fusion was significantly more effective in the expression of AID, NACHT, and hFTN-L. This result indicates that there




Figure 1
***E. coli* proteome profiles under non-stress and GdnHCl-stress condition.** (A) 2-DE gel image of *E. coli* proteome under the non-stress condition. (Arrow indicates RpoS spot under non-stress condition.) (B) 2-DE gel image of *E. coli* proteome under GdnHCl-stress condition (Arrow indicates RpoS spot under GdnHCl-stress condition). (Figure in a box presents relative spot intensities of RpoS, analyzed under non-stress and GdnHCl-stress conditions.)

Table 1: Result of proteome-wide finding of some aggregation-resistant *E. coli* proteins

Gene name ^a	Access No. ^a	Protein name ^a	pI/MW(kDa)		Sequence coverage ^d	Score ^d	Fold change
			Theoretical ^b	Experimental ^c			
rpoS	PI3445	RNA polymerase sigma factor rpoS	4.89/37.97	5.08/38.95	19	36	6.21
groL	P0A6F5	HSP60 chaperonin	4.85/57.20	4.90/58.63	16	34	2.13
htpG	P0A6Z3	Chaperone protein htpG	5.09/71.38	5.14/66.37	9	27	3.10
nusA	P0AFF6	Transcription elongationprotein nusA	4.53/54.87	4.69/59.09	16	44	3.46
gyrA	P0AES4	DNA gyrase subunit A	5.09/96.96	5.20/92.23	21	29	1.95
pflB	P09373	Formate acetyltransferase I	5.69/85.23	5.68/83.51	19	39	3.26
sdhA	P0AC41	Succinate dehydrogenase flavoprotein subunit	5.85/64.42	5.70/65.73	25	43	2.16

^a Gene name, accession number, and protein name were obtained from ExPASy Proteomics Server [35].

^b Theoretical values of pI and Mw were calculated using "Compute pI/Mw tool" [36].

^c Experimental values of pI and Mw were estimated through 2-DE gel image analysis in the present study.

^d Sequence coverage and score values were calculated using "ALDENTE : PEPTIDE MASS FINGERPRINTING TOOL" [37].

are no universal helpers to solve all protein solubility issues. Table 3 also shows that the results of direct- and fusion expression of heterologous proteins are highly reproducible.

Moreover, the plasmid vector for the expression of poly-histidine-tagged fusion mutant of G-CSF [(His)6::RpoS::(D4K)::G-CSF] was constructed (Fig. 2) to

purify fusion-free G-CSF. After (His)6::RpoS::(D4K)::G-CSF was bound onto the ProBond resin (Ni+2) column, the enterokinase proteolysis was carried out in a batch mode, and subsequently the digested product was collected and centrifuged. SDS-PAGE and Western blot analyses of the supernatant show that the recombinant G-CSF was easily released from the Rpos-fusion protein and was

Table 2: Primers used for the cloning of genes encoding various heterologous proteins

Heterologous proteins	Primer sequences			
		Direct expression		Fusion expression
mp-INS	Sense	cat atg ttt gtc aac caa cat		ctc gag ttt gtc aac caa cat
	Antisense	aag ctt tta gtt aca gta gtt c		aag ctt tta gtt aca gta gtt c
EGF	Sense	cat atg aac tct gac tcc gaa tgc		ctc gag aac tct gac tcc gaa tgc
	Antisense	aag ctt tta acg cag ttc cca cca		aag ctt tta acg cag ttc cca cca
ppGRN	Sense	cat atg ggc tcc agc ttc ctg		ctc gag ggc tcc agc ttc ctg
	Antisense	aag ctt tca ctt gtc ggc t		aag ctt tca ctt gtc ggc t
hIL-2	Sense	cat atg gca cct act tca agt		ctc gag gca cct act tca agt
	Antisense	aag ctt tta tca agt cag tgt		aag ctt tta tca agt cag tgt
AID	Sense	cat atg gac agc ctc ttg atg aac		ctc gag gac agc ctc ttg atg aac
	Antisense	aag ctt tca taa caa aag tcc ca		aag ctt tca taa caa aag tcc ca
GAD ₄₄₈₋₅₈₅	Sense	cat atg cgc cac gtt gat gt		ctc gag cgc cac gtt gat gt
	Antisense	atc gat tta taa atc ttg tcc		atc gat tta taa atc ttg tcc
CUT	Sense	cat atg gct ccc ctg ccg gat ac		ctc gag gct ccc ctg ccg gat ac
	Antisense	aag ctt tta aag ccc gcg gcg ct		aag ctt tta aag ccc gcg gcg ct
				aag ctt atg atg gtg gtg atg atg tta aag ccc gcg gcg ct (for metal affinity purification)
hFTN-L	Sense	cat atg agc tcc cag att cgt		ctc gag agc tcc cag att cgt
	Antisense	aag ctt tta gtc gtg ctt gag agt		aag ctt tta gtc gtg ctt gag agt
G-CSF	Sense	cat atg act cca ctc gga cct g		ctc gag acc ccc ctg ggc cct gcc
				ctc gag gac gat gac gat aaa acc ccc ctg ggc cct gcc (for enterokinase digestion)
	Antisense	aag ctt tca tgg ctg tgc aag		aag ctt tca tgg ctg tgc aag
NACHT	Sense	cat atg act gtg gtg ttc cag		ctc gag act gtg gtg ttc cag
	Antisense	aag ctt tca cag cag gta gta c		aag ctt tca cag cag gta gta c




Figure 2

The plasmid vector constructions for direct and fusion expression of heterologous proteins in *E. coli*. (A) Direct expression vector, (B) RpoS- and GST-fusion expression vector, (C) Hybrid vector for metal (Ni²⁺) affinity purification of RpoS-CUT-(His)₆, (D) Hybrid vector for metal (Ni²⁺) affinity purification of (His)₆-RpoS-D₄K-G-CSF, followed by enterokinase cleavage.

present in the form of soluble protein in the supernatant (Fig. 4).

Bioactivity of the recombinant fusion mutant of cutinase, RpoS::CUT

Cutinase has been used as a lipolytic enzyme in the composition of laundry and dishwashing detergents to more efficiently remove immobilized fats [23,24]. In addition, the oleochemistry industries [25], and pollutant degradation [26,27] represent other potential uses of cutinase. In recent years, the esterification and transesterification properties of cutinase have been intensively exploited and could be applied usefully in other chemical synthesis processes [28]. Because of these extensive potential applications, we were particularly interested in the production of a bioactive recombinant cutinase. Thus, we cloned the

cutinase gene from the genome of *Pseudomonas putida* and expressed it in *E. coli* using the fusion of RpoS. Cutinase is known for its hydrolytic activity for a variety of esters ranging from soluble *p*-nitrophenyl esters to insoluble long-chain triglycerides. The hydrolytic activity of cutinase, especially on *p*-nitrophenyl esters of fatty acids, is extremely sensitive to fatty acid chain length. Previously, Lin *et al.* [29] reported that microbial cutinase lacked a large hydrophobic surface around the active site, in contrast to other lipases and esterases. This structural characteristic of cutinase may be strongly related to high substrate-specificity, i.e. the extremely low activity on *p*-nitrophenyl ester of a long chain fatty acid such as *p*-nitrophenyl palmitate (PNP).

Table 3: Solubility of the expressed recombinant proteins

Heterologous proteins	Solubility* of the expressed recombinant proteins (%)		
	RpoS-fusion expression	GST-fusion expression	Direct expression
G-CSF	87.4 ± 2.1	91.5 ± 1.5	4.6 ± 0.5
CUT	44.9 ± 2.7	26.2 ± 1.1	3.1 ± 0.2
AID	18.1 ± 3.5	41.7 ± 1.5	8.7 ± 1.6
hFTN-L	47.8 ± 2.1	87.6 ± 2.8	1.7 ± 0.4
NACHT	36.4 ± 4.3	74.8 ± 2.2	8.4 ± 1.3
GAD ₄₄₈₋₅₈₅	59.0 ± 1.8	7.8 ± 2.7	1.9 ± 0.5
hIL-2	60.1 ± 2.4	85.7 ± 1.7	1.3 ± 0.3
EGF	80.4 ± 0.9	92.9 ± 1.2	3.0 ± 0.9
ppGRN	89.2 ± 3.5	90.1 ± 2.3	7.6 ± 0.6
mp-INS	75.9 ± 2.8	13.3 ± 1.0	1.3 ± 0.4

* The solubility was defined as the fraction of the soluble recombinant protein compared to the synthesized total (soluble + insoluble) recombinant protein. Average and standard deviation values were calculated based on the results of repeated triplicate experiments.




Figure 3
Results of direct and fusion expression of heterologous proteins. SDS-PAGE analyses of the RpoS-fusion expressed proteins (A), directly (non-fusion) expressed proteins (B), and GST-fusion expressed proteins (C).




Figure 4
Results of SDS-PAGE and Western blot analysis of RpoS-fusion and fusion-free G-CSF. -SDS-PAGE analysis (lane M-3): lane M, molecular markers; lane 1, supernatant of recombinant *E. coli* cell lysates containing recombinant (His)₆-RpoS-D₄K-G-CSF, which was loaded onto ProBond resin (Ni²⁺) column for metal affinity purification; lane 2, soluble fraction of enterokinase(EK)-digested product of (His)₆-RpoS-D₄K-G-CSF, containing RpoS and fusion-free G-CSF (indicated by an arrow); lane 3, purified soluble fusion-free G-SCF. -Western blot analysis (lane 4): result of immunoblotting analysis of purified soluble fusion-free G-SCF (loaded onto lane 3).

We assayed the enzymatic activity of our fusion mutant of cutinase (RpoS::CUT) and demonstrated the same selective bioactivity as native cutinase to degrade *p*-nitrophenyl butyrate (PNB) but not to degrade PNP (Fig. 5A and 5B). We also purified RpoS::CUT-His₆ (Fig. 6A) through Ni²⁺ affinity chromatography and analyzed the purified RpoS::CUT-His₆ using reversed phase HPLC (Fig. 6B). As shown in Figure 6B, RpoS::CUT-His₆ was analyzed as a single peak, which seems to indicate that the crafted mutant molecules of cutinase have uniform and correctly folded conformation due probably to the help of fusion partner, *E. coli* RpoS.

Conclusion

Using 2-dimensional gel electrophoresis, we found that *E. coli* RpoS, RNA polymerase sigma factor was GdnHCl stress-responsive and highly effective as a strong solubility enhancer when used as fusion partner for the expression of aggregation-prone heterologous proteins in *E. coli* BL21(DE3). The results of these findings may, therefore, be useful in the production of other biologically active industrial enzymes, as successfully demonstrated by cutinase.

Methods

Bacterial strain and plasmids

E. coli strain BL21(DE3) (F-ompT hsdS_B(rB⁻ mB⁻)) was selected under both non-stress and GdnHCl-stress conditions for 2-dimensional gel electrophoresis analysis. Through PCR amplification using appropriate primers (Table 2), the genes encoding mp-INS, EGF, ppGRN, hIL-2 [30], AID, GAD₄₄₈₋₅₈₅ [31], CUT, hFTN-L [32], G-CSF, and NACHT were cloned using previously cloned heterologous genes, except for CUT gene that was cloned from chromosomal DNA of *Pseudomonas putida* (ATCC 53552). The gene clones of mp-INS, ppGRN, AID, G-CSF, and NACHT were kindly donated by other researchers, as acknowledged (see Acknowledgements). The EGF gene was cloned using pCMV6-XL vector (OriGene Technologies, USA). Each of the recombinant genes above and various fusion/hybrid genes [rpoS(or GST gene)::each heterologous gene] were inserted into the *Nde*I-*Hind*III site of the same plasmid pT7-7 to construct the expression vectors. All the heterologous genes above were fused directly to the RpoS gene (cloned from the chromosomal DNA of *E. coli* BL21(DE3)) or GST gene [cloned from the GST-fusion vector pET42a(+)] (Novagen, USA) without any linker sequence. Therefore, each expression vector has no enzymatic cleavage site between rpoS (or GST gene) and heterologous gene, except for the case of G-CSF. For the purification of fusion-free recombinant G-CSF, the D₄K sequence for enterokinase digestion was inserted between RpoS and G-CSF genes to synthesize the polyhistidine-tagged fusion mutant of G-CSF, i.e. (His)₆::RpoS::(D₄K)::G-CSF. For the purification of RpoS::CUT, hexahistidine (His₆) was added to the C-terminus of CUT by PCR (Fig. 2). For this, the sequence coding for N-RpoS::CUT-His₆-C was inserted into the *Nde*I-*Hind*III site of plasmid pT7-7.

After complete DNA sequencing of all gel-purified hybrid plasmids, the *E. coli* strain BL21(DE3) (F-ompT hsdS_B(rB⁻ mB⁻)) was transformed with the hybrid plasmids, and ampicillin-resistant transformants were subsequently selected using LB-agar plates supplemented with ampicillin (100 mg/l).

Recombinant *E. coli* culture, gene expression, and recombinant protein purification

For shake flask experiments, 250 ml Erlenmeyer flasks containing 50 ml LB media and ampicillin at 100 mg/l of culture (37°C and 150 rpm) was used. When the culture turbidity (OD₆₀₀) reached 0.5, gene expression was induced with the addition of IPTG (1 mM), and after a further 4 h of cultivation the all recombinant cells (80 mg wet cell mass) were harvested by centrifugation (13,000 rpm (MICRO17TR, Hanil Science Industrial, Korea) × 5 min) and cell pellets resuspended in 5 ml lysis buffer (10 mM Tris-HCl, pH 7.5, 10 mM EDTA). Cell disruption was




Figure 5
Bioactivity of recombinant fusion mutant, RpoS::CUT. Assay results using cell-free supernatants from (A) *E. coli* BL21 (DE3) host and (B) recombinant *E. coli* BL21 (DE3) [pT7-RpoS-CUT] producing RpoS::CUT. Both PNB (●) and PNP (○) were used as substrates for the cutinase activity assay. (Concentrations: PNB = 6.6 mM; PNP = 6.6 mM).

achieved using a Branson Sonifier (Branson Ultrasonics Corp., Danbury, CT). The cell-free supernatant and insoluble protein aggregates were separated at 13,000 rpm (MICRO17TR, Hanil Science Industrial, Korea) for 10 min. The isolated inclusion bodies, if any, were washed twice with 1 % Triton X-100. Cell-free supernatants and the washed inclusion bodies were subjected to polyacrylamide (14%) gel electrophoresis (PAGE) analysis. Coomassie-stained protein bands were ultimately scanned, and the intensity of each recombinant protein band was estimated using densitometry (Duoscan T1200, Bio-Rad, Hercules, CA). The solubility of recombinant proteins was determined by analyzing the fraction of the soluble recombinant protein compared to the synthesized total (soluble + insoluble) recombinant protein. Average

and standard deviation values of the solubility were calculated based on the results of repeated triplicate experiments.

The purification of recombinant G-SCF was accomplished using metal affinity chromatography. That is, polyhistidine-tagged fusion mutant of G-CSF [(His)₆::RpoS::(D₄K)::G-CSF] (Fig. 4) were loaded onto ProBond resin (Ni²⁺) column. Prior to sample loading, the resin was washed twice with 10 column volumes of binding buffer (50 mM potassium phosphate, 300 mM KCl, 20 mM imidazole, pH 7.0). Binding buffer contains 20 mM imidazole to minimize non-specific binding of untagged protein contaminants, and binding was carried out in a batch mode at 4°C. Afterwards the resin was




Figure 6

Results of purified RpoS::CUT-6xHis analysis. (A) Results of SDS-PAGE analysis of RpoS::CUT-6xHis in soluble fraction of *E. coli* cell lysates (lane 1) and of purified RpoS::CUT-6xHis (lanes 2, 3). (M: molecular marker) (Arrow indicates recombinant RpoS::CUT-6xHis). (B) Result of reversed-phase HPLC analysis of the affinity-purified RpoS::CUT-6xHis. (Peaks in red rectangle and dotted circle correspond to the purified RpoS::CUT-6xHis and PBS buffer, respectively.)

washed twice with 5–8 ml Tris-HCl (10 mM Tris, pH 8.0) prior to enterokinase digestion step. The enterokinase digestion was carried out in a batch mode at 4°C for 10 h using 5-unit enterokinase (Invitrogen, CA, USA). Then, the proteolytic product was collected and centrifuged [3,000 rpm (MICRO17TR, Hanil Science Industrial, Korea) for 10 min], and the supernatant fraction was subjected to being analyzed by SDS-PAGE and western blotting.

Purification and HPLC analysis of cutinase fusion mutant

For the purification of RpoS::CUT-His₆, the soluble fraction was separated after cell disruption by centrifugation (13,000 rpm (MICRO17TR, Hanil Science Industrial, Korea)) for 30 min. The cell-free supernatant containing RpoS::CUT-His₆ was loaded onto the ProBond resin (Ni²⁺) column (Invitrogen) for affinity purification. Before the sample loading, the resin was washed twice with ten column volumes of binding buffer (pH 8.0, 50 mM sodium phosphate, 300 mM NaCl, 10 mM imidazole). Binding was carried out in a batch mode at 4°C. Afterwards, the resin was washed twice with 8 ml washing buffer (pH 8.0, 50 mM sodium phosphate, 300 mM NaCl,

50 mM imidazole) and eluted with elution buffer (pH 8.0, 50 mM sodium phosphate, 300 mM NaCl, 250 mM imidazole). The elution buffer in the eluted solution containing the purified RpoS::CUT-His₆ was changed with PBS buffer (137 mM NaCl, 2.7 mM KCl, 10 mM Na₂HPO₄, 2 mM KH₂PO₄, pH 7.4) using Amicon Ultra-4 centrifugal filter (Millipore, Ireland). Analysis of the purified RpoS::CUT-His₆ was performed with high-performance liquid chromatography (HPLC) (LC-20A Prominence, Shimadzu Co. Ltd., Japan). A Shim-pack CLC-NH2 column (60 × 150 mm, Shimadzu Co. Ltd., Japan) was equilibrated with 10 % acetonitrile containing 0.1 % trifluoroacetic acid at a flow rate of 1 ml/min. The sample was loaded onto the column and eluted with 75 % acetonitrile. The elution profile was monitored at 215 nm.

Sample preparation for proteome analysis and 2-dimensional gel electrophoresis

Flask culture conditions were identical to those for recombinant gene expressions. Cells were grown at 37°C and then 100 mM GdnHCl was added for the GdnHCl-induced proteome response when culture turbidity (OD₆₀₀) reached 0.5 in the LB media. After further 3 h cul-

tivation, the cells were harvested by centrifugation at 6,000 rpm (MEGA17R, Hanil Science Industrial, Korea) for 15 min (4°C) and then washed twice with 40 mM Tris buffer (pH 8.0). Cell pellets were resuspended in 500 µl of lysis buffer (8 M urea, 4% (w/v) CHAPS, 40 mM Tris, protease inhibitor cocktail; Roche Diagnostics GmbH, Mannheim, Germany) and disrupted by sonication. After sonication (Branson Sonifier 450, USA), the cell debris and the aggregated proteins were removed by centrifugation at 12,000 rpm (MICRO17TR, Hanil Science Industrial, Korea) for 60 min (4°C), and only soluble proteins were obtained. 2-dimensional gel electrophoresis was performed as described previously Kim *et al.* [33]. Stained gels were scanned using a UMAX powerlook 1100 scanner and Image Master software v 4.01 (Amersham Biosciences, Uppsala, Sweden) was used for gel image analysis, including quantification of spot intensities performed on volume bases (*i.e.* values calculated from the integration of spot optical intensity over spot area).

MALDI-TOF-MS analysis and protein identification

Samples for MALDI-TOF MS analysis were prepared through the extraction from silver-stained protein spots according to the previous protocol [33]. Enzymatic digestion was performed with 10 mg/ml of sequencing grade modified trypsin (Promega, WI, USA) in 25 nM ammonium bicarbonate (pH 8.0) for overnight at 37°C in a stationary incubator. The trypsin-digested protein spots were analyzed using a MALDI-TOF-MS system (Voyager DE-STR, PE Biosystem, Framingham, MA, U.S.A.) by the Korea Basic Science Institute (Seoul, Republic of Korea), and peptide mass fingerprinting for protein identification was performed using MS-Fit [34]. Spectra were calibrated using a matrix and tryptic autodigestion ion peaks as internal standards. Peptide mass fingerprints were analyzed using the MS-Fit [34]. The identification of a protein with respect to theoretical parameters (pI, molecular mass, etc.) was accepted if the peptide mass matched within a mass tolerance of 10 ppm.

Bioactivity assay

A hydrolytic enzyme activity of the recombinant cutinase fusion mutant was assessed as described below. The hydrolysis reactions occurred in 96-well microplates at 37°C for 15 min where each well contained 200 µl enzyme/substrate solution comprised of 106.7 µl phosphate buffer (0.1 M, pH 8.0), 13.3 µl Triton X-100 solution (4 g/l), cutinase solution. The reaction was initiated by adding 66.7 µl of substrate reagent solution to each well in the 96-well microplate. Absorbance changes ($\Delta A_{415\text{nm}}$ per min) were measured using a Bio-Rad microplate reader (Tecan, Austria), and solution A with no enzyme solution was used as blank. From the absorbance

changes measured at each column, an average absorbance for a specific reaction condition could then be calculated.

Abbreviations

2-DE, 2-dimensional gel electrophoresis; AID, human activation induced cytidine deaminase; CUT, *Pseudomonas putida* cutinase; EGF, human epidermal growth factor; GAD₄₄₈₋₅₈₅, deletion mutant of human glutamate decarboxylase; G-CSF, human granulocyte colony-stimulating factor; GdnHCl, Guanidine hydrochloride; GST, glutathione-S-transferase; hFTN-H, human ferritin heavy chain; hFTN-L, human ferritin light chain; hIL-2, human interleukin-2; MBP, maltose binding protein; mp-INS, mini-pro-insulin; NACHT, human cold autoinflammatory syndrome 1 protein (NALP3) NACHT domain; PNB, *p*-nitrophenyl butyrate; PNP, *p*-nitrophenyl phamate; ppGRN, human prepro-ghrelin; Trx, thioredoxin.

Authors' contributions

JSP, KYH, and JAS carried out all experiments. SHS and KYA carried out 2-dimensional gel electrophoresis and its analysis. JHL and SWK performed HPLC analysis. JSP and SJS drafted and revised the manuscript in collaboration with JL. JL also participated in the planning, design, and coordination of the research. All authors read and approved the final manuscript.

Acknowledgements

We thank Professor Hang Chul Shin at Soongsil University for kindly providing the gene clones of mpINS and G-CSF. We also appreciate Professors Won Tae Lee and Hyun Soo Cho at Yonsei University for the kind donation of ppGRN, AID, and NACHT clones, respectively. This study was supported by the National Research Laboratory Project of the Ministry of Science and Technology (grant no. ROA-2007-000-20084-0) of the Republic of Korea. This work was also supported by the Korea Health 21 R&D Project of the Ministry of Health & Welfare (grant no. A050750), by grant 031-061-029 of the Ecotechnopia 21 project of the Ministry of Environment, and by the Second Brain Korea 21 Project. Further supports from the Korea Science and Engineering Foundation (grant no. R01-2005-000-10355-0) and the Korea Research Foundation (grant no. KRF-2004-041-D00180) are also appreciated.

References

1. Finley JB, Qui SH, Luan CH, Luo M: **Structural genomics for *Caenorhabditis elegans*: high throughput protein expression analysis.** *Protein Expr Purif* 2004, **34**:49-55.
2. Himanen JP, Rajashankar KR, Lackmann M, Cowan CA, Henkemeyer M, Nikolov DB: **Crystal structure of an Eph receptor-ephrin complex.** *Nature* 2001, **414**:933-938.
3. de Marco A, Deuerling E, Mogk A, Tomoyasu T, Bukau B: **Chaperone-based procedure to increase yields of soluble recombinant proteins produced in *E. coli*.** *BMC Biotechnol* 2007, **7**:32.
4. Qiu J, Swartz JR, Georgiou G: **Expression of active human tissue-type plasminogen activator in *Escherichia coli*.** *Appl Environ Microbiol* 1998, **64**:4891-4896.
5. Tan WS, Dyson MR, Murray K: **Hepatitis B virus core antigen: enhancement of its production in *Escherichia coli*, and interaction of the core particles with the viral surface antigen.** *Biol Chem* 2003, **384**:363-371.
6. Braun P, Hu Y, Shen B, Halleck A, Koundinya M, Harlow E, LaBaer J: **Proteome-scale purification of human proteins from bacteria.** *Proc Natl Acad Sci USA* 2002, **99**:2654-2659.

7. Hammarstrom M, Hellgren N, van Den Berg S, Berglund H, Hard T: **Rapid screening for improved solubility of small human proteins produced as fusion proteins in *Escherichia coli*.** *Protein Sci* 2002, **11**:313-321.
8. Bedouelle H, Duplay P: **Production in *Escherichia coli* and one-step purification of bifunctional hybrid proteins which bind maltose. Export of the Klenow polymerase into the periplasmic space.** *Eur J Biochem* 1998, **171**:541-549.
9. LaVallie ER, DiBlasio EA, Kovacic S, Grant KL, Schendel PF, McCoy JM: **A thioredoxin gene fusion expression system that circumvents inclusion body formation in the *E. coli* cytoplasm.** *Biotechnology (N. Y.)* 1993, **11**:187-193.
10. Ahn JY, Choi H, Kim YH, Han KY, Park JS, Han SS, Lee J: **Heterologous gene expression using self-assembled supra-molecules with high affinity for HSP70 chaperone.** *Nucleic Acids Res* 2005, **33**:3751-3762.
11. Smith DB, Johnson KS: **Single-step purification of polypeptides expressed in *Escherichia coli* as fusions with glutathione-S-transferase.** *Gene* 1988, **67**:31-40.
12. Hammarstrom M, Woestenenk EA, Hellgren N, Hard T, Berglund H: **Effect of N-terminal solubility enhancing fusion proteins on yield of purified target protein.** *J Struct Funct Genomics* 2006, **7**:1-14.
13. Sakane I, Hongo K, Motojima F, Murayama S, Mizobata T, Kawata Y: **Structural stability of covalently linked GroES heptamer: Advantages in the formation of oligomeric structure.** *J Mol Biol* 2007, **367**:1171-1185.
14. Eaglestone SS, Ruddock LW, Cox BS, Tuite MF: **Guanidine hydrochloride blocks a critical step in the propagation of the prion-like determinant [PSI(+)] of *Saccharomyces cerevisiae*.** *Proc Natl Acad Sci USA* 2000, **97**:240-244.
15. Park JS, Ahn JY, Lee SH, Lee H, Han KY, Seo HS, Ahn KY, Min BH, Sim SJ, Choi IS, Kim YH, Lee J: **Enhanced stability of heterologous proteins by supramolecular self-assembly.** *Appl Microbiol Biotechnol* 2007, **75**:347-355.
16. Weber A, Kögl SA, Jung K: **Time-dependent proteome alterations under osmotic stress during aerobic and anaerobic growth in *Escherichia coli*.** *J Bacteriol* 2006, **188**:7165-7175.
17. Chen YC, Chen LA, Chen SJ, Chang MC, Chen TL: **A modified osmotic shock for periplasmic release of a recombinant creatinase from *Escherichia coli*.** *Biochem Eng J* 2004, **19**:211-215.
18. Ewis HE, Lu CD: **Osmotic shock: A mechanosensitive channel blocker can prevent release of cytoplasmic but not periplasmic proteins.** *FEMS Microbiol Lett* 2005, **253**:295-301.
19. Hirsch M, Elliott T: **Stationary-phase regulation of RpoS translation in *Escherichia coli*.** *J Bacteriol* 2005, **187**:7204-7213.
20. Wei B, Shin S, LaPorte D, Wolfe AJ, Romeo T: **Global regulatory mutations in *csr A* and *rpoS* cause severe central carbon stress in *Escherichia coli* in the presence of acetate.** *J Bacteriol* 2000, **182**:1632-1640.
21. Muffler A, Barth M, Marschall C, Hengge-Aronis R: **Heat shock regulation of sigmaS turnover: a role for DnaK and relationship between stress responses mediated by sigmaS and sigma32 in *Escherichia coli*.** *J Bacteriol* 1997, **179**:445-452.
22. Stenström CM, Jin H, Major LL, Tate WP, Isaksson LA: **Codon bias at the 3'-side of the initiation codon is correlated with translation initiation efficiency in *Escherichia coli*.** *Gene* 2001, **263**:273-284.
23. Flipsen JAC, Appel ACM, Van der Hijden HTWM, Verrips CT: **Mechanism of removal of immobilized triacylglycerol by lipolytic enzymes in a sequential laundry wash process.** *Enzyme Microb Technol* 1998, **23**:274-280.
24. Murphy CA, Cameron JA, Huang SJ, Vinopal RT: **Fusarium polyacrylate lactone depolymerase is cutinase.** *Appl Environ Microbiol* 1996, **62**:456-460.
25. Cristina MLC, Maria RAB, Joaquim MSC: **Cutinase: From molecular level to bioprocess development.** *Biotechnol Bioeng* 1999, **66**:17-34.
26. Kim YH, Lee J, Ahn JY, Gu MB, Moon SH: **Enhanced degradation of an endocrine-disrupting chemical, butyl benzyl phthalate, by *Fusarium oxysporum* f. sp. pisi cutinase.** *Appl Environ Microbiol* 2002, **68**:4684-4688.
27. Martinez C, De Geus P, Lauwereys M, Matthyssens G, Cambillau C: ***Fusarium solani* cutinase is a lipolytic enzyme with a catalytic serine accessible to solvent.** *Nature* 1992, **356**:615-618.
28. Gerard HC, Fett WF, Osman SF, Moreau RA: **Evaluation of cutinase activity of various industrial lipases.** *Biotechnol Appl Biochem* 1993, **17**:181-189.
29. Lin TS, Kolattukudy PE: **Structural studies on cutinase, a glycoprotein containing novel amino acids and glucuronic acid amide at the N terminus.** *Eur J Biochem* 1980, **106**:341-351.
30. Kim DY, Lee J, Saraswat V, Park YH: **Glucagon-induced self-association of recombinant proteins in *Escherichia coli* and affinity purification using a fragment of glucagon receptor.** *Biotechnol Bioeng* 2000, **69**:418-428.
31. Choi H, Ahn JY, Sim SJ, Lee J: **Glutamate decarboxylase-derived IDDM autoantigens displayed on self-assembled protein nanoparticles.** *Biochem Biophys Res Commun* 2005, **327**:604-608.
32. Kim SW, Kim YH, Lee J: **Thermal stability of human ferritin: concentration dependence and enhanced stability of an N-terminal fusion mutant.** *Biochem Biophys Res Commun* 2001, **289**:125-129.
33. Kim YH, Han KY, Lee K, Heo JH, Kang HY, Lee J: **Comparative proteome analysis of *Hansenula polymorpha* DL1 and A16.** *Proteomics* 2004, **4**:2005-2013.
34. MS Fit [<http://prospector.ucsf.edu/>]
35. ExPASy Proteomics Server [<http://www.expasy.org/>]
36. Compute pI/Mw tool [http://www.expasy.org/tools/pi_tool.html]
37. Aldente: peptide mass fingerprinting tool [<http://au.expasy.org/tools/aldente/>]

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

