

# Comparing the phosphorus use efficiency of pre-treated (organically) rock phosphate with soluble P fertilizers in maize under calcareous soils

Maria Mussarat<sup>1</sup>, Hazrat Ali<sup>1</sup>, Dost Muhammad<sup>1</sup>, Ishaq Ahmad Mian<sup>1</sup>, Shadman Khan<sup>1</sup>, Muhammad Adnan<sup>2</sup>, Shah Fahad<sup>3,4</sup>, Fazli Wahid<sup>2</sup>, Khadim Dawar<sup>1</sup>, Shamsheer Ali<sup>5</sup>, Afia Zia<sup>6</sup>, Manzoor Ahmad<sup>7</sup>, Sowm Khan<sup>1</sup>, Wajid Ali Shah<sup>7</sup>, Muhammad Romman<sup>8</sup>, Rainaz Parvez<sup>9</sup>, Manzer H. Siddiqui<sup>10</sup>, Abid Khan<sup>11</sup>, Depeng Wang<sup>12</sup> and Xue Jiang<sup>13</sup>

<sup>1</sup> Department of Soil and Environmental Science, The University of Agriculture Peshawar, Peshawar, Khyber Pakhtunkhwa, Pakistan

<sup>2</sup> Department of Agriculture, The University of Swabi, Swabi, Khyber Pakhtunkhwa, Pakistan

<sup>3</sup> Department of Agronomy, University of Haripur, Haripur, Khyber Pakhtunkhwa, Pakistan

<sup>4</sup> Hainan Key Laboratory for Sustainable Utilization of Tropical Bioresource, College of Tropical Crops, Hainan University, Hainan, Haikou, China

<sup>5</sup> Department of Soil and Environmental Science (AMK) campus, The University of Agriculture Peshawar, Mardan, Khyber Pakhtunkhwa, Pakistan

<sup>6</sup> Department of Agricultural Chemistry, The University of Agriculture Peshawar, Peshawar, Khyber Pakhtunkhwa, Pakistan

<sup>7</sup> Department of Agriculture, Bacha Khan University, Charsadda, Khyber Pakhtunkhwa, Pakistan

<sup>8</sup> Department of Botany, University of Chitral, Chitral, KP, Pakistan

<sup>9</sup> Department of Botany, Government Girls Degree College, Dargai Malakand, KP, Pakistan

<sup>10</sup> Botany and Microbiology Department, College of Science, King Saud University, Riyadh, Saudi Arabia

<sup>11</sup> Department of Horticulture, The University of Haripur, Haripur, KPK, Pakistan

<sup>12</sup> College of Life Science, Linyi University, Linyi City, Shandong Province, China

<sup>13</sup> College of Life Sciences and Technology, Xinjiang University, Urumqi City, Xinjiang Province, China

Submitted 5 January 2021

Accepted 23 April 2021

Published 24 May 2021

Corresponding authors

Muhammad Adnan,

madnanses@gmail.com

Depeng Wang,

dwyandywang@163.com

Academic editor

Muhammad Riaz

Additional Information and  
Declarations can be found on  
page 15

DOI 10.7717/peerj.11452

© Copyright

2021 Mussarat et al.

Distributed under

Creative Commons CC-BY 4.0

**OPEN ACCESS**

## ABSTRACT

In calcareous soils, phosphorus (P) availability to plant is impaired due to the formation of insoluble complexes with calcium and magnesium. Therefore, this study was executed to compare the P use efficiency (PUE) of four different P sources [rock phosphate (RP), acidulated rock phosphate (ARP), single super phosphate (SSP) and di ammonium phosphate (DAP)] alone or pre-treated with organic amendments (farm yard manure (FYM) enriched compost, simple compost and humic acid (HA)) along with control in maize crop under calcareous soils.

All treatments irrespective of P sources received 90 kg P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup>. Phosphorus application regardless of its sources and combination with organic amendments significantly improved maize growth, yield as well as P uptake and PUE. Rock phosphate when applied alone was recorded inferior but its performance significantly improved with compost or its pre-addition with FYM and HA, that further enhanced upon acidulation. Maize grain yield increased by 21, 22.2, 67.9 and 94% with RP, ARP, ARP enriched compost and ARP+ compost respectively, over control. Similarly, PUE of DAP improved from 31.7 to 43.1 and 39 with sample and enriched compost correspondingly. Post-harvest soil and grain P were at par for SSP, ARP and

DAP alone or in conjugation with organic amendments when averaged across the amendments. These results suggested that pretreatment of P sources with organic amendments is an economical and more feasible approach to improve maize yield and PUE. Moreover, on-farm acidulation of RP may give at par results with SSP and DAP with cheaper rate and hence recommended for P management in maize in alkaline calcareous soils.

**Subjects** Agricultural Science, Plant Science, Soil Science

**Keywords** Acidulated rock phosphate, Enrich compost, Humic acid, Calcareous soils, Phosphorus use efficiency

## INTRODUCTION

Plants need essential nutrients in the optimum range to complete their life cycle. These nutrients are either supplied through rain, weathering of soil minerals or irrigation, fertilization, or organic matter decay. The fastest route of nutrient availability is via decomposition of organic matter (*Akcura et al., 2019; Bohn, Mcneal & O'Conner, 1985*). However, nutrients availability not only depends on soil properties but also on climatic conditions. Soils of arid and semi-arid regions are low in organic matter and thus can't provide adequate amounts of nutrient to plants for the completion of growth cycle. Under such conditions, soil is supplemented with organic and mineral fertilizers (*Naeem et al., 2021; Ludibeth, Marina & Vicenta, 2012; Johnson, Swedell & Rothman, 2012*). These practices not only overcome the grain shortages but also reduce financial stress on the farmers by overwhelming the cost of fertilizers.

Alkaline soils are most deficient in nitrogen (N) and phosphorus (P), and as a result the yield of edible cereal like maize and wheat is far lower than the potential under these soils (*Adnan et al., 2018*). An ample supplementation of P in early growth stages is important for higher crop yield, as restricted supply of P limits the crop growth (*Grant et al., 2001*). The mineral P added as chemical fertilizers also makes insoluble complexes either with aluminum in acidic soils or calcium and magnesium in alkaline soil (*Makinde, 2007; Ayeni, Adeleye & Adejumo, 2012*), and become unavailable to plants (*Merino et al., 2010*). However, soil supplementation with organic manures in integration with mineral fertilizers will not only enhance nutrients availability; but will also improve soil conditions and crop yield. It is evident from the studies that blended application of organic and in-organic P sources proved to be an effective technique in boosting up inorganic P fertilizers (*Klammsteiner et al., 2020; Whalen & Chang, 2001*). The locally prepared low cost organo-mineral fertilizers have been reported to improve soil health both in terms of fertility and physical conditions (*Hussain, Abid & Raza, 2004*). However, it is a challenging task to locally develop sustainable low input agricultural systems that not only improve soil fertility but also enhance the crop yield.

Humic acid (HA) consists of carbon, oxygen, nitrogen and hydrogen (*Seleiman et al., 2021; Jerzykiewicz et al., 2002; Senesi, 1990*). It may also contain sulphur and P in variable amounts depending on the source. Like most natural polymers, HA have complex structures and contain a number of biologically active functional groups including

carboxylic acid, phenolic, hydroxyl, methoxyl and amine groups (Sposito, 2008). In soil, HA is formed through the chemical and biological humification of plant and animal matter. There is conclusive evidence that quite small amounts of certain organic substances (highly dispersed humic acid, some aromatic compounds and organic acids) have a definite, positive effect on the growth and development of the plant (Fahad et al., 2019a, 2019b; Rafiullah et al., 2021; Kononova, 2013). Similarly, composting is a biological process in which microorganisms convert organic wastes into soil like materials called humus. Compost is a rich source of soil nutrients especially in N and P. It is truly beneficial to soil health as it improves in soil structure, water storage, nutrient supply, soil microbial activity and also improves in soil fertility and productivity (Turan, 2020; Allison, 1973; Johnston, 1986).

Majority of experiments that have been conducted to date on the effect of different P sources over yield but very little is known about the pre-treating/blending of different P sources with organic amendments like HA and composts. Thus, the present study was conducted to compare the economic feasibility and effectiveness of commercial (SSP and DAP) and natural (RP and ARP) P fertilizers alone or in pre-treatment with HA, enriched and sample compost in relation to soil PUE and yield of maize crop under calcareous soils.

## MATERIALS AND METHODS

### Experimental site

A field experiment was conducted to study the effect of different P sources (RP, ARP, SSP and DAP) alone and pretreated with organic amendments (HA, Enriched compost and compost) along with control on yield and P uptake of maize at agronomic research farm (34.02, 71.48), The University of Agriculture, Peshawar. The soil of the experimental site was alkaline (pH = 7.4), calcareous (16.9%), non-saline (EC = 0.17 d Sm<sup>-1</sup>) in nature and silt loam in texture. It was low in organic matter (0.76%) and AB-DTPA extractable P (3.33 mg kg<sup>-1</sup>).

### Preparation of P fertilizers

#### *FYM compost*

Fresh animal dung having 50% moisture was mounded on the ground and covered with a plastic sheet. It was mixed after each 15 days and if necessary some amount of water was added to it until the final odorless FYM compost with black color was formed. It was tested for pH, EC, organic C, nitrogen and phosphorus content.

#### *Rock phosphate and its acidulation*

The ground RP was obtained from Hattar Industrial state, Haripur and was analyzed for total P content. This RP was utilized as such or in the acidulated form called acidulated rock phosphate (ARP) equivalent to SSP. The acidulation of RP was done by treating 100 kg RP with 30 kg H<sub>2</sub>SO<sub>4</sub> in the presence of 60 L water and ARP equivalent to commercial SSP fertilizer was formed (Matiullah & Rafique, 2012; Naseer & Muhammad,

2014). The paste of RP and acids were then retained for 14 days to complete of the reaction. After that the paste was passed through a sieve to get wet granules. These granules were then dried at room temperature for later use.

### ***Extraction of humic acid from coal***

Humic acid (HA) was extracted from coal according to a modified procedure of *Hai & Mir (1998)* and *Khattak, Ul-Ain & Iqbal (2013)*. A 5 kg ground coal was treated with 1.5 L concentrated  $\text{HNO}_3$ , 0.5 L  $\text{H}_2\text{SO}_4$  and 20 L water in a plastic bucket for 24 h and was stirred intermittently with a rod. In order to settle down all suspended particles, reactants (mixture of acid and coal) were allowed to stand for another 24 h. The supernatant was decanted. The pH of the reactants was then raised by the gradual addition of 0.5 N NaOH until it reached 11.5. The solution was then allowed to stand overnight and then centrifuged next day. Acidification of the filtrate was done by adding 0.5 N HCl until the pH reached 2.0. The solution containing HA suspended particles were again centrifuged. The semi-solid HA was dried at 50 °C in an oven to prepare dry solid humic acid.

### ***Enriched composts***

For preparation of enriched compost, a fresh farmyard manure (FYM) was collected from dairy farm of the University of Agriculture Peshawar and was spread on land for two days in order to evaporate excess water. Black polyethylene sheet was spread on a 3 m<sup>2</sup> clean plain area and 100–150 kg of FYM was mixed at the rate of 2% P with RP, ARP, SSP or DAP for preparing enriched composts of the given P sources. The manures and additives were mixed homogeneously and were covered by black polyethylene sheet in order to conserve heat produced during thermophilic stage of compost and to reduce the odor produced during composting.

### ***Preparation of humic acid-complexed P fertilizers***

A solution of humic acid (HA) was prepared in slight alkaline condition to facilitate its dissolution and was sprayed on the granules of ARP, SSP and DAP. The P sources and HA were mixed in such ratios that the final product would supply 90 kg  $\text{P}_2\text{O}_5$  and 5 kg HA ha<sup>-1</sup>. The coated ARP, SSP and DAP were left to dry and absorb the HA, and then applied to soil at the time of sowing.

## **Experimental description**

The experiment was laid out in randomized complete block design (RCBD) with factorial arrangement. Sixteen (16) treatments along with control were replicated three times. Azam variety of maize was sown at the rate of 30 kg ha<sup>-1</sup> on 15<sup>th</sup> June, 2019 in 3 × 5 m<sup>3</sup> plots and then plants were thinned after 3 weeks of emergence. All treatments were treated with basal dose of 150 kg N ha<sup>-1</sup> as a urea inclusive of N obtained from organic amendments. Hoeing and weeding were done manually. However, weedicides were also sprayed for controlling weeds. Standard agronomic practices were applied to all experimental units uniformly throughout the whole growing season.

## Agronomic attributes

### **Biological yield (kg ha<sup>-1</sup>)**

Biological yield was quantified by weighing dry plants harvested from whole plot and then converted into kg ha<sup>-1</sup> as follows (Eq. (1)):

$$\text{Biological yield (kg ha}^{-1}\text{)} = \frac{\text{Yield per plot (kg)}}{\text{Area of experimental plot}} \times \frac{100 \text{ m} \times 100 \text{ m}}{\text{ha}}$$

### **Grain yield (kg ha<sup>-1</sup>)**

Grain yield was measured by the shelling of dry ears obtained from whole plot and then it was converted into kg ha<sup>-1</sup> (Eq. (2)).

$$\text{Grain yield (kg ha}^{-1}\text{)} = \frac{\text{Yield per plot (kg)}}{\text{Area of experimental plot}} \times \frac{100 \text{ m} \times 100 \text{ m}}{\text{ha}}$$

### **Harvesting index (%)**

For determination of harvest index the following formula was used (Eq. (3)):

$$\text{Harvest index (HI \%)} = \frac{\text{Grain yield (kg ha}^{-1}\text{)}}{\text{Biological yield (kg ha}^{-1}\text{)}} \times 100$$

## Soil and plant analysis

Surface soil samples (0–20 cm) were collected at three stages (before sowing, at boot stage and postharvest stage) for evaluating the treatments role in P status of the soil. Soil organic matter was determined by the procedure according to [Nelson & Sommers \(1983\)](#).

Extractable P in soil was quantified on spectrophotometer according to the procedure of [Soltanpour & Schwab \(1977\)](#). The pH and EC of the soil was determined in 1:5 soil water suspensions by the methods of [MeClean \(1982\)](#), and [Richards \(1954\)](#), respectively. Plant P content was determined by using the protocol described by [Jones, Wolf & Mills \(1991\)](#) While, P use efficiency was determined by the method described by [Johnston & Syers \(2009\)](#) via the following formula (Eq. (4)):

$$\text{P use efficiency (kg kg}^{-1}\text{)} = \frac{\text{Grain weight (mg plant}^{-1}\text{)}}{\text{P available in soil (mg plant}^{-1}\text{)}}$$

## Statistical analysis

Statistical analysis of data was performed through analysis of variance (ANOVA) technique as suggested by [Steel & Torrie \(1981\)](#) according to factorial randomized complete block design (RCBD). The means were compared with least significant difference (LSD) at  $p < 0.05$  via computer software Statistix 8.1.

**Table 1** Biological yield of maize as affected by various sources of P fertilizers applied alone or in combination with different organic amendments.

Amendment	Phosphorus source				Mean
	RP	ARP	SSP	DAP	
Control	-	-	-	-	6,099 d
No amendment	7,144	7,625	7,895	8,426	7,773 c
Compost	8,800	9,316	9,606	10,047	9,442 a
Enriched compost	8,623	9,102	8,940	9,508	9,043 a
Humic acid	7,935	8,436	8,370	8,619	8,340 b
Mean	8,125 c	8,620 b	8,703 b	9,150 a	

**Note:**

LSD for P and S levels at  $p < 0.05$  is  $423 \text{ kg ha}^{-1}$  whereas for interaction is NS. RP, ARP, SSP and DAP stand for rock phosphate, acidulated rock phosphate, single super phosphate and di-ammonium phosphate, respectively. The means followed by different letters in each column are significantly different from each other at  $\alpha = 0.05$ .

## RESULTS AND DISCUSSION

### Biological yield ( $\text{kg ha}^{-1}$ )


Application of different phosphate fertilizers significantly increased biological yield of maize as shown in Table 1. Maximum biological yield of  $9,150 \text{ kg ha}^{-1}$  was produced by DAP which was statistically different from the rest of inorganic fertilizers. The SSP and ARP produced statistically similar biological yield of  $8,703 \text{ kg ha}^{-1}$  and  $8,620 \text{ kg ha}^{-1}$  respectively. Among the chemical fertilizers RP produced minimum biological yield but it was higher than control plot ( $6,099 \text{ kg ha}^{-1}$ ).

When averaged across amendments, the maximum biological yield of  $9,442 \text{ kg ha}^{-1}$  was observed for simple compost treated plots which slightly declined with P enriched compost. It should be noted that for P sources with compost the amount of P applied was  $10 \text{ t ha}^{-1}$  whereas, in case of enriched compost its amount was  $2 \text{ t ha}^{-1}$ . Thus, it is certified that the bulk of compost could be reduced by five times through its enrichment without any remarkable reduction in plant growth and yield. HA combined with chemicals fertilizers improved biological yield but their effectiveness was not enough like compost and enriched compost.

In case of integrated use of inorganic and organic fertilizers the remarkable biological yield of  $10,047 \text{ kg ha}^{-1}$  was obtained at DAP + compost treated plots which was significantly higher than the rest of treatment combination. The SSP + compost and ARP + compost receiving plots have also produced the biological yield of  $9,606$  and  $9,316 \text{ kg ha}^{-1}$  respectively (Fig. 1). The RP applied alone was not effective but its effectiveness was pronounced when applied in combination with compost or P enriched composts.

### Grain yield ( $\text{kg ha}^{-1}$ )

Grain yield of maize significantly increased with both sole application of chemical fertilizers and their combination with organic amendments (Table 2). The plots treated with sole DAP produced grain yield of  $3,256 \text{ kg ha}^{-1}$  that was higher than control ( $1,839 \text{ kg ha}^{-1}$ ) but produced superior grain yield of  $3,877 \text{ kg ha}^{-1}$  when DAP was applied with compost. This treatment (DAP + compost) showed the highest grain yield among all



**Figure 1** Increase in biological yield (%) of maize over control as affected by various sources of P fertilizers applied alone or in combination with different organic amendments.

Full-size DOI: 10.7717/peerj.11452/fig-1

**Table 2** Grain yield ( $\text{kg ha}^{-1}$ ) of maize as affected by various sources of P fertilizers applied alone or in combination with different organic amendments.

Amendment	Phosphorus source				Mean
	RP	ARP	SSP	DAP	
Control	-	-	-	-	1,839 e
No amendment	2,229	2,448	2,677	2,861	2,554 d
Compost	2,827	3,569	3,153	3,877	3,357 a
Enriched compost	2,878	3,088	3,105	3,514	3,146 b
Humic acid	2,621	2,840	2,807	2,772	2,760 c
Mean	2,639 c	2,986 b	2,935 b	3,256 a	


**Note:**

LSD for P and S levels at  $p < 0.05$  is  $192.3 \text{ kg ha}^{-1}$  whereas for interaction is NS. RP, ARP, SSP and DAP stand for rock phosphate, acidulated rock phosphate, single super phosphate and di-ammonium phosphate, respectively. The means followed by different letters in each column are significantly different from each other at  $\alpha = 0.05$ .

the applied treatments indicating that the growth and yield of the present crops is associated with solubility of P from the applied fertilizers. The ARP and SSP produced statistically similar grains yield of  $2,986 \text{ kg ha}^{-1}$  and  $2,935 \text{ kg ha}^{-1}$  respectively, those were 62.36% higher than control. The grain yield further increased when ARP and SSP were applied with different types of compost. The sole application of SSP increased grain yield by 40% but rose up to 60% when applied in combination with compost (Fig. 2). Similarly ARP applied alone showed 30% increase in grain yield over control but it was further enhanced when applied with compost suggesting that local acidification is as effective as SSP fertilizer.

### Harvesting index (%)

Analysis of variance regarding harvest index (HI) showed significant differences for phosphorus sources applied alone or in combination with organic amendments (Table 3). When data were averaged across inorganic p sources, maximum HI (35.4%) was noted for



**Figure 2** Increase in grains yield (%) of maize over control as affected by various sources of P fertilizers applied alone or in combination with different organic amendments.

Full-size DOI: 10.7717/peerj.11452/fig-2

**Table 3** Harvesting Index of maize as affected by various sources of P fertilizers applied alone or in combination with different organic amendments.

Amendment	Phosphorus source				Mean
	RP	ARP	SSP	DAP	
Control	-	-	-	-	30.2 c
No amendment	31.2	32.2	34.0	34.1	32.9 b
Compost	32.3	33.9	37.2	38.5	35.5 a
Enriched compost	33.3	34.0	34.8	37.0	34.8 ab
Humic acid	33.1	33.6	33.5	32.1	33.1 ab
Mean	32.5 b	33.4 ab	34.9 ab	35.4 a	

**Note:**

LSD for P and S levels at  $p < 0.05$  is 1.12 whereas for interaction it has also NS, RP, ARP, SSP and DAP stand for rock phosphate, acidulated rock phosphate, single super phosphate and di-ammonium phosphate, respectively. The means followed by different letters in each column are significantly different from each other at  $\alpha = 0.05$ .

DAP followed by ARP (34.9%) and SSP (33.4%) but minimum was calculated for RP applied alone (32.6%) which was better than control (30.2%). Similarly, data revealed that in case of organic P amendments higher HI (35.5%) was observed for compost followed by enriched compost (34.8%). Plots treated with HA gave a HI of 33.1% which was comparatively lower than compost and enriched compost.

### Cost benefit ratio

The benefit cost ratio (CBR) analysis of maize crop and fertilizers are presented in (Table 4). Maximum benefit of Rs. 98,762 and 98,762  $\text{ha}^{-1}$  were obtained from DAP + enriched compost and DAP + compost, respectively. However, their cost benefit ratio were too low (3.1 and 5) compared to ARP enriched compost having net return of 95,540 and CBR of 10.3. The sole RP though had a higher BCR (46.4) but its net return was as low as Rs. 74,637  $\text{ha}^{-1}$ . Any treatment combination who's CBR is greater than 3.0 has



**Table 4** Cost benefit ratio as affected by various sources of P fertilizers applied alone or in combination with different organic amendments.

Treatments	Yield		Income			Cost			Benefit	C:B ratio	
	Stover yield	Grain yield	Stover	Grains	Total	fertilizer	amed.	Total			
RP	0	4,916	2,229	4,916	71,328	76,244	1,607	0	1,607	74,637	46.4
RP	Compost	5,973	2,827	5,973	90,464	96,437	1,607	16,070	17,677	78,760	4.5
RP	E. Com	5,745	2,878	5,745	92,096	97,841	1,607	4,018	5,625	92,216	16.4
RP	H. Acid	5,314	2,621	5,314	83,872	89,186	1,607	2,000	3,607	85,579	23.7
ARP	0	5,177	2,448	5,177	78,336	83,513	5,262	0	5,262	78,251	14.9
ARP	Compost	4,653	3,569	4,653	114,219	118,872	5,262	16,070	21,332	97,540	4.6
ARP	E. Com	6,014	3,088	6,014	98,805	104,820	5,262	4,018	9,280	95,540	10.3
ARP	H. Acid	5,596	2,840	5,596	90,880	96,476	5,262	2,000	7,262	89,214	12.3
SSP	0	5,218	2,677	5,218	85,664	90,882	12,500	0	12,500	78,382	6.3
SSP	Compost	6,453	3,153	6,453	100,896	107,349	12,500	16,070	28,570	78,779	2.8
SSP	E. Com	5,835	3,105	5,835	99,349	105,184	12,500	4,018	16,518	88,667	5.4
SSP	H. Acid	5,563	2,807	5,563	89,813	95,377	12,500	2,000	14,500	80,877	5.6
DAP	0	5,565	2,861	5,565	91,552	97,117	15,652	0	15,652	81,465	5.2
DAP	Compost	6,170	3,877	6,170	124,053	130,224	15,652	16,070	31,722	98,502	3.1
DAP	E. Com	5,994	3,514	5,994	112,437	118,432	15,652	4,018	19,670	98,762	5.0
DAP	H. Acid	5,847	2,772	5,847	88,693	94,540	15,652	2,000	17,652	76,888	4.4

**Note:**

Calculated valued based on: stover Rs. 1/kg, maize grains Rs. 32/kg, RP Rs. 5/kg, compost Rs. 2,000/ton, HA Rs.120/kg H<sub>2</sub>SO<sub>4</sub> Rs. 70/l, DAP Rs. 80/kg, SSP Rs. 25/kg.

considered economical but its benefit should be also kept in mind. The two fertilizers having the same benefit net return then the fertilizer with high CBR would be selected. Similarly, the difference in net return between DAP and ARP enriched compost was too low that's why, ARP enriched compost is recommended under existing soil and climatic conditions.

### Soil organic matter (%)

Data regarding soil organic matter (SOM) as affected by sole application of chemical fertilizers or in combination with different amendments are shown in Table 5. Higher SOM was observed at DAP (0.94%) followed by ARP (0.83%) and SSP (0.82%) while minimum was observed in control plot (0.43%). When data were averaged across the P amendments, significantly higher SOM was observed in plots treated with simple compost. The SOM observed for enriched compost was 0.93% while for HA treated plots was 0.66%. The ARP + compost produced at par SOM with other conventional fertilizers.

### Soil phosphorus concentration at boot stage and after harvest (mg kg<sup>-1</sup>)

The ABDTPA extractable phosphorus in soil at boot stage and after harvesting the crop as influenced by the application of organic and inorganic fertilizers are shown in Tables 6 and 7, respectively. There were no significant differences among the sources for soil phosphorus at boot stage but they were higher than control. When the data were averaged across P amendment maximum soil P concentration (6.28 mg kg<sup>-1</sup>) at boot stage was

**Table 5** Soil organic matter (%) of harvested soil as affected by various sources of P fertilizers applied alone or in combination with different organic amendments.

Amendment	Phosphorus source				Mean
	RP	ARP	SSP	DAP	
Control	-	-	-	-	0.43 d
No amendment	0.52	0.56	0.57	0.59	0.56 c
Compost	0.87	1.19	1.16	1.37	1.15 a
Enriched compost	0.83	0.95	0.85	1.10	0.93 b
Humic acid	0.65	0.60	0.71	0.68	0.66 c
Mean	0.72 b	0.83 ab	0.82 ab	0.94 a	

**Note:**

LSD for P and S levels at  $p < 0.05$  for amendment is 0.16% and for interaction it has NS. RP, ARP, SSP and DAP stand for rock phosphate, acidulated rock phosphate, single super phosphate and di-ammonium phosphate, respectively. The means followed by different letters in each column are significantly different from each other at  $\alpha = 0.05$ .

**Table 6** Soil Phosphorus ( $\text{mg kg}^{-1}$ ) at boot stage as affected by various sources of P fertilizers applied alone or in combination with different organic amendments.

Amendment	Phosphorus source				Mean
	RP	ARP	SSP	DAP	
Control	-	-	-	-	3.42 c
No amendment	4.87	5.21	5.39	5.47	5.23 b
Compost	5.28	6.93	6.34	6.54	6.27 a
Enriched compost	6.27	6.03	6.52	6.07	6.22 a
Humic acid	4.92	5.67	5.63	5.23	5.37 b
Mean	5.34 a	5.96 a	5.97 a	5.83 a	

**Note:**

LSD for P and S levels at  $p < 0.05$  for amendment is  $0.81 \text{ mg kg}^{-1}$  and for interaction it has NS. RP, ARP, SSP and DAP stand for rock phosphate, acidulated rock phosphate, single super phosphate and di-ammonium phosphate, respectively. The means followed by different letters in each column are significantly different from each other at  $\alpha = 0.05$ .

**Table 7** Soil Phosphorus ( $\text{mg kg}^{-1}$ ) of harvested soil as affected by various sources of P fertilizers applied alone or in combination with different organic amendments.

Amendment	Phosphorus source				Mean
	RP	ARP	SSP	DAP	
Control	-	-	-	-	2.17 e
No amendment	3.73	4.07	3.77	4.99	4.14 d
Compost	5.17	5.91	5.90	6.15	5.78 a
Enriched compost	4.96	5.43	5.57	5.63	5.40 b
Humic acid	4.70	4.76	4.50	4.41	4.59 c
Mean	4.64 b	5.04 a	4.93 ab	5.29 a	

**Note:**

LSD for P and S levels at  $p < 0.05$  for amendment is  $0.39 \text{ mg kg}^{-1}$  and for interaction it has NS. RP, ARP, SSP and DAP stand for rock phosphate, acidulated rock phosphate, single super phosphate and di-ammonium phosphate, respectively. The means followed by different letters in each column are significantly different from each other at  $\alpha = 0.05$ .

**Table 8** Phosphorus concentration (%) in grains of maize as affected by various sources of P fertilizers applied alone or in combination with different organic amendments.

Amendment	Phosphorus source				Mean
	RP	ARP	SSP	DAP	
Control	-	-	-	-	0.11 c
No amendment	0.13	0.15	0.13	0.16	0.14 b
Compost	0.17	0.20	0.20	0.22	0.20 a
Enriched compost	0.17	0.21	0.19	0.16	0.18 a
Humic acid	0.16	0.14	0.13	0.16	0.15 b
Mean	0.16 a	0.17 a	0.16 a	0.17 a	

**Note:**

LSD for P and S levels at  $p < 0.05$  for amendment is 0.20 mg per kg and for interaction it has NS. RP, ARP, SSP and DAP stand for rock phosphate, acidulated rock phosphate, single super phosphate and di-ammonium phosphate, respectively. The means followed by different letters in each column are significantly different from each other at  $\alpha = 0.05$ .

observed for plot receiving compost followed by P enriched composts ( $6.22 \text{ mg kg}^{-1}$ ). These treatments were statistically different from HA treated plot having P contents of  $5.37 \text{ mg kg}^{-1}$ .

Post-harvest soil AB-DTPA extractable P was significantly altered by applied treatments. The soil P significantly increased with the sole application of chemical fertilizers or applied with different organic amendments. When data were averaged across fertilizer types, maximum P ( $5.29 \text{ mg kg}^{-1}$ ) was noted for DAP followed by ARP ( $5.04 \text{ mg kg}^{-1}$ ). Minimum P concentration ( $2.17 \text{ mg kg}^{-1}$ ) was observed for control. Similarly, highest P contents of  $5.59 \text{ mg kg}^{-1}$  was observed with the addition of compost followed by different P enriched composts ( $5.57 \text{ mg kg}^{-1}$ ). Addition of HA did not improve soil P compared to composts. Interactive effect demonstrated that soil P was highly improved ( $6.02 \text{ mg kg}^{-1}$ ) with DAP + compost followed by DAP enriched compost. As DAP is considered highly soluble then other P sources (mineral) hence, it added more P to soil with the combination of compost.

### Phosphorus concentrations in grains and stalk (%)

Data regarding phosphorus concentration in grains and stalk as affected by organic and inorganic P fertilizers are presented in Tables 8 and 9 correspondingly. Chemical fertilizers didn't show any significant effect for grain P concentration in maize. However, when data were averaged across different P amendments significant differences were observed. Higher P content (0.20%) was observed for the compost treated plots which was statistically higher than that observed for P enriched composts (0.18%).

Significant differences were observed among the P sources for stalk P content in maize. Averaging the data across fertilizers, maximum P concentration of 0.26% was observed at DAP followed by SSP (0.24%) which were statistically different from treatments fertilized with ARP (0.21%) and RP (0.18%), however, they were higher than control plots (0.13%). When data were averaged across P amendments, compost treated maize plants absorbed optimum P (0.27%) than other P amendments and it was

**Table 9** Phosphorus concentration (%) in stalk of maize as affected by various sources of P fertilizers applied alone or in combination with different organic amendments.

Amendment	Phosphorus source				Mean
	RP	ARP	SSP	DAP	
Control	-	-	-	-	0.13 d
No amendment	0.15	0.18	0.20	0.19	0.18 c
Compost	0.22	0.25	0.27	0.32	0.27 a
Enriched compost	0.17	0.21	0.23	0.27	0.22 b
Humic acid	0.17	0.19	0.25	0.24	0.21 bc
Mean	0.18 c	0.21 bc	0.24 ab	0.26 a	

**Note:**

LSD for P and S levels at  $p < 0.05$  for amendment is 0.039% and for interaction it has NS. RP, ARP, SSP and DAP stand for rock phosphate, acidulated rock phosphate, single super phosphate and di-ammonium phosphate, respectively. The means followed by different letters in each column are significantly different from each other at  $\alpha = 0.05$ .

**Table 10** Phosphorus use efficiency ( $\text{kg kg}^{-1}$ ) of maize as affected by various sources of P fertilizers applied alone or in combination with different organic amendments.

Amendment	Phosphorus source				Mean
	RP	ARP	SSP	DAP	
Control	-	-	-	-	22.30 e
No amendment	25.03	27.20	29.74	31.78	28.44 d
Compost	31.41	35.03	39.66	43.07	37.29 a
Enriched compost	31.97	34.30	34.49	39.04	34.95 b
Humic acid	29.12	31.56	31.18	30.79	30.66 c
Mean	29.38 c	32.02 b	33.77 b	36.17 a	

**Note:**

LSD for P and S levels at  $p < 0.05$  for amendment is  $2.19 \text{ kg kg}^{-1}$  and for interaction it has NS. RP, ARP, SSP and DAP stand for rock phosphate, acidulated rock phosphate, single super phosphate and di-ammonium phosphate, respectively. The means followed by different letters in each column are significantly different from each other at  $\alpha = 0.05$ .

statistically higher than the rest of treatment combination. In Enriched compost fertilized plots, plants absorbed slightly higher P (0.23%) than sole chemical fertilizers.

### Phosphorus use efficiency (%)

Phosphorus use efficiency (PUE) was significantly affected by various P sources either applied alone or in combination with different amendments (Table 10). Maximum PUE was observed for DAP ( $36.17 \text{ kg kg}^{-1}$ ) followed by SSP ( $33.7 \text{ kg kg}^{-1}$ ) and ARP ( $32.02 \text{ kg kg}^{-1}$ ) while the minimum PUE ( $23.12 \text{ kg kg}^{-1}$ ) was observed for control. When data were averaged across organic source, maximum PUE was observed for sample compost ( $37.29 \text{ kg kg}^{-1}$ ) which was statistically different from all other amendments. Although P enriched compost had closely similar PUE value  $34.95 \text{ kg kg}^{-1}$  suggesting that chemical fertilizers enriched with compost has responsive effect because they may release much phosphorus to the soil which is the key factor in organic agriculture for improving yield and yield related components. The HA treated plots resulted a PUE of  $30.66 \text{ kg kg}^{-1}$  indicating that it had also additional beneficial P value when applied with chemical fertilizers, it may be meaningful in releasing more P from in-organic sources.

## DISCUSSION

Biological and grain yield of maize showed a significant response to the application of mineral phosphorus and in combination with compost except rock phosphate. The increment in the yield of cereals crops was recorded due to the application of P fertilizers (Bilen, Bilen & Turan, 2019; Huang et al., 2011). Amano (1989) reported similar findings of higher harvest index of maize with the application of phosphorus fertilizer. However, Osinama (1995) reported no significant differences in grain yield, stover yield and harvest index with different application of the P fertilizers. This could happen due to the slow release of P from the composts (Zhang et al., 2013) and the availability of enough P to plant for long time can ensure better root system resulted in maximum growth, yield and P uptake (Table 2). Alkhader, Rayyan & Rusan (2013) also applied four different levels of phosphorus fertilizers on lettuce both in field as well as in glass house experiments and found that among all the treatments RP produced lowest plant growth and yield as compared to others P sources. When data were averaged across the organic amendments grain yield was higher than chemical fertilizers. As we know that compost and P enriched compost are considered soil conditioner which can improve soil health. Good and optimal supply of phosphorus can enhanced root activities which help plants in exploring soil nutrients and retained more soil moisture. That is why, control plot produced uneconomical grain yield which haven't utilize sufficient nutrients required by the plants for its proper nourishment. Jamal et al. (2018) summarized from their experiment that 90 kg ha<sup>-1</sup> P applied along with 500 g of HA resulted optimum yield of wheat crop. With the increase of biological yield, grain yield was also enhanced because of the optimum levels of nutrients were taken up by the crop (Alkhader, Rayyan & Rusan, 2013). Our results are also supported by Qin et al. (2010) and Jiang & Xie (2009), who concluded that zero tillage with phosphorus amendments has a significant increase in the grain yield of crop compared to conventional tillage. These findings are also aligned with Mashori et al. (2013) suggesting that RP integrated with SSP and FYM enhanced maize growth, biological yield, leaf P content and uptake. The production of rice was positively affected by mineral fertilizers amendments (Turan, 2019; Lan et al., 2012).

The application of mineral fertilizers enhanced the P fraction on the surface of soil (Adnan et al., 2020; Jiang & Xie, 2009). That's why, post-harvest soil and grain P concentrations were statistically similar for SSP, ARP and DAP when averaged across the amendments. Similarly, the HA compost and enriched compost gave at par results when averaged across the fertilizer types, indicating that any phosphate fertilizer applied with amendments could improve the soil P level and its uptake by plant (Tables 7 & 8). These results were in line with Mashori et al. (2013), who concluded that RP integrated with SSP and FYM enhanced maize growth, dry matter, leaf P content and uptake. Our results are also in agreement to many researchers (Turan et al., 2019; Jiang & Xie, 2009). With the maintaining of organic matter on the surface of soil enhanced the fertility of soil. Organic fraction of soil significantly enhanced the acidifying nature of soil, thus, enhanced the availability of soluble P (Huang et al., 2012). Organic materials increased the

soil organic matter from 1.17 to 2.85 kg ha<sup>-1</sup> and reduced the oxidation stability of SOM (Wang, Miao & Li, 2015; Li et al., 2017; Sayyed et al., 2019; Adnan et al., 2019). Organic section of soil enhanced soil fertility (Zhang et al., 2011). Our result contrasts with the result of Huang & Yuan (1994) who reported the decrease of organic phosphorus with the amendments of mineral P and no tillage surface. Moreover, we observed an increase in soil P resulting from an increase in soil organic P, which is also in line with Tiecher, Dos Santos & Calegari (2012).

The soil P content at boot stage of the maize crop has shown an influential effect with the application of mineral P. This agrees with the findings of Zhang et al. (2006). Lan et al. (2012) and Lee, Lee & Kim (2004) reported that the application of fertilizer significantly enhanced the accumulation of P fraction in soil. Our results are also supported by Sharif, Khattak & Sarir (2002) who reported that HA addition to the soil generally reduce P fixation and increase the recovery of P. Similarly, Zhang et al. (1994) observed that in the rice field, organic P linearly increase with the amendments of P fertilizers. Improvement in grain P concentration of maize crops is in line with Khan et al. (2020) who reported that as the level of organic substitution increased from 25–75%, the phosphorus content exhibit a progressive increase. Roy & Tripathi (1987) also reported increase in nutrient content and uptake with increasing application of P. Our results oppose the findings of Ibrahim et al. (2011) who reported linear decrease in P uptake with the increments of fertilizer P. According to Jamal et al. (2018), application of different levels of P and HA either applied alone or in combination significantly increased soil P concentration over control. Our results contrasts with Zhang et al. (2006) who determined the limitation of extractable P in paddy soil under calcareous condition, but in line with Adeyemi et al. (2020) and Lan et al. (2012) who observed significant increase in labile organic P and total organic P concentration with application of P. These findings were in consistency with Waldrip-Dail et al. (2009) who reported that P values increased as the crop gradually matured. Phosphorous use efficiency showed a remarkable result for the integrated application of mineral and organic P and these findings were also studied by Ademba et al. (2015) who reported that integrated use of phosphate fertilizers and manure applications significantly improved maize yield and PUE. The excess P application can increase the risk of P movement to surface and groundwater (Grant et al., 2005) and thus PUE decreases.

## CONCLUSIONS

ARP performed similar to SSP and DAP either applied alone or in combination with composts and HA for most of studied traits. The performance of sole RP was inferior but improved by addition of compost, HA or it's pretreatment with organic amendments—(i.e. sample and RP-enriched compost). Phosphate fertilizer (DAP/SSP) in integration with composts significantly improved the yield and PUE of maize as compared to other treatments. Plant growth and yield though was comparatively lower in different P enriched compost treated plots but still it could be considered economical and feasible as the bulk of compost was reduced by 10 times. These findings imply that, the PUE and economic feasibility of natural/chemical P fertilizers can be improved by either

their integrated application or pre-treatment with organic amendment and shall be practiced for improved crop productivity under calcareous soils.

## ADDITIONAL INFORMATION AND DECLARATIONS

### Funding

This research was funded by The National Key Research and Development Program of China (2018YFD0300606). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

### Grant Disclosures

The following grant information was disclosed by the authors:

National Key Research and Development Program of China: 2018YFD0300606.

### Competing Interests

The authors declare that they have no competing interests.

### Author Contributions

- Maria Mussarat conceived and designed the experiments, prepared figures and/or tables, and approved the final draft.
- Hazrat Ali conceived and designed the experiments, prepared figures and/or tables, and approved the final draft.
- Dost Muhammad conceived and designed the experiments, analyzed the data, prepared figures and/or tables, and approved the final draft.
- Ishaq Ahmad Mian performed the experiments, prepared figures and/or tables, and approved the final draft.
- Shadman Khan performed the experiments, analyzed the data, prepared figures and/or tables, and approved the final draft.
- Muhammad Adnan performed the experiments, analyzed the data, prepared figures and/or tables, and approved the final draft.
- Shah Fahad performed the experiments, analyzed the data, prepared figures and/or tables, and approved the final draft.
- Fazli Wahid performed the experiments, prepared figures and/or tables, and approved the final draft.
- Khadim Dawar performed the experiments, prepared figures and/or tables, and approved the final draft.
- Shamsher Ali performed the experiments, prepared figures and/or tables, and approved the final draft.
- Afia Zia performed the experiments, authored or reviewed drafts of the paper, and approved the final draft.
- Manzoor Ahmad performed the experiments, authored or reviewed drafts of the paper, and approved the final draft.
- Sowm Khan performed the experiments, authored or reviewed drafts of the paper, and approved the final draft.

- Wajid Ali Shah performed the experiments, authored or reviewed drafts of the paper, and approved the final draft.
- Muhammad Romman conceived and designed the experiments, authored or reviewed drafts of the paper, and approved the final draft.
- Rainaz Parvez conceived and designed the experiments, authored or reviewed drafts of the paper, and approved the final draft.
- Manzer H. Siddiqui conceived and designed the experiments, authored or reviewed drafts of the paper, and approved the final draft.
- Abid Khan conceived and designed the experiments, prepared figures and/or tables, and approved the final draft.
- Depeng Wang conceived and designed the experiments, analyzed the data, authored or reviewed drafts of the paper, and approved the final draft.
- Xue Jiang conceived and designed the experiments, prepared figures and/or tables, and approved the final draft.

### Data Availability

The following information was supplied regarding data availability:

Raw data are available as a [Supplemental File](#).

### Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.11452#supplemental-information>.

## REFERENCES

- Ademba JS, Kwach JK, Esilaba AO, Ngari SM. 2015. The effects of phosphate fertilizers and manure on maize yields in South Western Kenya, Egypt. *East African Agricultural and Forestry Journal* **81**(1):1 DOI 10.1080/00128325.2015.1040640.
- Adeyemi O, Keshavarz-Afshar R, Jahanzad E, Battaglia ML, Luo Y, Sadeghpour A. 2020. Effect of wheat cover crop and split nitrogen application on corn yield and nitrogen use efficiency. *Agronomy* **10**(8):1081 DOI 10.3390/agronomy10081081.
- Adnan M, Fahad S, Khan IA, saeed M, Saud S, Ihsan MZ, Raiz M, Wang D, Wu C. 2019. Integration of poultry manure and phosphate solubilizing bacteria improved availability of Ca bound P in calcareous soils. *3 Biotech* **9**(10):1–10 DOI 10.1007/s13205-019-1894-2.
- Adnan M, Fahad S, Zamin M, Shah S, Mian IA, Danish S, Zafar-ul-Hye M, Battaglia ML, Naz RMM, Saeed B, Saud S, Ahmad I, Yue Z, Brtnicky M, Holatko J, Datta R. 2020. Coupling phosphate-solubilizing bacteria with phosphorus supplements improve maize phosphorus acquisition and growth under lime induced salinity stress. *Plants* **9**(7):900 DOI 10.3390/plants9070900.
- Adnan M, Shah Z, Sharif M, Rahman H. 2018. Liming induces carbon dioxide (CO<sub>2</sub>) emission in PSB inoculated alkaline soil supplemented with different phosphorus sources. *Environmental Science and Pollution Research* **25**(10):9501–9509 DOI 10.1007/s11356-018-1255-4.
- Akcura M, Turan V, Kokten K, Kaplan M. 2019. Fatty acid and some micro element compositions of cluster bean (*Cyamopsis tetragonoloba*) genotype seeds growing under Mediterranean climate. *Industrial Crops and Products* **128**(2):140–146 DOI 10.1016/j.indcrop.2018.10.062.



- Alkhader A, Rayyan A, Rusan M. 2013.** The effect of phosphorus fertilizers on the growth and quality of lettuce (*Lactuca sativa* L.) under greenhouse and field conditions. *Journal of Food, Agriculture & Environment* **11**(2):777–783.
- Allison FE. 1973.** *Soil organic matter and its role in crop production*. Amsterdam: Elsevier.
- Amano LO. 1989.** Comparative productivity of corn and sorghum as affected by population density and nitrogen fertilization. *Philippine Agriculture* **72**:247–254.
- Ayeni LS, Adeleye EO, Adejumo JO. 2012.** Comparative effect of organic, organomineral and mineral fertilizers on soil properties, nutrient uptake, growth and yield of maize (*Zea mays*). *International Research Journal of Agricultural Science and Soil Science* **2**(11):493–497.
- Bilen S, Bilen M, Turan V. 2019.** Relationships between cement dust emissions and soil properties. *Polish Journal of Environmental Studies* **28**(5):3089–3098 DOI [10.15244/pjoes/92521](https://doi.org/10.15244/pjoes/92521).
- Bohn HL, Mcneal BL, O'Conner GA. 1985.** *Soil chemistry*. Second Edition. New York: John Wiley and sons.
- Fahad S, Adnan M, Hassan S, Saud S, Hussain S, Wu C, Wang D, Hakeem KR, Alharby HF, Turan V, Khan MA, Huang J. 2019a.** Rice responses and tolerance to high temperature. In: *Advances in Rice Research for Abiotic Stress Tolerance*, Amsterdam: Elsevier, 201–224.
- Fahad S, Rehman A, Shahzad B, Tanveer M, Saud S, Kamran M, Ihtisham M, Khan SU, Turan V, Rahman MH. 2019b.** Rice responses and tolerance to metal/metalloid toxicity. In: *Advances in Rice Research for Abiotic Stress Tolerance*, Amsterdam: Elsevier, 299–312.
- Grant C, Bittman S, Montreal M, Plenchette C, Morel C. 2005.** Soil and fertilizer phosphorus: effects on plant P supply and mycorrhizal development. *Canadian Journal of Plant Science* **85**(1):3–14 DOI [10.4141/P03-182](https://doi.org/10.4141/P03-182).
- Grant K, Cravens DW, Low GS, Moncrief WC. 2001.** The role of satisfaction with territory design on the motivation, attitudes, and work outcomes of salespeople. *Journal of the Academy of Marketing Science* **29**(2):165–178 DOI [10.1177/03079459994533](https://doi.org/10.1177/03079459994533).
- Hai SM, Mir S. 1998.** The lignitic coal derived HA and the prospective utilization in Pakistan agriculture and industry. *Science Technology* **17**(3):32–40.
- Huang JG, Yuan L. 1994.** Effects of tillage on P status in paddy soils. *Journal of Southwest Agriculture University* **16**:475–477.
- Huang M, Zou Y, Feng Y, Cheng Z, Mo Y, Ibrahim M, Xia B, Jiang P. 2011.** No-tillage and direct seeding for super hybrid rice production in rice–oilseed rape cropping system. *European Journal of Agronomy* **34**(4):278–286 DOI [10.1016/j.eja.2011.02.005](https://doi.org/10.1016/j.eja.2011.02.005).
- Huang M, Zou Y, Jiang P, Xia B, Feng Y, Cheng Z, Mo Y. 2012.** Effect of tillage on soil and crop properties of wet-seeded flooded rice. *Field Crops Research* **129**:28–38 DOI [10.1016/j.fcr.2012.01.013](https://doi.org/10.1016/j.fcr.2012.01.013).
- Hussain N, Abid M, Raza I. 2004.** Response of wheat to phosphorus in the presence of farmyard manure. *Indus Journal Plant Science* **3**:298–302.
- Ibrahim M, Yamin M, Sarwar G, Anayat A, Habib F, Ullah S. 2011.** Tillage and farm manure affect root growth and nutrient uptake of wheat and rice under semi-arid conditions. *Applied Geochemistry* **26**:194–197 DOI [10.1016/j.apgeochem.2011.03.102](https://doi.org/10.1016/j.apgeochem.2011.03.102).
- Jamal A, Hussain I, Sarir M, Sharif M, Fawad M. 2018.** Investigating combination and individual impact of phosphorus and humic acid on yield of wheat and some soil properties. *Türk Tarım ve Doğa Bilimleri Dergisi* **5**(4):492–500.
- Jerzykiewicz M, Jezierski A, Czechowski F, Drozd J. 2002.** Influence of metal ions binding on free radical concentration in humic acids: a quantitative electron paramagnetic resonance study. *Organic Geochemistry* **33**(3):265–268 DOI [10.1016/S0146-6380\(01\)00158-9](https://doi.org/10.1016/S0146-6380(01)00158-9).

- Jiang X-J, Xie D-T. 2009.** Combining ridge with no-tillage in lowland rice-based cropping system: long-term effect on soil and rice yield. *Pedosphere* **19**(4):515–522  
DOI [10.1016/S1002-0160\(09\)60144-2](https://doi.org/10.1016/S1002-0160(09)60144-2).
- Johnson CA, Swedell L, Rothman JM. 2012.** Feeding ecology of olive baboons (*Papio anubis*) in Kibale National Park, Uganda: preliminary results on diet and food selection. *African Journal of Ecology* **50**(3):367–370.
- Johnston AE. 1986.** Soil organic matter, effects on soils and crops. *Soil Use and Management* **2**(3):97–105 DOI [10.1111/j.1475-2743.1986.tb00690.x](https://doi.org/10.1111/j.1475-2743.1986.tb00690.x).
- Johnston AE, Syers JK. 2009.** A new approach to assessing phosphorus use efficiency in agriculture. *Better Crops* **93**(3):14–16.
- Jones ML, Wolf B, Mills HA. 1991.** *A practical sampling, preparation, analysis, and interpretation guide: plant analysis handbook*. Athens: Micro-Macro Publishing, Inc.
- Khan MA, Mahmood-ur-Rahman, Ramzani PMA, Zubair M, Rasool B, Khan MK, Ahmed A, Khan SA, Turan V, Iqbal M. 2020.** Associative effects of lignin-derived biochar and arbuscular mycorrhizal fungi applied to soil polluted from Pb-acid batteries effluents on barley grain safety. *Science of the Total Environment* **710**(4):136294 DOI [10.1016/j.scitotenv.2019.136294](https://doi.org/10.1016/j.scitotenv.2019.136294).
- Khattak MA, Ul-Ain Q, Iqbal N. 2013.** Impact of role ambiguity on job satisfaction, mediating role of job stress. *International Journal of Academic Research in Accounting, Finance and Management Sciences* **3**(3):28–39 DOI [10.6007/IJARAFMS/v3-i3/33](https://doi.org/10.6007/IJARAFMS/v3-i3/33).
- Klammsteiner T, Turan V, Fernández-Delgado Juárez M, Oberegger S, Insam H. 2020.** Suitability of black soldier fly frass as soil amendment and implication for organic waste hygienization. *Agronomy* **10**(10):1578 DOI [10.3390/agronomy10101578](https://doi.org/10.3390/agronomy10101578).
- Kononova MM. 2013.** *Soil organic matter: its nature, its role in soil formation and in soil fertility*. Amsterdam: Elsevier.
- Lan ZM, Lin XJ, Wang F, Zhang H, Chen CR. 2012.** Phosphorus availability and rice grain yield in a paddy soil in response to long-term fertilization. *Biology and Fertility of Soils* **48**(5):579–588 DOI [10.1007/s00374-011-0650-5](https://doi.org/10.1007/s00374-011-0650-5).
- Lee CH, Lee IB, Kim PJ. 2004.** Effects of long-term fertilization on organic phosphorus fraction in paddy soil. *Soil Science and Plant Nutrition* **50**(4):485–491  
DOI [10.1080/00380768.2004.10408504](https://doi.org/10.1080/00380768.2004.10408504).
- Li S, Li J, Li G, Li Y, Yuan J, Li D. 2017.** Effect of different organic fertilizers application on soil organic matter properties. *Compost Science & Utilization* **25**(1):31–36  
DOI [10.1080/1065657X.2017.1344160](https://doi.org/10.1080/1065657X.2017.1344160).
- Ludibeth SM, Marina IE, Vicenta EM. 2012.** Vermicomposting of sewage sludge: earthworm population and agronomic advantages. *Compost Science & Utilization* **20**(1):11–17  
DOI [10.1080/1065657X.2012.10737016](https://doi.org/10.1080/1065657X.2012.10737016).
- Makinde EA. 2007.** Effects of an organo-mineral fertilizer application on the growth and yield of maize. *Journal of Applied Sciences Research* **3**(10):1152–1155.
- Mashori NM, Memon M, Memon KS, Kakar H. 2013.** Maize dry matter yield and P uptake as influenced by rock phosphate and single super phosphate treated with farm manure. *Soil & Environment* **32**(2):1.
- Matiullah MF, Rafique M. 2012.** Indoor radon monitoring near an in situ leach mining site in DG Khan, Pakistan. *Journal of Radiological Protection* **32**(4):427–437  
DOI [10.1088/0952-4746/32/4/427](https://doi.org/10.1088/0952-4746/32/4/427).
- McClellan EO. 1982.** Soil pH and lime requirement. In: Page AL, Miller RH, Keeny DR, eds. *Method of soil Analysis: Part 22nd Edition*. Vol. 9. Madison: American Society of, 199–208.

- Merino G, Alberdi M, Ivanov AG, Diaz MR. 2010.** Al<sup>+3</sup> ca<sup>+2</sup> interaction in plants growing in acid soils: Al phyto-toxicity response to calcareous amendments. *Journal of Soil Science & Plant Nutrition* **0**(3):217–243.
- Naeem I, Masood N, Turan V, Iqbal M. 2021.** Prospective usage of magnesium potassium phosphate cement combined with *Bougainvillea alba* derived biochar to reduce Pb bioavailability in soil and its uptake by *Spinacia oleracea* L. *Ecotoxicology and Environmental Safety* **208**(1–4):111723 DOI [10.1016/j.ecoenv.2020.111723](https://doi.org/10.1016/j.ecoenv.2020.111723).
- Naseer M, Muhammad D. 2014.** Direct and residual effect of hazara rock phosphate (HRP) on wheat and succeeding maize in alkaline calcareous soils. *Pakistan Journal of Botany* **46**(5):1755–1761.
- Nelson DW, Sommers L. 1983.** Total carbon, organic carbon, and organic matter. *Methods of soil analysis: Part 2 Chemical and Microbiological Properties* **9**:539–579.
- Osinama OA. 1995.** Evaluation of three P sources for maize in South Western Nigeria. *Tropical Agriculture* **62**:33–37.
- Qin J, Hu F, Li D, Li H, Lu J, Yu R. 2010.** The effect of mulching, tillage and rotation on yield in non-flooded compared with flooded rice production. *Journal of Agronomy and Crop Science* **196**(6):397–406 DOI [10.1111/j.1439-037X.2010.00430.x](https://doi.org/10.1111/j.1439-037X.2010.00430.x).
- Rafiullah, Khan MJ, Muhammad D, Mussarat M, Huma, Adnan M, Fazli W, Arif M, Amanullah. 2021.** Foliar versus soil phosphorus (P) application for improving P use efficiency in wheat and maize in calcareous soils. *Journal of Plant Nutrition* **42**(19):2423–2434 DOI [10.1080/01904167.2019.1659344](https://doi.org/10.1080/01904167.2019.1659344).
- Richards LA. 1954.** *Diagnosis and improvement of saline and Alkali soils (USDA): Handbook*. Vol. 60. Washington: US. Govt., Printing Office.
- Roy RK, Tripathi RS. 1987.** Effect of irrigation and fertilizers on yield, water-use efficiency and nutrient concentration in winter maize. *Indian Journal of Agronomy* **32**(4):314–318.
- Sayed MI, Akman F, Turan V, Araz A. 2019.** Evaluation of radiation absorption capacity of some soil samples. *Radiochimica Acta* **107**(1):83–93 DOI [10.1515/ract-2018-2996](https://doi.org/10.1515/ract-2018-2996).
- Seleiman MF, Almutairi KF, Alotaibi M, Shami A, Alhammad BA, Battaglia ML. 2021.** Nano fertilization as an emerging fertilization technique: why modern agriculture can benefit from its use? *Plants* **10**(1):2 DOI [10.3390/plants10010002](https://doi.org/10.3390/plants10010002).
- Senesi N. 1990.** Application of electron spin resonance (ESR) spectroscopy in soil chemistry. In: *Advances in Soil Science*. New York, NY: Springer, 77–130.
- Sharif M, Khattak RA, Sarir MS. 2002.** Wheat yield and nutrients accumulation as affected by humic acid and chemical fertilizers. *Sarhad Journal of Agriculture* **18**(3):323–329.
- Soltanpour PN, Schwab AP. 1977.** A new soil test for simultaneous extraction of macro-and micro-nutrients in alkaline soils. *Communications in Soil Science and Plant Analysis* **8**(3):195–207 DOI [10.1080/00103627709366714](https://doi.org/10.1080/00103627709366714).
- Sposito G. 2008.** *The chemistry of soils*. Oxford: Oxford University Press.
- Steel RG, Torrie JH. 1981.** *Principles and procedures of statistics*. New York: McGraw-Hill.
- Tiecher T, Dos Santos DR, Calegari A. 2012.** Soil organic phosphorus forms under different soil management systems and winter crops, in a long term experiment. *Soil and Tillage Research* **124**:57–67 DOI [10.1016/j.still.2012.05.001](https://doi.org/10.1016/j.still.2012.05.001).
- Turan V. 2019.** Confident performance of chitosan and pistachio shell biochar on reducing Ni bioavailability in soil and plant plus improved the soil enzymatic activities, antioxidant defense system and nutritional quality of lettuce. *Ecotoxicology and Environmental Safety* **183**(4):109594 DOI [10.1016/j.ecoenv.2019.109594](https://doi.org/10.1016/j.ecoenv.2019.109594).

- Turan V. 2020.** Potential of pistachio shell biochar and dicalcium phosphate combination to reduce Pb speciation in spinach, improved soil enzymatic activities, plant nutritional quality, and antioxidant defense system. *Chemosphere* **245**(4):125611  
DOI [10.1016/j.chemosphere.2019.125611](https://doi.org/10.1016/j.chemosphere.2019.125611).
- Turan V, Schröder P, Bilen S, Insam H, Fernández-Delgado Juárez M. 2019.** Co-inoculation effect of Rhizobium and *Achillea millefolium* L. oil extracts on growth of common bean (*Phaseolus vulgaris* L.) and soil microbial-chemical properties. *Scientific Reports* **9**(1):15178  
DOI [10.1038/s41598-019-51587-x](https://doi.org/10.1038/s41598-019-51587-x).
- Waldrip-Dail H, He Z, Erich SM, Honeycutt WC. 2009.** Soil phosphorus dynamics in response to poultry manure amendment. *Soil Science* **174**(4):195–201 DOI [10.1097/SS.0b013e31819cd25d](https://doi.org/10.1097/SS.0b013e31819cd25d).
- Wang ZH, Miao YF, Li SX. 2015.** Effect of ammonium and nitrate nitrogen fertilizers on wheat yield in relation to accumulated nitrate at different depths of soil in drylands of China. *Field Crops Research* **183**(7):211–224 DOI [10.1016/j.fcr.2015.07.019](https://doi.org/10.1016/j.fcr.2015.07.019).
- Whalen JK, Chang C. 2001.** Phosphorus accumulation in cultivated soils from long-term annual applications of cattle feedlot manure. *Journal of Environmental Quality* **30**(1):229–237  
DOI [10.2134/jeq2001.301229x](https://doi.org/10.2134/jeq2001.301229x).
- Zhang ZS, Cao CG, Cai ML, Li CF. 2013.** Crop yield, P uptake and soil organic phosphorus fractions in response to short-term tillage and fertilization under a rape-rice rotation in central China. *Journal of Soil Science and Plant Nutrition* **13**(4):871–882.
- Zhang Q, Wang GH, Feng YK, Sun QZ, Witt C, Dobermann A. 2006.** Changes in soil phosphorus fractions in a calcareous paddy soil under intensive rice cropping. *Plant and Soil* **288**(1–2):141–154 DOI [10.1007/s11104-006-9100-9](https://doi.org/10.1007/s11104-006-9100-9).
- Zhang YS, Werner W, Scherer HW, Sun X. 1994.** Effect of organic manure on organic phosphorus fractions in two paddy soils. *Biology and Fertility of Soils* **17**(1):64–68  
DOI [10.1007/BF00418674](https://doi.org/10.1007/BF00418674).
- Zhang JS, Zhang FP, Yang JH, Wang JP, Cai ML, Li CF, Cao CG. 2011.** Emissions of N<sub>2</sub>O and NH<sub>3</sub>, and nitrogen leaching from direct seeded rice under different tillage practices in central China. *Agriculture Ecosystem and Environment* **140**(1–2):164–173  
DOI [10.1016/j.agee.2010.11.023](https://doi.org/10.1016/j.agee.2010.11.023).