



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



ABO blood group and link to COVID-19: A comprehensive review of the reported associations and their possible underlying mechanisms

Eric Pereira^a, Stela Felipe^a, Raquel de Freitas^a, Valdevane Araújo^a, Paula Soares^a, Jannison Ribeiro^b, Luiz Henrique dos Santos^a, Juliana Osório Alves^a, Natália Canabrava^c, Mauricio van Tilburg^c, Maria Izabel Guedes^c, Vânia Ceccatto^{a,*}

^a Superior Institute of Biomedical Sciences, State University of Ceará, Dr. Silas Munguba Av., Fortaleza, 60714-903, Ceará, Brazil

^b Hematology and Hemotherapy Center of Ceará, José Bastos Av., Fortaleza, 60431-086, Ceará, Brazil

^c Biotechnology and Molecular Biology Laboratory, State University of Ceará, Dr. Silas Munguba Av., Fortaleza, 60714-903, Ceará, Brazil

ARTICLE INFO

Keywords:

ABO blood Group system
ABO blood Types
COVID-19
SARS-CoV-2
ABO antibodies
Glycobiology

ABSTRACT

ABO blood group is long known to be an influencing factor for the susceptibility to infectious diseases, and many studies have been describing associations between ABO blood types and COVID-19 infection and severity, with conflicting findings. This narrative review aims to summarize the literature regarding associations between the ABO blood group and COVID-19. Blood type O is mostly associated with lower rates of SARS-CoV-2 infection, while blood type A is frequently described as a risk factor. Although results regarding the risk of severe outcomes are more variable, blood type A is the most associated with COVID-19 severity and mortality, while many studies describe O blood type as a protective factor for the disease progression. Furthermore, genetic associations with both the risk of infection and disease severity have been reported for the ABO locus. Some underlying mechanisms have been hypothesized to explain the reported associations, with incipient experimental data. Three major hypotheses emerge: SARS-CoV-2 could carry ABO(H)-like structures in its envelope glycoproteins and would be asymmetrically transmitted due to a protective effect of the ABO antibodies, ABH antigens could facilitate SARS-CoV-2 interaction with the host' cells, and the association of non-O blood types with higher risks of thromboembolic events could confer COVID-19 patients with blood type O a lower risk of severe outcomes. The hypothesized mechanisms would affect distinct aspects of the COVID-19 natural history, with distinct potential implications to the disease transmission and its management.

1. Introduction

The Severe Acute Respiratory Syndrome-Coronavirus 2 (SARS-CoV-2), the causing agent of COVID-19, emerged in late 2019 and rapidly spread worldwide, becoming a global health threat that, in one year, accounted for about 70 million cumulative cases [1]. Its high infectivity [2], with the occurrence of asymptomatic transmission [3], may have contributed to such a scenario. Mainly characterized by symptoms like fever, headache, cough, fatigue, breathing difficulties, anosmia, and ageusia; COVID-19 can either remain asymptomatic or develop to severe and critical manifestations, like dyspnea, hypoxia, acute respiratory distress syndrome, and multiorgan failure [4]. Age and co-morbidities, like diabetes and cardiovascular diseases, increase the risk of severe outcomes [5]. However, given the poorly understood pathogenesis of

COVID-19 and the variability in clinical presentation and outcomes, there has been a search for other biological markers that could predict the susceptibility to SARS-CoV-2 infection and the risk of severe outcomes [4,6].

The ABO blood group, which comprises distinct carbohydrate epitopes at the surface of blood cells and other tissues, like the respiratory and gastrointestinal epithelium [7], has been considered as an influencing factor for the susceptibility to infectious diseases. For instance, associations between the ABO blood type and the risk or severity of infections have been described for the Human Immunodeficiency Virus (HIV) [8], hepatitis B [8,9], West Nile virus (WNV) [10], and *Helicobacter pylori* [11,12]. Blood group antigens can influence infections directly, acting as receptors or co-receptors for microorganisms and toxins; or indirectly, through the anti-blood group antibodies, which can

* Corresponding author.

E-mail address: vania.ceccatto@uece.br (V. Ceccatto).

<https://doi.org/10.1016/j.micpath.2022.105658>

Received 29 March 2022; Received in revised form 22 June 2022; Accepted 22 June 2022

Available online 25 June 2022

0882-4010/© 2022 Elsevier Ltd. All rights reserved.




Fig. 2. - Structural representations of the ABO blood group antigens. (1.5-column fitting image).

A and O blood type individuals produce anti-B, B and O individuals produce anti-A, and AB individuals are devoid of both anti-A and anti-B [31]. The origin of ABO antibodies is still controversial, however, their synthesis seems to be mostly triggered by either microbiota or pathogenic microorganisms bearing ABO(H)-like antigens [32,33]. The ABO antibodies are able to induce complement-mediated lysis of RBCs and are mainly of IgM class [31].

3. ABO blood group and link to infections

ABO and FUT2 are clearly polymorphic genes and their products are highly present on epithelial cells in contact with the external environment. They are maximally expressed in mucins, which are thought to have a protective role in binding to microorganisms and are found in the body's exposed surfaces, like in the lungs and the gastrointestinal tract. Such features are suggestive of a possible function for ABO antigens in the interaction with environmental pathogens [7,34]. In addition, phylogeny of the ABO determinants showed that their firstly appeared as tissular antigens [35].

The widespread distribution of carbohydrates on cells' surfaces and their structural diversity place them as informational molecules able to act in cell recognition [36]. Indeed, pathogenic microorganisms and toxins can require binding to specific carbohydrate antigens to trigger infection and disease [37]. This was observed for *H. pylori*, whose infection was shown to depend on the expression of the Lewis b (Le^b) antigen in the gastric mucosa [38], and for the uropathogenic *E. coli* R45 strain, which binds to galactosyl-globoside - a precursor of the type 1 chain exposed at the membrane of non-secretor uroepithelial cells - and its sialylated derivate [39]. Furthermore carbohydrate antigens akin to that found on the surface of hosts' cells can be synthesized by bacteria in a co-evolving process suggestive of molecular mimicry, in which pathogenic or symbiotic microorganisms mimic carbohydrate motifs found at the host species to evade their innate immune system [37]. Indeed, Georg et al. screened 282 gram-negative strains isolated from humans, of which 48% showed ABO blood group activity [40].

Likewise, enveloped viruses can also acquire carbohydrate antigens

from their previous host as part of their structures and be asymmetrically transmitted due to a protective effect of the naturally occurring antibodies [41]. During their synthesis, the viral proteins are glycosylated in the Golgi apparatus and the endoplasmic reticulum of the host cells by their glycosylation enzymes [42]. Hence, viral glycoproteins can bear the glycosylation pattern of the cell in which they were synthesized. This was experimentally demonstrated for the Human T-cell Lymphotropic Virus (HTLV) iii_b, which was neutralized by monoclonal anti-A antibodies after propagation in peripheral blood mononuclear cells (PBMC) from A-type individuals, but not after propagation in PBMC of B or O individuals [42]. A similar effect was observed for the Measles virus produced in cells transfected with cDNA encoding human A, B, or null "O" transferases. The viruses produced in A- or B-type cells were partially neutralized, in a complement-dependent manner, by human pre-immune serum containing anti-A or anti-B antibodies, respectively, while the viruses produced in type O cells were not neutralized [43]. Moreover, anti-A monoclonal antibodies and type O plasma inhibited the adhesion of transfected CHO cells co-expressing the A antigen and the Spike (S) protein of SARS-CoV to ACE2-expressing Vero cells [44].

The link between ABO polymorphism and infectious disease has been extensively studied and is detailed reviewed elsewhere [13]. Associations have been reported for instance, with HIV [8], hepatitis B [9], West Nile Virus [10], and malaria [45]. Thus, the ABO polymorphism has been considered as an adaptive trait that could narrow the spread of infections within a given species. In turn, pathogenic microorganisms have been considered as a selective pressure generating diversity in the distribution of blood types among different populations, according to the exposition to distinct pathogens [37,46]. It is worth mentioning that associations between the ABO blood type and infectious diseases are most reliant on statistics, with few experimental data, and that illness may be multifactorial in nature, with the blood type constituting one of the multiple predisposing factors [34].

4. Associations between ABO blood types and COVID-19

4.1. Susceptibility to SARS-CoV-2 infection

Soon after the onset of the COVID-19 pandemic, an association between the ABO blood group and SARS-CoV-2 infection was firstly reported by Zhao et al., in which blood type A was significantly related to higher risk of COVID-19 infection in comparison to non-A blood types; while blood type O was significantly related to lower risk of COVID-19 infection compared to non-O blood types [14]. Subsequently, several studies have been reporting significant associations between blood type A and higher susceptibility to SARS-CoV-2 infection and/or blood type O and lower susceptibility [15,47–58]. A summary of the findings is described in Table 1.

As can be seen from Table 1, blood type A is frequently associated with higher risks of acquiring COVID-19, while blood type O is mostly described as a protective factor. Nonetheless, there are several disagreements, especially regarding the higher susceptibility. For instance, Muñiz-Diaz et al. evaluated the distribution of ABO blood types in two cohorts – convalescent plasma donors, with regular donors as the control

group, and hospitalized COVID-19 patients, with general patients as the control group -. In the former, blood type A was significantly over-represented compared to non-A blood types, while blood type O was significantly underrepresented compared to non-O types. However, no significant association was found in the later cohort [18].

Indeed, in a multiethnic cohort enrolling hospitalized COVID-19 patients - with general patients as the control group - no significant difference in ABO phenotype distribution was found [19]. In this study, the authors challenged the use of regular blood donors as the control group in studies concerning associations between ABO blood types and COVID-19 susceptibility. Blood type O individuals are preferentially recruited for blood donation and might be overrepresented among regular donors, which may have biased some previous results towards O-related protection. However, with a similar cohort choice, Wu et al. found significant associations between ABO blood group and the risk of SARS-CoV-2 infection, in which blood type A was associated with higher susceptibility to SARS-CoV-2 infection, and blood type O, to lower susceptibility [16]. Furthermore, studies using blood bank data do not always associate blood type O to lower risk of SARS-CoV-2 infection [59–62].

Table 1

Studies on associations between ABO blood types and the risk of SARS-CoV-2 infection.

Reference	Country	Study Design ^b	Cases		Controls		ABO Risk ^a			
			Definition ^c	N	Definition ^d	N	A	B	AB	O
Zhao et al. [14]	China	I	I	2.173	I	27.080	↑	ns	ns	↓
Li et al. [15]	China	II	II	265	I	3.694	↑	ns	ns	↓
Muñiz-Diaz et al. [18]	Spain	II	III	854	II	75.870	↑	ns	ns	↓
Franchini et al. [53]	Italy	III	III	447	II	16.911	ns	ns	ns	↓
Dzik et al. [19]	USA	II	II	957	III	5.840	ns	ns	ns	ns
Wu et al. [16]	China	II	II	187	III	1.991	↑	ns	ns	↓
Padhi et al. [69]	India	IV	IV	–	I	–	ns	↑	ns	ns
Khalil; Feghali; Hassoun [20]	Lebanon	II	II	146	III	6.497	ns	ns	ns	ns
Liu et al. [74]	International	I	I	9.383	IV	54.218	↑	↑	ns	↓
Fan et al. [52]	China	II	II	105	III	103	↑	ns	ns	ns
Solmaz; Araç [54]	Turkey	V	II	1.667	I	127.091	↑	ns	ns	↓
Barnkob et al. [47]	Denmark	II	I	7.422	I	2.204.742	↑	ns	↑	↓
Boudin et al. [76]	France	II	I	1.279	IV	409	ns	ns	ns	ns
Latz et al. [68]	US	II	I	1.289	V	7.648	ns	↑	↑	↓
Levi et al. [66]	Brazil	II	I	2.037	VI	1.813.237	ns	ns	ns	ns
Yalaoui et al. [55]	Tunisia	III	I	51	II or IV	63.375 or 1.506	↑	ns	ns	ns
Göker et al. [56]	Turkey	III	V	186	II	1.881	↑	ns	ns	↓
Almadhi et al. [59]	Bahrain	V	I	2.334	II	4.985	ns	↑	↓	ns
Lehrer; Rheinstein [65]	UK	II	I	717	I	11.855	ns	ns	ns	ns
Mullins et al. [67]	US	II	II	227	I	–	ns	ns	ns	ns
Negro et al. [78]	Italy	II	VI	129	VII	38	↑	↓	ns	↓
Ray et al. [70]	Canada	II	I	7.071	IV	218.485	ns	↑	↑	↓
El-Shitany et al. [51]	Saudi Arabia/Egypt	V	I	726	IV	707	ns	ns	ns	↓
Kibler et al. [77]	France	II	VII	22	VIII	680	↑	ns	ns	Ref
Leaf et al. [63], ^e	US	II	VIII	2.033	II	3.100.000	↑	ns	ns	↓
Niles et al. [64], ^e	US	II	I	34.178	IV	242.358	ns	ns	ns	↓
Al-Youha et al. [71]	Kuwait	II	I	3.304	I	3.730.027	ns	↑	↑	↓
Muñoz-Culla et al. [57]	Spain	II	VI	412	II	17.796	↑	↑	↑	↓
Kotila et al. [72]	Nigeria	II	I	302	II	9.138	ns	↑	↑	↓
Mahmud et al. [58]	Bangladesh	II	V	438	I	–	↑	ns	ns	ns
Schetelig et al. [48]	Germany	V	I	6.257	IV	118.868	↑	ns	↑	Ref
Ozkarafakili; Gareayaghi & Kara [60]	Turkey	II	II	492	II	51.966	ns	ns	ns	ns
Gamboa-Aguilar et al. [61]	Mexico	III	V	2.416	II	5.000	↓	ns	↓	↑
Enguita-Gérman et al. [49]	Spain	II	IX	87.090	–	–	↑	ns	ns	↓
Jawdat et al. [73]	Saudi Arabia	II	I	373	I	9.939	ns	↑	↑	↓
Kander et al. [75]	Sweden	II	X	338	IX	29.174	ns	ns	↑	ns
Gurung et al. [62]	Nepal	III	I	1091	II	2.182	ns	ns	↑	ns
Jukić et al. [50]	Croatia	II	VIII	780	II, V and X	614.673	↑	ns	ns	↓

^a vs. other blood types. Upward arrow: increased risk, downward arrow: decreased risk; ns: non-significant = $p > 0.05$.

^b I: Meta-analysis, II: Cohort study, III: Case-control study, IV: Mixed method, V: Cross-sectional study.

^c I: SARS-CoV-2+ individuals (Mostly RT-PCR), II: COVID-19 hospitalized patients, III: Convalescent plasma-donors, IV: Officially reported cases, V: General COVID-19 patients, VI: Symptomatic cases, VII: COVID-19 patients underwent TAVR (Transaortic valvular replacement), VIII: Critically ill patients, IX: General population, X: COVID-19 patients admitted to ICU (Intensive care unit).

^d I: Local anthropological data, II: Blood donors, III: Non-COVID-19 hospitalized patients, IV: SARS-CoV-2- individuals, V: Individuals tested for SARS-CoV-2, VI: Patients in general, VII: Asymptomatic COVID-19 cases, VIII: Patients underwent TAVR, IX: Non-COVID-19 patients admitted to ICU, X: Pregnant women with ABO typing.

^e Adjusted for ethnicity.

Other studies enrolling multiethnic cohorts in the U.S reported no significant association between ABO blood group and the risk of acquiring COVID-19 [63] or significantly higher odds for blood type O [64]. Nonetheless, after adjusting for ethnicity, blood type A was significantly associated with a higher risk of SARS-CoV-2 infection [63], and blood type O, with lower risk [63,64], among white non-Hispanic participants, but not in the Hispanic and black non-Hispanic groups. Indeed, a more recent study conducted in Mexico described a significantly higher risk of SARS-CoV-2 infection among blood type O individuals, while blood type A was significantly related to lower risk [61]. Ethnicity constitutes, therefore, a potential confounding factor in studies concerning ABO blood types and the risk of SARS-CoV-2 infection and should be considered in the interpretation of their results.

While several studies failed to reproduce significant associations between ABO blood-type and SARS-CoV-2 infection [20,60,65–67], others reached significant associations for blood types other than A and O. Blood type B was related to significantly higher risk of SARS-CoV-2 infection in some studies [59,68–73]; and a meta-analysis indicated that blood type B is associated with higher risk of SARS-CoV-2 infection in Asian, but not in European cohorts [74]. AB blood type, in turn, has been significantly related to either higher [47,62,68,70–73,75] or lower [59,61] risk of acquiring COVID-19.

Moreover, some authors studied the association between ABO blood types and the risk of SARS-CoV-2 infection in more specific backgrounds. Boudin et al. found no significant difference in ABO phenotype distribution between infected and uninfected crewmembers exposed to

Table 2

- Studies on associations between ABO blood types and the risk of severe COVID-19 outcomes.

Reference	Country	Study Design ^b	Cases		Type of Association	ABO Risk ^a			
			Definition ^c	N		A	B	AB	O
Muñiz-Diaz et al. [18]	Spain	I	I	965	Mortality	↑	ns	ns	↓
Dzik et al. [19]	USA	I	II	957	Mortality	ns	ns	ns	ns
Padhi et al. [69]	India	II	III	–	Mortality	ns	↑	ns	↓
Khalil; Feghali; Hassoun [20]	Lebanon	I	IV	146	Severity and mortality	ns	ns	ns	ns
Solmaz; Araç [54]	Turkey	III	II	1.667	Admission to ICU ^d and death	ns	ns	ns	ns
Barnkob et al. [47]	Denmark	I	V	7.422	Hospitalization and death	ns	ns	ns	ns
Latz et al. [68]	US	I	IV	1289	Hospitalization, admission to ICU, intubation, and death	ns	ns	ns	ns
Almadhi et al. [59]	Bahrain	III	IV	2.334	Admission to ICU	ns	ns	ns	ns
Lehrer; Rheinstein [65]	UK	I	V	717	Mortality	ns	ns	ns	ns
Mullins et al. [67]	US	I	II	227	Severity, admission to ICU, intubation, dialysis, hospital or ICU LOS ^e , and death	ns	ns	ns	ns
Negro et al. [78]	Italy	I	V	167	Severity and mortality	ns	ns	ns	ns
Ray et al. [70]	Canada	I	V and VI	218.485	Severity and mortality	ns	ns	ns	↓
El-Shitany et al. [51]	Saudi Arabia/ Egypt	III	IV	726	Oxygen saturation, artificial respiration, myalgia, and recovery time	↓	↓	ns	↑
Garibaldi et al. [85]	Brazil	I	VII	72	Severity and risk of death (SOFA ^f and SAPS-3 ^g)	↑	ns	ns	↓
Hoiland et al. [86]	Canada	I	VIII	95	Mechanical ventilation, CRRT ^h , and ICU LOS	↑	↓	↑	↓
Kibler et al. [77]	France	I	IX	22	Hospitalization and death	↑	ns	ns	ns
Sardu et al. [87]	Italy	I	X	164	Cardiac injury and death	ns	ns	ns	↓
Solhpour et al. [17]	Iran	I	XI	93	Severity	↑	ns	ns	↓
Valenti et al. [80]	Italy	I	VII	505	Mechanical ventilation and admission to ICU	ns	ns	ns	ref
Göker et al. [56]	Turkey	IV	IV	186	Admission to ICU, intubation, and death	ns	ns	ns	ns
Al-Youha et al. [71]	Kuwait	I	V	3.304	Respiratory support, admission to ICU, and death	ns	ns	ns	ns
Badedi et al. [79]	Saudi Arabia	I	V	404	Symptom presentation, hospital admission, disease duration and severity	ns	ns	ns	ns
Muñoz-Culla et al. [57]	Spain	I	VII, XII, XIII	213	Severity	ns	ns	ns	ns
Garg et al. [84]	India	I	IV	383	Severity	↑	↑	↓	↓
Szymanski et al. [88]	US	I	IV	4.968	Mortality	↑	ns	ns	ref
Ozkarafakili, Gareayaghi & Kara [60]	Turkey	I	II	492	Admission to ICU and death	ns	ns	ns	ns
Gamboa-Aguilar et al. [61]	Mexico	IV	IV	2.416	Hospitalization and death	ns	ns	ns	ns
Enguita-Gérman [49]	Spain	I	V	6.771	Hospitalization and death	ns	ns	ns	ns
Jawadat et al. [73]	Saudi Arabia	I	V	373	Hospitalization and admission to ICU	ns	ns	ns	ns
Nafakhi et al. [89]	Iraq	V	VII and XIII	169	Respiratory support, admission to ICU, death, and long-term symptoms	ns	↑	↑	ref
Gursoy & Avcı [81]	Turkey	I	II	969	Hospital LOS, admission to ICU, ICU LOS, and death	ns	ns	ns	ns
Bokhary et al. [82]	Saudi Arabia	I	II	316	Admission to ICU and death	ns	ns	ns	ns
Jelinek et al. [83]	United Arab Emirates	III	VII	193	Severity	ns	ns	ns	↓
Alabsi et al. [90]	Saudi Arabia	III	XIV	100	Chronic chemosensitive dysfunction	–	–	–	↑

^a Risk of severity and death vs. other blood types. Upward arrow: increased risk, downward arrow: decreased risk, ns: non-significant = $p > 0.05$.

^b I: Cohort study, II: Mixed method, III: Cross-sectional study, IV: Case-control study, V: Longitudinal observational study.

^c I: COVID-19 patients transfused during hospitalization, II: Hospitalized COVID-19 patients, III: Officially reported death rates, IV: General COVID-19 patients, V: SARS-CoV-2-positive individuals (Mostly RT-PCR), VI: SARS-CoV-2-negative individuals, VII: Severe COVID-19 patients, VIII: Patients admitted to intensive care unit (ICU), IX: COVID-19 patients underwent TAVIR, X: Hypertensive COVID-19 patients, XI: COVID-19 patients with severe hypoxia, XII: light COVID-19, XIII: Mild COVID-19, XIV: Patients recovered from COVID-19.

^d Intensive Care Unit.

^e Length of Stay.

^f Sequential Organ Failure Assessment.

^g Simplified Acute Physiology Score 3.

^h Continuous Renal Replacement Therapy.

SARS-CoV-2 during an outbreak in an aircraft carrier [76]. Conversely, blood type A was significantly associated with higher risk of developing COVID-19 among patients who underwent transcatheter aortic valve replacement (TAVR) [77]. Furthermore, after no association between ABO phenotype and SARS-CoV-2 positivity was found, Negro et al. sought to investigate whether ABO blood type correlates with the symptomatic disease in SARS-CoV-2-positive individuals. A significant positive association was found between blood type A and the development of symptomatic COVID-19, while blood types B and O were significantly related to lower risks of developing symptomatic infection [78].

Although studies concerning ABO blood types and the risk of SARS-CoV-2 infection are highly heterogeneous, we were unable to find any trend in susceptibility according to the study design or the type and size of the case and control groups. However, among the studies summarized here, a dichotomy between blood types A – higher risk – and O – lower risk – was more clearly observed in European and East Asian settings, while the involvement of blood types B or AB has been mostly observed in Middle Eastern, Indian, and North American settings (Table 1). Data from Africa and Latin America are scarce, but it is noteworthy that in both the settings, blood type A was not associated yet with higher susceptibility to SARS-CoV-2 infection [61,66,72] and, in the case of Latin America, blood type O was not associated yet with lower susceptibility [61,66].

4.2. COVID-19 severity and outcomes

In addition to the risk of SARS-CoV-2 infection or symptomatic disease, the ABO blood group has also been associated with severe outcomes. Severity is usually defined as hospitalization, admission to Intensive Care Unit (ICU), mechanical ventilation requirement, or death. The association of ABO blood types with COVID-19 severity is generally performed by comparing the blood type distribution between distinct subgroups of patients. Studies on this subject were mostly designed to uncover associations for both the risk of infection and severity, with fewer studies designed to evaluate associations for severity only. A summary of the findings is described in Table 2.

As can be seen from Table 2, results regarding severity and outcomes are more variable than those regarding the risk of infection. Several studies that reported associations between ABO blood type and the risk of SARS-CoV-2 infection failed to describe any association for severity [47,49,54,56,57,59,61,68,71,73,78] or reached distinct associations [51,70]. Other studies also reported no association between ABO blood types and severe COVID-19 outcomes but in accordance with the finds reported for the risk of infection [19,20,60,65,67]. Further studies also failed to report any association between the ABO blood group and severe COVID-19 outcomes [79–82].

Nonetheless, blood type A has been the most associated with severe outcomes, while blood type O frequently appears as a protective factor against the disease progression [17,18,69,70,77,83–88]. For instance, although failing to describe a significant difference in ABO phenotype distribution between hospitalized COVID-19 patients and the control group, Muñoz-Díaz et al. described a significant positive association between blood type A and mortality, and a significant negative association for blood type O [18]. Blood type A was significantly associated with hospitalization and mortality among COVID-19 patients who underwent TAVR [77] and to severe hypoxia among young COVID-19 patients without co-morbidities [17]. Blood type O, in turn, was significantly associated with lower risks of cardiac injury and death, and with lower pro-thrombotic indexes, among hypertensive COVID-19 patients [87].

Enrolling COVID-19 patients admitted to ICU, Hoiland et al. described significantly higher odds of requiring mechanical ventilation and Continuous Renal Replacement Therapy (CRRT), as well as a higher ICU length of stay (LOS), for A and AB groups; while B and O groups were negatively associated with these outcomes [86]. Conversely,

El-Shitany et al. reported an association between blood type O and unfavorable outcomes, such as lower oxygen saturation and mechanical ventilation requirement, while blood type A was linked to lower odds of requiring artificial ventilation, and blood type B, to shorter recovery periods [51]. Significant associations with severity [84] and mortality [69,89] were also reported for blood type B, while blood type AB has been linked to either favorable [84], or unfavorable [89] outcomes. More recently, some studies sought to investigate whether ABO blood types are associated with long-term symptoms presentation after COVID-19 recovery. Blood type AB was significantly associated with long-term palpitation and dizziness [89], and blood type O, with subjective loss of smell [90].

Studies regarding associations between ABO blood types and COVID-19 severity and outcomes are even more heterogeneous than those concerning the risk of infection. The studies evaluated distinct outcomes and some of them had a unique design, which makes it more difficult to establish trends. Among the studies summarized here (Table 2), the majority did not reach statistical significance, and we were unable to identify any trend in the outcomes according to the size of the enrolled groups. However, it is possible to observe a trend towards significance when the enrolled group consisted of critically ill COVID-19 patients – like those requiring ICU admission [86] or blood transfusion [18] and those presenting severe hypoxia [17] - or patients with cardiovascular co-morbidities [77,87]. Taken together, in these groups of patients, blood type A was the most associated with unfavorable outcomes and blood type O constituted a protective factor. Conversely, in broader enrolled groups, like general hospitalized COVID-19 patients or individuals with a positive test, there was a trend towards non-significant results (Table 2).

4.3. Genetic link with susceptibility and severity

Several studies have been designed to uncover single nucleotide polymorphisms (SNPs) that could predict COVID-19 susceptibility or severity, and associations have been reported for the ABO locus. In a meta-analysis of two Genome-Wide Association Studies (GWAS) enrolling COVID-19 patients with respiratory failure, Ellinghaus et al. analyzed 8,582,968 SNPs and described a cross-replicating association with the Reference SNP (rs)657152 at locus 9q34, which corresponds to the ABO gene [6]. Other GWAS also reported associations for the ABO gene. An association with rs9411378 at chromosome 9q34.2 was identified by Shelton et al. in contrasting SARS-CoV-2-positive and –negative individuals, while no association with this SNP was identified when comparing severity phenotypes [91]. In this study, blood type O was shown to be protective regarding susceptibility, while blood types A, B, and AB did not differ from each other. An association with rs91280525 for susceptibility was replicated in a meta-analysis of previous GWAS [92]. Nonetheless, a GWAS enrolling critically ill COVID-19 patients failed to identify any association with the ABO locus [93].

More recently, Vargas-Alarcón et al. evaluated the allele distribution of six SNPs located at the ABO gene among COVID-19 patients and healthy hospital staff members. Two SNPs located at the coding region - rs8176740 and rs512770 – were found to be associated with the risk of developing COVID-19, with the risk alleles being related to O blood type [94]. The rs8176740 polymorphism was also associated with high platelet blood levels [94]. In contrast, after finding significantly lower levels of blood type O among critical COVID-19 patients (vs. non-critical patients), Jelinek et al. performed a candidate gene approach based on previous GWAS data and examined 240 SNPs in the ABO gene for association with critical COVID-19 presentation. 8 SNPs located at the ABO 3' untranslated region were significantly associated with disease severity, with significantly lower frequencies of the risk alleles in non-critical patients with blood type O [83].

Furthermore, some studies sought to genotype the ABO and/or FUT2 genes in COVID-19 patients and control groups. AB-related alleles were significantly associated with SARS-CoV-2 infection and O-related alleles

were significantly less represented among the patients' group, while no association was described for FUT2 [95]. Conversely, a study found no significant difference in the distribution of the rs8176719 deletion allele – the main O blood type-determinant – between patients and controls nor between severe and non-severe patients [96], while the rs601338 G>A variant – the main non-secretor determinant – was associated with lower risks of ICU admission, mechanical ventilation and decease, especially in blood type A-individuals [80]. Another study described associations of the A/B-related rs8176747 with higher susceptibility to COVID-19 infection and the O-related rs8176719 with lower susceptibility, failing to describe any association with the FUT2-related rs1047781. Nonetheless, when comparing ABO secretor individuals vs. O and non-secretors, significant results were obtained, with the former group being more prone to COVID-19 onset [97].

Data from previous GWAS have been integrated to protein (p) or expression (e) quantitative trait loci (QTL) and colocalization analysis in Summary-Based Mendelian Randomization studies. In a multi-omics-based study, a protein-increasing allele of the ABO gene (rs505922) co-localized with susceptibility and severity-associated loci in both lungs and blood; and increased levels of blood ABO proteins was associated with higher risk of COVID-19 and its severity [98]. Likewise, in a Mendelian Randomization study based on previous GWAS, the blood level of ABO proteins was associated with hospitalization, respiratory support requirement, and decease [99]. Further QTL-based studies have also been reporting associations between the expression of ABO proteins (rs505922) and COVID-19 severity phenotypes [100,101]; however, pQTL for rs505922 was also associated with the levels of several other proteins [100,101], which suggests potential pleiotropic effects. Interestingly, among the proteins associated with rs505922, CD209 antigen

was recently described as a receptor for SARS-CoV-2 S protein in lung and kidney epithelium and endothelial tissue [102]. Moreover, genetically predicted ABO levels were significantly associated with COVID-19 clinical phenotypes driven by the circulatory system, such as deep vein thrombosis, pulmonary heart disease, and acute pulmonary infarction [103]. Genetic associations between the ABO locus and COVID-19 are summarized in Table 3.

5. Hypothesis linking ABO blood type and COVID-19: underlying mechanisms and their contribution to the interpretation of reported associations

Since an association between the ABO blood group and COVID-19 was firstly reported, several underlying mechanisms have been hypothesized [22,23]. The proposed hypotheses are most reliant on what is already known for other enveloped viruses, including SARS-CoV [21, 44], or in the previous associations between ABO blood group and other COVID-19 risk factors and co-morbidities [23,104]. The hypothesized mechanisms could affect either the risk of infection or the disease severity and outcomes, with distinct potential implications to the transmission dynamics or disease management.

5.1. ABO antibodies and host's glycosylation pattern

Upon entry in the human body, SARS-CoV-2 can infect and replicate in the upper respiratory tract [105,106], several cell types of the bronchial epithelium, and the alveolar type 1 pneumocytes [107], as well as in the enterocytes [108]. Epithelial cells of both the human airways and digestive tract are known to express host factors used by SARS-CoV-2,

Table 3
- Genetic associations of the ABO locus with COVID-19 infection and severity.

Reference	Country	Study Design	Sample size (n)	SNP	Outcomes
Ellinghaus et al. [6]	Italy/Spain	COVID-19 with RF ^a vs. blood donors, and non-COVID-19 controls	1.610 cases vs.2.205 controls	rs657152	Severe COVID-19 with respiratory failure
Pairo-Castineira et al. [93]	UK	Critically ill COVID-19 vs. matched controls	2.244 cases vs.1.157.277 controls	–	None
Cordero et al. [98]	International	Integrative genomics	–	rs505922	Susceptibility and severity
Valenti et al. [80]	Italy	Non-O vs. O COVID-19 with RF	Non-O 256 vs.O 153	rs601338 (FUT2)	Mechanical ventilation, ICU ^a admission, and mortality
Matzhold et al. [95]	Austria	Hospitalized COVID-19 patients vs. blood donors	336 patients vs. 250.298 controls	Several SNPs corresponding to variants of ABO alleles, FUT-2 rs601338, FUT3	Susceptibility to infection, hospitalization, ICU admission, mechanical ventilation, and mortality
Coto et al. [96]	Spain	Hospitalized COVID-19 patients vs. matched healthy controls	318 patients vs. 350 controls	rs8176719	Susceptibility and severity
Zhou et al. [100]	International	Hospitalized COVID-19 patients vs. healthy controls.	14.134 cases vs.1.2 million controls	rs505922	Severity
Zhu et al. [101]	International	Hospitalized COVID-19 patients vs. healthy controls.	6.492 patients vs.1.012.809 controls	rs505922	Susceptibility, hospitalization and severity.
Pathak et al. [103]	International	COVID-19 hospitalized patients vs.controls	7.885 cases vs. 961.804 controls	–	Severity
CHGI [92]	International (multiethnic)	COVID-19 patients vs. matched controls	49.562 cases vs. 2.000.000 controls	rs91280525	Susceptibility and severity
Shelton et al. [91]	US/UK	SARS-CoV-2 positive vs. controls	12.972 cases vs. 101.268 controls	rs9411378	Susceptibility
Vargas-Alarcón et al. [94]	Mexico	COVID-19 patients vs. healthy hospital staff members	415 cases vs. 288 controls	rs8176740 and rs512770	Susceptibility and platelets concentration
Jelinek et al. [83]	United Arab Emirates	Critical vs. non-critical COVID-19 patients	193 cases vs. 453 controls	rs199969472, rs34266669, rs76700116, rs7849280, rs34039247, rs10901251, rs9411475, and rs13291798	Severity
Nishida et al. [97]	Japan	COVID-19 patients vs. healthy controls	461 cases vs. 1.193 controls	rs8176719 and rs8176747 (ABO) and rs1047781 (FUT2)	Susceptibility
Palmos et al. [99]	International	Mendelian Randomization	–	Several SNPs related to the ABO gene.	Severity

^a Intensive Care Unit.

like ACE2 and TMPRSS2 [109–111], in addition to A/B transferases in secretor individuals [7,112].

During the replication cycle of SARS-CoV-2, viral RNA is synthesized within double-membrane vesicles that bud from the host endoplasmic reticulum following extensive remodeling [113], and the structural proteins assemble to the viral genome at the reticulum-Golgi intermediate compartment (ERGIC) to form progeny virions, which are released from the cell surface by exocytosis [114]. The passage of virions through the secretory pathway raises questions concerning the processing of the viral surface glycoproteins [115], due to the presence of glycosylation enzymes at this pathway [116].

Indeed, the SARS-CoV-2 spike (S) protein is a highly glycosylated trimer, with 22 canonical N-linked glycosylation sites per protomer [117]. Glycomics-oriented studies revealed a high occupancy of N-linked sequons and the presence of O-linked glycosylation sites with low occupancy in recombinant S protein of SARS-CoV-2 produced in either mammalian [115,117–119] or insect [120] cell lines. In addition, mass spectrometry approaches showed that the glycosylation profile of the S protein is highly heterogeneous and cell type-dependent. In recombinant S protein produced in insect Sf9 cells, a predominance of less processed glycans, like oligomannose, was found [120]; while a more complex glycan processing, like fucosylation, galactosylation, and sialylation, was shown to take place in mammalian HEK 293 cells [115, 117,119]. Nonetheless, these glycans do not fully represent those found at the envelope glycoproteins of virions emitted from the human respiratory or digestive tract and cannot be decorated with ABH antigens, since HEK293 cells do not synthesize ABH transferases [25].

Aiming to uncover whether the S protein can be decorated with ABH antigens when synthesized in ABO- and FUT2-expressing cells, Deleers et al. produced recombinant S protein in transfected CHO cells expressing either the A or B transferase and the FUT2 gene. This approach showed that the S protein can carry ABH antigens according to the glycosyltransferase repertoire of the previous host cell [121]. Hence, ABO antibodies could constitute a barrier to SARS-CoV-2 infection according to the rules of blood transfusion. When transmitted in ABO-incompatible situations, SARS-CoV-2 could be partially neutralized by anti-A or anti-B antibodies, which can explain the lower incidence of COVID-19 in type O individuals, who display both anti-A and anti-B. Nonetheless, such protection would only be initial as, once the infection is established, the virions would carry the host's ABO profile and the autologous ABO antibodies would no longer be effective [122]. Therefore, this possible underlying mechanism can only explain associations between the ABO blood type and the risk of SARS-CoV-2 infection and does not apply to the reported associations concerning severity and outcomes.

The ABO antibodies hypothesis has some implications for the dynamics of COVID-19 transmission. Firstly, the reduced risk of viral infections in case of ABO-incompatible transmission does not constitute individual protection, but rather collective protection that could delay the spread of newly emerging viruses and slow down the epidemics, in addition to the required behavioral changes [44]. Nonetheless, as the epidemic progresses, more and more ABO-compatible transmission might occur, and the final number of cases may not be reduced to a great extent [25]. This could explain, for instance, the lack of association reported by Boudin et al. [76], in which the study participants consisted of crewmembers in close enough contact within an aircraft carrier that may have facilitated ABO-compatible transmission and reduced any O-related protection.

Moreover, the protection provided by ABO antibodies in ABO-incompatible transmission would critically depend on the relative frequency of ABO phenotypes in a given population. Such protection would be more effective in populations with low coefficients of variation between the ABO phenotypes and is expected to be inefficient in populations with high coefficients of variation, in which more ABO-compatible transmission may occur [25]. Importantly, although blood type O is likely more protective to SARS-CoV-2 infection due to the

presence of both anti-A and anti-B antibodies, this advantage seems to disappear in populations in which this blood type is highly predominant, like in some Central and South American countries [25].

Following this rationale, differences in ABO phenotypes distribution between distinct cohorts can explain some interstudy discrepancies. For instance, the frequency of the B phenotype is increased in Asia at the expense of both A and O phenotypes, which, together, are largely predominant in Western Europe, with the Middle East displaying an intermediary distribution [25]. Such difference could explain the association between blood type B and the increased risk of SARS-CoV-2 infection reported by some Indian or Middle Eastern studies [59,69, 71,73]. Likewise, the lack of association between ABO blood type and the risk of SARS-CoV-2 infection [19,63] or a significant positive association with blood type O [64] in U.S. multiethnic cohorts before adjusting for ethnicity can be explained by the fact that ethnic groups among whom blood type O is predominant, like Hispanics and black non-Hispanics, have been disproportionately affected by the COVID-19 pandemic in the U.S. [19,64]. In addition, the association between blood type O and the risk of COVID-19 onset described in a Mexican study can be explained by the high prevalence of this blood type in the Mexican population [61].

More recently, Boukhari et al. sought to investigate the impact of ABO compatibility on a secondary attack rate among the spouses of hospital staff members who were symptomatic for COVID-19. The risk of symptomatic COVID-19 transmission was significantly lower for ABO-incompatible couples, while ABO compatibility was associated with higher transmissibility regardless of the ABO blood type of the partner. This supports that all blood types are equally susceptible to COVID-19, and the protection provided by ABO-incompatibility may be due to ABO antibodies [123].

Furthermore, one should consider the variability of the ABO antibodies titer. The levels of ABO antibodies decrease with improving hygiene and tend to be low in developed countries [124], which may pose a challenge to its protective effect against viral infections and constitutes a potential confounding factor in association studies. Indeed, a study enrolling COVID-19 patients and asymptomatic controls showed that, for both A, B, and O blood types, the patients' group displayed significantly lower IgM agglutination scores compared to the control group [121]. In this way, some authors have been suggesting the promotion of anti-glycan antibodies through dedicated microbiota to enhance the natural anti-viral protection and slow down the spread of newly emergent viruses [121,125].

5.2. S protein priming and SARS-CoV-2-host cells interaction

The S protein of SARS-CoV-2 harbors a multibasic S1/S2 cleavage site, which constitutes a target for furin, a proprotein convertase. The furin-mediated S1/S2 cleavage generates two subunits – S1 and S2 – and is pivotal for the S protein's activation. S1 binds to cellular receptors, while S2 is further cleaved by TMPRSS2 and promotes virus-cell membrane fusion. The furin cleavage site was shown to be required for SARS-CoV-2 infection of the human lung cells Calu-3 and for the S protein-mediated cell-cell fusion, which promotes a systemic viral spread [126]. Although no correlation was yet reported for the furin levels and ABO polymorphism, an association between blood type O and lower levels of the furin-related proprotein convertase subtilisin-kexin type 9 (PCSK9) was previously reported [127]. Therefore, Abdelmassih et al. hypothesized that furin levels may follow the same pattern as PCSK9 and be decreased in blood type O individuals, which might reduce S protein priming and SARS-CoV-2 entry into host cells [128].

S1, in turn, is subdivided into the S1A and S1B domains. The former corresponds to the amino-terminal portion of the polypeptide chain, namely NTD (from N-terminal domain), and, in some coronaviruses, interacts with glycolipids and glycoproteins displaying terminal sialic acid motifs at the host cells' surface [129]. The latter constitutes the Receptor-Binding Domain (RBD) and binds to the ACE2 receptor [130].

In silico studies based on molecular docking simulations [131] and electronic density mapping surface [132] predict the existence of a sialic acid-binding pocket in the SARS-CoV-2 NTD similar to that found in MERS-CoV, which might confer to SARS-CoV-2 the ability to infect human cells in a dual manner, resulting in a higher diffusion speed [132].

ABH antigens are known to modulate the clustering of sialylated glycans at the cell's surface through carbohydrate-carbohydrate interactions and to regulate their interaction with cognate-binding proteins. A model was proposed in which antigen A is peripherally located at the sialylated glycan clusters, allowing for continuous clusters with increased sialic acids concentrations. Conversely, antigen B is mostly located at the center of the cluster, which may pose a steric hindrance for sialic acids-binding proteins, while in type O cells, the clusters are small and lack H antigens [133]. Following this rationale, it was hypothesized that ABO blood group may correlate with COVID-19 susceptibility and severity through carbohydrate-carbohydrate interactions between ABH antigens and sialylated glycans at the cells surface. According to such hypothesis, antigen A maximizes the interaction of SARS-CoV-2 with the target cells by promoting the clustering of sialylated glycans and increasing the likelihood of NTD binding [134].

Moreover, in a glycan microarray with ABH antigens, the SARS-CoV-2 RBD bound preferentially to the type I structure of antigen A, like the SARS-CoV RBD, which suggests a possible role for antigen A in coronavirus entry into respiratory or digestive tract cells [48]. A similar result was described by Wu et al., who also demonstrated sequence similarities between the SARS-CoV-2 RBD and galectins, a family of carbohydrate-binding proteins [135]. This could explain, for instance, the genetic association reported by Valenti et al. between non-secretor status and lower risk of severe outcomes among blood type A individuals [80]. Conversely, Boukhari et al. failed to detect any preferential binding of recombinant RBD to both A type I saccharide and lung tissue from A and O donors, as well as to salivary mucin from secretor-phenotype saliva [123]. Furthermore, although individuals with blood type A1 display more than fourfold higher A epitope density compared to those with A2 subtype, comparable rates of infection between A1 and A2 participants were reported by Schetelig et al., which suggests that the A antigen itself does not facilitate viral entry [48].

Hypotheses linking the ABO blood type to S protein priming or SARS-CoV-2 cellular entry remain largely elusive and need further studies. These hypotheses could explain both the higher SARS-CoV-2 infection rates and higher disease severity reported for blood type A, as well as the lower rates of infection and lower disease severity reported for blood type O. Unlike the ABO antibodies hypothesis, the possible role of ABO antigens in regulating virus-cell interaction may be mostly reliant on the antigen itself and may not depend on the ABO coefficient of variation at a population level. In addition, the possible pleiotropy effect of the ABO gene (rs505922) over the CD209 receptor gene [101] could also be considered into the framework of these hypotheses.

5.3. ABO blood group and cardiovascular risk

Associations between non-O blood types and increased risk of cardiovascular disorders are well documented and are detailed reviewed elsewhere [136–138]. Associations have been reported, for instance, with venous thromboembolism, myocardial infarction, peripheral vascular disease, cerebrovascular stroke, and pulmonary embolism. A possible explanation relies on the ~30% lower levels of von Willebrand factor (vWF) and coagulation factor VIII (FVIII) in blood type O individuals [139,140]. vWF mediates platelet aggregation and adhesion and can be released in response to inflammatory stimuli, like infections. It also plays a role in hemostasis by complexing with FVIII and stabilizing it at the circulation [141]. As a glycoprotein, vWF carries ABO (H)-like structures in Asp-linked sugar chains [142], which are likely to mediate its homeostasis. Indeed, glycosylation is likely to mediate the clearance of vWF, which is largely performed by lectin receptors [143,

144]. A and B antigens were shown to protect vWF from proteolysis by the ADAMTS13 metalloprotease [145]. Furthermore, the ABO blood group indirectly modulates the circulating levels of FVIII by affecting the clearance of the vWF/FVIII complex, accounting for about ~30% of its variation [140].

Likewise, some adhesion molecules, like the intercellular cell adhesion molecule 1 (ICAM1) and P-selectin, are protected by antigen A from enzymatic cleavage, prompting the attachment of leucocytes to them at the vascular wall, which increases inflammation and decreases circulation [23]. In addition, a study enrolling COVID-19 hypertensive patients reported significantly higher values of pro-thrombotic indexes - vWF, FVIII, D-dimer, and activated pro-thrombin time - in non-O vs O patients, which was consistent with the higher rates of cardiac injury and death in the former group [87]. Blood type O individuals are therefore less prone to thromboembolic events that constitute relevant outcomes in COVID-19 disease, especially among ICU-admitted patients [146, 147].

The link between ABO blood group and cardiovascular disorders does not affect the susceptibility to SARS-CoV-2 infection and cannot explain the reported associations between ABO blood type and the risk of infection. However, it constitutes a possible underlying mechanism for the lower risk of severe outcomes associated with blood type O and, at a lesser extent, the higher risk related to blood type A in some studies (Table 2), especially among patients with cardiovascular co-morbidities [77,87]. Into the framework of this hypothesis, the reported associations rely on constitutive tracts related to ABO blood type, which is in line with a study that reported associations between genetically-driven ABO expression levels and COVID-19 clinical phenotypes driven by the circulatory system [103]. Following this rationale, a risk stratification for cardiovascular complications in COVID-19 patients based on the ABO phenotype has been suggested, as well as the introduction of early anticoagulant therapy in non-O hypertensive patients [87].

6. Conclusions

A literature survey of epidemiological studies suggests that the ABO blood type may be an influencing factor for both SARS-CoV-2 infection and COVID-19 severity. Although the findings are conflicting and quite variable, blood type O is mostly associated with lower risk of SARS-CoV-2 infection, while blood type A, with higher risk. Blood types B and AB were also associated with higher risk of infection in some studies. Although the majority of the studies regarding COVID-19 severity and outcomes failed to report any association with the ABO blood group, associations with lower risk of severe outcomes have been frequently described for blood type O, while blood type A has been frequently described as a risk factor for severe outcomes. Moreover, genetic links with both susceptibility to infection and severity have been reported for the ABO locus.

Driven by the previous knowledge about ABO blood group biology and its role in the pathogenesis of infectious diseases, some underlying mechanisms that could explain the reported associations emerge; like a protective effect of the ABO antibodies against viral infections, a role for ABO(H) antigens in facilitating viral entry into host cells, and the association between non-O blood types and higher risk of thromboembolic events. These underlying mechanisms may affect different phases of the natural history of COVID-19, with distinct implications to the SARS-CoV-2 transmission dynamics and to COVID-19 outcomes. Nonetheless, there are little experimental data on this subject and more studies are needed to elucidate the mechanisms underlying the reported associations, which is pivotal for a deeper understanding of the relationship between the ABO blood group and COVID-19 and whether or not it can be translated to disease prevention or management strategies.

Funding

This research did not receive any specific grant from funding

agencies in the public, commercial, or non-profit sectors.

Ethics approval

Not applicable.

Consent to participate

Not applicable.

Consent to publish

Not applicable.

Code availability

Not applicable.

CRediT authorship contribution statement

Eric Pereira: Writing – original draft, Formal analysis, Data curation, Conceptualization. **Stela Felipe:** Writing – review & editing, Conceptualization. **Raquel de Freitas:** Writing – review & editing. **Valdevane Araújo:** Data curation, Conceptualization. **Paula Soares:** Data curation, Conceptualization. **Jannison Ribeiro:** Supervision, Data curation, Conceptualization. **Luiz Henrique dos Santos:** Conceptualization. **Juliana Osório Alves:** Writing – review & editing. **Natália Canabrava:** Conceptualization. **Mauricio van Tilburg:** Supervision, Conceptualization. **Maria Izabel Guedes:** Supervision, Conceptualization. **Vânia Ceccatto:** Writing – review & editing, Supervision, Data curation, Conceptualization.

Declaration of competing interest

The authors have no financial or non-financial competing interests to disclose.

Data availability

No data was used for the research described in the article.

References

- [1] World Health Organization, COVID-19 Weekly Epidemiological Update [Internet], 2020. Available from: <https://www.who.int/publications/m/item/weekly-epidemiological-update—15-december-2020>. (Accessed 18 December 2020).
- [2] B. Hu, H. Guo, P. Zhou, Z.L. Shi, Characteristics of SARS-CoV-2 and COVID-19, *Nat. Rev. Microbiol.* 19 (2021) 141–154, <https://doi.org/10.1038/s41579-020-00459-7>.
- [3] D. Han, R. Li, Y. Han, R. Zhang, J. Li, Covid-19: insight into the asymptomatic sars-cov-2 infection and transmission, *Int. J. Biol. Sci.* 16 (2020) 2803–2811, <https://doi.org/10.7150/ijbs.48991>.
- [4] T.P. Velavan, S.R. Pallerla, J. Rütter, Y. Augustin, P.G. Kreamsner, S. Krishna, et al., Host genetic factors determining COVID-19 susceptibility and severity, *EBioMedicine* (2021) 72, <https://doi.org/10.1016/j.ebiom.2021.103629>.
- [5] Z. Wu, J.M. McGoogan, Characteristics of and important lessons from the coronavirus disease 2019 (COVID-19) outbreak in China: summary of a report of 72314 cases from the Chinese center for disease control and prevention, *JAMA, J. Am. Med. Assoc.* 323 (2020) 1239–1242, <https://doi.org/10.1001/jama.2020.2648>.
- [6] D. Ellinghaus, F. Degenhardt, L. Bujanda, M. Buti, A. Albillos, P. Invernizzi, et al., Genome-wide association study of severe covid-19 with respiratory failure, *N. Engl. J. Med.* 383 (2020) 1522–1534, <https://doi.org/10.1056/nejmoa2020283>.
- [7] S. Marionneau, A. Cailleau-Thomas, J. Rocher, B. Le Moullac-Vaidye, N. Ruvoën, M. Clément, et al., ABH and Lewis histo-blood group antigens, a model for the meaning of oligosaccharide diversity in the face of a changing world, *Biochimie* 83 (2001) 565–573, [https://doi.org/10.1016/S0300-9084\(01\)01321-9](https://doi.org/10.1016/S0300-9084(01)01321-9).
- [8] Z. Batool, S.H. Durrani, S. Tariq, Association of abo and Rh blood group types to hepatitis B, hepatitis C, hiv and Syphilis infection, A five Year* experience in healthy blood donors in A tertiary care hospital, *J. Ayub Med. Coll. Abbottabad* 29 (2017) 90–92.
- [9] W. Jing, S. Zhao, J. Liu, M. Liu, ABO blood groups and hepatitis B virus infection: a systematic review and meta-analysis, *BMJ Open* 10 (2020), <https://doi.org/10.1136/bmjopen-2019-034114>.
- [10] Z. Kaidarova, M.D. Bravo, H.T. Kamel, B.S. Custer, M.P. Busch, M.C. Lanteri, Blood group A and D negativity are associated with symptomatic West Nile virus infection, *Transfusion* 56 (2016) 1699–1706, <https://doi.org/10.1111/trf.13622>.
- [11] L.C. De Mattos, J.R. Cintra, F.E. Sanches, R.D.C.M. Alves Da Silva, M.A. Ruiz, H. W. Moreira, ABO, Lewis, secretor and non-secretor phenotypes in patients infected or uninfected by the *Helicobacter pylori* bacillus, *Sao Paulo Med. J.* 120 (2002) 55–58, <https://doi.org/10.1590/s1516-31802002000200006>.
- [12] L.C. Martins, T.C. de Oliveira Corvelo, H.T. Oti, R. do Socorro Pompeu Loiola, D. C.F. Aguiar, K.A. dos Santos Barile, et al., ABH and Lewis antigen distributions in blood, saliva and gastric and H pylori infection in gastric ulcer patients, *World J. Gastroenterol.* 12 (2006) 1120–1124, <https://doi.org/10.3748/wjg.v12.i7.1120>.
- [13] L. Cooling, Blood groups in infection and host susceptibility, *Clin. Microbiol. Rev.* 28 (2015) 801–870, <https://doi.org/10.1128/CMR.00109-14>.
- [14] J. Zhao, Y. Yang, H. Huang, D. Li, D. Gu, X. Lu, et al., Relationship between the ABO blood group and the COVID-19 susceptibility, *Clin. Infect. Dis.* 73 (2021) 328–331, <https://doi.org/10.1093/cid/ciaa1150>.
- [15] J. Li, X. Wang, J. Chen, Y. Cai, A. Deng, M. Yang, Association between ABO blood groups and risk of SARS-CoV-2 pneumonia, *Br. J. Haematol.* 190 (2020) 24–27, <https://doi.org/10.1111/bjh.16797>.
- [16] Y. Wu, Z. Feng, P. Li, Q. Yu, Relationship between ABO blood group distribution and clinical characteristics in patients with COVID-19, *Clin Chim Acta. Netherlands* 509 (2020) 220–223, <https://doi.org/10.1016/j.cca.2020.06.026>.
- [17] A. Solhpour, A. Jafari, M. Pourhoseingholi, F. Soltani, Corona Covid-19 virus and severe hypoxia in young patients without underlying disease: high prevalence rate with blood group A, *Trends Anaesth Crit Care* 34 (2020) 63–64, <https://doi.org/10.1016/j.tacc.2020.08.005>.
- [18] E. Muñiz-Diaz, J. Llopis, R. Parra, I. Roig, G. Ferrer, J. Grifols, et al., Relationship between the ABO blood group and COVID-19 susceptibility, severity and mortality in two cohorts of patients, *Blood Transfus. Italy* 19 (2021) 54–63, <https://doi.org/10.2450/2020.0256-20>.
- [19] S. Dzik, K. Eliason, E.B. Morris, R.M. Kaufman, C.M. North, COVID-19 and ABO blood groups, *Transfusion* 60 (2020) 1883–1884, <https://doi.org/10.1111/trf.15946>.
- [20] A. Khalil, R. Feghali, M. Hassoun, The Lebanese COVID-19 cohort; A challenge for the ABO blood group system, *Front. Med.* 7 (2020) 1–7, <https://doi.org/10.3389/fmed.2020.585341>.
- [21] Y. Cheng, G. Cheng, C.H. Chui, F.Y. Lau, E. Al, ABO blood group and susceptibility to severe acute respiratory syndrome to, *J. Am. Med. Assoc.* 293 (2005) 1450–1451, <https://doi.org/10.3122/jabfm.19.6.648>.
- [22] C. Gérard, G. Maggipinto, J.M. Minon, COVID-19 and ABO blood group: another viewpoint, *Br. J. Haematol.* 190 (2020) e93–e94, <https://doi.org/10.1111/bjh.16884>.
- [23] X. Dai, ABO blood group predisposes to COVID-19 severity and cardiovascular diseases, *Eur. J. Prev. Cardiol.* 27 (2020) 1436–1437, <https://doi.org/10.1177/2047487320922370>.
- [24] L. Dean, The ABO blood group, in: L. Dean (Ed.), *Blood Groups and Red Cell Antigens*, National Center for Biotechnology Information, Bethesda, 2005, pp. 31–39.
- [25] J. Le Pendu, A. Breiman, J. Rocher, M. Dion, N. Ruvoën-Clouet, ABO blood types and COVID-19: Spurious, Anecdotal, or truly important relationships? A reasoned review of Available data, *Viruses* 13 (2021) 1–20, <https://doi.org/10.1016/b978-0-08-022449-7.50017-x>.
- [26] W.M. Watkins, Some genetical aspects of the biosynthesis of human blood group substances, in: C. Wolstenholme, C. O'Connor (Eds.), *Biochemistry of Human Genetics*, Little, Brown and Company, Boston, 1959, pp. 217–238.
- [27] F. Yamamoto, Review: ABO blood group system—ABH oligosaccharide antigens, anti-A and anti-B, A and B glycosyltransferases, and ABO genes, *Immunohematol* 20 (2004) 3–22.
- [28] F.I. Yamamoto, H. Clausen, T. White, J. Marken, S.I. Hakomori, Molecular genetic basis of the histo-blood group ABO system, *Nature* 345 (1990) 229–233, <https://doi.org/10.1038/345229a0>.
- [29] W.M. Watkins, The ABO blood group system: historical background, *Transfus. Med.* 11 (2001) 243–265, <https://doi.org/10.1046/j.1365-3148.2001.00321.x>.
- [30] S.I. Patenaude, N.O.L. Seto, S.N. Borisova, A. Szpacenko, S.L. Marcus, M. M. Palcic, et al., The structural basis for specificity in human abo(h) blood group biosynthesis, *Nat. Struct. Biol.* 9 (2002) 685–690, <https://doi.org/10.1038/nsb832>.
- [31] D.R. Branch, Anti-A and anti-B: what are they and where do they come from? *Transfusion* 55 (2015) S74–S79, <https://doi.org/10.1111/trf.13087>.
- [32] G.F. Springer, R.E. Horton, Blood group isoantibody stimulation in man by feeding blood group-active bacteria, *J. Clin. Invest.* 48 (1969) 1280–1291, <https://doi.org/10.1172/JCI106094>.
- [33] D. Bello-Gil, C. Audebert, S. Olivera-Ardid, M. Pérez-Cruz, G. Even, N. Khasbiullina, et al., The formation of glycan-specific natural antibodies repertoire in Galt-KO mice is determined by gut microbiota, *Front. Immunol.* 10 (2019) 1–13, <https://doi.org/10.3389/fimmu.2019.00342>.
- [34] P. Greenwell, Blood group antigens: molecules seeking a function? *Glycoconj. J.* 14 (1997) 159–173, <https://doi.org/10.1023/A:1018581503164>.
- [35] R. Oriol, J. Le Pendu, R. Mollicone, Genetics of ABO, H, Lewis, X and related antigens, *Vox Sang.* 51 (1986) 161–171, <https://doi.org/10.1111/j.1423-0410.1986.tb01946.x>.
- [36] J. Geisel, M.K. Steuer, H.L. Ko, J. Beuth, The role of ABO blood groups in infections induced by *Staphylococcus saprophyticus* and *Pseudomonas*

- aeruginosa, Zentralblatt für Bakteriologie 282 (1995) 427–430, [https://doi.org/10.1016/S0934-8840\(11\)80714-0](https://doi.org/10.1016/S0934-8840(11)80714-0).
- [37] P. Gagneux, A. Varki, Evolutionary considerations in relating oligosaccharide diversity to biological function, *Glycobiology* 9 (1999) 747–755, <https://doi.org/10.1093/glycob/9.8.747>.
- [38] T. Borén, P. Falk, K.A. Roth, G. Larson, S. Normark, Attachment of *Helicobacter pylori* to human gastric epithelium mediated by blood group antigens, *Science* 262 (1993), <https://doi.org/10.1126/science.1229933>.
- [39] A. Stapleton, E. Nudelman, H. Clausen, S.I. Hakomori, W.E. Stamm, Binding of uropathogenic *Escherichia coli* R45 to glycolipids extracted from vaginal epithelial cells is dependent on histo-blood group secretor status, *J. Clin. Invest.* 90 (1992) 965–972, <https://doi.org/10.1172/JCI115973>.
- [40] F. Georg, M.D. Springer, P. Williamson, W. Brandes, Blood group Activity of gram-negative bacteria, *J. Exp. Med.* 113 (1961) 1077–1093, <https://doi.org/10.1046/j.1537-2995.1988.28388219149.x>.
- [41] R.M. Seymour, M.J. Allan, A. Pomiankowski, K. Gustafsson, Evolution of human ABO polymorphism by two complementary selective pressures, *Proc. R. Soc. B Biol. Sci.* 271 (2004) 1065–1072, <https://doi.org/10.1098/rspb.2004.2674>.
- [42] M. Arendrup, J.-E.S. Hansen, H. Clausen, C. Nielsen, L.R. Mathiesen, J.O. Nielsen, Antibody to histo-blood group A antigen neutralizes HIV produced by lymphocytes from blood group A donors but not from blood group B or O donors, *AIDS* 5 (1991) 441–444, <https://doi.org/10.1097/00002030-199104000-00014>.
- [43] A.F. Preece, K.M. Strahan, J. Devitt, F.I. Yamamoto, K. Gustafsson, Expression of ABO or related antigenic carbohydrates on viral envelopes leads to neutralization in the presence of serum containing specific natural antibodies and complement, *Blood* 99 (2002) 2477–2482, <https://doi.org/10.1182/blood.V99.7.2477>.
- [44] P. Guillon, M. Clément, V. Sébille, J.G. Rivain, C.F. Chou, N. Ruvoën-Clouet, et al., Inhibition of the interaction between the SARS-CoV Spike protein and its cellular receptor by anti-histo-blood group antibodies, *Glycobiology* 18 (2008) 1085–1093, <https://doi.org/10.1093/glycob/cwn093>.
- [45] J.A. Rowe, I.G. Handel, M.A. Thera, A.M. Deans, K.E. Lyke, A. Koné, et al., Blood group O protects against severe *Plasmodium falciparum* malaria through the mechanism of reduced rosetting, *Proc. Natl. Acad. Sci. USA* 104 (2007) 17471–17476, <https://doi.org/10.1073/pnas.0705390104>.
- [46] L. Ségurel, Z. Gao, M. Przeworski, Ancestry runs deeper than blood: the evolutionary history of ABO points to cryptic variation of functional importance, *Bioessays* 35 (2013) 862–867.
- [47] M.B. Barnkob, A. Pottegård, H. Støvring, T.M. Haunstrup, K. Homburg, R. Larsen, et al., Reduced prevalence of SARS-CoV-2 infection in ABO blood group O, *Blood Adv.* 4 (2020) 4990–4993, <https://doi.org/10.1182/BLOODADVANCES.2020002657>.
- [48] J. Schetelig, H. Baldauf, S. Wendler, F. Heidenreich, R. Real, M. Kolditz, et al., Blood group A epitopes do not facilitate entry of SARS-CoV-2, *J. Intern. Med.* 290 (2021) 223–226, <https://doi.org/10.1111/joim.13256>.
- [49] M. Enguita-Gérman, J. Librero, L. Leache, M. Gutiérrez-Valencia, I. Tamayo, C. Jericó, et al., Role of the ABO blood group in COVID-19 infection and complications: a population-based study, *Transfus. Apher. Sci.* (2022), <https://doi.org/10.1016/j.transci.2022.103357>.
- [50] I. Jukić, A. Hećimović, T. Vuk, M. Vinković, T. Kereš, M. Lampalo, et al., Prevalence of ABO and RhD blood group phenotypes in the Croatian population and in patients with severe COVID-19 in Croatia, *Blood Transfus.* (2022), <https://doi.org/10.2450/2022.0311-21>.
- [51] N.A. El-Shitany, M. El-Hamamsy, A.A. Alahmadi, B.G. Eid, T. Neamatallah, H. S. Almkadi, et al., The impact of abo blood grouping on covid-19 vulnerability and seriousness: a retrospective cross-sectional controlled study among the Arab community, *Int. J. Environ. Res. Publ. Health* 18 (2021) 1–19, <https://doi.org/10.3390/ijerph18010276>.
- [52] Q. Fan, W. Zhang, B. Li, D.-J. Li, J. Zhang, F. Zhao, Association between ABO blood group system and COVID-19 susceptibility in Wuhan, *Front. Cell. Infect. Microbiol.* 10 (2020) 1–10, <https://doi.org/10.3389/fcimb.2020.00404>.
- [53] M. Franchini, C. Glingani, C. Del Fante, M. Capuzzo, V. Di Stasi, G. Rastrelli, et al., The protective effect of O blood type against SARS-CoV-2 infection, *Vox Sang.* 116 (2021) 249–250, <https://doi.org/10.1111/vox.13003>.
- [54] İ. Solmaz, S. Araç, ABO blood groups in COVID-19 patients; Cross-sectional study, *Int. J. Clin. Pract.* 75 (2021), <https://doi.org/10.1111/ijcp.13927>.
- [55] S. Yalouji, R. Fakhfakh, F. Tritar, N. Chaouch, T. Mestiri, M. Besbes, et al., Groupes sanguins ABO et risque d'atteinte par le covid-19, *Tunis. Med.* 98 (2020) 888–891.
- [56] H. Göker, E. Aladağ-Karakulak, H. Demiroğlu, C.M. Ayaz, Y. Büyükaşık, A. Ç. İnkaya, et al., The effects of blood group types on the risk of COVID-19 infection and its clinical outcome, *Turk. J. Med. Sci.* 50 (2020) 679–683, <https://doi.org/10.3906/sag-2005-395>.
- [57] M. Muñoz-Culla, A. Roncancio-Clavijo, B. Martínez, M. Gorostidi-Aicua, L. Pineiro, A. Azkune, et al., O group is a protective factor for COVID19 in Basque population, *PLoS One* 16 (2021) 3–6, <https://doi.org/10.1371/journal.pone.0249494>.
- [58] R. Mahmud, M.A. Rassel, F.B. Monayem, S.K.J.B. Sayeed, M.S. Islam, M.M. Islam, et al., Association of ABO blood groups with presentation and outcomes of confirmed SARS CoV-2 infection: a prospective study in the largest COVID-19 dedicated hospital in Bangladesh, *PLoS One* 16 (2021) 1–10, <https://doi.org/10.1371/journal.pone.0249252>.
- [59] M.A. Almadhi, A. Abdulrahman, A. Alawadhi, A.A. Rabaan, S. Atkin, M. AlQahtani, The effect of ABO blood group and antibody class on the risk of COVID-19 infection and severity of clinical outcomes, *Sci. Rep.* 11 (2021) 19–23, <https://doi.org/10.1038/s41598-021-84810-9>.
- [60] M.A. Ozkarafakili, N. Gareayaghi, Z.M.Y. Kara, Relationship between ABO blood types and COVID-19 severity, *Med. Bull. Sisli Hosp.* 56 (2022) 1–20, <https://doi.org/10.14744/SEMB.2021.15045>.
- [61] J. Gamboa-aguilera, Á.C. Zamorano-montaño, A. Enríquez-osorio, W. Torres-cubillas, J.L. López-arroyo, A. Mata, et al., Abo blood group, Atherothrombotic comorbidities, and COVID-19: a case-control study of their association in the Mexican population, *Arch. Med. Res.* 53 (2022) 100–108, <https://doi.org/10.1016/j.arcmed.2021.09.005>.
- [62] S. Gurung, N.B. Mahotra, L. Shrestha, A. Sheralpi, S.P. Joshi, G. Shrestha, et al., Association of ABO blood group with susceptibility to SARS-CoV-2 infection in Rupandehi district of Nepal, *Sage Open Med.* 10 (2022) 1–5, <https://doi.org/10.1177/20503121221095413>.
- [63] R.K. Leaf, H. Al-Samkari, S.K. Brenner, S. Gupta, D.E. Leaf, ABO phenotype and death in critically ill patients with COVID-19, *Br. J. Haematol.* 190 (2020) 204–208, <https://doi.org/10.1111/bjh.16984>.
- [64] J.K. Niles, H.E. Karnes, J.S. Dlott, H.W. Kaufman, Association of ABO/Rh with SARS-CoV-2 positivity: the role of race and ethnicity in a female cohort, *Am. J. Hematol.* 96 (2021) 1–3, <https://doi.org/10.1002/ajh.26019>.
- [65] S. Lehrer, P.H. Rheinstein, ABO blood groups, COVID-19 infection and mortality, *Blood Cells, Mol. Dis.* 89 (2021) 1–4, <https://doi.org/10.1016/j.bcmd.2021.102571>.
- [66] J.E. Levi, P.R. Telles, H. Scrivani, G. Campana, Lack of association between ABO blood groups and susceptibility to SARS-CoV-2 infection, *Vox Sang.* 116 (2021) 251–252, <https://doi.org/10.1111/vox.13015>.
- [67] J. Mullins, A.H. Al-Tarbsheh, H. Chieng, P. Chaukiyal, S. Ghalib, E. Jain, et al., The association of ABO blood type with the risk and severity of COVID-19 infection, *Am. J. Blood Res.* 11 (2021) 53–58.
- [68] C.A. Latz, C. DeCarlo, L. Boitano, C.Y.M. Png, R. Patell, M.F. Conrad, et al., Blood type and outcomes in patients with COVID-19, *Ann. Hematol. Annals of Hematology* 99 (2020) 2113–2118, <https://doi.org/10.1007/s00277-020-04169-1>.
- [69] S. Padhi, S. Suvankar, D. Dash, V.K. Panda, A. Pati, J. Panigrahi, et al., ABO blood group system is associated with COVID-19 mortality: an epidemiological investigation in the Indian population, *Transfus. Clin. Biol.* 27 (2020) 253–258, <https://doi.org/10.1016/j.tracu.2020.08.009>.
- [70] J.G. Ray, M.J. Schull, M.J. Vermeulen, A.L. Park, Association between ABO and Rh blood groups and SARS-CoV-2 infection or severe COVID-19 illness: a population-based cohort study, *Ann. Intern. Med.* 174 (2021) 308–315, <https://doi.org/10.7326/M20-4511>.
- [71] S.A. Al-Youha, W. Alduaij, A. Al-Serri, S.M. Almazzeedi, M. Al-Haddad, M. H. Jamal, et al., The impact of ABO blood groups on clinical outcomes and susceptibility to COVID-19: a retrospective study in an unselected population, *Transfusion* 61 (2021) 1631–1641, <https://doi.org/10.1111/trf.16365>.
- [72] T.R. Kotila, T.O. Alonge, A. Fowotade, O.I. Famuyiwa, A.S. Akinbile, Association of the ABO blood group with SARS-CoV-2 infection in a community with low infection rate, *Vox Sang.* 116 (2021) 910–915, <https://doi.org/10.1111/vox.13077>.
- [73] D. Jawdat, A. Hajeer, S. Massadeh, N. Aljawini, M.S. Abedalthagafi, M. Alaamery, Correlation between ABO blood group phenotype and the risk of COVID-19 infection and severity of disease in a Saudi Arabian cohort, *J. Epidemiol. Glob. Health* 12 (2022) 85–91, <https://doi.org/10.1007/s44197-021-00023-3>.
- [74] N. Liu, T. Zhang, L. Ma, H. Zhang, H. Wang, W. Wei, et al., The impact of ABO blood group on COVID-19 infection risk and mortality: a systematic review and meta-analysis, *Blood Rev.* 48 (2021), <https://doi.org/10.1016/j.blre.2020.100785>.
- [75] T. Kander, M.F. Bjurström, A. Frigyesi, M. Jöud, C.U. Nilsson, ABO and RhD blood group are not associated with mortality and morbidity in critically ill patients: a multicentre observational study of 29 512 patients, *BMC Anesthesiol.* 22 (2022) 1–8, <https://doi.org/10.1186/s12871-022-01626-4>.
- [76] L. Boudin, F. Janvier, O. Bylickei, F. Dutasta, ABO blood groups are not associated with the risk of acquiring SARS-CoV-2 infection in young adults, *Haematologica* 105 (2020) 2841–2843, <https://doi.org/10.3810/psm.1999.02.687>.
- [77] M. Kibler, A. Carmona, B. Marchandot, K. Matsushita, A. Trimaille, M. Kanso, et al., Risk and severity of COVID-19 and ABO blood group in transcatheter aortic valve patients, *J. Clin. Med.* 9 (2020) 1–12, <https://doi.org/10.1101/2020.06.13.20130211>.
- [78] P. Negro, M. Congedo, A. Zizza, M. Guido, G. Sacquegna, G. Pulito, et al., Role of ABO blood system in COVID-19: findings from a southern Italian study, *Transfus. Med.* 1–5 (2021), <https://doi.org/10.1111/tme.12797>.
- [79] M. Badedi, A. Alnami, H. Darraj, A. Alrajhi, D.A. Mutawwam, M. Somaily, et al., Clinical characteristics and ABO blood groups in COVID-19 patients, *Saudi Arabia, Medicine* 100 (2021) 1–6, <https://doi.org/10.1097/MD.00000000000026738>.
- [80] L. Valenti, S. Villa, G. Baselli, R. Temporiti, A. Bandera, L. Scudeller, et al., Association of ABO blood group and secretor phenotype with severe COVID-19, *Transfusion* 60 (2020) 3067–3070, <https://doi.org/10.1111/trf.16130>.
- [81] V. Gursoy, S. Avci, Effect of ABO blood groups on length of hospital stay according to age in Covid-19 patients, *Hematol Transfus Cell Ther* 44 (2022) 7–12, <https://doi.org/10.1016/j.htct.2021.08.013>.
- [82] D.H. Bokhary, N.H. Bokhary, L.E. Seadawi, A.M. Moafa, H.H. Khairallah, A. A. Bakhs, Variation in COVID-19 disease severity and clinical outcomes between different ABO blood groups, *Cureus* 12 (2022), <https://doi.org/10.3389/fmicb.2021.799519>.
- [83] H.F. Jelinek, M. Mousa, N. Alkaabi, E. Alefishat, G. Daw Elbait, H. Kannout, et al., Allelic variants within the ABO blood group phenotype confer protection against

- critical COVID-19 hospital presentation, *Front. Med.* 8 (2022), <https://doi.org/10.3389/fmed.2021.759648>.
- [84] I. Garg, S. Srivastava, V. Dogra, M. Bargotyia, S. Bhattar, U. Gupta, et al., Potential association of COVID-19 and ABO blood group: an Indian study, *Microb. Pathog.* 158 (2021) 1–7, <https://doi.org/10.1016/j.micpath.2021.105008>.
- [85] P.M.M. Garibaldi, L.C. Oliveira, B.A. da Fonseca, M.A. Martins, C.H. Miranda, C. E.L. Almado, et al., Histo-blood group A is a risk factor for severe COVID-19, *Transfus. Med.* 1–4 (2021), <https://doi.org/10.1111/tme.12796>.
- [86] R.L. Hoiland, N.A. Fergusson, A.R. Mitra, D.E.G. Griesdale, D.V. Devine, S. Stukas, et al., The association of ABO blood group with indices of disease severity and multiorgan dysfunction in COVID-19, *Blood Adv.* 4 (2020) 4981–4989, <https://doi.org/10.1182/BLOODADVANCES.2020002623>.
- [87] C. Sardu, R. Marfella, P. Maggi, V. Messina, P. Cirillo, V. Codella, et al., Implications of ABO blood group in hypertensive patients with covid-19, *BMC Cardiovasc. Disord.* 20 (2020) 1–10, <https://doi.org/10.1186/s12872-020-01658-z>.
- [88] J. Szymanski, L. Mohrmann, J. Carter, R. Nelson, S. Chekuri, A. Assa, et al., ABO blood type association with SARS-CoV-2 infection mortality: a single-center population in New York City, *Transfusion* 61 (2021) 1064–1070, <https://doi.org/10.1111/trf.16339>.
- [89] A. Nafakhi, I.S. Rabeaa, R. Al-Darraj, H. Nafakhi, A. Mechi, A. Al-Khalidi, et al., Association of ABO blood group with in-hospital adverse outcome and long term persistent symptoms of COVID-19 infection: a single-center longitudinal observational study, *Heal Sci Reports* 5 (2022) 1–6, <https://doi.org/10.1002/hsr.2.656>.
- [90] R.A.M. Alabsi, N.C. Sandeepa, R.T. Misfer, M.M. Alraqdi, M.I.M. Hamdi, Correlation between post-COVID-19, chemosensitive function, blood group, and oral health-related quality of life, *Int. J. Dentistry.* 2022 (2022) 1–8, <https://doi.org/10.1155/2022/8715777>.
- [91] J.F. Shelton, A.J. Shastri, C. Ye, C.H. Weldon, T. Filshstein-Sonmez, D. Coker, et al., Trans-ancestry analysis reveals genetic and nongenetic associations with COVID-19 susceptibility and severity, *Nat. Genet.* 53 (2021) 801–808, <https://doi.org/10.1038/s41588-021-00854-7>.
- [92] CHGI, Mapping the human genetic architecture of COVID-19, *Nature* 600 (2021) 472–477, <https://doi.org/10.1038/s41586-021-03767-x>.
- [93] E. Pairo-Castineira, S. Clohisey, L. Klaric, A.D. Bretherick, K. Rawlik, D. Pasko, et al., Genetic mechanisms of critical illness in COVID-19, *Nature* 591 (2021) 92–98, <https://doi.org/10.1038/s41586-021-03767-x>.
- [94] G. Vargas-Alarcón, J. Ramírez-Bello, R. Posadas-Sánchez, G. Rojas-Velasco, A. López-Reyes, L. Martínez-Gómez, et al., The rs8176740 T/A and rs512770 T/C genetic variants of the ABO gene increased the risk of COVID-19, as well as the plasma concentration platelets, *Biomolecules* 12 (2022) 1–11, <https://doi.org/10.3390/biom12040486>.
- [95] E.M. Matzhold, A. Berghold, M.K.B. Bemelmans, C. Banfi, E. Stelzl, H.H. Kessler, et al., Lewis and ABO histo-blood types and the secretor status of patients hospitalized with COVID-19 implicate a role for ABO antibodies in susceptibility to infection with SARS-CoV-2, *Transfusion* 61 (2021) 2736–2745, <https://doi.org/10.1111/trf.16567>.
- [96] E. Coto, G.M. Albaiceta, M.G. Clemente, J. Gómez, Lack of association between SNPsrs8176719 (O blood group) and COVID-19: data from Spanish age matched patients and controls, *Transfusion* 61 (2021) 654–656, <https://doi.org/10.1111/trf.16206>.
- [97] N. Nishida, M. Sugiyama, Y. Kawai, I. Naka, N. Iwamoto, T. Suzuki, et al., Genetic association of IL17 and the importance of ABO blood group antigens in saliva to COVID-19, *Sci. Rep.* 12 (2022) 1–12, <https://doi.org/10.1038/s41598-022-07856-3>.
- [98] A.I. Hernández Cordero, X. Li, S. Milne, C.X. Yang, Y. Bossé, P. Joubert, et al., Multi-omics highlights ABO plasma protein as a causal risk factor for COVID-19, *Hum. Genet.* 140 (2021) 969–979, <https://doi.org/10.1007/s00439-021-02264-5>.
- [99] A.B. Palmos, V. Millischer, D.K. Menon, T.R. Nicholson, L.S. Taams, B. Michael, et al., Proteome-wide Mendelian randomization identifies causal links between blood proteins and severe COVID-19, *PLoS Genet.* 18 (2022) 1–23, <https://doi.org/10.1371/journal.pgen.1010042>.
- [100] S. Zhou, G. Butler-Laporte, T. Nakanishi, D.R. Morrison, J. Afilalo, M. Afilalo, et al., A Neanderthal OAS1 isoform protects individuals of European ancestry against COVID-19 susceptibility and severity, *Nat. Med.* 27 (2021) 659–667, <https://doi.org/10.1038/s41591-021-01281-1>.
- [101] J. Zhu, C. Wu, L. Wu, Associations between genetically predicted protein levels and COVID-19 severity, *J. Infect. Dis.* 223 (2021) 19–22, <https://doi.org/10.1093/infdis/jiaa660>.
- [102] R. Amraei, W. Yin, M.A. Napoleon, E.L. Suder, J. Berrigan, Q. Zhao, et al., CD209L/L-SIGN and CD209/DC-SIGN act as receptors for SARS-CoV-2, *ACS Cent. Sci.* 7 (2021) 1156–1165, <https://doi.org/10.1021/acscentsci.0c01537>.
- [103] G.A. Pathak, K. Singh, T.W. Miller-Fleming, F.R. Wendt, N. Ehsan, K. Hou, et al., Integrative genomic analyses identify susceptibility genes underlying COVID-19 hospitalization, *Nat. Commun.* 12 (2021) 1–11, <https://doi.org/10.1038/s41467-021-24824-z>.
- [104] T. Koster, A. Blann, E. Briët, J. Vandenbroucke, F. Rosendaal, Role of clotting factor VIII in effect of von Willebrand factor occurrence of deep-vein thrombosis, *Lancet* 345 (1995) 152–155, [https://doi.org/10.1016/s0140-6736\(95\)90166-3](https://doi.org/10.1016/s0140-6736(95)90166-3).
- [105] L. Zou, F. Ruan, M. Huang, L. Liang, H. Huang, Z. Hong, SARS-CoV-2 viral load in upper respiratory specimens of infected patients, *N. Engl. J. Med.* 382 (2020) 1177–1179, <https://doi.org/10.1056/nejmc2000231>.
- [106] R. Wölfel, V.M. Corman, W. Guggemos, M. Seilmaier, S. Zange, M.A. Müller, et al., Virological assessment of hospitalized patients with COVID-2019, *Nature* 581 (2020) 465–469, <https://doi.org/10.1038/s41586-020-2196-x>.
- [107] K.P.Y. Hui, M.C. Cheung, R.A.P.M. Perera, K.C. Ng, C.H.T. Bui, J.C.W. Ho, et al., Tropism, replication competence, and innate immune responses of the coronavirus SARS-CoV-2 in human respiratory tract and conjunctiva: an analysis in ex-vivo and in-vitro cultures, *Lancet Respir. Med.* 8 (2020) 687–695, [https://doi.org/10.1016/S2213-2600\(20\)30193-4](https://doi.org/10.1016/S2213-2600(20)30193-4).
- [108] R. Zang, M.F.G. Castro, B.T. McCune, Q. Zeng, P.W. Rothlauf, N.M. Sonnek, et al., TMPRSS2 and TMPRSS4 mediate SARS-CoV-2 infection of human small intestinal enterocytes, *Sci. Immunol.* 3582 (2020) 1–14, <https://doi.org/10.1101/2020.04.21.054015>.
- [109] S. Lukassen, R.L. Chua, T. Trefzer, N.C. Kahn, M.A. Schneider, T. Muley, et al., SARS-CoV-2 receptor ACE 2 and TMPRSS 2 are primarily expressed in bronchial transient secretory cells, *EMBO J.* 39 (2020) 1–15, <https://doi.org/10.15252/emboj.20105114>.
- [110] W. Sungnak, N. Huang, C. Bécaivin, M. Berg, R. Queen, M. Litvinukova, et al., SARS-CoV-2 entry factors are highly expressed in nasal epithelial cells together with innate immune genes, *Nat. Med.* 26 (2020) 681–687, <https://doi.org/10.1038/s41591-020-0868-6>.
- [111] C.G.K. Ziegler, S.J. Allon, S.K. Nyquist, I.M. Mbanjo, V.N. Miao, C.N. Tzouanas, et al., SARS-CoV-2 receptor ACE2 is an interferon-stimulated gene in human airway epithelial cells and is detected in specific cell subsets across tissues, *Cell* 181 (2020) 1016–1035, <https://doi.org/10.1016/j.cell.2020.04.035>.
- [112] V. Ravn, E. Dabelsteen, Tissue distribution of histo-blood group antigens, *Apmis* 108 (2000) 1–28, <https://doi.org/10.1034/j.1600-0463.2000.d01-1.x>.
- [113] M. Bouhaddou, D. Memon, B. Meyer, K. White, V. Rezelj, The global phosphorylation landscape of SARS-CoV-2 infection, *Cell* 182 (2020) 685–712, <https://doi.org/10.1016/j.cell.2020.06.034>.
- [114] C.A.M. de Haan, P.J.M. Rottier, Molecular interactions in the assembly of coronaviruses, *Adv. Virus Res.* 64 (2005) 165–230, [https://doi.org/10.1016/S0065-3527\(05\)64006-7](https://doi.org/10.1016/S0065-3527(05)64006-7).
- [115] P. Zhao, J.L. Praissman, O.C. Grant, Y. Cai, T. Xiao, K.E. Rosenbalm, et al., Virus-receptor interactions of glycosylated SARS-CoV-2 spike and human ACE2 receptor, *Cell Host Microbe* 28 (2020) 586–601, <https://doi.org/10.1101/2020.06.25.172403>.
- [116] M. Ujike, F. Taguchi, Incorporation of spike and membrane glycoproteins into coronavirus virions, *Viruses* 7 (2015) 1700–1725, <https://doi.org/10.3390/v7041700>.
- [117] Y. Watanabe, J.D. Allen, D. Wrapp, J.S. McLellan, M. Crispin, Site-specific glycan analysis of the SARS-CoV-2 spike, *Science* 369 (2020) 330–333, <https://doi.org/10.1126/science.abb9983>.
- [118] M. Sanda, L. Morrison, R. Goldman, N-and O-glycosylation of the SARS-CoV-2 spike protein, *Anal. Chem.* 93 (2021), <https://doi.org/10.1021/acs.analchem.0c03173>, 2003–9.
- [119] A. Shajahan, N.T. Supekar, A.S. Gleinich, P. Azadi, Deducing the N- and O-glycosylation profile of the spike protein of novel coronavirus SARS-CoV-2, *Glycobiology* 30 (2020) 981–988, <https://doi.org/10.1093/glycob/cwaa042>.
- [120] Z. Sun, K. Ren, X. Zhang, J. Chen, Z. Jiang, J. Jiang, et al., Mass spectrometry analysis of newly emerging coronavirus HCoV-19 spike protein and human ACE2 reveals camouflaging glycans and unique post-translational modifications, *Engineering* 7 (2021) 1441–1451, <https://doi.org/10.1016/j.eng.2020.07.014>.
- [121] M. Deleers, A. Breiman, V. Daubie, C. Maggetto, I. Barreau, T. Besse, et al., Covid-19 and blood groups: ABO antibody levels may also matter, *Int. J. Infect. Dis.* 104 (2021) 242–249, <https://doi.org/10.1016/j.ijid.2020.12.025>.
- [122] F.Z. Zaidi, A.R.Z. Zaidi, S.M. Abdullah, S.Z.A. Zaidi, COVID-19 and the ABO blood group connection, *Transfus. Apher. Sci.* 59 (2020) 1–2, <https://doi.org/10.1016/j.transci.2020.102838>.
- [123] R. Boukhari, A. Breiman, J. Jazat, N. Ruvoën-Clouet, S. Martinez, A. Damais-Cepitelli, et al., ABO blood group incompatibility protects against SARS-CoV-2 transmission, *Front. Microbiol.* 12 (2022), <https://doi.org/10.3389/fmicb.2021.799519>.
- [124] T. Mazda, R. Yabe, O. NaThalang, T. Thammavong, K. Tadokoro, Differences in ABO antibody levels among blood donors: a comparison between past and present Japanese, Laotian, and Thai populations, *Immunohematol* 23 (2007) 38–41, <https://doi.org/10.21307/immunohematology-2019-261>.
- [125] A. Breiman, N. Ruven-Clouet, J. Le Pendu, Harnessing the natural anti-glycan immune response to limit the transmission of enveloped viruses such as SARS-CoV-2, *PLoS Pathog.* 16 (2020) 8–11, <https://doi.org/10.1371/journal.ppat.1008556>.
- [126] M. Hoffmann, H. Kleine-Weber, S. Pöhlmann, A multibasic cleavage site in the spike protein of SARS-CoV-2 is essential for infection of human lung cells, *Mol. Cell* 78 (2020) 779–784, <https://doi.org/10.1016/j.molcel.2020.04.022>.
- [127] S. Li, R.X. Xu, Y.L. Guo, Y. Zhang, C.G. Zhu, J. Sun, et al., ABO blood group in relation to plasma lipids and proprotein convertase subtilisin/kexin type 9, *Nutr. Metabol. Cardiovasc. Dis.* 25 (2015) 411–417, <https://doi.org/10.1016/j.numecd.2014.10.015>.
- [128] A. AbdelMassih, R. Mahrous, A. Taha, A. Saud, A. Osman, B. Kamel, et al., The potential use of ABO blood group system for risk stratification of COVID-19, *Med. Hypotheses* 145 (2020) 1–6, <https://doi.org/10.1016/j.mehy.2020.110343>.
- [129] M. Alejandra Tortorici, A.C. Walls, Y. Lang, C. Wang, Z. Li, D. Koerhuis, et al., Structural basis for human coronavirus attachment to sialic acid receptors, *Nat. Struct. Mol. Biol.* 26 (2019) 481–489, <https://doi.org/10.1038/s41594-019-0233-y>.

- [130] Y. Renhong, Z. Yuanyuan, L. Yaning, X. Lu, G. Yingying, Z. Qiang, Structural basis for the recognition of SARS-CoV-2 by full-length human ACE2, *Science* 367 (2020) 1444–1448, <https://doi.org/10.1126/science.abb2762>.
- [131] M. Awasthi, S. Gulati, D.P. Sarkar, S. Tiwari, S. Kateriya, P. Ranjan, et al., The sialoside-binding pocket of SARS-CoV-2 spike glycoprotein structurally resembles MERS-CoV, *Viruses* 12 (2020) 1–10, <https://doi.org/10.3390/v12090909>.
- [132] E. Milanetti, M. Miotto, L. Di Rienzo, M. Nagaraj, M. Monti, T.W. Golbek, et al., In-silico evidence for a two receptor based strategy of SARS-CoV-2, *Front. Mol. Biosci.* 8 (2021) 1–11, <https://doi.org/10.3389/fmolb.2021.690655>.
- [133] M. Cohen, N. Hurtado-Ziola, A. Varki, ABO blood group glycans modulate sialic acid recognition on erythrocytes, *Blood* 114 (2009) 3668–3676, <https://doi.org/10.1182/blood-2009-06-227041>.
- [134] J.C. Silva-filho, C.G.F. de Melo, J.L. de Oliveira, The influence of ABO blood groups on COVID-19 susceptibility and severity: a molecular hypothesis based on carbohydrate-carbohydrate interactions, *Med. Hypotheses* 144 (2020) 1–6, <https://doi.org/10.1016/j.mehy.2020.110155>.
- [135] S.C. Wu, C.M. Arthur, J. Wang, H. Verkerke, C.D. Josephson, D. Kalman, et al., The SARS-CoV-2 receptor-binding domain preferentially recognizes blood group A, *Blood Adv.* 5 (2021) 1305–1309, <https://doi.org/10.1182/bloodadvances.2020003259>.
- [136] S.R. Stowell, C.P. Stowell, Biologic roles of the ABH and Lewis histo-blood group antigens part II: thrombosis, cardiovascular disease and metabolism, *Vox Sang.* 114 (2019) 535–552, <https://doi.org/10.1111/vox.12786>.
- [137] O. Wu, N. Bayoumi, M.A. Vickers, P. Clark, ABO(H) blood groups and vascular disease: a systematic review and meta-analysis, *Thromb. Haemostasis* 6 (2007) 62–69, <https://doi.org/10.1160/TH08-05-0302>.
- [138] F. Dentali, A.P. Sironi, W. Ageno, S. Turato, C. Bonfanti, F. Frattini, et al., Non-O blood type is the commonest genetic risk factor for VTE: results from a meta-analysis of the literature, *Semin. Thromb. Hemost.* 38 (2012) 535–547, <https://doi.org/10.1055/s-0032-1315758>.
- [139] F. Yamamoto, E. Cid, M. Yamamoto, A. Blancher, ABO research in the modern Era of Genomics, *Transfus. Med. Rev.* 26 (2012) 103–118, <https://doi.org/10.1016/j.tmr.2011.08.002>.
- [140] P. Kamphuisen, J. Eikenboom, R. Bertina, Elevated factor VIII levels and risk of venous thrombosis, *Arterioscler. Thromb. Vasc. Biol.* 21 (2001) 731–738, <https://doi.org/10.1111/j.1365-2141.2012.09134.x>.
- [141] K. Kiouptsi, C. Reinhardt, Physiological roles of the von Willebrand factor-factor VIII interaction, in: U. Hoeger, J.R. Harris (Eds.), *Vertebrate and Invertebrate Respiratory Proteins, Lipoproteins and Other Body Fluid Proteins*, Springer, Cham, 2020, pp. 437–464.
- [142] T. Matsui, Y. Fujimura, S. Nishida, K. Titani, Human plasma α 2-macroglobulin and von Willebrand factor possess covalently linked ABO(H) blood group antigens in subjects with corresponding ABO phenotype, *Blood* 82 (1993) 663–668, <https://doi.org/10.1182/blood.v82.2.663.bloodjournal822663>.
- [143] P.K. Grewal, S. Uchiyama, D. Ditto, N. Varki, D.T. Le, V. Nizet, et al., The Ashwell receptor mitigates the lethal coagulopathy of sepsis, *Nat. Med.* 14 (2008) 648–655, <https://doi.org/10.1038/nm1760>.
- [144] N. Rydz, L.L. Swystun, C. Notley, A.D. Paterson, J.J. Riches, K. Sponagle, et al., The C-type lectin receptor CLEC4M binds, internalizes, and clears von Willebrand factor and contributes to the variation in plasma von Willebrand factor levels, *Blood* 121 (2013) 5228–5237, <https://doi.org/10.1182/blood-2012-10-457507>.
- [145] D.J. Bowen, An influence of ABO blood group on the rate of proteolysis of von Willebrand factor by ADAMTS13, *J. Thromb. Haemostasis* 1 (2003) 33–40, <https://doi.org/10.1046/j.1538-7836.2003.00007.x>.
- [146] B.K. Tan, S. Mainbourg, A. Friggeri, L. Bertolotti, M. Douplat, Y. Dargaud, et al., Arterial and venous thromboembolism in COVID-19: a study-level meta-analysis, *Thorax* 76 (2021) 970–979, <https://doi.org/10.1136/thoraxjnl-2020-215383>.
- [147] V. Mai, B. Kim, S. Mainbourg, F. Potus, M. Cucherat, J.C. Lega, et al., Venous thromboembolism in COVID-19 compared to non-COVID-19 cohorts: a systematic review with meta-analysis, *Vasc. Pharmacol.* 139 (2021) 1–8, <https://doi.org/10.1016/j.vph.2021.106882>.