

RESEARCH ARTICLE

Open Access

c-kit expression profile and regulatory factors during spermatogonial stem cell differentiation

Lei Zhang^{1,2,3}, Jiangjing Tang^{1,2,4}, Christopher J Haines², Huai Feng⁵, Liangxue Lai⁴, Xiaoming Teng^{1*} and Yibing Han^{1,2*}

Abstract

Background: It has been proven that *c-kit* is crucial for proliferation, migration, survival and maturation of spermatogenic cells. A periodic expression of *c-kit* is observed from primordial germ cells (PGCs) to spermatogenetic stem cells (SSCs). However, the expression profile of *c-kit* during the entire spermatogenesis process is still unclear. This study aims to reveal and compare *c-kit* expression profiles in the SSCs before and after the anticipated differentiation, as well as to examine its relationship with retinoic acid (RA) stimulation.

Results: We have found that there are more than 4 transcripts of *c-kit* expressed in the cell lines and in the testes. The transcripts can be divided into short and long categories. The long transcripts include the full-length canonical *c-kit* transcript and the 3' end short transcript. Short transcripts include the 3.4 kb short transcript and several truncated transcripts (1.9-3.2 kb). In addition, the 3.4 kb transcript (starting from intron 9 and covering exons 10 ~ 21) is discovered to be specifically expressed in the spermatogonia. The extracellular domain of Kit is obtained in the spermatogonia stage, but the intracellular domain (50 kDa) is constantly expressed in both SSCs and spermatogonia. The *c-kit* expression profiles in the testis and the spermatogonial stem cell lines vary after RA stimulation. The wave-like changes of the quantitative expression pattern of *c-kit* (increase initially and decrease afterwards) during the induction process are similar to that of the *in vivo* male germ cell development process.

Conclusions: There are dynamic transcription and translation changes of *c-kit* before and after SSCs' anticipated differentiation and most importantly, RA is a significant upstream regulatory factor for *c-kit* expression.

Keywords: *c-kit*, Expression profile, Spermatogonial stem cells, Differentiation, Regulatory factor

Background

Spermatogenesis starts from diploid spermatogonial stem cells (SSCs). The SSCs, also known as type A single (A_s) Spg, are located on the basement membrane of the seminiferous tubules. The A_s Spg can self-renew or produce the type A paired (A_p) Spg. After successive divisions, A_p Spg differentiates, forms chains of 4, 8 or 16 aligned Spg (A_{al}) and migrates along the basement membrane. A_{al} Spg differentiates into A1 Spg that further divides and differentiates into A2, A3, A4 intermediates and B Spg, which undergoes meiosis after a final mitosis stage [1]. The "undifferentiated" (A_s , A_p and A_{al}) and the "differentiating"

(A1, A2, A3, A4, intermediate and B) Spg differ in the expression profiles of *c-kit* [2]. *c-kit* is allelic to the W locus on mouse chromosome 5 [3]. The 21-exon gene encodes for a 5150 bp transcript, which is translated into a product of 145 kDa protein with 979 amino acid residues. This product is known as Kit [4]. Kit transduces growth regulatory signals across the plasma membrane and has three main functional regions, the extracellular, the transmembrane and the intracellular domains [5,6]. Its transcription process is only activated after binding with Kitl expressed by the Sertoli cells. The Kit/Kitl pathway is considered to be crucial for the proliferation, migration, survival and maturation of the germ cells [7-18]. In spite of the 5.1 Kb full-length canonical transcript, two alternative mRNAs of *c-kit*, 3.2 and 2.3 kb in length, exist in the haploid cells of the mouse testis [19]. With an Open Reading Frame (ORF) that starts in the intron 16 of the mouse *c-kit*, an alternative spermatid-specific *c-kit* transcript contains all of the

* Correspondence: xiaomingteng@hotmail.com; ybhan@yahoo.com
¹Shanghai first maternity and infant health hospital, Tongji University, Shanghai, China

²Department of Obstetrics and Gynaecology, Prince of Wales hospital, The Chinese University of Hong Kong, Hong Kong, China
Full list of author information is available at the end of the article

downstream exons (including 12 hydrophobic amino acids followed by the last 190 carboxyl terminal residues), encodes for Tr-Kit (~30 kDa) [7,20,21]. The 30 kDa Tr-Kit is found in the residual sperm cytoplasm and it has evident functions in the activation of oocyte during fertilization in mice [21,22].

c-kit has been a marker for SSCs pluripotency lost and its expression continues until meiosis is initiated [2,18]. The expression of protein Kit in the male germ cells is contradictory to those of gene *c-kit*. In early studies, Kit expression is detected in type A (A1–A4), intermediate, type B spermatogonia, as well as preleptotene spermatocytes, but not in the undifferentiated spermatogonia [2,18]. More recent studies demonstrate that Kit is also expressed A_s, A_{pr} and A_{al}. Therefore, whether Kit is expressed in spermatogonia and whether Kit/Kitl activation is a prerequisite for differentiation or not remain to be a question [23–28]. Even though the inactivation of *c-kit* by its specific inhibitor Imatinib results in Spg self-renewal impairment [29], both Kit⁻ and Kit⁺ spermatogonia have exhibited stem cell activities as evaluated by intra-seminiferous transplantation [1,24,30]. The POU5F1⁺/Kit⁺ subset of mouse SSCs can differentiate into several lines of somatic cells except for sperm cells [31].

We hypothesize that the expression profiles of *c-kit* in the male germ cells during spermatogenesis are dynamically changed before and after the expected differentiation, and these changes are important for their functional responses to the spermatogenesis-related genes. In this study, we have investigated the expression of *c-kit* in the immortal cell lines representing the SSCs, the differentiating spermatogonia and spermatocytes in hopes of understanding its natural expression patterns. We have also compared the *c-kit* expression patterns in those cell lines with their corresponding stage testes. The cell line c18-4 and 5 dpp mouse testes (before the initiation of spermatogonia differentiation) represent the undifferentiated spermatogonia. CRL-2053 and 10 dpp mouse testes (after the initiation of spermatogonia differentiation) represent the differentiating spermatogonia. CRL-2196 cells represent primary spermatocytes. The 60 dpp testes represent a mixture of the undifferentiated, the differentiating, the maturing and the matured germ cells.

RA, an active metabolite of vitamin A, is a vital signaling molecule for normal fetal development, pattern formation, cell proliferation, differentiation and apoptosis [32,33]. RA is considered to be crucial for germ cells to undergo meiosis in both male and female [34,35]. Testes of adult vitamin A-deficient mice/rat have seminiferous tubules that only contain Sertoli cells, type A spermatogonia and few preleptotene spermatocytes. With a reduced *c-kit* expression or without *Stra8* expression, the type A spermatogonia will arrest before differentiation (before A1 stage spermatogonia) [36]. Administration of

vitamin A to these animals results in a synchronized spermatogenesis emerging from type A spermatogonia and an enhanced expression of *c-kit* [37]. Therefore, RA is a key regulatory factor for *c-kit* expression.

Methods

Cell lines and animals

The c18-4 cell line represents the mouse SSCs [38]. CRL-2053 (ATCC) is a type B spermatogonia cell line [39]. CRL-2196 (GC-2spd(ts), ATCC) is a spermatocyte cell line [40,41]. C57/BL6 mice at different ages were purchased from laboratory animal service center (LASEC), The Chinese University of Hong Kong. All procedures were approved by the Animal Research Ethics Committee of the University.

Cell culture

All cells were cultured in the Dubecco modified eagle medium/F12 (DMEM/F12, Invitrogen, Carlsbad, CA, USA) supplemented with 10% fetal bovine serum (FBS, Invitrogen, Carlsbad, CA, USA). A subcultivation ratio of 1:6 to 1:10 was applied. Media were renewed 1 to 2 times per week. The cells were frozen in complete growth medium supplemented with 5% (v/v) DMSO and stored in liquid nitrogen.

Mouse testes collection

Mice at 5 days post partum (dpp), 10 dpp and 60 dpp were sacrificed by cervical dislocation. For RNA extraction, testes were washed twice with phosphate buffered saline (PBS) and then immersed in “RNA-later” stabilization reagent (Qiagen, Valencia, CA, USA). Before protein extraction, testes were washed twice with PBS, transported in iceboxes and stored in -80°C. Three batches of animals were used for each experiment.

In vitro tissue culture and RA induction

In vitro tissue culture was carried out according to the methods described by previous study [42]. Testes from 5 dpp, 10 dpp and 60 dpp mice were detunicated, cut into small pieces per testis, placed on Millicell CM filters (Millipore, Bedford, MA, USA) floating on the surface of medium and covered with drops of medium (DMEM/F12 + 10% FBS). RA (Sigma-Aldrich Co., Saint Louis, MO, USA) diluted in ethanol was added to the culture medium to make a final a concentration of 0.7 μM or 2 μM. Tissues were harvested after 24 hours of RA treatment. Total RNA was isolated using the RNeasy mini kit (Qiagen, Valencia, CA, USA).

In vitro cultured germ cells and RA induction

For germ cell exclusive induction assay, 2 × 10⁶ c18-4 or CRL-2063 cells were pre-seeded into T25 cell culture flasks separately (2 flasks each group) overnight before

the treatment in full medium (DMEM/F12 + 10% FBS). Induction media (DMEM/F12 + 10% FBS) with a final concentration of 2 μ M RA dissolved in ethanol were used in the treatment (induction) group. The same amount of ethanol without RA medium was set up as the control group. After 24 hours of induction, the induction media was removed, cells were washed with PBS twice, and cells were collected and stored at -80°C until analysis. Three independent replications were carried out for each experiment.

Methods for RNA preparation, electrophoresis and Northern blot

Total RNA from cells and testes was isolated using the RNeasy mini kit (Qiagen, Valencia, CA, USA) following the manufacturer's instructions. Sizes of RNA were estimated by comparing with 2 μ g RNA Millennium size markers (Ambion, Austin, TX, USA) by measuring the distance from each band to the loading well.

DNA fragments corresponding to exons 10–12 and exons 18–20 of the full-length *c-kit* transcript were obtained by PCR with *c-kit* specific primers using the 60 dpp mouse testis cDNA as the template. Primers sequences are shown in Table 1. Amplified DNA fragments were inserted into the Topo-TA vector (Invitrogen, Carlsbad, CA, USA). The plasmids were then extracted by QIAprep spin miniprep kit (Qiagen, Valencia, CA, USA) and were sent to commercial company for sequencing.

RNA probes were prepared by MAXIscript kit (Ambion, Austin, TX, USA) following the manufacturer's instructions. mRNA-complementary (antisense) transcripts were synthesized in a 20 μ l *in vitro* transcription system containing 1 μ g DNA template, 2 μ l 10 \times transcription buffer, 1 μ l 10 mM ATP, 1 μ l 10 mM CTP, 1 μ l 10 mM GTP, 5 μ l 800 Ci/mmol [α - ^{32}P] UTP at a concentration of 10 mCi/mL (Perkinelmer, San Jose, CA, USA) and 2 μ l T3 enzyme mix. After purification with NucAway Spin columns (Ambion, Austin, TX, USA), the RNA probes were hybridized with the blots with RNA samples in the ULTRAhyb ultrasensitive hybridization buffer (Ambion, Austin, TX, USA) at 68°C overnight. The same blot was stripped and re-probed with α - ^{32}P -labeled beta-actin RNA probe as internal control. Northern hybridization was performed twice with probes and membranes that were made independently. The sequences of PCR primers and RNA probes are shown in Table 1.

Rapid amplification of cDNA ends (RACE), cloning and sequencing

The number and size of *c-kit* mRNA expressed in mice cell lines and testis were determined by the Northern blot, the existence of these transcripts were further confirmed by RACE and sequencing. We used the BD-Smarter RACE protocol from BD Biosciences Clontech

Table 1 PCR primers, real-time PCR primers and RNA probe sequence of mouse *c-kit* gene

Probe name	Probe sequence (5' \rightarrow 3')
PCR primers	
Exons 10-12	Sense 5' -TGGGGATCATTGTGATGGT-3' Anti-sense 5'-ATGGCAGCATCCGACTTAAT-3'
Exons 18-20	Sense 5' - CCTCTGGGAGCTCTTCTCCT-3' Anti-sense 5'- GCTGTCCGAGATCTGCTTCT-3'
Real-time PCR primers	
Exon 7-8	AACGTTTACGT GAACACAAAACCCAG
Exon 20-21	GCACCAAGCACATTTACTCCAACCTT
Exon 21+	CTGATATGTTGTCCAACCTGTTGACA
Exons 10-12 probe (extracellular domain)	ATGGCAGCATCCGACTTAATCAAGCCA TATGCAGTGGCCTCAACGACC TTCCCGAAGGCCACCAGCTCCCAATGTC TTTCCAAAACCTCAGCCTGTTTC TGGGAAACTCCCATTTGTGATCATAAGG AAGTTGCGTCGGGTCTATGT AAACATAATTGTTTCCATTATCTCCTCG ACAACCTTCCATTGTACTTC ATACATGGGTTTCTGCAAATATTTGTAGG TGAGCAC CATCACAAATG ATCCCCAT
Exons 18-20 probe (intracellular domain)	GCTGTCCGAGATCTGCTTCTCAATAAGTT GGACAACCTGCTTGAATGT TGGCCTTTTCAAGGGGTGAGCGTCCAG CAAGTCTTCATGAC GTCATACATTTCCGCGAGCG CGTGCTCCGG GCTGACCATC CGGAAGCCTTCTTGATCATCTTGAGAACT GGAGTCGACCGGCATC CCTGGGTAGGGGCTGCTTCTAAGGAGAAG AGCTCCAGAGG

Exons 10-12 probe hybridizes to *c-kit* extracellular domain coding area. Exons 18-20 probe hybridizes to *c-kit* intracellular domain coding area.

(Paloalto, CA, USA) in RACE analysis. The full-length cDNAs was made by joint action of the SMARTer II A Oligonucleotide and SMARTScribe Reverse Transcriptase (a variant of MMLV RT) in reverse transcription reactions. The first strand of cDNA synthesis was obtained from 1 μ g total RNA. PCR amplification was done with specific primers hit exons 10–12 and exons 18–21 on the full-length *c-kit* transcript (Table 2) in conjunction with universal primers that were provided in the kit. Advantage 2 PCR kit (Clontech, Paloalto, CA, USA) was used for the 5' and 3' PCR amplification. Nested PCR and touchdown PCR were used to safeguard the specificity of the amplification. Electrophoresis of the PCR products, bands cutting and gel extraction (QIAquick gel extraction kit; QIAGEN, Valencia, CA, USA) were performed. All of the clear RACE PCR product gel extractions were cloned to TA vector (TOPO TA cloning kit for sequencing, Invitrogen,

Table 2 RACE primers

Name	5' or 3'	Sequence (5' → 3')	No. of bases	Exons hitting	Position on NM_021099
e11	5'	CAGCCTGTTTCTGGG AAACTCCCATTG	27	Exon 11	1825-1798
e12	5'	GCAACTGTCATGGC AGCATCCGACTT	26	Exon 12	1920-1895
e18	5'	TGCTCTCTGGTGCCA TCCACTTCAC	25	Exon 18	2552-2756
e20	5'A	GGTCAGCGTCCCAG CAAGTCTTCAT	25	Exon 20	2786-2762
e20	5'B	AAGGGGTCAGCGTC CCAGCAAGTCT	25	Exon 20	2790-2766
e20	5'C	TGCTTGGTGCTGTCC GAGATCTGCT	25	Exon 20	2856-2832
e21	5'	GGGGTTGCAGTTTG CCAAGTTGGAG	25	Exon 21	2887-2863
e10	3'	AAATCCAGGCCAC ACTCTGTTCCAGC	26	Exon 10	1602-1627
e11	3'	TGGGAGTTTCCCAG AAACAGGCTGAG	26	Exon 11	1802-1827
e18	3'	CCGTGAAGTGATG GCACCAAGAGAG	25	Exon 18	2550-2574
e19	3'A	AGGAAGCAGCCCT ACCCAGGGATG	25	Exon 19	2650-2674
e19	3'B	GGGATGCCGGTTCGA CTCCAAGTTCT	25	Exon 19	2669-2693
e20	3'A	TGACCCCTTGA AAA GGCCAACATTCA	26	Exon 20	2782-2807
e20	3'B	GCAGATCTCGGACA GCACCAAGCAC	25	Exon 20	2833-2857

Requirement of a good gene specific primer for RACE: It should be 23-28 nt, has 50-70% GC and Tm > 70°C, and does not complement to the 3' of the Universal Primer Mix.

Carlsbad, CA, USA) and sent to commercial company for sequencing. 5' and 3' RACE results were combined to obtain the full-length *c-kit* transcripts sequence information.

Quantitative real-time RT-PCR

Total RNA (2 µg) was treated with DNase I (Sigma, Saint Louis, USA) for 15 minutes at room temperature and then reversely transcribed by High Capacity cDNA Reverse Transcription Kit (Applied Biosystems, Foster City, CA, USA).

Real-time RT-PCR analysis of *c-kit* was performed with Taqman universal PCR master mix and Taqman gene expression assays on the ABI Prism 7900HT Real Time PCR System, according to the manufacturer's instructions (Applied Biosystems, Foster City, CA, USA). The relative expression level of each target gene was calculated by the comparative CT method and was normalized to glyceraldehyde 3-phosphate dehydrogenase (GAPDH) expression. Three *c-kit* gene-specific probes that hit different parts of the full-length transcript (exon 7–8, exon 20–21 and exon 21⁺) were used.

Real-time RT-PCR analysis for other genes were performed with Power SYBR PCR master mix and gene specific primers on the ABI Prism 7900HT Real Time PCR System, according to the manufacturer's instructions (Applied Biosystems, Foster City, CA, USA). The relative expression level of each target gene was calculated by the comparative CT method and was normalized to GAPDH expression. The primers of the candidate genes are list in Table 3.

Each RT-PCR analysis was repeated 3 times after GAPDH normalization.

Western blot

Cells and testis tissues were lysed on ice in RIPA buffer containing 1% freshly added protease inhibitors. Protein electrophoresis and gel blotting were performed with NuPAGE electrophoresis system (Invitrogen, Carlsbad, CA, USA) following the manufacturer's instructions. The blotted PCDF membranes were blocked with 5% (wt/vol) non-fat dry milk (RT, 60 minutes) and probed for Kit at 4°C overnight, using either 1 µg/ml of a monoclonal antibody (rat anti-mouse; NOVUS, Littleton, CO, USA) directed against the extracellular domain of the Kit or a polyclonal antibody (rabbit anti-human, mouse, rat; NOVUS, Littleton, CO, USA) directed against the amino acid near S715 of the human Kit (1 µg/ml) followed by the HRP-conjugated secondary antibodies (Santa Cruz, Santa Cruz, CA, USA) staining.

Protein lysate from Kit expressed in human megakaryoblast cell lines (ATCC no. CRL-2021) was set up as positive control and protein lysate from Kit negative mouse myoblast cell line (ATCC no. CRL-1772) was set up as negative control. The same blot was stripped and re-probed with mouse beta-actin primary antibody (Santa Cruz, Santa Cruz, CA, USA) as internal control.

Immunofluorescence staining of cells

The cover slips with the cells were washed 3 times with PBS again and were incubated with 5% normal goat serum (Santa Cruz, Santa Cruz, CA, USA) in PBS for

Table 3 Gene specific primers of the candidate genes

Name	Forward	Reverse
BMP4	TTCTGGTAACCGAATGCTGA	CCTGAATCTCGGCGACTTTTT
Cyp26b1	GCAAGATCTACTGGGCGAAC	TTGGGCGAGTAGCTCTCAAGT
DAZL	GTCCTTACATGTACCATT TGTGAC	GACTCCAACAAAACAGC AGACAA
EGR 3	AGCTGAACTGGGCTGTGTCT	AATGGGGAGTGGGTATGTGA
Kitl	TCTGCGGAATCCTGTGACT	TGGAAGATTGCCACCAGTTT
PLZF	GCAAGAACAGCGTCAAGACA	TGGGATCACGTGAAGCTATG
RARα	TCCGAAGAGATAGTACCCAGC	AAAGCAAGGCTTGTAGATGCC
Stra8	GTTTCTGCGTGTCCACAAG	CACCCGAGGCTCAAGCTTC

30 minutes before being incubated with the primary antibody overnight at 4°C. The cells were then incubated with the secondary antibody and mounted with UltraCruz™ Mounting Medium with DAPI (SantaCruz, Santa Cruz, CA, USA). The following antibodies were used in this study: the FITC monoclonal rat-anti-mouse Kit extracellular domain (1:200, 105805, BioLegend, San Diego, CA, USA), the monoclonal rat-anti-mouse Kit extracellular domain (1:200, NBP1-43359, NOVUS, Littleton, CO, USA), the monoclonal rat-anti-mouse Kit extracellular domain (1:100, KJ-14, Santa Cruz, Santa Cruz, CA, USA); the polyclonal rabbit-anti-human/mouse/rat Kit intracellular domain (1:200, NBP1-19865, NOVUS, Littleton, CO, USA), the polyclonal goat-anti-mouse Kit C-terminus (1:100, M14, Santa Cruz, USA) and the polyclonal rabbit-anti-human/mouse Kit C-terminus (1:100, C19, Santa Cruz, Santa Cruz, CA, USA). The secondary antibodies used in this study included: the Alexa 488-conjugated goat-anti-rat IgG (1:500; Invitrogen, Carlsbad, CA, USA); the Alexa 594-conjugated goat-anti-rabbit IgG (1:500; Invitrogen, Carlsbad, CA, USA) and the Texas red-conjugated donkey-anti-goat IgG (1:100; Santa Cruz, Santa Cruz, CA, USA).

Statistical analysis

Statistical analysis was performed by unpaired two-tail student t test using SPSS software (Version 17.0). All experiments were performed for at least three independent times and a P value of less than 0.05 was considered statistically significant.

Results

Transcription of *c-kit* in cell lines and testes

Northern blots revealed at least 4 transcripts in the cells and testes (Figure 1B and C). Even more *c-kit* transcripts expressed in c18-4, CRL-2053 and 5 dpp, 10 dpp and 60 dpp testes were assayed by RACE (Figure 2A, B, C, D). The 1.5 kb transcript expressed in the c18-4 cells was not shown by RACE (Figure 1C). Four representative transcripts (Type A, B, C and D) were illustrated in Figure 3 and a multiple blast of their sequences was shown in Additional file 1. Quantitative expression discrepancies (either in the 5' end or in the 3' end) of *c-kit* among the cell lines and testes of different stages existed. Multiple blast assays demonstrated that exons 17–21 region was highly conserved in the *c-kit* transcripts. The four representative transcripts included the full-length canonical transcript (transcript A, 5.1 kb, expressed in 10 and 60 dpp testes), the 3' end short transcript (transcript B, 3.9 kb, expressed in c18-4, CRL-2053, 5 and 10 dpp testes), the short transcript (transcript C, 3.4 kb, expressed in CRL-2053) and the truncated transcripts (transcripts D, 1.9-3.2 kb, no expression detected by Northern blot analysis) respectively (Figure 3).




Figure 1 *c-kit* northern hybridization. (A) RNA electrophoresis. 10 µg total RNA from different samples were loaded. RNA sizes were marked with 2 µg (2 µl) RNA Millennium size markers (Ambion). All RNA samples were in good qualities. (B) Northern hybridization with *c-kit* probe hit exons 10-12. Short transcripts (purple arrowhead) with a size between 4 and 5 kb could be observed in c18-4, CRL-2053 and different aged testes. Long transcripts (blue arrowhead) with a size between 5 and 6 kb were seen in 10 dpp and 60 dpp testes. (C) Northern hybridization with *c-kit* probe hit exons 18-20. A 1.5 kb short transcript (red arrowhead) was observed in c18-4 and a 2.7 kb short transcript (green arrowhead) was observed in CRL-2053. Short transcripts (purple arrowhead) with a size between 4 and 5 kb could be observed in 5 dpp, 10 dpp and 60 dpp testes. Long transcripts (blue arrowhead) with a size between 5 and 6 kb were seen in 10 dpp and 60 dpp testes. (D) Northern hybridization of beta-actin (internal control).

Quantitative analysis of different transcript expressions in testes and germ cell lines

Three sets of specific primers for RT-PCR were designed to detect *c-kit* transcripts (e7-8, e20-21 and e21⁺ as indicated by triangles in Figure 3). Products from the e7-8 primers represented the long transcripts (transcripts A + B). Products from the e20-21 primers represented the total transcripts (transcripts A + B + C + D). The ratio of the long transcripts could be calculated as (e7-8/e20-21) × 100%.





Figure 2 RACE PCR products gel electrophoresis. (A) 5' and 3' RACE with primer sets hit exons 10-12. c18-4 had two 5' ends (1.8 kb and 1.4 kb). CRL-2053 and 60 dpp testes had one 5' end (1.8 kb), c18-4 had one 3 kb 3' end (3 kb). CRL-2053 and 60 dpp testes had two 3' end (3 kb and 1.8 kb). **(B)** 5' and 3' RACE with primer sets hit exons 10-12. 5 dpp and 10 dpp testis had one 5' ends (2 kb). 5 dpp and 10 dpp testes had two 3' end (3 kb and 2.1 kb). CRL-2053 and 60 dpp testes had two 3' end (3 kb and 1.8 kb). **(C)** 5' and 3' RACE with primer sets hit exons 18-21. c18-4 had two 5' ends (2.9 kb and 1.5 kb). CRL-2053 had three 5' ends (2.9 kb, 1.4 kb and 1.6 kb). 5 dpp, 10 dpp and 60 dpp testes had the same one 5' end (2.9 kb); c18-4 had one 3' end (2.9 kb). CRL-2053 5 dpp and 10 dpp and 60 dpp testes had two 3' end (3 kb and 1.8 kb). **(D)** Summary of c-kit RACE PCR fragments from cell lines and testis.

The ratio of the short transcripts (Transcript C + D) was consequently $(1 - \text{ratio}_{\text{long transcripts}}) \times 100\%$. Those produced by e21⁺ primers represented all transcripts with a long 3' end except for transcript B (Transcripts A + C + D). The ratio of the long 3' UTR transcripts was $(e21^+/e20-21) \times 100\%$. The ratio of the short 3' UTR transcripts was $(1 - \text{ratio}_{\text{long 3' UTR transcripts}}) \times 100\%$.

From 5 dpp to 10 dpp, expressions of both transcripts A + B and transcripts C + D increased significantly (Figure 4A, B). The ratios of long and short transcripts remained constant (22.47% in 5 dpp and 22.45% in 10 dpp, Figure 4B). The ratio of long 3' transcripts jumped

from 2.8% to 4.01% (Figure 4C). From 10 dpp to 60 dpp, transcripts A + B + C + D increased moderately but transcripts A + C + D decreased. Hence, it can be deduced that transcript B must have been enhanced significantly (Figure 4A). Consequently, since the relative amount of transcripts A + B did not change, the quantity of transcript A must have declined dramatically. The ratio of short transcripts progressed from 22.45% in 10 dpp to 54.91% in 60 dpp (Figure 4B). Long 3' transcripts dropped from 4.01% (10 dpp) to 0.60% (60 dpp) (Figure 4C).

Primers e21⁺ did not detect any bands in all three types of cell lines. As a result, all transcripts in these cell lines



lacked of the 3' UTR ends. The relative quantity of transcript A was equivalent with transcript B in the cell lines. The expression of type A + B multiplied significantly in CRL-2053 (about 20 folds of that in c18-4) and then diminished (about 5 folds of that in c18-4) in CRL-2196 (Figure 5A). The ratio of type A + B transcripts was 65%, 92.9% and 56.4% in c-18-4, CRL-2053 and CRL-2196 respectively (Figure 5B).


Translation of *c-kit* in the testes and male germ cell lines

It was shown that full-length Kit (145 KDa) was expressed in CRL-2053 cells and testes except in c18-4 cells (Figure 6A). With an anti-intracellular antibody, a 50 KDa Kit was expressed in the c18-4, CRL-2053 and all testes (Figure 6B). From 5 dpp to 60 dpp, the 145 KDa Kit increased, but the 50 KDa Kit remained homogenous in the testes (Figure 6). Both the c18-4 and CRL-2053 cells expressed the intracellular domain of Kit (nuclear region) (Figure 7B and D). Unlike the CRL-2053 cells, the c18-4 cells did not express the extracellular domain (membrane region) (Figure 7A). In the 5 dpp testes, expression of the extracellular domain was very minimal in the germ cells (Figure 8A). In the 10 dpp testes, a portion of the germ cells adjacent to the basement membrane of the seminiferous tubules began to express the membrane domain

(Figure 8B). In the 60 dpp testes, some spermatogonia and spermatocytes were tested positive for membrane domain (Figure 8C). In contrast, the nuclear domain was expressed in all stages (5 dpp, 10 dpp and 60 dpp) in the spermatogonia, spermatocyte and spermatids, but not in the mature spermatozoon (Figure 8D, E and F). Leydig cells highly expressed the nuclear domain of Kit in the 60 dpp testes (Figure 8F).

Expression changes of *c-kit* and other differentiation-related genes in the testes after RA stimulation


5 dpp, 10 dpp and 60 dpp mouse testes were treated with either 0.7 μ M or 2 μ M RA *in vitro* for 24 h. Quantitative expression of *c-kit* and SSCs differentiation related genes were determined by Real-time PCR using three pairs of primers (e7-8, e20-21 and e21⁺ as indicated by arrow heads in Figure 3). The total *c-kit* mRNA level increased following the RA treatment and exhibited a concentration-dependent pattern in 5 dpp and 10 dpp testes (Figure 9A, C). After RA stimulation, the 60 dpp testes did not display concentration-dependent increases any more (Figure 9E). In 5 dpp and 10 dpp testes, expressions of *Cyp26b1* and *Stra8* were significantly up-regulated (Figure 9B, D). Expressions of *Dazl* and *Kitl* were enhanced moderately in 5 dpp and 60 dpp testes




(Figure 9B, F). *Bmp4* had a different response to RA treatment in testes of different ages (Figure 9B, D, F). The effects of RA on the expressions of *Bmp4*, *Cyp26b1* and *Stra8* were more significant in the 5 dpp testes than that in the 10 dpp testes. Expressions of *RARα* and *Egr3* were not altered in either stage (Figure 9B, D, F).


Expression dynamics of *c-kit* and differentiation-related genes in germ cell lines after RA stimulation

c-kit and other differentiation-related genes changes were analyzed in C18-4 and CRL-2053 cells after RA treatment (2 μM RA for 24 hours). Unlike in the testes, the expressions of all transcripts declined in the c18-4 cells after RA stimulation (Figure 10A). In CRL-2053 cells, the amount of the long transcripts (e7-8 in Figure 10B) declined to about 50% but the quantity of total transcripts (e20-21 in Figure 10B) escalated to approximately 2 folds,



indicating that the short transcripts must have increased even more significantly (Figure 10B). Expressions of *Cyp26b1* and *Stra8* increased significantly in both c18-4 and CRL-2053 cells ($P < 0.01$) (Figure 10C, 10D). The two germ cell marker genes, *Dazl* and *Pou5f1* had distinctive responses to RA. *Dazl* was boosted in 5 dpp and 60 dpp testes (Figure 9B, F), but not in either cell lines (Figure 10C, D). *Pou5f1* diminished in c18-4 cells but did





not change in CRL-2053 cells (Figure 10C and D). *Egr3* (an early growth response gene) was suppressed in c18-4 and CRL-2053 cells (Figure 10C, 10D), but it was not altered in the testes (Figure 9B, D, F). *Rara* (an RA receptor gene) did not respond to RA stimulation.


Discussion

A. *c-kit* transcripts during spermatogenesis

In addition of the conventional full-length *c-kit* and Tr-*kit* discovered, we also found more than 3*c-kit* mRNA transcripts in the SSCs and spermatogonia. CRL-2053 had the highest amount of transcript A (Figure 5A). Though its quantity increased, the percentages of Transcript A declined in CRL-2196 and 60 dpp testes when compared with that in c18-4 cells and testes at 5 dpp (Figure 4A, B, Figure 5A, B). The 145 KDa Kit was also absent in c18-4 cells (Figure 6A). On this basis, acquisition of this transcript marked the start of the transition from SSCs to spermatogonia. The percentage of this transcript decreased in CRL-2196 cells and 60 dpp testes. This was caused by the emergence of new short transcripts, which were important for later stage spermatogenesis.

Expression of transcript B was the highest in the 10 dpp testis (Figure 4A). The 3.9 kb short 3' UTR transcript was composed of 21 exons, identical to the full-length transcript. The only difference was that the 3.9 kb short 3' UTR transcript had a 1.2 kb shorter 3' UTR than the full-length transcript. Combining its abundance in the testes (95.9% ~ 99.4%, Figure 4C) with the strong positive staining of Leydig cells in the 60 dpp testes (Figure 8F), it could be inferred that this transcript might be a somatic form. Functions of 3' UTR included supplying binding sites for microRNAs and post-transcriptional regulation. Absence of the 3' end UTR in all transcripts in the immortal germ cells (Figure 5A) indicated that the 3' UTR modification was lost during immortalization and it might be controlled by testicular somatic factors.

Transcript C, encoding the 50 KDa Kit, was stably expressed in the testes (Figure 6B). The percentage of the short transcripts (Transcript C + D) was the highest in the 60 dpp testes (54.91%) (Figure 4B) and CRL-2196 (43.6%) (Figure 5B). Consequently, these transcripts might have significant roles in later stage spermatogenesis beginning from spermatocytes. Multiple sequence



alignment of *c-kit* transcripts was shown in Additional file 1.

B. Kit profile during spermatogenesis

Two forms of Kit were discovered in this study: the 145 kDa and 50 kDa Kit. The 145 kDa Kit was located in the cytoplasm/membrane domain in CRL-2053 cells (Figure 7C) but not in c18-4 cells. Its expression escalated accordingly in the 5 dpp, 10 dpp, 15 dpp, 40 dpp and 60 dpp testes (Figure 6A). Unlike the 145 kDa full-length Kit, the 50 kDa Kit, possibly the product of transcript C, was expressed in both nuclear and cytoplasm/membrane domains in CRL-2053 cells (Figure 6). The

50 kDa Kit was stably expressed in the testes (Figure 6B). Therefore, the 145 kDa was indeed the marker for spermatogonia [43]. We highlighted here that its location shifted from the nucleus to the cytoplasm and then to the membrane domain. This might be vital for the initiation of spermatogenesis in SSCs. Expression of the extracellular full-length Kit on membranes, not in the nucleus or in the cytoplasm, endorsed the cells the ability to correspond with Kitl signals. Hence, this expression played important roles in differentiation initiation. We also demonstrated here that SSCs did not have full-length transcript (transcript A), nor the full-length Kit (145 kDa), which agreed with other studies indicating




Figure 9 Expression of *c-kit* and other germ cell differentiation-related genes in the 5 dpp, 10 dpp and 60 dpp mouse testes stimulated by RA. 5 dpp, 10 dpp and 60 dpp testes were treated with 2 μM RA diluted in ethanol for 24 h *in vitro*. Testes treated with the same concentration of ethanol without RA were the control group. Realtime PCR was carried out for quantitative determination of the expression of *c-kit* and its potential regulatory genes (BMP4, Cyp26b1, DAZL, EGR3, Kitl, RARα and Stra8). Values of the vertical axis represented the expression fold change comparing with the control group. The results were normalized to GAPDH values. Error bars represent the S.E.M. Values with ** represented a significance with a $P < 0.01$ whereas values with * represented a significance with a $P < 0.05$ comparing with the control group. **(A)** *c-kit* expression in the 5 dpp testes. **(B)** Expression of germ cell differentiation-related genes in the 5 dpp testes. **(C)** *c-kit* expression in the 10 dpp testes. **(D)** Expression of germ cell differentiation-related genes in the 10 dpp testes. **(E)** *c-kit* expression in the 60 dpp testes. **(F)** Expression of germ cell differentiation-related genes in the 60 dpp testes.

that the activation of the Kit/Kitl signaling pathway was not required for SSCs' self-renewal [23,24]. Kit was initially expressed in the nucleus, and then ventured out to the cytoplasm and then to the membrane domain when SSCs became spermatogonia. ORF finder comparison of the *c-kit* putative proteins sequences were shown in Additional file 2.

C. RA responses in germ cells and testes

Some studies showed that RA directly acts on spermatogenic cells by stimulating *Stra8* and *c-kit* gene expression, whereas some studies testified that exogenous RA

could not stimulate *c-kit* expression [16,42,44]. We demonstrated that RA enhanced *c-kit* expression (Figure 9A, C and E) in testes. *Stra8* (gene stimulated by RA) was also significantly amplified in both cell lines and 5 dpp/10 dpp testes (Figure 9B, D). However, the expression pattern of *c-kit* in the cell lines was different from that in the testes (Figure 10 vs Figure 9). Both long and short transcripts were reduced in c18-4 cells after RA stimulation (Figure 10A). On the other hand, the long transcripts were reduced whereas the short transcripts were promoted in CRL-2053 cells (Figure 10B). We concluded that RA indirectly impacted upon *c-kit* expression in




Figure 10 Expression of *c-kit* and other germ cell differentiation-related genes in the c18-4 and CRL-2053 cells stimulated by RA.

The c18-4 and CRL-2053 cells were treated with 2 μM RA diluted in ethanol for 24 h *in vitro*. Cells treated with the same concentration of ethanol without RA were the control group. Realtime PCR was carried out for quantitative determination of the expression of *c-kit* and its potential regulatory genes (*Bmp4*, *Cyp26b1*, *Dazl*, *Egr3*, *Pou5f1* and *Stra8*). Values of the vertical axis represented the expression fold change comparing with the control group. The results were normalized to GAPDH values. Error bars represent the S.E.M. Values with ** represented a significance with a $P < 0.01$ whereas values with * represented a significance with a $P < 0.05$ comparing with the control group. (A) *c-kit* expression in the c18-4 cells. (B) *c-kit* expression in the CRL-2053 cells. (C) Expression of germ cell differentiation-related genes in the c18-4 cells. (D) Expression of germ cell differentiation-related genes in the CRL-2053 cells.

male germ cells, while some unknown factors from the testes somatic cells might be involved. It agreed with previous works that RA indirectly controlled the timing of meiosis by juxtacrine of Sertoli cells [45].

Suppression of *Bmp4* by RA was obvious. RA reduced *Bmp4* (a SSCs pluripotential maintenance gene) expression in 5 dpp testes (Figure 9B) and did not alter *Bmp4* expression in 10 dpp and 60 dpp testes (Figure 9D, F). *BMP4* was also reduced in c18-4 cells after RA stimulation (Figure 10C). Excessive exogenous RA would push SSCs into abnormal differentiation and finally apoptosis [16]. Our results validated that after 24 hours of 2 μM RA treatment, expression of *Cyp26b1* (a RA degradation gene) was stimulated. This would degrade excessive RA into the inactive form in both cell lines and testes (Figure 9B, C, Figure 10C, D). *Cyp26b1* was not altered in 60 dpp testes (Figure 9F). The increase of the protective gene also varied in CRL-2053 cells (~20 folds increase of *Cyp26b1*) and in c18-4 cells (~6 folds increase of *Cyp26b1*). We confirmed here, again, that *Stra8* was the most immediate responsive gene after RA stimulation. The germ cell marker genes (*Dazl* and *Pou5f1*), early growth response gene (*Egr3*), and the

RA receptor gene (*Rara*) did not respond to RA, especially not when RA was added to testes tissue culture.

Conclusions

There are dynamic transcription and translation changes of *c-kit* before and after SSCs' anticipated differentiation. These changes differ between in the cell lines and in the testis. The responses to RA stimulation are different between the cell lines and testis too. As a significant upstream regulatory factor for *c-kit* expression, RA might play with other unknown factors to precisely regulate the expression profiles of *c-kit* in order to regulate normal spermatogenesis.

Additional files

Additional file 1: Multiple sequence alignment of *c-kit* transcripts.

Additional file 2: Multiple sequence alignment of ORF finder predicted *c-kit* proteins.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

LZ and YH carried out the molecular genetic studies, participated in the RACE, Northern blot, sequence alignment and drafted the manuscript. JT carried out the immunoassays. CH and HF participated in the design of the study. LL participated online mRNA and protein prediction analysis. XT performed the statistical analysis. All authors read and approved the final manuscript.

Acknowledgement

This study is funded by Hong Kong RGC grant (464809) to YB Han, Key project No. 12JC1407602 by Science and Technology commission of Shanghai municipality to YB Han.

Author details

¹Shanghai first maternity and infant health hospital, Tongji University, Shanghai, China. ²Department of Obstetrics and Gynaecology, Prince of Wales hospital, The Chinese University of Hong Kong, Hong Kong, China. ³Diagnosics and Reproductive Health Center BGI, Beishan Industrial ZoneYantian District, Shenzhen 518083, China. ⁴Guangzhou Institute of Biomedicine and Health Chinese Academy of Sciences, Guangzhou, China. ⁵Department of Obstetrics and Gynecology, New York University Medical College, Manhasset, NY, USA.

Received: 8 May 2013 Accepted: 16 October 2013

Published: 27 October 2013

References

- Barroca V, Lassalle B, Coureuil M, Louis JP, Le PF, Testart J, Allemand I, Riou L, Fouchet P: **Mouse differentiating spermatogonia can generate germinal stem cells in vivo.** *Nat Cell Biol* 2009, **11**:190–196.
- Schrans Stassen BH, De Kant HJ V, De Rooij DG, Van PAM: **Differential expression of c-kit in mouse undifferentiated and differentiating type A spermatogonia.** *Endocrinology* 1999, **140**:5894–5900.
- Chabot B, Stephenson DA, Chapman VM, Besmer P, Bernstein A: **The proto-oncogene c-kit encoding a transmembrane tyrosine kinase receptor maps to the mouse W locus.** *Nature* 1988, **335**:88–89.
- Yarden Y, Kuang WJ, Yang Feng T, Coussens L, Munemitsu S, Dull TJ, Chen E, Schlessinger J, Francke U, Ullrich A: **Human proto-oncogene c-kit: a new cell surface receptor tyrosine kinase for an unidentified ligand.** *EMBO J* 1987, **6**:3341–3351.
- Blechman JM, Lev S, Barg J, Eisenstein M, Vaks B, Vogel Z, Givol D, Yarden Y: **The fourth immunoglobulin domain of the stem cell factor receptor couples ligand binding to signal transduction.** *Cell* 1995, **80**:103–113.
- Blechman JM, Lev S, Givol D, Yarden Y: **Structure-function analyses of the kit receptor for the steel factor.** *Stem Cells* 1993, **11**(Suppl 2):12–21.
- Albanesi C, Geremia R, Giorgio M, Dolci S, Sette C, Rossi P: **A cell- and developmental stage-specific promoter drives the expression of a truncated c-kit protein during mouse spermatid elongation.** *Development* 1996, **122**:1291–1302.
- Dym M, Jia MC, Dirami G, et al: **Expression of c-kit receptor and its autophosphorylation in immature rat type A spermatogonia.** *Biol Reprod* 1995, **52**(1):8–19.
- Gu Y, Runyan C, Shoemaker A, Surani A, Wylie C: **Steel factor controls primordial germ cell survival and motility from the time of their specification in the allantois, and provides a continuous niche throughout their migration.** *Development* 2009, **136**(8):1295–1303.
- Guerif F, Cadoret V, Rahal-Perola V, et al: **Apoptosis, onset and maintenance of spermatogenesis: evidence for the involvement of Kit in Kit-haplodeficient mice.** *Biol Reprod* 2002, **67**(1):70–79.
- Ohta H, Yomogida K, Dohmae K, Nishimune Y: **Regulation of proliferation and differentiation in spermatogonial stem cells: the role of c-kit and its ligand SCF.** *Development* 2000, **127**(10):2125–2131.
- Orth JM, Qiu J, Jester WF Jr, Pilder S: **Expression of the c-kit gene is critical for migration of neonatal rat gonocytes in vitro.** *Biol Reprod* 1997, **57**(3):676–683.
- Packer AI, Besmer P, Bachvarova RF: **Kit ligand mediates survival of type A spermatogonia and dividing spermatocytes in postnatal mouse testes.** *Mol Reprod Dev* 1995, **42**(3):303–310.
- Prabhu SM, Meistrich ML, McLaughlin EA, Roman SD, Warne S, Mendis S, Itman C, Loveland KL: **Expression of c-Kit receptor mRNA and protein in the developing, adult and irradiated rodent testis.** *Reproduction* 2006, **131**:489–499.
- Runyan C, Schaible K, Molyneux K, Wang Z, Levin L, Wylie C: **Steel factor controls midline cell death of primordial germ cells and is essential for their normal proliferation and migration.** *Development* 2006, **133**(24):4861–4869.
- Snyder EM, Davis JC, Zhou Q, Evanoff R, Griswold MD: **Exposure to Retinoic Acid in the Neonatal but Not Adult Mouse Results in Synchronous Spermatogenesis.** *Biol Reprod* 2011, **84**(5):886–893.
- Yan W, Suominen J, Toppari J: **Stem cell factor protects germ cells from apoptosis in vitro.** *J Cell Sci* 2000, **113**(Pt 1):161–168.
- Yoshinaga K, Nishikawa S, Ogawa M, Hayashi S, Kunisada T, Fujimoto T, Nishikawa S: **Role of c-kit in mouse spermatogenesis: identification of spermatogonia as a specific site of c-kit expression and function.** *Development* 1991, **113**:689–699.
- Sorrentino V, Giorgi M, Geremia R, Besmer P, Rossi P: **Expression of the c-kit proto-oncogene in the murine male germ cells.** *Oncogene* 1991, **6**:149–151.
- Rossi P, Marziali G, Albanesi C, Charlesworth A, Geremia R, Sorrentino V: **A novel c-kit transcript, potentially encoding a truncated receptor, originates within a kit gene intron in mouse spermatids.** *Dev Biol* 1992, **152**:203–207.
- Sette C, Bevilacqua A, Bianchini A, Mangia F, Geremia R, Rossi P: **Parthenogenetic activation of mouse eggs by microinjection of a truncated c-kit tyrosine kinase present in spermatozoa.** *Development* 1997, **124**:2267–2274.
- Sette C, Dolci S, Geremia R, Rossi P: **The role of stem cell factor and of alternative c-kit gene products in the establishment, maintenance and function of germ cells.** *Int J Dev Biol* 2000, **44**:599–608.
- Kubota H, Avarbock MR, Schmidt JA, Brinster RL: **Spermatogonial stem cells derived from infertile Wv/Wv mice self-renew in vitro and generate progeny following transplantation.** *Biol Reprod* 2009, **81**:293–301.
- Morimoto H, Kanatsu Shinohara M, Takashima S, Chuma S, Nakatsujii N, Takehashi M, Shinohara T: **Phenotypic plasticity of mouse spermatogonial stem cells.** *PLoS One* 2009, **4**:e7909.
- Nakagawa T, Sharma M, Nabeshima Y, Braun RE, Yoshida S: **Functional hierarchy and reversibility within the murine spermatogenic stem cell compartment.** *Science* 2010, **328**:62–67.
- Shinohara T, Orwig KE, Avarbock MR, Brinster RL: **Spermatogonial stem cell enrichment by multiparameter selection of mouse testis cells.** *Proc Natl Acad Sci U S A* 2000, **97**:8346–8351.
- Suzuki H, Sada A, Yoshida S, Saga Y: **The heterogeneity of spermatogonia is revealed by their topology and expression of marker proteins including the germ cell-specific proteins Nanos2 and Nanos3.** *Dev Biol* 2009, **336**:222–231.
- Yoshida S, Sukeo M, Nakagawa T, Ohbo K, Nagamatsu G, Suda T, Nabeshima Y: **The first round of mouse spermatogenesis is a distinctive program that lacks the self-renewing spermatogonia stage.** *Development* 2006, **133**:1495–1505.
- Heim C, Minniear K, Dann CT: **Imatinib has deleterious effects on differentiating spermatogonia while sparing spermatogonial stem cell self renewal.** *Reprod Toxicol* 2011, **31**:454–463.
- Trefil P, Bakst MR, Yan H, Hejnar J, Kalina J, Mucksova J: **Restoration of spermatogenesis after transplantation of c-Kit positive testicular cells in the fowl.** *Theriogenology* 2010, **74**:1670–1676.
- Izadyar F, Pau F, Marh J, Slepko N, Wang T, Gonzalez R, Ramos T, Howerton K, Sayre C, Silva F: **Generation of multipotent cell lines from a distinct population of male germ line stem cells.** *Reproduction* 2008, **135**:771–784.
- Jones Villeneuve EM, McBurney MW, Rogers KA, Kalnins VI: **Retinoic acid induces embryonal carcinoma cells to differentiate into neurons and glial cells.** *J Cell Biol* 1982, **94**:253–262.
- Livera G, Rouiller-Fabre V, Pairault C, Levacher C, Habert R: **Regulation and perturbation of testicular functions by vitamin A.** *Reproduction* 2002, **124**:173–180.
- Bowles J, Knight D, Smith C, Wilhelm D, Richman J, Mamiya S, Yashiro K, Chawengsaksophak K, Wilson MJ, Rossant J, et al: **Retinoid signaling determines germ cell fate in mice.** *Science* 2006, **312**:596–600.
- Koubova J, Menke DB, Zhou Q, Capel B, Griswold MD, Page DC: **Retinoic acid regulates sex-specific timing of meiotic initiation in mice.** *Proc Natl Acad Sci U S A* 2006, **103**:2474–2479.
- Morales C, Griswold MD: **Retinol-induced stage synchronization in seminiferous tubules of the rat.** *Endocrinology* 1987, **121**(1):432–434.

37. Van PAM, De Rooij DG: Retinoic acid is able to reinitiate spermatogenesis in vitamin A-deficient rats and high replicate doses support the full development of spermatogenic cells. *Endocrinology* 1991, **128**:697–704.
38. Hofmann MC, Braydich-Stolle L, Dettin L, Johnson E, Dym M: Immortalization of mouse germ line stem cells. *Stem Cells* 2005, **23**:200–210.
39. Hofmann MC, Narisawa S, Hess RA, Millan JL: Immortalization of germ cells and somatic testicular cells using the SV40 large T antigen. *Exp Cell Res* 1992, **201**:417–435.
40. Hofmann MC, Abramian D, Millan JL: A haploid and a diploid cell coexist in an in vitro immortalized spermatogenic cell line. *Dev Genet* 1995, **16**:119–127.
41. Hofmann MC, Hess RA, Goldberg E, Millan JL: Immortalized germ cells undergo meiosis in vitro. *Proc Natl Acad Sci U S A* 1994, **91**:5533–5537.
42. Zhou Q, Li Y, Nie R, Friel P, Mitchell D, Evanoff RM, Pouchnik D, Banasik B, McCarrey JR, Small C, *et al*: Expression of stimulated by retinoic acid gene 8 (Stra8) and maturation of murine gonocytes and spermatogonia induced by retinoic acid in vitro. *Biol Reprod* 2008, **78**:537–545.
43. Shinohara T, Avarbock MR, Brinster RL: beta1- and alpha6-integrin are surface markers on mouse spermatogonial stem cells. *Proc Natl Acad Sci U S A* 1999, **96**:5504–5509.
44. Wang Y, Culty M: Identification and distribution of a novel platelet-derived growth factor receptor beta variant: effect of retinoic acid and involvement in cell differentiation. *Endocrinology* 2007, **148**(5):2233–2250.
45. Pellegrini M, Filipponi D, Gori M, *et al*: ATRA and KL promote differentiation toward the meiotic program of male germ cells. *Cell Cycle* 2008, **7**(24):3878–3888.

doi:10.1186/1471-213X-13-38

Cite this article as: Zhang *et al.*: c-kit expression profile and regulatory factors during spermatogonial stem cell differentiation. *BMC Developmental Biology* 2013 **13**:38.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

