

Review Article

Vitamin C versus Cancer: Ascorbic Acid Radical and Impairment of Mitochondrial Respiration?

Rumiana Bakalova ^{1,2}, Zhivko Zhelev,^{3,4} Thomas Miller,⁵ Ichio Aoki,^{1,2} and Tatsuya Higashi ¹

¹Department of Molecular Imaging and Theranostics, National Institute of Radiological Sciences (NIRS), National Institutes for Quantum and Radiological Science and Technology (QST), Chiba, Japan

²Group of Quantum-State Controlled MRI, Institute for Quantum Life Science, National Institutes for Quantum and Radiological Science and Technology (QST), Chiba, Japan

³Institute of Biophysics and Biomedical Engineering, Bulgarian Academy of Sciences, Sofia, Bulgaria

⁴Medical Faculty, Trakia University, Stara Zagora, Bulgaria

⁵IC-MedTech Co., San Diego, CA, USA

Correspondence should be addressed to Rumiana Bakalova; bakalova.rumiana@qst.go.jp

Received 21 July 2019; Revised 14 September 2019; Accepted 28 September 2019; Published 10 January 2020

Academic Editor: Francisco J. Romero

Copyright © 2020 Rumiana Bakalova et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Vitamin C as a cancer therapy has a controversial history. Much of the controversy arises from the lack of predictive biomarkers for stratification of patients, as well as a clear understanding of the mechanism of action and its multiple targets underlying the anticancer effect. Our review expands the analysis of cancer vulnerabilities for high-dose vitamin C, based on several facts, illustrating the cytotoxic potential of the ascorbyl free radical (AFR) via impairment of mitochondrial respiration and the mechanisms of its elimination in mammals by the membrane-bound NADH:cytochrome b5 oxidoreductase 3 (Cyb5R3). This enzyme catalyzes rapid conversion of AFR to ascorbate, as well as reduction of other redox-active compounds, using NADH as an electron donor. We propose that vitamin C can function in “protective mode” or “destructive mode” affecting cellular homeostasis, depending on the intracellular “steady-state” concentration of AFR and differential expression/activity of Cyb5R3 in cancerous and normal cells. Thus, a specific anticancer effect can be achieved at high doses of vitamin C therapy. The review is intended for a wide audience of readers—from students to specialists in the field.

1. Introduction

In the April 2019 issue of *Nature Reviews Cancer*, Ngo et al. have described cancer vulnerabilities for high-dose vitamin C therapy [1]. The authors note that vitamin C as a cancer therapy has a controversial history, and much of the controversy arises from the lack of predictive biomarkers for stratification of patients, as well as a clear understanding of the mechanism of action and its multiple targets underlying the anticancer effect [2].

Our article links empirical and experimental observations from clinical practice that may, at first glance, seem disjointed. However, common characteristics of the pathologies discussed herein include mitochondrial dysfunction and

severe inflammation. These are important targets for vitamin C and hallmarks of cancer. In particular, we expand the analysis of cancer vulnerabilities for high-dose vitamin C, based on several significant and indicative facts, illustrating the cytotoxic potential of the ascorbyl free radical (AFR) via impairment of mitochondrial respiration and the mechanisms of its elimination in mammals:

- (i) AFR is an intermediate product from the oxidation of ascorbate by one-electron reduction of transition metal ions and/or interaction with free radicals [1]. AFR does not readily react with oxygen or other molecules to generate more reactive radicals and may temporarily accumulate in cells [3]

- (ii) AFR is a by-product of some vital biochemical reactions. Ascorbate participates in the recovery of hemoglobin from methemoglobin in erythrocytes, as well as in the enzymatic hydroxylation (e.g., during the synthesis of noradrenalin in the chromaffin granules of dopamine-synthesizing cells) [4–6]. Both processes are accompanied by AFR production
- (iii) NADH:cytochrome b5 oxidoreductase 3 (Cyb5R3) catalyzes rapid conversion of AFR to ascorbate, as well as reduction of other redox-active compounds, using NADH as an electron donor [7, 8]. Thus, Cyb5R3 eliminates AFR, restores the ascorbate “pool,” and helps maintain the NAD^+/NADH ratio in cells. This enzyme is ubiquitously expressed in all mammalian cells, implying an important role in cell homeostasis
- (iv) Lack or downregulation of Cyb5R3 leads to development of recessive hereditary methemoglobinemia (RHM), accompanied by neuropathies and impairment of mitochondrial function [7, 9]. Ascorbate treatment has beneficial effects on methemoglobinemia, especially in patients with ascorbate deficiency. However, ascorbate deficiency does not explain the neuropathy and mitochondrial impairment associated with this pathology
- (v) Downregulation of membrane-bound Cyb5R3 in cells leads to lower NAD^+/NADH ratio, impaired mitochondrial respiration rate, and decreased ATP production, which is associated with higher sensitivity to oxidative stress [9]
- (vi) We propose that the accumulation of high concentrations of AFR in cells during oxidative stress and/or some specific metabolic activity is responsible for the impairment of mitochondrial respiration in the absence or downregulation of Cyb5R3, particularly the enzyme located on the outer mitochondrial membrane (OMM Cyb5R3). This can induce cytotoxicity in mammalian cells. Perhaps this is why enzymes found in plants that convert ascorbate to AFR (ascorbate oxidase and ascorbate peroxidase) did not evolve in animals and humans [3]

Is AFR harmful for mitochondria and can pharmacological doses of vitamin C attack cancer cells by inhibiting membrane-bound Cyb5R3 and impairing mitochondrial respiration?

2. Cyb5R3/VDAC1 Is a Redox-Cycling System That Converts AFR to Ascorbate

The discovery of enzyme activity of Cyb5R3 dates back to 1951, when Lehninger reported that isolated rat liver mitochondria catalyze rapid oxidation of NADH in the presence of added cytochrome c. This mitochondrial NADH:cytochrome c oxidoreductase activity was found to be independent of the respiratory chain. Thirty years later, it was found that this enzyme is NADH:semi-dehy-

droascorbic acid reductase, which uses a cytochrome b5-like hemoprotein of the outer mitochondrial membrane (OMM cytochrome b5), which is distinct from microsomal cytochrome b5 [10]. A similar enzyme was also found in *Saccharomyces cerevisiae*—mitochondrial NADH:cytochrome b5 reductase, which catalyzes the reduction of D-erythroascorbic acid radical [11].

Recently, OMM NADH:cytochrome b5 reductase was classified as OMM Cyb5R3 (NADH:ferricytochrome b5 oxidoreductase 3; EC 1.6.2.2.). *Cyb5R3 belongs to the class of cancer-related and disease-related genes. Cyb5R3 enzyme belongs to the class of potential drug targets and predicted membrane proteins (according to The Human Protein Atlas).* The Cyb5R3 protein is encoded in two isoforms: (i) soluble, exclusively expressed in erythrocytes, and (ii) membrane-bound, expressed in all other cells and anchored to the outer mitochondrial membrane, endoplasmic reticulum, and plasmatic membrane [8]. Deficiency of cytosolic Cyb5R3 causes type I RHM, which is a benign condition with mild cyanosis, fatigue, and shortness of breath upon exertion [9, 12]. Deficiency of membrane-bound Cyb5R3 causes incurable type II RHM, accompanied by severe neuropathy, permanent cyanosis, and impairment of mitochondrial respiration [9, 12].

OMM Cyb5R3 is functionally connected to the voltage-dependent anion channel 1 (VDAC1), the most abundant protein of the outer mitochondrial membrane [8]. Initially, it was believed that VDAC1 possesses NADH-dependent oxidoreductase activity. However, VDAC1 does not contain flavin, heme, and/or metal ions in its active center, nor any other catalytic motif of the known NAD(P)H-dependent oxidoreductases. Recombinant VDAC1 and VDAC1 purified from comigrating proteins either lack oxidoreductase activity or exhibit very low activity [8, 13, 14]. The purity of VDAC1 preparations studied usually reaches only 90–95%. In all cases, other proteins of equal molecular weights (32–35 kDa) are present, which affect the efficiency of separation [15–18]. It was reported that Cyb5R3 consists of 5–10% of the purified VDAC1 [15, 18]. The oxidoreductase activity ascribing to VDAC1 is inhibited by nonspecific VDAC blockers [15, 16, 19], as well as by specific Cyb5R3 inhibitors [19]. This is direct evidence that oxidoreductase activity of VDAC1 is due to the presence of Cyb5R3 in the samples. It seems that both proteins cofunction and could be considered as one “redox-cycling system” of the outer mitochondrial membrane. Recent studies on cells treated with paraquat (considered to be carcinogenic) indicate that VDAC1 maintains membrane integrity and keeps mitochondria intact, while Cyb5R3 acts as an oxidoreductase [8, 20].

3. The Cyb5R3/VDAC1 Redox-Cycling System Reduces AFR and Helps Maintain Mitochondrial Homeostasis

The crucial role of Cyb5R3 in regeneration of AFR to ascorbate and mitochondrial homeostasis has been demonstrated by Shirabe et al. on human dermal fibroblasts (HDFs), derived from a patient with type II RHM with a point mutation in the Cyb5R3 gene [21]. This mutation results

in a complete absence of membrane-bound Cyb5R3 in the cells. The authors have reported severely decreased cellular NADH:semidehydroascorbic acid reductase activity and rotenone-insensitive NADH:cytochrome c reductase activity, compared to neonatal HDFs. These data suggest that OMM Cyb5R3 and its NADH:semidehydroascorbic acid reductase activity might contribute to the protection and normal functioning of the cell, as well as protection of its mitochondrial electron transport chain (ETC).

Siendones et al. have found that membrane-bound Cyb5R3-deficient HDFs, derived from two patients with similar mutations, are characterized by the decrease of several parameters: NAD^+/NADH ratio, mitochondrial respiration rate, activities of mitochondrial Complexes I-IV, and ATP production [22]. This is associated with higher sensitivity of Cyb5R3-deficient HDFs to oxidative stress and accelerated senescence, compared to neonatal HDFs. The role of Cyb5R3 in the modulation of mitochondrial function has also been evaluated in Cyb5R3-siRNA-silenced MRC-5 normal fibroblasts [22]. Downregulation of Cyb5R3 is accompanied by a significant decrease of the NAD^+/NADH ratio and mitochondrial respiration rate. Cyb5R3-siRNA-silenced MRC-5 cells and type II RHM HDFs displayed low proliferation rates and increased cell death. This is consistent with impairment of mitochondrial metabolism and reduction of oxidative phosphorylation, based on analysis of oxygen consumption. The authors suggest that impairment of Cyb5R3 expression and/or its AFR reductase activity may result not only in NADH accumulation but also in defects of mitochondrial ETC due to increased oxidative damage, ultimately leading to senescence.

Recently, it was reported that Cyb5R3 is essential for cardiomyocyte function [23]. Cardiomyocyte-specific inducible Cyb5R3-knockout mice ($\text{Myh6-Cre}^{\text{ERT2}}\text{-flox/flox}$) show a loss of mitochondrial biogenesis, associated with 30% loss of total ATP, 50% loss of Complex IV activity, and 25% loss of Complex IV protein quantity [23]. RNA_{seq} pathway analysis indicated significantly decreased expression of Complexes I, III, and IV after downregulation of membrane-bound Cyb5R3 in cardiomyocytes. This study shows that Cyb5R3 is essential for normal mitochondrial function and cell survival.

The studies described above suggest that the presence of OMM Cyb5R3 is essential for the following parameters at normal conditions: activity of mitochondrial ETC, oxygen consumption, ATP production, and resistance to oxidative stress.

In vitro studies on isolated mitochondria indicate that the OMM Cyb5R3/VDAC1 complex is responsible for the transfer of electrons from cytosolic NADH into mitochondria and that the process is dependent to Complex IV [24, 25]. This is accompanied by oxygen uptake, proton pumping, and generation of mitochondrial potential, supported by small catalytic amounts of external (mitochondrial) cytochrome c. Their data also demonstrate that cytochrome c-dependent NADH oxidation is strongly inhibited by dextran sulfate (500 kDa)—an inhibitor of VDAC1. They have suggested that in physiological conditions, cytochrome c might be transferred in a very limited amount outside the mitochondria so as to promote the activation of the

Cyb5R3-dependent electron transport pathway. The process is highly dependent on induction of contact sites in the outer and inner mitochondrial membranes (membrane remodeling), as well as on inhibitors of VDAC1. The activity of this Cyb5R3/VDAC1-dependent system becomes functional in removing the excess cytosolic NADH and is essential for cell survival under impairment of the ETC at the level of one of the first respiratory complexes. Since the Cyb5R3-catalyzed electron transfer generates an electrochemical membrane potential associated with the activity of the cytochrome oxidase [24, 26, 27], it represents an additional pathway for providing energy to cells. This aspect is essential for the cells in which oxidative phosphorylation activity is not properly supported by the respiratory chain.

The same authors also demonstrate a nonenzymatic induction of this alternative pathway for ATP synthesis in mitochondria [24, 25]. They added ascorbate (instead of NADH) and small catalytic amounts of external cytochrome c to intact mitochondria and detected oxygen uptake, accompanied by cytochrome c reduction and ascorbate oxidation [24, 25]. These processes are dependent on Complex IV and the induction of contact sites in mitochondrial membrane and are sensitive to inhibitors of VDAC1. It can be assumed that oxidation of ascorbate by small catalytic amounts of external cytochrome c leads also to AFR production. AFR may enter the mitochondria via VDAC1 and affect mitochondrial respiration at Complexes III-IV—a hypothesis, which is described below in the context of the anticancer effect of high doses of vitamin C.

4. The Cyb5R3/VDAC1 Redox-Cycling System in Cancer

Studies discussed above suggest that OMM Cyb5R3/VDAC1 is vital for mitochondrial homeostasis, protection against oxidative stress, prevention of cell senescence, and cellular longevity. These events clearly relate to AFR elimination and maintenance of the cytosolic NAD^+/NADH ratio—crucial factors for cell survival.

It has been shown that VDAC1 is highly expressed in all cells as a consequence of exposure to various toxic substances and plays a crucial role in the protection against intoxication [8, 28, 29]. VDAC1 interacts with both proapoptotic and antiapoptotic factors, which makes it a gatekeeper for mitochondria-mediated cell death or survival signaling pathways [30, 31]. This illustrates the complexity of VDAC functions in normal and cancer cells. It has been observed that overexpression of VDAC1 in cancer cells is associated with high metastatic potential, low therapeutic efficiency, and poor prognosis [30, 31]. In this case, VDAC1 appears to be involved in protecting mitochondria from reactive oxygen species (ROS), functioning as a prosurvival pathway [28–30].

Cyb5R3 is also recognized as a carcinogen detoxification gene [32, 33], although its role in carcinogenesis is not well studied. Single reports on the role of OMM Cyb5R3 in cancer have appeared over the past 10 years [32, 34, 35]. It was found that the enzyme is overexpressed in cancer cells, protecting them against oxidative stress and induction of apoptosis [32, 34, 35].

Rajcevic et al. have analyzed the proteomic profile of metabolic proteins in the invasive glioblastoma phenotype by applying a functional analysis, using isobaric peptide tagging chemistry (iTRAQ) combined with bioinformatics analysis (Ingenuity® Pathway Analysis) [35]. Cyb5R3 is identified as a key protein in a canonical pathway of the cancerous phenotype associated with mitochondrial dysfunction.

Two groups have reported that membrane-bound Cyb5R3 overexpression and polymorphism are associated with breast cancer risk in women [32, 34]. Blanke et al. have found that one variant of Cyb5R3 is significantly overexpressed in African American women with breast cancer, which is accompanied by impaired detoxification of aromatic and heterocyclic amine mammary carcinogens in cigarette smoke [32]. The authors suggest that overexpression of membrane-bound Cyb5R3 should be considered as a valuable marker in etiologic studies of prostate, bladder, and colon cancers, which are also associated with exposure to arylamine and heterocyclic amine carcinogens.

A large cohort study on patients with estrogen receptor-negative (ER^{neg}) and progesterone receptor-negative (PR^{neg}) breast tumors reveals a central role of membrane-bound Cyb5R3 in extravasation/colonization of cancer cells and metastasis formation at distinct sites (e.g., lung) [34]. Immunohistochemical analysis has identified a significant correlation between high Cyb5R3 expression and poor disease-free (DFS; $p = 0.02$) and overall survival (OS; $p = 0.04$). Multivariate analysis revealed about 2-fold higher risk of poor outcome for ER^{neg}/PR^{neg} patients exhibiting strong staining of Cyb5R3, compared to those expressing weak staining. Cyb5R3 gene knockdown using siRNA in metastatic cells leads to a significantly decreased tumor burden in lungs when injected intravenously in immunodeficient mice. The cellular effects of Cyb5R3-siRNA knockdown showed signaling alterations associated with extravasation, such as transforming growth factor beta (TGF β) and hypoxia-inducible factor alpha (HIF α) pathways, and strong suppression of proliferation. It is interesting to note that Cyb5R3-siRNA-silenced cancer cells stop proliferation and metastasis, but they continue to survive [34].

5. AFR in Mitochondrial Homeostasis at Low and High Doses of Vitamin C

Our hypothesis is focused on the intracellular generation of AFR at low and high doses of vitamin C, together with its effect on mitochondrial respiration, intracellular redox balance, and energy metabolism mediated by the OMM Cyb5R3/VDAC1.

We propose that vitamin C can function in “protective mode” or “destructive mode” affecting cellular homeostasis, depending on the OMM Cyb5R3 expression/activity and intracellular “steady-state” concentration of AFR. The hypothesis is focused on the effects of AFR in the mitochondrial intermembrane space and the cytosol.

Membrane-bound Cyb5R3 has several characteristics that could be crucial to mitochondrial homeostasis:

- (i) The redox potential of the OMM-bound cytochrome b5 varies from -160 mV to -272 mV, depending on the NAD⁺/NADH ratio [8, 13]. Therefore, OMM Cyb5R3 can catalyze the reduction of a variety of redox cyclers in addition to AFR, such as quinone-like compounds, bioreductive xenobiotics/drug, and nitrocompounds [8, 17]. These redox cyclers can influence the intracellular redox state
- (ii) Membrane-bound Cyb5R3 helps maintain the cytosolic NAD⁺/NADH ratio, supplying cells with reducing equivalents by transferring electrons from cytosolic NADH to the respective membrane components (electron acceptors and carriers) [7, 8, 10, 11, 22]
- (iii) The membrane-bound Cyb5R3 is considered to be one of the major enzymes that maintain the NAD⁺/NADH ratio, by using the coenzyme Q (CoQ) “pools” of the outer mitochondrial membrane and endoplasmic reticulum [8, 10, 11]

CoQ is most likely the main acceptor of excess electrons, as it is the most abundant membrane-bound redox-active compound in the cell. CoQ is also a key electron carrier in mitochondria. Therefore, outside the inner mitochondrial membrane, CoQ can serve as a “buffer” for excess-reducing equivalents, accepting electrons from excess cytosolic NADH via Cyb5R3.

We also assume that the ascorbate could serve as a “buffer” of excess-reducing equivalents in the intracellular aqueous phase of cancer cells due to their oxidative environment, because ascorbate is one of the most abundant cytosolic redox-active compounds. Steady-state levels of ascorbate are maintained by Cyb5R3 [6–8], and they are significantly higher in cancer cells compared to normal cells [1, 36].

Figures 1 and 2 represent two modes of action of vitamin C on mitochondrial homeostasis via generation of AFR and its elimination by the OMM Cyb5R3/VDAC1: (i) “protective mode” at low/normal (steady-state) doses and (ii) “destructive mode” at high doses of vitamin C.

Ascorbate and AFR exist as anions, which imply their penetration from the cytosol into the mitochondrial intermembrane space and back through the VDAC1 (Figure 1).

In normal cells (Figure 1), low concentrations of AFR should be generated due to the low (steady-state) intracellular levels of vitamin C, as a result of (i) relatively low expression of vitamin C transport proteins [37] and (ii) low steady-state levels of ROS and normal levels of reducing equivalents [38]. Both factors are responsible for the limited production of AFR, which is a product of one-electron oxidation of ascorbate. Since AFR is relatively stable, it could be rapidly eliminated by the OMM Cyb5R3/VDAC1. This prevents effects on mitochondrial respiration, and all complexes will operate in a “normal mode.” Electrons are transferred from Complex III’s Qo site of the CoQ “pool” to cytochrome c, creating a proton gradient and leading to normal ATP synthesis. In this case, the CoQ “pool” is balanced and the Qo site of Complex III transfers one electron to




FIGURE 1: "Protective mode" of action of the OMM Cyb5R3/VDAC1 at low (steady-state) concentrations of vitamin C in normal cells (at normoxia). OMM Cyb5R3/VDAC1 converts ascorbyl free radical (AFR; $Asc^{\bullet-}$) to ascorbate ($AscH^-$), using cytosolic NADH as an electron donor. This prevents effects on mitochondrial respiration and all complexes operate in a "normal mode," creating a proton gradient and leading to normal ATP synthesis. The CoQ "pool" is balanced [39], and the Q_o site of Complex III transfers one electron to cytochrome c and the second electron to ubiquinone in the Q_i site. VDAC1: voltage-dependent anion channel 1; Cyb5R3: NADH:cytochrome b5 oxidoreductase 3; OMM: outer mitochondrial membrane; Cyt. c: cytochrome c.

cytochrome c and the second electron to ubiquinone while bound in the Q_i site [39]. This model is typical for normal (noncancerous) cells.

In cancer cells, relatively high "steady-state" levels of vitamin C can be achieved due to overexpression of its transport proteins [1, 37, 40]. Cancer cells are also characterized by permanent oxidative stress and redox imbalance, accompanied mainly by overproduction of superoxide, hypoxic environment, and overproduction of cytosolic NADH due to upregulation of the aldehyde dehydrogenase pathway [41, 42]. These distinct characteristics suggest increased production of intracellular AFR at "steady-state" levels of vitamin C in cancer cells, compared to normal cells. On the other hand, AFR should be rapidly eliminated in the mitochondrial environment of cancer cells due to overexpressed Cyb5R3/VDAC1 [30, 32, 34, 35] and high cytosolic levels of NADH [38, 40, 41] as an electron donor. Overexpression of the Cyb5R3/VDAC1 appears to be a compensatory mechanism that protects cancer cells and their mitochondria by

regulating AFR accumulation at "steady-state" conditions, while also maintaining the cytosolic $NAD^+/NADH$ ratio.

6. What Processes Can Be Induced in Cancerous Mitochondria at High "Therapeutic" Intracellular Concentrations of Vitamin C?

As well described by Ngo et al. and others, the first consequence of the high-dose "therapeutic" vitamin C is to selectively elevate its concentration in cancer cells [1, 36, 43]. Chen et al. have reported that vitamin C in pharmacological doses selectively generates AFR and hydrogen peroxides in the extracellular fluids of cancer-bearing patients [44]. Keshari et al. have studied ascorbate/dehydroascorbate ratios in solid tumors in vivo as a marker of oxidative stress at different expressions of glucose transporters [45, 46]. The authors have detected AFR as an intermediate product, using time-resolved MR spectroscopy. These studies show that




FIGURE 2: “Destructive mode” of action of ascorbyl free radical via the OMM Cyb5R3/VDAC1 at high (“therapeutic”) concentrations of vitamin C in cancer cells (at normoxia). High intracellular concentration of ascorbate may induce Cyb5R3 end-product inhibition, accompanied by elevated levels of ascorbyl free radical (AFR) in the mitochondrial intermembrane space and decrease of the NAD⁺/NADH ratio in the cytosol. AFR may transfer one electron to oxidized cytochrome c, causing a partial (or complete) arrest of electron flow between Complex III and Complex IV. Rapid reduction of cytochrome c by AFR effectively competes with and perturbs the proton pumping at Complex III, as well as the temporal electron transport provided by CoQ cycles of Complex III. When Complex III is blocked, succinate (from the citric acid cycle) builds up, mitochondrial membrane potential rises, and the CoQ “pool” becomes unbalanced. Reverse electron transport (RET) to Complex I is driven, which causes a superoxide “burst” [95, 96]. RET is also accompanied by synthesis of succinate and NADH. We propose that the “destructive mode” caused by high-dose vitamin C may contribute to RET not only during tissue reperfusion at angiogenesis but also during tumor hypoxia and normoxia in cancer cells. In addition, the reduction of cytochrome c by AFR results in production of DHA. DHA is converted to ascorbate by glutathione, which provokes a depletion of reducing equivalents in cancer cells—a crucial factor for their survival. VDAC1: voltage-dependent anion channel 1; Cyb5R3: NADH:cytochrome b5 oxidoreductase 3; OMM: outer mitochondrial membrane; Cyt. c: cytochrome c; RET: reverse electron transport; DHA: dehydroascorbate; AscH⁻: ascorbate in anion form; Asc^{•-}: ascorbyl free radical.

AFR is also generated inside the cells. It should be clarified that high intracellular levels of AFR can only be achieved with intravenous vitamin C administration, because ascorbate concentrations in the plasma and tissues are tightly controlled as a function of the oral dose [47].

We suppose that high “therapeutic” intracellular concentration of ascorbate in cancer cells is then likely to induce “enzyme (Cyb5R3) end-product inhibition”—a negative feedback, regulating any enzymatic pathway (Figure 2). The

inhibition of OMM Cyb5R3 will result in the accumulation of high amounts of AFR under the permanent oxidative stress within these cells.

In the mitochondrial intermembrane space, AFR may transfer one electron to cytochrome c, causing a partial (or complete) arrest of electron flow between Complex III and Complex IV. It has been shown that the rate of reduction of cytochrome c by AFR, with production of DHA, is orders of magnitude faster than the reduction of cytochrome c by ascorbate [48, 49].

Rapid reduction of cytochrome *c* by AFR effectively competes with the electron transport from the CoQ cycle of Complex III to the cytochrome *c*, which perturbs the proton pumping. Competitive inhibition by AFR, bypassing Complex III to cytochrome *c*, drives the CoQ “pool” to a more reduced (overcharged) state. If this overcharging phenomenon occurs, there will be insufficient ubiquinone, available for binding at the Qi site. This unbalances the Qo/Qi cycle. It also increases the lifetime of semiubiquinone in the Qo pocket, allowing oxygen to accept the second electron from ubiquinol during the second step of the CoQ cycle, yielding superoxide (Figure 2) [50].

Briefly, we propose that a high level of AFR in the intermembrane space of mitochondria blocks Complex III, at least partially, via unbalancing the CoQ “pool” and impairing mitochondrial respiration in vitamin C-treated cancer cells. Precisely timed redox sequences responsible for the efficient flow of electrons from Complex III to oxidized cytochrome *c* are vulnerable to AFR flux. This drives the CoQ “pool” to a more reduced (overcharged) form. This also causes the lifetime of semiubiquinone in the Qo pocket to lengthen, allowing oxygen to accept the second electron from ubiquinol during the second step of the CoQ cycle, yielding superoxide. This can interfere with the timing of the Qo/Qi cycle and consequently the CoQ “pool” redox balance, impairing mitochondrial respiration. A highly reduced CoQ “pool” inhibits proton pumping, yielding up to 80% lower ATP plus higher levels of superoxide. We further propose that the damaging effect of high-dose vitamin C on cancer cells is similar to the mechanism of ischemia-reperfusion injury (IRI) or reoxygenation damage [51]. In IRI, the mitochondrial CoQ “pool” becomes highly reduced by stopped flow through Complex III.

It is interesting to note that this mechanism can influence the mitochondrial ATP production in different directions depending on the cell type. In the case of cancer cells, this mechanism should result in a significant reduction of mitochondrial ATP and increase of superoxide production (Figure 2).

It has been generally accepted that cancer cells use glycolysis as a source of ATP (Warburg effect) and do not rely on mitochondrial respiration for ATP production. However, recent studies demonstrate that the majority of ATP in cancer cells is produced by mitochondria and some tumors show heavy dependence on oxidative phosphorylation [52–56]. Many solid tumors are poorly perfused and have a limited supply of glucose, but enough oxygen to generate mitochondrial ATP [57]. The ETC is able to function optimally at oxygen levels as low as 0.5% [58]. Therefore, blocking mitochondrial ATP production will induce cell death in poorly perfused tumors.

Even ignoring the dependence of cancer cells on mitochondrial ATP, a function of the Krebs cycle (at Complex II) is necessary for the supply of metabolites for synthesis of nucleic acids, fatty acids, etc. Blocking the ETC will stop this supply and suppress proliferation. Krebs cycle metabolites (such as succinate, fumarate, and itaconate) are also coupled with nonmetabolic signaling in cancer and immune cells, which is crucial for cancer progression and invasion [59, 60].

The key consideration in targeting mitochondria by high-dose vitamin C (via AFR) is that normal cells rely mostly on mitochondrial respiration for ATP production. As was mentioned above, high concentrations of AFR cannot be reached in normal cells, especially *in vivo* in the absence of oxidative stress, due to lower expression of glucose/ascorbate transport proteins [36, 47] and low amounts and normal functioning of Cyb5R3 [32, 33]. However, normal cells should also be vulnerable to high concentrations of AFR generated under oxidative stress. Perhaps, this is a part of the dual effect of vitamin C, which can act as an antioxidant or prooxidant, depending on the environment [61].

It is shown that in the case of noncancer cells with mitochondrial deficiency, high-dose vitamin C could even increase ATP production. Eleff et al. reported that administration of vitamin C and quinone-like provitamin menadione on the skeletal muscles of a 17-year-old patient with a severe defect in Complex III of the mitochondrial ETC increased the recovery rate, compared to the recovery rate of the young female controls [62]. This rare defect included a stop-codon mutation (G15242A) in the mtDNA-encoded cytochrome *b* gene, which was accompanied by an effective prevention of aerobic metabolism and oxidative phosphorylation. The authors suppose that both substances bypass the deficient Complex III as electron transfer mediators to carry the electrons from CoQ to cytochrome *c*. Thus, they increased ATP production from mitochondria compared to the initial level. It is interesting to note that redox cycles of vitamin C and menadione are mediated by Cyb5R3/VDAC1 [8, 17] and the enzyme activity could be important for this bypass. In this patient, the combination of vitamin C and menadione resulted in the production of more ATP than in the case of vitamin C applied alone. The authors suggest that both substances can directly reduce cytochrome *c*, since the reduction potential of cytochrome *c* is over +200 mV—more positive than those of ascorbate and menadione [62–65].

7. What Could Be the Consequences of AFR-Mediated Electron Transfer in Cancerous Mitochondria from High-Dose Vitamin C Therapy?

- (i) Induction of reverse electron transport in the respiratory chain and acceleration of superoxide production at Complex I, as well as Complex III (Figure 2) [50, 65–67]. Degradation of Complex I when the CoQ “pool” is unbalanced, and Complex III is dysfunctional [67]. In hypoxic conditions (common to many types of cancer cells), these redox reactions occur even faster, exponentially increasing superoxide production [50, 65–67].
- (ii) Superoxide-mediated destruction of Fe/S clusters from the complexes and induction of Fenton’s reactions in mitochondria [68–70], in the presence of AFR and/or ascorbate

- (iii) Mitochondria within cancer cells are vulnerable to a “destructive mode” of action from high-dose vitamin C, which leads to their collapse

In addition, the conversion of AFR to DHA and subsequent reduction of DHA to ascorbate by glutathione- and NADPH-dependent pathways will promote a depletion of glutathione and reducing equivalents in the cancer cells leading to severe oxidative stress—crucial factors for induction of apoptosis and cell death [38].

An important consequence of the reduction of cytochrome c by AFR is the transfer of electrons from cytoplasmic NADH to Complex IV, which is accompanied by production of 1 mol ATP per 1 mol NADH—suggesting the likelihood of thermogenesis [71, 72]. Cancer cells often express heat shock proteins and are very sensitive to the elevated temperature of their environment [38, 73]. This may also contribute to converting “cold” tumors into “hot” tumors, an important cell death-facilitating factor potentially from combining high-dose vitamin C therapy with applications in immunotherapy [74].

It is also reported that NADH reduces conduction of VDAC by partial channel block and/or modulation of its activity [75, 76]. As ATP and many other vital substances pass from the mitochondria into the cytosol through VDAC [77, 78], partial inhibition of the VDAC channel would also contribute to the impairment of mitochondrial respiration and decrease of cell viability.

Recent studies show that Cyb5R3/VDAC1 can activate bioreductive antitumor drugs, which in combination with vitamin C may affect mitochondrial activity, exhibiting additive or synergistic anticancer effects [17, 79, 80]. For example, vitamin C can sensitize cancer cells to redox-responsive chemotherapeutics (e.g., furanonaphthoquinones) after their bioreductive activation via Cyb5R3/VDAC1 [81, 82]. In this case, the “therapeutic” intravenous dose of vitamin C could be decreased to more tolerant values. It has been reported that high doses of vitamin C exhibit selective antitumor effects in combination with menadione—an analogue of 1,4-naphoquinone [83–85]. A recent study demonstrates that high/tolerable concentrations of menadione inhibit Cyb5R3 [86]. It can be assumed that menadione-mediated inhibition of Cyb5R3 is involved in potentiation of the anticancer effect of high-dose vitamin C via overproduction of AFR and subsequent impairment of mitochondrial respiration (as we assume with our “model”). Moreover, vitamin C/menadione could achieve a selective cytotoxicity against cancer cells due to the rapid UBIAD1-mediated conversion of menadione to vitamin K2 in normal cells and downregulation of UBIAD1 in the majority of cancers leading to strong inhibition of this conversion [87, 88]. These assumptions could be a prerequisite/basis for the design of future studies aimed to elucidate the molecular mechanisms of the synergistic anticancer effects of vitamin C and vitamin C/menadione with conventional chemotherapeutics, described in the literature [89]. Clarifying the effects of conventional anticancer drugs and high-dose vitamin C on Cyb5R3 may also be a prerequisite for predicting the effectiveness of vitamin C in adjuvant settings and stratification of patients with good/poor prognosis.

8. Concluding Remarks and Outstanding Questions

The high concentrations of cytochrome c in mitochondria [90], as well as the possible existence of dedicated pools of CoQ and cytochrome c (as supported by the “plasticity model” of mitochondria) [90, 91], could limit the proposed direct effects of AFR on oxidative phosphorylation, as described in Figure 2. In fact, reported concentrations of vitamin C in cells under normal conditions are significantly higher than those of cytochrome c (~10–30 nmol/mg protein versus ~0.1–1 nmol/mg protein) [90, 92]. In this case, ascorbate and AFR concentrations in cancer cells after treatment with high doses of vitamin C should be much higher than cytochrome c concentrations [93]. This supports the possibility of direct electron transfer from AFR to cytochrome c at Complex IV. Remodeling of the mitochondrial membrane and, in particular, the contact sites of Cyb5R3/VDAC1/Complex IV to optimize performance can explain the different efficiency and selectivity of high-dose vitamin C therapy when combining with different anticancer drugs (depending on their mechanisms). In addition, the abovementioned consequences of AFR-mediated electron transfer in cancerous mitochondria at high-dose vitamin C therapy may explain the induction of apoptosis even without strong inhibition of oxidative phosphorylation and ATP production. The mechanism, described in Figure 2, may not cause strong cancer cell death, but it can damage the mitochondria, stop proliferation, and keep cancer cells latent, so they become vulnerable to the immune system of the intact organism.

The analysis of the literature raises several outstanding questions that need experimental verification:

- (1) Does AFR impair mitochondrial respiration by reducing cytochrome c, causing an arrest of electron flow between Complexes III and IV, a reverse electron transport, and an induction of oxidative stress in the absence or downregulation of Cyb5R3/VDAC1
- (2) Does vitamin C in low/normal doses work in “protective mode” in normal and cancer cells? If yes, is this related to maintaining the balance of the CoQ “pool” and cytosolic reducing equivalents
- (3) Does vitamin C in high doses work in “destructive mode” in cancer cells, but not in normal cells? If yes, is this related to maintaining the imbalance of the CoQ “pool” and cytosolic reducing equivalents
- (4) What are the levels of mitochondrial superoxide, succinate, and ATP in normal and cancer cells of the same origin, treated with low, moderate, and high doses of vitamin C? Is there a difference in the levels of mitochondrial superoxide, succinate, and ATP in cells with different proliferative index, treated with vitamin C
- (5) Does the effect of pharmacological vitamin C relate to the expression and activity of membrane-bound Cyb5R3/VDAC1? Can high doses of vitamin C attack

cancer cells only by inhibiting Cyb5R3/VDAC1 and specific destruction of cancer mitochondria

- (6) What is the impact/significance of Cyb5R3/VDAC1 as a predictive biomarker for stratification of cancer-bearing patients for high-dose vitamin C therapy
- (7) Does pharmacological vitamin C have a potential as adjuvant cancer therapy in combination with redox-responsive drugs

Our article is intended to provoke the design of new experiments (in vitro and in vivo) that can provide evidence Pro and Con for the assumptions made, to change the “status quo” in the controversial field of “pharmacological vitamin C.”

If vitamin C can clinically exploit the pathway presented here, we suggest it offers a new, exciting rationale for cancer studies. Perhaps this orthogonal mechanism of action can safely and more effectively compliment current cancer treatments in the adjuvant setting and help to let the phoenix fly [94].

Abbreviations

AFR:	Ascorbyl free radical
CoQ:	Coenzyme Q
Cyb5R3:	NADH:cytochrome b5 oxidoreductase 3
DHA:	Dehydroascorbic acid
HDF:	Human dermal fibroblast
OMM:	Outer mitochondrial membrane
RHM:	Recessive hereditary methemoglobinemia
VDAC1:	Voltage-dependent anion channel 1.

Conflicts of Interest

No potential conflicts of interest are disclosed.

Acknowledgments

This study was partially supported by the IC-MedTech Co. (San Diego, CA, USA) and the Japan Agency for Medical Research and Development (AMED) (Project for Cancer Research and Therapeutic Evolution, P-CREATE, no. 16 cm0106202h0001).

References

- [1] B. Ngo, J. van Riper, L. C. Cantley, and J. Yun, “Targeting cancer vulnerabilities with high-dose vitamin C,” *Nature Reviews Cancer*, vol. 19, no. 5, pp. 271–282, 2019.
- [2] A. Unly, O. Kirca, and M. Ozdogan, “High-dose vitamin C and cancer,” *Journal of Oncological Sciences*, vol. 1, pp. 10–12, 2015.
- [3] N. Smirnoff, “Ascorbic acid metabolism and functions: a comparison of plants and mammals,” *Free Radical Biology & Medicine*, vol. 122, pp. 116–129, 2018.
- [4] A. Allen, C. Fisher, A. Premawardhana et al., “Methemoglobinemia and ascorbate deficiency in hemoglobin E β -thalassaemia: metabolic and clinical implications,” *Blood*, vol. 120, no. 15, pp. 2939–2944, 2012.
- [5] E. J. Diliberto Jr. and P. L. Allen, “Mechanism of dopamine β -hydroxylation: semidehydroascobate as the enzymic oxidation product of ascorbate,” *The Journal of Biological Chemistry*, vol. 256, 1981.
- [6] F. S. Menniti, J. Knoth, and Diliberto EJ Jr, “Role of ascorbic acid in dopamine beta-hydroxylation. The endogenous enzyme cofactor and putative electron donor for cofactor regeneration,” *The Journal of Biological Chemistry*, vol. 261, no. 36, pp. 16901–16908, 1986.
- [7] R. de Cabo, E. Siendones, R. Minor, and P. Navas, “Cyb5R3: a key player in aerobic metabolism and aging?,” *Aging*, vol. 2, no. 1, pp. 63–68, 2009.
- [8] A. B. Nikiforova, N.-E. L. Saris, and A. G. Kruglov, “External mitochondrial NADH-dependent reductase of redox cyclers: VDAC1 or Cyb5R3?,” *Free Radical Biology & Medicine*, vol. 74, pp. 74–84, 2014.
- [9] E. Siendones, M. Ballesteros, and P. Navas, “Cellular and molecular mechanisms of recessive hereditary Methaemoglobinemia type II,” *Journal of Clinical Medicine*, vol. 7, no. 10, p. 341, 2018.
- [10] A. Ito, S. Hayashi, and T. Yoshida, “Participation of a cytochrome b₅-like hemoprotein of outer mitochondrial membrane (OM cytochrome b) in NADH-semidehydroascorbic acid reductase activity of rat liver,” *Biochemical and Biophysical Research Communications*, vol. 101, no. 2, pp. 591–598, 1981.
- [11] J. S. Lee, W. K. Huh, B. H. Lee et al., “Mitochondrial NADH-cytochrome b₅ reductase plays a crucial role in the reduction of d-erythroascorbyl free radical in *Saccharomyces cerevisiae*,” *Biochimica et Biophysica Acta (BBA) - General Subjects*, vol. 1527, no. 1-2, pp. 31–38, 2001.
- [12] M. J. Percy and T. R. Lappin, “Recessive congenital methaemoglobinemia: cytochrome b₅ reductase deficiency,” *British Journal of Haematology*, vol. 141, no. 3, pp. 298–308, 2008.
- [13] S. Kimura, H. Nishida, and T. Iyanagi, “Effects of flavin-binding motif amino acid mutations in the NADH-cytochrome b₅ reductase catalytic domain on protein stability and catalysis,” *Journal of Biochemistry*, vol. 130, no. 4, pp. 481–490, 2001.
- [14] M. Gonzalez-Gronow, R. Ray, F. Wang, and S. V. Pizzo, “The voltage-dependent anion channel (VDAC) binds tissue-type plasminogen activator and promotes activation of plasminogen on the cell surface,” *The Journal of Biological Chemistry*, vol. 288, no. 1, pp. 498–509, 2013.
- [15] A. Deniaud, C. Rossi, A. Berquand et al., “Voltage-dependent anion channel transports calcium ions through biomimetic membranes,” *Langmuir*, vol. 23, no. 7, pp. 3898–3905, 2007.
- [16] V. Shoshan-Barmatz, D. Ben-Hail, L. Admoni, Y. Krelin, and S. S. Tripathi, “The mitochondrial voltage-dependent anion channel 1 in tumor cells,” *Biochimica et Biophysica Acta (BBA) - Biomembranes*, vol. 1848, no. 10, pp. 2547–2575, 2015.
- [17] E. Simamura, H. Shimada, Y. Ishigaki, T. Hatta, N. Higashi, and K. Hirai, “Bio-reductive activation of quinone anti-tumor drugs by mitochondrial voltage-dependent anion channel 1,” *Anatomical Science International*, vol. 83, no. 4, pp. 261–266, 2008.
- [18] M. Tamura, T. Yubisui, and M. Takeshita, “Microsomal NADH-cytochrome b₅ reductase of bovine brain: purification and properties,” *Journal of Biochemistry*, vol. 94, no. 5, pp. 1547–1555, 1983.
- [19] M. Baker, J. D. Ly, and A. Lawen, “Characterization of VDAC1 as a plasma membrane NADH-oxidoreductase,” *BioFactors*, vol. 21, no. 1-4, pp. 215–221, 2004.

- [20] H. Shimada, K. Hirai, E. Simamura et al., "Paraquat toxicity induced by voltage-dependent anion channel 1 acts as an NADH-dependent oxidoreductase," *The Journal of Biological Chemistry*, vol. 284, no. 42, pp. 28642–28649, 2009.
- [21] K. Shirabe, M. T. Landi, M. Takeshita, G. Uziel, E. Fedrizzi, and N. Borgese, "A novel point mutation in a 3' splice site of the NADH-cytochrome b5 reductase gene results in immunologically undetectable enzyme and impaired NADH-dependent ascorbate regeneration in cultured fibroblasts of a patient with type II hereditary methemoglobinemia," *American Journal of Human Genetics*, vol. 57, no. 2, pp. 302–310, 1995.
- [22] E. Siendones, S. SantaCruz-Calvo, A. Martín-Montalvo et al., "Membrane-bound Cyb5R3 is a common effector of nutritional and oxidative stress response through FOXO3a and Nrf2," *Antioxidants & Redox Signaling*, vol. 21, no. 12, pp. 1708–1725, 2014.
- [23] N. T. Carew, H. M. Altmann, J. C. Galley et al., "Abstract 20733: cytochrome b5 reductase 3 is essential for cardiomyocyte function," *Circulation*, vol. 136, Supplementary 1, 2017.
- [24] D. Marzulli, G. la Piana, E. Fransvea, and N. E. Lofrumento, "Modulation of cytochrome c-mediated extramitochondrial NADH oxidation by contact site density," *Biochemical and Biophysical Research Communications*, vol. 259, no. 2, pp. 325–330, 1999.
- [25] G. La Piana, D. Marzulli, V. Gorgoglione, and N. E. Lofrumento, "Porin and cytochrome oxidase containing contact sites involved in the oxidation of cytosolic NADH," *Archives of Biochemistry and Biophysics*, vol. 436, no. 1, pp. 91–100, 2005.
- [26] M. E. Bodrova, V. I. Dedukhova, E. N. Mokhova, and V. P. Skulachev, "Membrane potential generation coupled to oxidation of external NADH in liver mitochondria," *FEBS Letters*, vol. 435, no. 2-3, pp. 269–274, 1998.
- [27] G. La Piana, D. Marzulli, M. I. Consalvo, and N. E. Lofrumento, "Cytochrome c-induced cytosolic nicotinamide adenine dinucleotide oxidation, mitochondrial permeability transition, and apoptosis," *Archives of Biochemistry and Biophysics*, vol. 410, no. 2, pp. 201–211, 2003.
- [28] V. Shoshan-Barmatz, Y. Krelin, A. Shteinfein-Kuzmine, and T. Arif, "Voltage-dependent anion channel 1 as an emerging drug target for novel anti-cancer therapeutics," *Frontiers in Oncology*, vol. 7, 2017.
- [29] A. K. S. Camara, Y. F. Zhou, P. C. Wen, E. Tajkhorshid, and W. M. Kwok, "Mitochondrial VDAC1: a key gatekeeper as potential therapeutic target," *Frontiers in Physiology*, vol. 8, 2017.
- [30] L. Leanza, M. Zoratti, E. Gulbins, and I. Szabo, "Mitochondrial ion channels as oncological targets," *Oncogene*, vol. 33, no. 49, pp. 5569–5581, 2014.
- [31] A. Magri, S. Reina, and V. De Pinto, "VDAC1 as pharmacological target in cancer and neurodegeneration: focus on its role in apoptosis," *Frontiers in Chemistry*, vol. 6, 2018.
- [32] K. L. Blanke, J. C. Sacco, R. C. Millikan, A. F. Olshan, J. Luo, and L. A. Trepanier, "Polymorphisms in the carcinogen detoxification genes *Cyb5A* and *Cyb5R3* and breast cancer risk in African American women," *Cancer Causes & Control*, vol. 25, no. 11, pp. 1513–1521, 2014.
- [33] A. Martín-Montalvo, Y. Sun, A. Diaz-Ruiz et al., "Cytochrome b₅ reductase and the control of lipid metabolism and healthspan," *Aging and Mechanisms of Disease*, vol. 2, 2016.
- [34] R. R. Lund, R. Leth-Larsen, T. D. Caterino et al., "NADH-cytochrome b5 reductase 3 promotes colonization and metastasis formation and is a prognostic marker of disease-free and overall survival in estrogen receptor-negative breast cancer," *Molecular & Cellular Proteomics*, vol. 14, no. 11, pp. 2988–2999, 2015.
- [35] U. Rajcevic, K. Petersen, J. C. Knol et al., "iTRAQ-based proteomics profiling reveals increased metabolic activity and cellular cross-talk in angiogenic compared with invasive glioblastoma phenotype," *Molecular & Cellular Proteomics*, vol. 8, no. 11, pp. 2595–2612, 2009.
- [36] M. Levine, S. J. Padayatty, and M. G. Espey, "Vitamin C: a concentration-function approach yields pharmacology and therapeutic discoveries," *Advances in Nutrition*, vol. 2, no. 2, pp. 78–88, 2011.
- [37] K. C. Sagun, J. M. Cárcamo, and D. W. Golde, "Vitamin C enters mitochondria via facilitative glucose transporter 1 (Glut1) and confers mitochondrial protection against oxidative injury," *The FASEB Journal*, vol. 19, no. 12, pp. 1657–1667, 2005.
- [38] D. Trachootham, J. Alexandre, and P. Huang, "Targeting cancer cells by ROS-mediated mechanisms: a radical therapeutic approach?," *Nature Reviews Drug Discovery*, vol. 8, no. 7, pp. 579–591, 2009.
- [39] M. Sarewicz and A. Osyczka, "Electronic connection between the quinone and cytochrome c redox pools and its role in regulation of mitochondrial electron transport and redox signaling," *Physiological Reviews*, vol. 95, no. 1, pp. 219–243, 2015.
- [40] A. A. Chen, C. J. Marsit, B. C. Christensen et al., "Genetic variation in the vitamin C transporter, *SLC23A2*, modifies the risk of HPV16-associated head and neck cancer," *Carcinogenesis*, vol. 30, no. 6, pp. 977–981, 2009.
- [41] J. Y. Kim-Muller, J. Fan, Y. J. R. Kim et al., "Aldehyde dehydrogenase 1a3 defines a subset of failing pancreatic β cells in diabetic mice," *Nature Communications*, vol. 7, 2016.
- [42] J. H. Kang, S. H. Lee, D. Hong et al., "Aldehyde dehydrogenase is used by cancer cells for energy metabolism," *Experimental & Molecular Medicine*, vol. 48, no. 11, article e272, 2016.
- [43] G. Nauman, J. Gray, R. Parkinson, M. Levine, and C. Paller, "Systematic review of intravenous ascorbate in cancer clinical trials," *Antioxidants*, vol. 7, no. 7, p. 89, 2018.
- [44] Q. Chen, M. G. Espey, A. Y. Sun et al., "Ascorbate in pharmacologic concentrations selectively generates ascorbate radical and hydrogen peroxide in extracellular fluid *in vivo*," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 104, no. 21, pp. 8749–8754, 2007.
- [45] K. R. Keshari, J. Kurhanewicz, R. Bok, P. E. Larson, D. B. Vigneron, and D. M. Wilson, "Hyperpolarized ¹³C dehydroascorbate as an endogenous redox sensor for *in vivo* metabolic imaging," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 108, no. 46, pp. 18606–18611, 2011.
- [46] K. R. Keshari, V. Sai, Z. J. Wang, H. F. VanBrocklin, J. Kurhanewicz, and D. M. Wilson, "Hyperpolarized [1-¹³C]dehydroascorbate MR spectroscopy in a murine model of prostate cancer: comparison with ¹⁸F-FDG PET," *Journal of Nuclear Medicine*, vol. 54, no. 6, pp. 922–928, 2013.
- [47] S. J. Padayatty, H. Sun, Y. Wang et al., "Vitamin C pharmacokinetics: implications for oral and intravenous use," *Annals of Internal Medicine*, vol. 140, no. 7, pp. 533–537, 2004.
- [48] I. Yamazaki, "The reduction of cytochrome c by enzyme-generated ascorbic free radical," *The Journal of Biological Chemistry*, vol. 237, pp. 224–229, 1962.

- [49] C. Paciolla and L. De Gara, "Reduction of cytochrome c by ascorbic free radical," *Bollettino della Società Italiana di Biologia Sperimentale*, vol. 67, no. 2, pp. 137–144, 1991.
- [50] L. Bleier and S. Drose, "Superoxide generation by complex III: from mechanistic rationales to functional consequences," *Biochimica et Biophysica Acta (BBA) - Bioenergetics*, vol. 1827, no. 11–12, pp. 1320–1331, 2013.
- [51] E. T. Chouchani, V. R. Pell, E. Gaude et al., "Ischaemic accumulation of succinate controls reperfusion injury through mitochondrial ROS," *Nature*, vol. 515, no. 7527, pp. 431–435, 2014.
- [52] P. Caro, A. U. Kishan, E. Norberg et al., "Metabolic signatures uncover distinct targets in molecular subsets of diffuse large B cell lymphoma," *Cancer Cell*, vol. 22, no. 4, pp. 547–560, 2012.
- [53] J. Fan, J. J. Kamphorst, R. Mathew et al., "Glutamine-driven oxidative phosphorylation is a major ATP source in transformed mammalian cells in both normoxia and hypoxia," *Molecular Systems Biology*, vol. 9, no. 1, p. 712, 2013.
- [54] R. Haq, J. Shoag, P. Andreu-Perez et al., "Oncogenic BRAF regulates oxidative metabolism via PGC1 α and MITF," *Cancer Cell*, vol. 23, no. 3, pp. 302–315, 2013.
- [55] F. Vazquez, J.-H. Lim, H. Chim et al., "PGC1 α expression defines a subset of human melanoma tumors with increased mitochondrial capacity and resistance to oxidative stress," *Cancer Cell*, vol. 23, no. 3, pp. 287–301, 2013.
- [56] S. E. Weinberg and N. S. Chandel, "Targeting mitochondria metabolism for cancer therapy," *Nature Chemical Biology*, vol. 11, no. 1, pp. 9–15, 2015.
- [57] R. K. Jain, L. L. Munn, and D. Fukumura, "Dissecting tumour pathophysiology using intravital microscopy," *Nature Reviews Cancer*, vol. 2, no. 4, pp. 266–276, 2002.
- [58] W. L. Rumsey, C. Schlosser, E. M. Nuutinen, M. Robiolio, and D. F. Wilson, "Cellular energetics and the oxygen dependence of respiration in cardiac myocytes isolated from adult rat," *The Journal of Biological Chemistry*, vol. 265, no. 26, pp. 15392–15402, 1990.
- [59] A. G. M. Tielens, C. Rotte, J. J. van Hellemond, and W. Martin, "Mitochondria as we don't know them," *Trends in Biochemical Sciences*, vol. 27, no. 11, pp. 564–572, 2002.
- [60] D. G. Ryan, M. P. Murphy, C. Frezza et al., "Coupling Krebs cycle metabolites to signalling in immunity and cancer," *Nature Metabolism*, vol. 1, pp. 16–33, 2019.
- [61] J. Du, J. J. Cullen, and G. R. Buettner, "Ascorbic acid: chemistry, biology and the treatment of cancer," *Biochimica et Biophysica Acta (BBA) - Reviews on Cancer*, vol. 1826, no. 2, pp. 443–457, 2012.
- [62] S. Eleff, N. G. Kennaway, N. R. Buist et al., "31P NMR study of improvement in oxidative phosphorylation by vitamins K3 and C in a patient with a defect in electron transport at complex III in skeletal muscle," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 81, no. 11, pp. 3529–3533, 1984.
- [63] G. C. Wagner, R. J. Kassner, and M. D. Kamen, "Redox potentials of certain vitamins K: implications for a role in sulfite reduction by obligately anaerobic bacteria," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 71, no. 2, pp. 253–256, 1974.
- [64] T. Matsui, Y. Kitagawa, M. Okumura, and Y. Shigeta, "Accurate standard hydrogen electrode potential and applications to the redox potentials of vitamin C and NAD/NADH," *The Journal of Physical Chemistry A*, vol. 119, no. 2, pp. 369–376, 2015.
- [65] A. P. West, G. S. Shadel, and S. Ghosh, "Mitochondria in innate immune responses," *Nature Reviews Immunology*, vol. 11, no. 6, pp. 389–402, 2011.
- [66] E. L. Robb, A. R. Hall, T. A. Prime et al., "Control of mitochondrial superoxide production by reverse electron transport at complex I," *The Journal of Biological Chemistry*, vol. 293, no. 25, pp. 9869–9879, 2018.
- [67] A. Guarás, E. Perales-Clemente, E. Calvo et al., "The CoQH₂/CoQ ratio serves as a sensor of respiratory chain efficiency," *Cell Reports*, vol. 15, no. 1, pp. 197–209, 2016.
- [68] J. C. Crack, J. Green, M. R. Cheesman, N. E. le Brun, and A. J. Thomson, "Superoxide-mediated amplification of the oxygen-induced switch from [4Fe-4S] to [2Fe-2S] clusters in the transcriptional regulator FNR," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 104, no. 7, pp. 2092–2097, 2007.
- [69] S. J. Dixon and B. R. Stockwell, "The role of iron and reactive oxygen species in cell death," *Nature Chemical Biology*, vol. 10, no. 1, pp. 9–17, 2014.
- [70] D. B. Zorov, M. Juhaszova, and S. J. Sollott, "Mitochondrial reactive oxygen species (ROS) and ROS-induced ROS release," *Physiological Reviews*, vol. 94, no. 3, pp. 909–950, 2014.
- [71] J. E. Silva, "Thermogenic mechanisms and their hormonal regulation," *Physiological Reviews*, vol. 86, no. 2, pp. 435–464, 2006.
- [72] C. Bell, N. R. Stob, and D. R. Seals, "Thermogenic responsiveness to β -adrenergic stimulation is augmented in exercising versus sedentary adults: role of oxidative stress," *The Journal of Physiology*, vol. 570, Part 3, pp. 629–635, 2006.
- [73] S. K. Calderwood and J. Gong, "Heat shock proteins promote cancer: it's a protection racket," *Trends in Biochemical Sciences*, vol. 41, no. 4, pp. 311–323, 2016.
- [74] J. B. A. G. Haanen, "Converting cold into hot tumors by combining immunotherapies," *Cell*, vol. 170, no. 6, pp. 1055–1056, 2017.
- [75] M. Zizi, M. Forte, E. Blachly-Dyson, and M. Colombini, "NADH regulates the gating of VDAC, the mitochondrial outer membrane channel," *The Journal of Biological Chemistry*, vol. 269, no. 3, pp. 1614–1616, 1994.
- [76] R. Diaz-Ruiz, M. Rigoulet, and A. Devin, "The Warburg and Crabtree effects: on the origin of cancer cell energy metabolism and of yeast glucose repression," *Biochimica et Biophysica Acta (BBA) - Bioenergetics*, vol. 1807, no. 6, pp. 568–576, 2011.
- [77] S. F. Okada, W. K. O'Neal, P. Huang et al., "Voltage-dependent anion channel-1 (VDAC-1) contributes to ATP release and cell volume regulation in murine cells," *The Journal of General Physiology*, vol. 124, no. 5, pp. 513–526, 2004.
- [78] D. Ben-Hail, R. Begas-Shvartz, M. Shalev et al., "Novel compounds targeting the mitochondrial protein VDAC1 inhibit apoptosis and protect against mitochondrial dysfunction," *The Journal of Biological Chemistry*, vol. 291, no. 48, pp. 24986–25003, 2016.
- [79] E. Guerriero, A. Sorice, F. Capone et al., "Vitamin C effect on mitoxantrone-induced cytotoxicity in human breast cancer cell lines," *PLoS One*, vol. 9, no. 12, article e115287, 2014.
- [80] E. Hatem, S. Azzi, N. el Banna et al., "Auranofin/vitamin C: a novel drug combination targeting triple-negative breast cancer," *JNCI: Journal of the National Cancer Institute*, vol. 111, no. 6, pp. 597–608, 2019.
- [81] E. Simamura, K. Hirai, H. Shimada, J. Koyama, Y. Niwa, and S. Shimizu, "Furanonaphthoquinones cause apoptosis of

- cancer cells by inducing the production of reactive oxygen species by the mitochondrial voltage-dependent anion channel,” *Cancer Biology & Therapy*, vol. 5, no. 11, pp. 1523–1529, 2006.
- [82] S. Vatolin, T. Radivoyevitch, and J. P. Maciejewski, “New drugs for pharmacological extension of replicative life span in normal and progeroid cells,” *npj Aging and Mechanisms of Disease*, vol. 5, no. 1, 2019.
- [83] J. Verrax, J. Stockis, A. Tison, H. S. Taper, and P. B. Calderon, “Oxidative stress by ascorbate/menadione association kills K562 human chronic myelogenous leukaemia cells and inhibits its tumour growth in nude mice,” *Biochemical Pharmacology*, vol. 72, no. 6, pp. 671–680, 2006.
- [84] M. Tomasetti, E. Strafella, S. Staffolani, L. Santarelli, J. Neuzil, and R. Guerrieri, “ α -Tocopheryl succinate promotes selective cell death induced by vitamin K3 in combination with ascorbate,” *British Journal of Cancer*, vol. 102, no. 8, pp. 1224–1234, 2010.
- [85] X. Ren, S. M. Santhosh, L. Coppo, F. T. Ogata, J. Lu, and A. Holmgren, “The combination of ascorbate and menadione causes cancer cell death by oxidative stress and replicative stress,” *Free Radical Biology & Medicine*, vol. 134, pp. 350–358, 2019.
- [86] J. T. Szilagyi, K. C. Fussell, Y. Wang et al., “Quinone and nitrofurantoin redox cycling by recombinant cytochrome b5 reductase,” *Toxicology and Applied Pharmacology*, vol. 359, pp. 102–107, 2018.
- [87] K. Nakagawa, Y. Hirota, N. Sawada et al., “Identification of UBIAD1 as a novel human menaquinone-4 biosynthetic enzyme,” *Nature*, vol. 468, no. 7320, pp. 117–121, 2010.
- [88] W. J. Fredericks, J. Sepulveda, P. Lal et al., “The tumor suppressor TERE1 (UBIAD1) prenyltransferase regulates the elevated cholesterol phenotype in castration resistant prostate cancer by controlling a program of ligand dependent SXR target genes,” *Oncotarget*, vol. 4, no. 7, pp. 1075–1092, 2013.
- [89] D. Ivanova, Z. Zhelev, D. Lazarova, P. Getsov, R. Bakalova, and I. Aoki, “Vitamins C and K3: a powerful redox system for sensitizing leukemia lymphocytes to everolimus and barasertib,” *Anticancer Research*, vol. 38, no. 3, pp. 1407–1414, 2018.
- [90] G. Lenaz and M. Genova, “Structural and functional organization of the mitochondrial respiratory chain: a dynamic super-assembly,” *The International Journal of Biochemistry & Cell Biology*, vol. 41, no. 10, pp. 1750–1772, 2009.
- [91] R. Acin-Perez and J. A. Enriquez, “The function of the respiratory supercomplexes: the plasticity model,” *Biochimica et Biophysica Acta (BBA) - Bioenergetics*, vol. 1837, no. 4, pp. 444–450, 2014.
- [92] K. J. Lenton, H. Therriault, A. M. Cantin, T. Fülöp, H. Payette, and J. R. Wagner, “Direct correlation of glutathione and ascorbate and their dependence on age and season in human lymphocytes,” *The American Journal of Clinical Nutrition*, vol. 71, no. 5, pp. 1194–1200, 2000.
- [93] A. C. Carr and J. Cook, “Intravenous vitamin C for cancer therapy – identifying the current gaps in our knowledge,” *Frontiers in Physiology*, vol. 9, p. 1182, 2018.
- [94] N. Shenoy, E. Creagan, T. Witzig, and M. Levine, “Ascorbic acid in cancer treatment: let the phoenix fly,” *Cancer Cell*, vol. 34, no. 5, pp. 700–706, 2018.
- [95] D. Speijer, “Oxygen radicals shaping evolution: why fatty acid catabolism leads to peroxisomes while neurons do without it,” *BioEssays*, vol. 33, no. 2, pp. 88–94, 2011.
- [96] H. Dubouchaud, L. Walter, M. Rigoulet, and C. Batandier, “Mitochondrial NADH redox potential impacts the reactive oxygen species production of reverse electron transfer through complex I,” *Journal of Bioenergetics and Biomembranes*, vol. 50, no. 5, pp. 367–377, 2018.