

# Does stopping left-right nodal flow mirror impaired mechanosensation in the left-right organizer?

R Jacinto\*, P Sampaio, M Roxo-Rosa, SS Lopes

From Cilia 2014 - Second International Conference  
Paris, France. 18-21 November 2014

## Background

In most vertebrates organ asymmetries arise in early development through the left-right organizer (LRO). LRO ciliated cells induce a leftward fluid flow responsible for the asymmetric expression of Nodal pathway genes in tissues that will originate internal organs. We have recently published that *charon/dand5* transcription is an early flow target in zebrafish by being expressed in the LRO, first symmetrically and later asymmetrically to the right where the fluid flow is weaker. We showed that absence of flow originates symmetric *charon* expression [1]. Pkd2 has been reported as part of a mechanosensor complex that senses flow and induces a calcium inward flux in kidney cells [2] and mouse LRO [3]. In agreement, mouse and zebrafish Pkd2 mutants have LR defects [4,5].

## Objective

Determine if impaired mechanosensing renders the same *charon/dand5* phenotype as absent flow.

## Methods

We injected two morpholinos in zebrafish embryos: a *dnah7*-MO to stop cilia movement [1] and a *pkd2*-MO. We also co-injected both. We recorded flow by particle analysis, quantified Pkd2 expression and correlated these with *charon/dand5* expression patterns by *in situ* hybridization and qRT-PCR.

## Results

50-60% of injected embryos have symmetric *charon/dand5* expression when flow is low and homogeneous or when

Pkd2 is absent from LRO ciliary membrane. We will show if *charon/dand5* quantification by qRT-PCR is also similar. When both flow and Pkd2 are impaired, 80% of embryos showed symmetric *charon/dand5*, which indicates a synergistic effect.

## Conclusion

Although no flow and no Pkd2 have virtually the same phenotype, there seems to be a positive epistasis when both are affected.

## Acknowledgements

Supported by FCT-ANR/BEX-BID/0153/2012 grant and PD/BD/52420/2013 PhD scholarship.

Published: 13 July 2015

## References

1. Sampaio P, Ferreira R, Guerrero A, Pintado P, Tavares B, Amaro J, Smith A, Montenegro-Johnson T, Lopes SS: **Left-right organizer flow Dynamics: how much cilia activity reliably yields laterality?** *Dev Cell* 2014, **29**(6):716-728.
2. Alenghat FJ, Luo Y, Williams E, Vassilev P, Li X, et al: **Polycystins 1 and 2 mediate mechanosensation in the primary cilium of kidney cells.** *Nat Genet* 2003, **33**(2):129-137.
3. McGrath J, Somlo S, Makova S, Tian X, Brueckner M: **Two populations of node monocilia initiate left-right asymmetry in the mouse.** *Cell* 2003, **114**(1):61-73.
4. Pennekamp P, Karcher C, Fisher A, Schweickert A, Skryabin B, Horst J, Blum M, Dworniczak B: **The ion channel polycystin-2 is required for left-right axis determination in mice.** *Current Biol* 2002, **12**(11):938-943.
5. Schottenfeld J, Sullivan-Brown J, Burdine R: **Zebrafish curly up encodes a Pkd2 ortholog that restricts left-side-specific expression of southpaw.** *Development* 2002, **134**(8):1605-1615.

doi:10.1186/2046-2530-4-S1-P39

Cite this article as: Jacinto et al.: Does stopping left-right nodal flow mirror impaired mechanosensation in the left-right organizer? *Cilia* 2015 **4**(Suppl 1):P39.