

ORIGINAL RESEARCH

Reduction of \dot{V}_{O_2} slow component by priming exercise: novel mechanistic insights from time-resolved near-infrared spectroscopy

Yoshiyuki Fukuoka^{1,2}, David C. Poole³, Thomas J. Barstow³, Narihiko Kondo⁴, Masato Nishiwaki⁵, Dai Okushima⁶ & Shunsaku Koga⁶

1 Environmental Physiology Laboratory, Prefectural University of Kumamoto, Kumamoto, Japan

2 Graduate School of Health and Sports Science, Doshisha University, Kyoto, Japan

3 Departments of Anatomy and Physiology and Kinesiology, Kansas State University, Manhattan, Kansas

4 Graduate School of Cultural Studies and Human Science, Kobe University, Kobe, Japan

5 Faculty of Engineering, Osaka Institute of Technology, Osaka, Japan

6 Applied Physiology Laboratory, Kobe Design University, Kobe, Japan

Keywords

Heavy exercise, hemoglobin concentration, muscle microcirculation, muscle O_2 diffusing capacity.

Correspondence

Yoshiyuki Fukuoka, Faculty of Health and Sports Science, Doshisha University, 1-3 Kyotanabe, Kyoto 610-0394, Japan.

Tel: +81-774-65-7530

Fax: +81-774-65-6029

E-mail: yfukuoka@mail.doshisha.ac.jp

Funding Information

This investigation was supported by a grant for scientific research from the Ministry of Education, Science, and Culture of Japan (Grant-in-Aid for Scientific Research 20650103, 24650401) to S. Koga.

Received: 30 January 2015; Revised: 12 May 2015; Accepted: 15 May 2015

doi: 10.14814/phy2.12432

Physiol Rep, 3 (6), 2015, e12432,
doi:10.14814/phy2.12432

Abstract

Novel time-resolved near-infrared spectroscopy (TR-NIRS), with adipose tissue thickness correction, was used to test the hypotheses that heavy priming exercise reduces the \dot{V}_{O_2} slow component ($\dot{V}_{O_{2SC}}$) (1) by elevating microvascular [Hb] volume at multiple sites within the quadriceps femoris (2) rather than reducing the heterogeneity of muscle deoxygenation kinetics. Twelve subjects completed two 6-min bouts of heavy work rate exercise, separated by 6 min of unloaded cycling. Priming exercise induced faster overall \dot{V}_{O_2} kinetics consequent to a substantial reduction in the $\dot{V}_{O_{2SC}}$ (0.27 ± 0.12 vs. 0.11 ± 0.09 L·min⁻¹, $P < 0.05$) with an unchanged primary \dot{V}_{O_2} time constant. An increased baseline for the primed bout [total (Hb + Mb)] (197.5 ± 21.6 vs. 210.7 ± 22.5 μ mol L⁻¹, $P < 0.01$), reflecting increased microvascular [Hb] volume, correlated significantly with the $\dot{V}_{O_{2SC}}$ reduction. At multiple sites within the quadriceps femoris, priming exercise reduced the baseline and slowed the increase in [deoxy (Hb + Mb)]. Changes in the inter-site coefficient of variation in the time delay and time constant of [deoxy (Hb + Mb)] during the second bout were not correlated with the $\dot{V}_{O_{2SC}}$ reduction. These results support a mechanistic link between priming exercise-induced increase in muscle [Hb] volume and the reduced $\dot{V}_{O_{2SC}}$ that serves to speed overall \dot{V}_{O_2} kinetics. However, reduction in the heterogeneity of muscle deoxygenation kinetics does not appear to be an obligatory feature of the priming response.

Introduction

Exercise performed above the lactate threshold in the heavy domain evinces a \dot{V}_{O_2} slow component ($\dot{V}_{O_{2SC}}$), superimposed upon the primary (fast) \dot{V}_{O_2} kinetics, that elevates pulmonary and muscle \dot{V}_{O_2} . This “extra” \dot{V}_{O_2} , which may exceed 1000 mL·min⁻¹, slows overall \dot{V}_{O_2} kinetics (Gerbi-

no et al. 1996) and is associated with compromised exercise tolerance (Murgatroyd and Wylde 2011; Grassi et al. 2015). Heavy priming exercise substantially reduces the $\dot{V}_{O_{2SC}}$ contribution to the achieved \dot{V}_{O_2} in the presence or absence of speeded fast component kinetics (Bangsbo et al. 2000; Rossiter et al. 2001; Tordi et al. 2003; Paterson et al. 2005; Sahlin et al. 2005; Poole and Jones 2012).

Resolution of the mechanistic bases for priming exercise-induced $\dot{V}_{O_{2SC}}$ reduction will significantly improve our understanding of muscle energetics. In this regard, current theories for the etiology of the reduced $\dot{V}_{O_{2SC}}$ include increased bulk and local blood flow and O_2 delivery (\dot{Q}_{O_2}) via residual vasodilation and acidemia-induced rightward shift of the hemoglobin (Hb) O_2 dissociation curve (Poole et al. 1991; Saitoh et al. 2009; Jones et al. 2011). These effects would (presumably) serve to improve matching of \dot{Q}_{O_2} -to- \dot{V}_{O_2} thereby raising muscle and microvascular oxygen pressure (PO_2) and enhancing blood-myocyte O_2 flux and mitochondrial control by raising intracellular PO_2 (McDonough et al. 2005). In support of this contention, several previous studies carried out by conventional (continuous wave, CW) near-infrared spectroscopy (NIRS) observed a priming exercise-induced slowing of muscle deoxygenation (i.e., [deoxy (Hb + Mb)]) kinetics (Rossiter 2011) and a reduced [deoxy (Hb + Mb)] at the baseline of the second exercise bout (Spencer et al. 2012). However, multichannel CW-NIRS found that the reduction in spatial heterogeneity across 10 sites in the quadriceps femoris muscle(s) did not correlate with the priming exercise-induced decrease in the $\dot{V}_{O_{2SC}}$ (Saitoh et al. 2009), suggesting that improved matching of \dot{Q}_{O_2} -to- \dot{V}_{O_2} was not responsible. One striking limitation of CW-NIRS is that it assesses relative rather than absolute [deoxy (Hb + Mb)] and thus exercise-induced alterations and variability in the optical physics (i.e., absorbance, scattering, path length) and also adipose tissue thickness (ATT) among subjects may have obscured the underlying response(s) (Koga et al. 2011). Consequently, it remains unknown to what extent prior heavy exercise influences the temporal and spatial profiles of the absolute [deoxy (Hb + Mb)] (ATT corrected) (Bowen et al. 2013) and consequently the mean muscle PO_2 (Koga et al. 2012) of the different muscle regions during subsequent heavy exercise.

To circumvent these problems we have established a state-of-the art method to quantify absolute [deoxy (Hb + Mb)] and its spatial heterogeneity within the quadriceps femoris (QF) muscle during cycle exercise using time-resolved (TR) NIRS with ATT correction (Chin et al. 2011; Koga et al. 2011, 2014; Bowen et al. 2013). Moreover, based upon our current understanding of muscle O_2 diffusion where the muscle O_2 diffusing capacity (D_mO_2) is determined by the number/volume of red blood cells (RBCs) in the capillary bed adjacent to muscle fibers at any instant (Federspiel and Popel 1986; Groebe and Thews 1990), we hypothesized that priming exercise-induced increases in [total (Hb + Mb)] (Burnley et al. 2002; DeLorey et al. 2007; Jones et al. 2008) would relate quantitatively to $\dot{V}_{O_{2SC}}$ reductions. In addition, if this represents a crucial mechanism for improving myo-

cyte O_2 delivery and metabolic control, decreased \dot{Q}_{O_2} -to- \dot{V}_{O_2} heterogeneity would not be an obligatory facet of the reduced $\dot{V}_{O_{2SC}}$ response. Accordingly, using TR-NIRS across multiple QF muscle sites during primed heavy exercise we tested the following hypotheses: (1) The $\dot{V}_{O_{2SC}}$ would be reduced in proportion to the greater [total Hb] volume present. (2) The faster overall \dot{V}_{O_2} kinetics would not be dependent upon a reduction in the heterogeneity of the absolute [deoxy (Hb + Mb)] kinetics.

Methods

Subjects

Twelve physically active healthy male subjects (age, 23.4 ± 4.3 years; height, 173.8 ± 4.8 cm; and weight, 62.5 ± 6.3 kg) participated in this study. The study was approved by the Human Subjects Committee of Kobe Design University, in accordance with the Declaration of Helsinki. Explanation of the experimental protocol, including any possible risks and benefits associated with the testing procedures, were discussed with the subjects, and written consent was obtained from all subjects before voluntary participation in this study.

Correction of amplitude of muscle deoxygenation for adipose tissue thickness

Measurement of ATT under each optode site over the vastus lateralis (VL) and the rectus femoris (RF) muscles at the distal- and proximal sites was made by B-mode ultrasound (model Logiq 400; GE-Yokogawa Medical Systems, Japan) with the subject in an upright position (Chin et al. 2011; Koga et al. 2011). In order to quantify the influence of ATT on the NIRS signal (Niwayama et al. 2000; Koga et al. 2011), we applied a linear regression of the relationship between [total (Hb + Mb)] and ATT that was established by a previous report (Bowen et al. 2013). Resting [total (Hb + Mb)] was determined at each site from a 2 min resting average in the upright-seated position. Measured [deoxy (Hb + Mb)] and [total (Hb + Mb)] values at each individual muscle site were corrected to a common ATT of 0 mm on this regression of [total (Hb + Mb)] and ATT. This normalization process allowed absolute values of both [deoxy (Hb + Mb)] and [total (Hb + Mb)] to be compared between subjects and muscle sites differing in ATT.

Exercise tests

On each data-collection day, subjects reported to the laboratory at least 2 h after their last meal. They were asked to avoid caffeine and alcohol ingestion and strenuous

exercise for 24 h before the test. The temperature and relative humidity of the laboratory were maintained at 25°C and 50%, respectively. On the first visit, seat height and handlebar position on an electronically braked cycle ergometer (Combi 232C, Tokyo, Japan) were recorded and reproduced on subsequent tests.

Incremental exercise tests

The first visit was used to familiarize the subjects with testing procedures and to determine the peak \dot{V}_{O_2} , gas exchange threshold (GET), and work rates for the constant work-rate tests. The protocol, which was designed to produce volitional exhaustion within 10–15 min, consisted of 4 min of unloaded exercise, followed by work rate increases of 25–30 W·min⁻¹ until the limit of tolerance (Chin et al. 2011; Koga et al. 2011). Pedaling frequency was held constant at 60 rpm for all exercise bouts with the aid of an audible metronome. The peak \dot{V}_{O_2} was defined as the highest \dot{V}_{O_2} achieved during the test averaged over a 20-sec interval. The GET was estimated from gas exchange measurements using the V-slope method, ventilatory equivalents, and end-tidal gas tensions (Beaver et al. 1986).

Constant work-rate exercise tests

Square-wave exercise transition tests were conducted on separate days. Each constant work-rate exercise test was performed for 6 min. The work rates for heavy exercise were calculated to elicit 40% of the difference between the GET \dot{V}_{O_2} and peak \dot{V}_{O_2} [$\Delta 40 \dot{V}_{O_2} = \text{GET } \dot{V}_{O_2} + 0.4 (\text{peak } \dot{V}_{O_2} - \text{GET } \dot{V}_{O_2})$], based on the \dot{V}_{O_2} /work rate with account taken of the lag in \dot{V}_{O_2} relative to the work rate during ramp exercise. Over the next two visits, repeated bouts of heavy cycling exercise were performed twice at approximately the same time of day, with 3 days between visits. Moreover, a subset of four subjects performed, on separate days, an additional heavy exercise bout designed to elicit 20% of the difference between the GET \dot{V}_{O_2} and peak \dot{V}_{O_2} ($\Delta 20 \dot{V}_{O_2}$). The constant work-rate exercise protocol consisted of 1 min of rest and 4 min of unloaded exercise, followed by two 6-min bouts of heavy exercise separated by 6 min of unloaded exercise at a pedal frequency of 60 rpm.

Measurements

Pulmonary \dot{V}_{O_2}

Subjects breathed through a low-resistance hot-wire flowmeter for measurement of inspiratory and expiratory flows. The flowmeter was calibrated repeatedly by

inputting known volumes of room air at various mean flows and flow profiles. Expired oxygen and carbon dioxide concentrations were determined by gas analysis (Minato-Medical AE-300S, Japan) from a sample drawn continuously from the mouthpiece. Alveolar gas exchange variables were calculated breath-by-breath gas exchange measurement system (Minato-Medical AE-300S, Japan) (Beaver et al. 1981) from time aligned gas volume and concentration signals.

Muscle deoxygenation

The absolute muscle [deoxy (Hb + Mb)], [oxy (Hb + Mb)], and [total (Hb + Mb)] profiles at 4 sites in the quadriceps of the dominant leg were measured by two time-resolved spectroscopy (TRS) NIRS systems (TRS-20 each with two channels, Hamamatsu Photonics K.K., Japan). This system measures the distribution of in vivo optical path lengths, thereby enabling the determination of absolute [Hb + Mb] concentration ($\mu\text{mol L}^{-1}$). Previous studies showed that the deoxygenation measured by the TRS correlated significantly with the oxyhemoglobin saturation in both the blood and a purified-hemoglobin phantom solution (Hamaoka et al. 2000).

The optodes were housed in black rubber holders that helped to minimize extraneous movement, thus ensuring that the position of the optodes was fixed and invariant. The distal optodes were placed on the lower third of the VL and the RF muscles parallel to the major axis of the thigh. The location of the distal optodes on the VL muscle was chosen to represent the single site NIRS measurement conducted by previous studies (e.g., (Ferreira et al. 2007, 2005; Wilkerson et al. 2004)). The proximal optode pairs on the VL and the RF muscles were located ~10–15 cm from the distal optode pairs. The interoptode spacing between emitter and receiver was 3 cm. The depth of the measured area was assumed to be approximately half of the distance between the emitter and the receiver, ~1.5 cm. The skin under the probes was carefully shaved. Pen marks were made on the skin to indicate the margins of the rubber holder to check for any downward sliding of the probe during cycling and for accurate probe repositioning on subsequent days. No sliding was observed in any subject at the end of each protocol. The principles of operation and algorithms utilized by the equipment have been described in detail elsewhere (Oda et al. 1999; Ohmae et al. 2006). Briefly, the TRS system consists of a pico-second (ps) light pulser that emits three wavelengths (760, 795, and 830 nm), with a repetition frequency of 5 MHz and a full width at half maximum (FWHM) of 100 ps. A time-correlated single-photon counting board was installed to acquire temporal profiles of photons. The laser diodes and photomultiplier tube are connected to a

lightweight plastic probe by optical fibers for single-photon detection. The time-correlated single-photon counting board is used for the parallel acquisition of time-resolved reflectance curves. In the present study, estimation of the optical path length, reduced scattering coefficient (μ_s') and absorption coefficient (μ_a) are achieved by fitting the receiver profile of photon counts over time to a function based on diffusion theory (Oda et al. 1999; Ohmae et al. 2006). The output frequency was selected as 0.5 Hz. We reasoned that, as [Mb] would not be expected to change within the sampled volumes, any increased [Hb + Mb] reflected that of [Hb] (Davis and Barstow 2013). Validation of the equipment was performed before each test, by measuring the instrument's responses when the input and receiving fibers faced each other through a neutral-density filter in a black tube.

Data analysis

Individual responses of pulmonary \dot{V}_{O_2} and [deoxy (Hb + Mb)] during the baseline (BL)-to-exercise transitions were time-interpolated to 1-sec intervals, and averaged across each transition for each subject. The response curve of \dot{V}_{O_2} was fit by a three-term exponential function (Eq. 1) that included amplitudes, time constants, and time delays, using nonlinear least-squares regression techniques (Gerbino et al. 1996; Fukuba et al. 2002; Grassi et al. 2003; Ferreira et al. 2005). The computation of best-fit parameters was chosen by the program (Kaleida Graph) so as to minimize the sum of the squared differences between the fitted function and the observed response.

The first exponential term started with the onset of exercise and the second and third terms began after independent time delays (Ma et al. 2010).

$$H(t) = 0 \text{ for } t < 0 \\ 1 \text{ for } t \geq 0$$

$$\begin{aligned} \dot{V}_{O_2}(t) = & \dot{V}_{O_2}(\text{BL}) + H \cdot (t) \cdot A_i \cdot (1 - e^{-t/\tau_i}) \\ & \text{phase 1 [initial (cardio dynamic) component]} \\ & + H \cdot (t - \text{TD}_p) \cdot A_p \cdot [1 - e^{-(t-\text{TD}_p)/\tau_p}] \\ & \text{phase 2 (primary component)} \\ & + H \cdot (t - \text{TD}_s) \cdot A_s \cdot [1 - e^{-(t-\text{TD}_s)/\tau_s}] \\ & \text{phase 3 (slow component, SC)} \end{aligned} \quad (1)$$

where the subscripts *i*, *p*, and *s* refer to initial, primary, and slow components, respectively; \dot{V}_{O_2} (BL) is the unloaded exercise baseline value; A_i , A_p , and A_s are the asymptotic amplitudes for the exponential terms; τ_i , τ_p , and τ_s are the time constants; and TD_p , and TD_s are the time delays. Mean response time (MRT_p) for the primary

phase of \dot{V}_{O_2} was defined as the sum of $\text{TD}_p + \tau_p$. The phase I \dot{V}_{O_2} at the start of phase II (i.e., at TD_p) was assigned the value for that time (A_i'). The physiologically relevant amplitude of the primary exponential component during phase II (A_p') was defined as the sum of $A_i' + A_p$. Because of concerns regarding the validity of using the extrapolated asymptotic value for the SC (A_s) for comparisons, we used the value of the slow exponential function at the end of exercise, defined as A_s' . Alternatively, the slow component of \dot{V}_{O_2} was calculated as the change from 3 to 6 min ($\dot{V}_{O_2}(6-3)$).

Subsequently, absolute [deoxy (Hb + Mb)] data were then fit from the time of initial and primary increases in [deoxy (Hb + Mb)] to 120 s with a two-exponential model of the form in Eq. 2 to determine the time course of muscle deoxygenation.

$$\begin{aligned} [\text{Deoxy(Hb + Mb)}](t) \\ = & [\text{deoxy(Hb + Mb)}](\text{BL}) + A_i \cdot [1 - e^{t/\tau_i}] \\ & \text{phase 1 (initial component)} + A_p \cdot [1 - e^{-(t-\text{TD})/\tau_p}] \\ & \text{phase 2 (primary component)}, \end{aligned} \quad (2)$$

where the subscripts *i* and *p* refer to initial and primary components, respectively; [deoxy (Hb + Mb)] (BL) is the unloaded exercise baseline value; A_i and A_p are the asymptotic amplitudes for the exponential terms; τ_i and τ_p are the time constants; and TD is the initial component duration from the onset of exercise to the onset of primary component of [deoxy (Hb + Mb)]. The absolute primary amplitude (A_a) was defined as the sum of $A_p + \text{BL}$. The TD and τ_p of the [deoxy (Hb + Mb)] response were summed (MRT_p) to provide an indication of the overall dynamics of the primary component (Koga et al. 2007).

The averaged slow components of both the [deoxy (Hb + Mb)] and [total (Hb + Mb)] (i.e., [deoxy (Hb + Mb)](6-3) and [total (Hb + Mb)](6-3)) were calculated as the changes from 3 to 6 min across the four sites. The priming-induced reductions in both [deoxy (Hb + Mb)](6-3) and [total (Hb + Mb)](6-3) from the first bout to the second bout were calculated to analyze the correlation with the reduced slow component of \dot{V}_{O_2} ($\dot{V}_{O_2}(6-3)$).

Statistics

Data are presented as mean \pm SD. Intersite coefficient of variation [CV (%); 100-SD/mean of the four sites values] for each subject was calculated to show spatial heterogeneity of the amplitude and kinetic profiles of the muscle deoxygenation (Saitoh et al. 2009). When the comparison for differences between the first and the second bouts was justified, a paired t-test was used. Significance

was accepted when $P < 0.05$. A two-way ANOVA [sites (distal/proximal) and muscles (VL/RF)] was performed to evaluate significant differences in the muscle deoxygenation profiles. When a significant difference was detected, this was further examined by post hoc *Scheffe* test. The relationship between two variables was analyzed by the correlation or linear regression analysis.

Results

Peak \dot{V}_{O_2} /body mass reflected the active lifestyle for these subjects ($52.5 \pm 6.2 \text{ mL}\cdot\text{min}^{-1}\cdot\text{kg}^{-1}$). The average work rates at the GET + $\Delta 40$ and $\Delta 20$ were $183 \pm 25 \text{ W}$ and $163 \pm 20 \text{ W}$, respectively.

Pulmonary O_2 uptake kinetics

Priming increased baseline \dot{V}_{O_2} and substantially reduced the $\dot{V}_{O_{2SC}}$ (A_s') 59% (both $P < 0.05$, Fig. 1 and Table 1) without altering end-exercise \dot{V}_{O_2} . In addition, priming caused a significant shortening of the primary component time delay (TD_p , $P < 0.05$) and consequently the mean response time (MRT_p , $\sim 7\%$, $P < 0.05$) but did not change either the time constant (τ_p) or the amplitude (A_p' , phase I + phase II).

[total (Hb + Mb)]

As shown in the [total (Hb + Mb)] profiles at each site in Figure 2, the averaged baseline [total (Hb + Mb)] response across the four muscle sites after priming was significantly greater than for the first bout of exercise (197.5 ± 21.6 vs. $210.7 \pm 22.5 \mu\text{mol L}^{-1}$, $P < 0.01$). The

averaged end-exercise [total (Hb + Mb)] across the four muscle sites after priming was also significantly elevated (218.8 ± 22.8 vs. $225.7 \pm 24.0 \mu\text{mol L}^{-1}$, $P < 0.05$), while the averaged amplitude of [total (Hb + Mb)] was significantly reduced after priming (23.3 ± 3.0 vs. $15.1 \pm 2.0 \mu\text{mol L}^{-1}$, $P < 0.05$). Specifically, at both proximal and distal RF sites the end-exercise values of the [total (Hb + Mb)] response were increased after priming, ($P < 0.05$) whereas those for the VL muscle were not (proximal, $P = 0.083$; distal, $P = 0.081$) despite a significantly greater baseline ($P < 0.05$).

At the $\Delta 40\dot{V}_{O_2}$ (GET + $\Delta 40$) exercise intensity the reduced $\dot{V}_{O_{2SC}}$ (Table 1) was correlated with an increased BL of [total (Hb + Mb)] across the four muscle sites ($r = -0.720$, $P < 0.01$). As expected, for $\Delta 20\dot{V}_{O_2}$ (GET + $\Delta 20$, $n = 4$) there was a much smaller reduction of $\dot{V}_{O_{2SC}}$ compared with $\Delta 40\dot{V}_{O_2}$ (i.e., ~ 60 vs. $160 \text{ mL}\cdot\text{min}^{-1}$, $P < 0.05$). For both $\Delta 20\dot{V}_{O_2}$ and $\Delta 40\dot{V}_{O_2}$ data sets the reduced $\dot{V}_{O_{2SC}}$ was correlated significantly with the increased [total (Hb + Mb)] BL across the four muscle sites ($r = -0.731$, $P < 0.01$, Figure 3, 95% confidential interval, $\pm 50.4 \text{ mL}\cdot\text{min}^{-1}$). In contrast, the reduction in [total (Hb + Mb)](6-3) or reduced amplitude of [total (Hb + Mb)] response from the first bout to the second bout was not related to the \dot{V}_{O_2} (6-3) reduction from the first bout to the second bout ($r = -0.228$, $P = 0.472$; $r = 0.06$, $P = 0.853$, respectively). Only the increased BL in [total (Hb + Mb)] was clearly shown after priming to be related to the reduction of $\dot{V}_{O_{2SC}}$.




Figure 1. Mean pulmonary O_2 uptake responses to the first (square) and second (triangle) bouts of heavy cycling exercise along with the best fitting three-term exponential function.

Table 1. Parameters of pulmonary \dot{V}_{O_2} kinetics following the onset of two sequential bouts of heavy exercise.

	1st bout	2nd bout
BL, $L\cdot\text{min}^{-1}$	0.47 ± 0.09	$0.58 \pm 0.08^*$
TD_p , s	20.9 ± 7.4	$16.8 \pm 4.2^*$
τ_p , s	21.1 ± 6.5	22.5 ± 7.4
MRT_p , s	42.1 ± 6.2	$39.3 \pm 7.1^*$
A_p' , $L\cdot\text{min}^{-1}$	1.88 ± 0.33	1.79 ± 0.33
A_a , $L\cdot\text{min}^{-1}$	2.35 ± 0.31	2.37 ± 0.31
TD_s , s	143.4 ± 64.7	176.1 ± 86.8
A_s' , $L\cdot\text{min}^{-1}$	0.27 ± 0.12	$0.11 \pm 0.09^*$
\dot{V}_{O_2} (6-3), $L\cdot\text{min}^{-1}$	0.22 ± 0.13	$0.10 \pm 0.03^*$

Values are mean \pm SD. \dot{V}_{O_2} , O_2 uptake; BL, baseline; TD_p , time delay of phase II; τ_p , time constant of phase II; MRT_p , mean response time of the primary component of the response ($MRT_p = TD_p + \tau_p$); A_p' , amplitude of phase I + phase II, not including BL; A_a , absolute primary amplitude (BL + A_p'); TD_s , time delay of phase III (slow component); A_s' , amplitude of phase III; \dot{V}_{O_2} (6-3), the change in \dot{V}_{O_2} from 3 min to 6 min.

* $P < 0.05$, compared to 1st bout.




Figure 2. Adipose tissue thickness corrected absolute mean concentrations of total hemoglobin/myoglobin [total (Hb + Mb)] response at each site between first bout (square) and second bout (triangle) of heavy exercise using mean values of all subjects (please note that error bars are omitted for clarity). The baseline of [total (Hb + Mb)] response at each site before the second bout of exercise was significantly higher than for the first bout of exercise ($P < 0.05$).

[Deoxy (Hb + Mb)] kinetics and their spatial heterogeneity

Priming heavy exercise significantly increased averaged τ_p [deoxy (Hb + Mb)] (i.e., slower kinetics) across the four muscle sites ($P < 0.01$) and significantly shortened the TD (Fig. 4 and Table 2). Regarding the spatial heterogeneity of [deoxy (Hb + Mb)], the intersite TD CVs became significantly greater after priming ($P < 0.05$), while intersite CVs for τ_p were reduced ($P < 0.05$) and intersite CVs for MRT_p unchanged.

Baseline [deoxy (Hb + Mb)] was decreased significantly by priming and the amplitude of the response increased across the four muscle sites ($P < 0.05$, Fig. 4 and Table 3) such that the A_a (the sum of A_p + BL) was unchanged.

The intersite CV for baseline and A_p were not significantly different after priming. The proximal RF muscle was characterized by a greater [deoxy (Hb + Mb)] baseline compared with the proximal and distal VL for both exercise bouts. Priming increased (i.e., slowed) τ_p , decreased (i.e., made faster) TD, and decreased BL in both proximal and distal sites of the VL (Table 4).

Relationship between [deoxy (Hb + Mb)] heterogeneity and \dot{V}_{O_2} kinetics

The priming exercise-induced decrease in the \dot{V}_{O_2SC} was not correlated with the altered muscle deoxygenation kinetics (as reflected in the shorter TD and slower [deoxy (Hb + Mb)] response). In addition, the averaged [deoxy




Figure 3. Relationship between the priming-induced reduction in \dot{V}_{O_2} slow component and the increased [total (Hb + Mb)] at baseline across the four sites ($r = -0.731$, $P < 0.01$). Dotted lines indicate the 95% confidence interval. Open circles indicate subjects who only performed at $\Delta 40\dot{V}_{O_2}$ exercise intensity whereas gray and black circles are the four subjects who performed heavy exercise at both $\Delta 20\dot{V}_{O_2}$ and $\Delta 40\dot{V}_{O_2}$. The number from 1 to 4 is each subject's number.

[Hb + Mb]] slow component was calculated as the change in [deoxy (Hb + Mb)] from 3 to 6 min across four sites. No correlation was found between the priming-induced small reduction in [deoxy (Hb + Mb)](6-3) and that of \dot{V}_{O_2} (6-3) ($r = -0.127$, $P = 0.694$).

Discussion

The original findings of this investigation were consistent with our two hypotheses. First, absolute [total (Hb + Mb)] was increased at baseline (Burnley et al. 2002; DeLorey et al. 2007; Jones et al. 2008) within and among muscles/regions and this response significantly correlated with the reduced \dot{V}_{O_2SC} . Second, neither the reduction in the baseline [deoxy (Hb + Mb)] nor the reduced intersite CVs of the τ_p kinetics correlated with the decreased \dot{V}_{O_2SC} or any other differences in the \dot{V}_{O_2} response (i.e., reduced MRT_p primary component). These results suggest that neither priming-induced increases in the local \dot{Q}_{O_2} relative to \dot{V}_{O_2} , nor its spatial variation (i.e., intersite CVs of [deoxy (Hb + Mb)] kinetics), contributed to the altered pulmonary \dot{V}_{O_2} kinetics. However, the sustained increase in [total (Hb + Mb)] at baseline may, by augmenting D_mO_2 and the intracellular consequences of such (i.e., elevated $P_{intramyocyte}O_2$ -induced

enhancement of mitochondrial oxidative regulation), reduce the \dot{V}_{O_2SC} .

Effects of priming on pulmonary \dot{V}_{O_2} kinetics

Our finding of the speeded 'overall' \dot{V}_{O_2} kinetics (reduced TD_p and MRT_p without change in the primary component τ_p and reduction in the \dot{V}_{O_2SC} amplitude) after prior heavy exercise are consistent with the majority of previous studies (e.g. Bearden and Moffatt 2001; Burnley et al. 2000; Koppo and Bouckaert 2000; MacDonald et al. 1997; Perrey et al. 2003; Scheuermann et al. 2001) with a few exceptions (Bangsbo et al. 2000; Rossiter et al. 2001; Tor-di et al. 2003; Paterson et al. 2005; Sahlin et al. 2005). A reduction in the \dot{V}_{O_2SC} by itself (i.e., when primary component τ_p and A_p are unchanged as in this study) does not reflect a speeding of phase 2 \dot{V}_{O_2} kinetics. As previously determined (Gerbino et al. 1996; Jones et al. 2003, 2006) the faster overall \dot{V}_{O_2} kinetics occurred consequent to a reduced relative contribution of the \dot{V}_{O_2SC} to the overall response.

Mechanistic basis for reduced \dot{V}_{O_2SC}

The primary determinant of D_mO_2 is considered to be the number of RBCs in the capillary bed adjacent to the contracting muscle fibers at any given instant (Federspiel and Popel 1986; Groebe and Thews 1990). Thus, from Fick's law [$\dot{V}_{O_2SC} = D_mO_2 (P_{capillary}O_2 - P_{intramyocyte}O_2)$], for any given \dot{V}_{O_2} , the greater the D_mO_2 the less capillary-to-myocyte O_2 pressure differential is required. Capillary (or microvascular) PO_2 ($P_{capillary}O_2$) is set by the $\dot{Q}_{O_2}/\dot{V}_{O_2}$ relationship and, accordingly, if D_mO_2 increases, the requisite \dot{V}_{O_2} can be achieved with less reduction in $P_{intramyocyte}O_2$ and hence improved mitochondrial oxidative phosphorylation. The present findings of elevated total [Hb + Mb] after priming, resulting presumably from increased [Hb], provide support for priming exercise elevating D_mO_2 which is expected to raise $P_{intramyocyte}O_2$, thereby "tightening" mitochondrial control (decreasing $\Delta[PCr]$, $\Delta[Pi]$, $\Delta[NADH]$, and $\Delta[ADP]_{free}$, (Hogan et al. 1992)) and potentially reducing the \dot{V}_{O_2SC} .

Effect of prior heavy exercise on muscle deoxygenation kinetics

The absolute concentrations at BL for [deoxy (Hb + Mb)] at the four different sites before subsequent heavy exercise were significantly reduced compared to the first exercise bout. This extends the findings of Spencer et al. (2012) from single site VL measurements during moderate exercise and supports that local muscle \dot{Q}_{O_2}




Figure 4. Adipose tissue thickness corrected absolute mean concentrations of deoxyhemoglobin/myoglobin [deoxy (Hb + Mb)] response at each site for first (square) and second (triangle) bouts (error bars are omitted for clarity). A significantly lower baseline at the proximal and distal sites of the vastus lateralis (VL) muscle was present after priming.

was augmented in the presence of elevated muscle \dot{V}_{O_2} (DeLorey et al. 2007). Muscle \dot{V}_{O_2} is increased for a period following exercise [i.e., excess postexercise O_2 consumption (EPOC)], while fractional O_2 extraction is reduced, reflecting a slower decrease in blood flow relative to \dot{V}_{O_2} (Ferreira et al. 2006). The present results suggest that the sustained elevations of blood flow (and \dot{Q}_{O_2}) during postexercise (recovery) that increase $P_{\text{microvascular}}O_2$ (and $P_{\text{intramyocyte}}O_2$), and additionally may benefit heat and metabolite removal, may be relatively homogeneous. Also, differences in the Bohr effect (i.e., a rightward shift in the O_2 -Hb dissociation curve) due to lactic acidosis (Bhambhani et al. 1997; Grassi et al. 2003) and increases in muscle temperature (Koga et al. 2013) across the different sites could contribute to the reduced spatial and temporal heterogeneity at baseline in the primed muscles.

During exercise the macroscopic heterogeneity of O_2 delivery and \dot{V}_{O_2} is undoubtedly impacted by differences in motor unit/muscle recruitment patterns as well as inherent heterogeneities in vascular and metabolic control. To elucidate these effects more fully further studies must exploit technology that can address both microscopic- and macroscopic heterogeneity, that is, the broad extremities of mismatching of O_2 delivery and \dot{V}_{O_2} within and among contracting muscle(s).

Relationship between heterogeneity of muscle deoxygenation and pulmonary \dot{V}_{O_2} kinetics

The diminished [deoxy (Hb + Mb)] baseline prior to the second bout might be attributed in part to the increased

Table 2. Kinetics of [deoxy (Hb + Mb)] across the four sites and intersite CVs for the primary component at the onset of two bouts of heavy exercise.

	1st bout	2nd bout
TD, s	12.5 ± 4.3	9.5 ± 4.6*
τ_p , s	11.7 ± 3.8	19.4 ± 7.6**
MRT _p , s	24.1 ± 6.9	28.4 ± 9.0
CV of TD, %	38.7 ± 25.0	62.6 ± 31.9*
CV of τ_p , %	72.7 ± 34.4	49.0 ± 20.9*
CV of MRT _p , %	39.9 ± 14.3	49.7 ± 29.1

Values are mean ± SD. [deoxy (Hb + Mb)], change in absolute concentration of deoxyhemoglobin + myoglobin; CV, intersite coefficient of variation.

*,***P* < 0.05, 0.01, compared to 1st bout.

Table 3. Amplitude of [deoxy (Hb + Mb)] across the four sites and intersite CVs for the primary component at the onset of two bouts of heavy exercise.

	1st bout	2nd bout
BL, $\mu\text{mol L}^{-1}$	55.0 ± 7.3	50.2 ± 6.7*
A_p , $\mu\text{mol L}^{-1}$	24.5 ± 10.2	30.2 ± 15.9*
A_a , $\mu\text{mol L}^{-1}$	79.5 ± 11.1	78.2 ± 19.2
A_i , $\mu\text{mol L}^{-1}$	-1.4 ± 3.9	0.1 ± 6.7
[deoxy (Hb + Mb)](6-3), $\mu\text{mol L}^{-1}$	2.6 ± 2.6	2.1 ± 4.4
CV of BL, %	19.1 ± 10.4	21.7 ± 10.5
CV of A_p , %	35.7 ± 13.5	30.3 ± 13.7

Values are mean ± SD. A_p , amplitude of primary component; A_a , sum of A_p and BL; A_i , amplitude of initial component; [deoxy (Hb + Mb)](6-3), the slow component calculating as the changes from 3 min to 6 min.

**P* < 0.05, compared to 1st bout.

bulk and local O₂ delivery relative to \dot{V}_{O_2} (Hernández et al. 2010). However, the τ_p of the \dot{V}_{O_2} kinetics was not decreased from the first to the second bout. This suggests that within and among the muscles, muscle \dot{V}_{O_2} kinetics during the primary phase were not dependent on O₂ delivery (i.e., the condition was to the right of the O₂ dependency ‘tipping point’) (Poole and Jones 2005; Koga et al. 2007).

There is substantial heterogeneity in muscle blood flow within an active muscle (Whipp et al. 1995; Piiper 2000; Kalliokoski et al. 2006; Heinonen et al. 2007) and the effects of heavy priming exercise on \dot{V}_{O_2} (e.g., reduced $\dot{V}_{O_{2SC}}$) might be related to greater homogeneity of \dot{Q}_{O_2} distribution and matching to \dot{V}_{O_2} (Germino et al. 1996; DeLorey et al. 2004, 2007; Jones et al. 2006; Kalliokoski et al. 2006). Indeed, Saitoh et al. (2009) reported that priming reduced the dynamic spatial heterogeneities of muscle deoxygenation matching of \dot{Q}_{O_2} to \dot{V}_{O_2} within the active muscle (as reflected in the shorter TD and slower

Table 4. Amplitude and kinetics of [deoxy (Hb + Mb)] and their spatial heterogeneity following the onset of heavy exercise.

	1st bout					2nd bout						
	Baseline, $\mu\text{mol L}^{-1}$	A_i , $\mu\text{mol L}^{-1}$	A_p , $\mu\text{mol L}^{-1}$	TD, s	τ_p , s	MRT _p , s	Baseline, $\mu\text{mol L}^{-1}$	A_i , $\mu\text{mol L}^{-1}$	A_p , $\mu\text{mol L}^{-1}$	TD, s	τ_p , s	MRT _p , s
Distal RF	56.8 ± 9.1	-2.6 ± 2.2	20.6 ± 8.0	14.8 ± 6.7	17.9 ± 13.6	32.3 ± 12.8	54.0 ± 7.7	-0.9 ± 4.8	30.1 ± 19.7	8.0 ± 4.3 [§]	22.3 ± 19.2	29.7 ± 20.8
Distal VL	46.1 ± 6.2**	-1.0 ± 2.0	22.8 ± 11.2	13.1 ± 9.1	6.3 ± 2.5	19.3 ± 10.6	40.2 ± 7.0** [§]	-0.6 ± 2.0	32.7 ± 15.1	9.0 ± 8.1 [§]	19.1 ± 9.6 [§]	28.1 ± 13.8
Proximal RF	67.9 ± 17.7	-2.2 ± 3.3	28.7 ± 13.0	13.5 ± 6.8	15.1 ± 8.9	29.2 ± 13.0	62.6 ± 15.1	-0.8 ± 3.4	33.3 ± 19.9	15.6 ± 10.6	18.8 ± 11.2	33.6 ± 17.9
Proximal VL	51.9 ± 6.0**	-2.4 ± 4.7	24.8 ± 16.1	9.6 ± 3.9	8.4 ± 6.1	17.9 ± 4.3	45.3 ± 6.8** [§]	-0.7 ± 4.4	29.1 ± 15.5	6.7 ± 3.5*	15.1 ± 6.6 [§]	21.8 ± 8.9

Values are mean ± SD (*n* = 12).

*,***P* < 0.05, 0.01 versus Proximal RF, #*P* < 0.05 versus Distal RF, [§]*P* < 0.05, 0.01 versus 1st bout.

[deoxy (Hb + Mb)] kinetics). However in their study, the dynamic spatial heterogeneities of muscle deoxygenation did not correlate with the kinetics of phase II \dot{V}_{O_2} . Murias et al. (2011) reported that at the reduced moderate intensity, $\tau\dot{V}_{O_2}$ after priming correlated with the reduction in the [deoxy (Hb + Mb)]/ \dot{V}_{O_2} ratio (proportional to the inverse of \dot{Q}_{O_2}): The interpretation being that for $\tau\dot{V}_{O_2}$'s $> \sim 20$ s in a young, healthy population, improved microvascular O_2 distribution within the active tissues may play an important role in the faster rate of adjustment in \dot{V}_{O_2} during moderate exercise following priming. In contrast, the intransigent phase II $\tau\dot{V}_{O_2}$ in the present investigation (~ 20 s) suggests that the spatial heterogeneity of microvascular \dot{Q}_{O_2} (in relation to \dot{V}_{O_2}) is not related to (i.e., does not determine) the primary phase of \dot{V}_{O_2} kinetics during heavy exercise in active young subjects.

Comparison between RF and VL for the different [deoxy (Hb + Mb)] kinetics

The proximal RF was characterized by a greater baseline [deoxy (Hb + Mb)] for both bouts compared to the VL. Histochemically, RF contains proportionally fewer slow twitch fibers in the surface region compared with the VL (Johnson et al. 1973). Judging from our observations (Table 4), the greater baseline [deoxy (Hb + Mb)] in the proximal RF versus the proximal VL is the result of lower \dot{Q}_{O_2} relative to \dot{V}_{O_2} (i.e., higher $\dot{V}_{O_2}/\dot{Q}_{O_2}$). Interestingly, Chin et al. (2011) reported that, compared with the VL, the RF [deoxy (Hb + Mb)] kinetics evidenced a “right-shifted” response throughout ramp exercise. This may be evidence for a lower activation (supported by EMG) rather than a higher $\dot{Q}_{O_2}/\dot{V}_{O_2}$ at a given \dot{V}_{O_2} especially as muscles comprised of fast twitch fibers typically have far lower $\dot{Q}_{O_2}/\dot{V}_{O_2}$ (and thus lower $P_{\text{microvascular } O_2}$) than their slow twitch counterparts (Richardson et al. 1998; Behnke et al. 2002; McDonough et al. 2005). The $\dot{V}_{O_{2SC}}$ is linked either directly or indirectly to the recruitment of type II muscle fibers at higher work rates and/or to metabolic changes occurring within the initially recruited fibers (Grassi et al. 2015). Furthermore, it is thought that the availability of O_2 may play an important role in regulating the recruitment of these high-threshold type II motor units (Bearden and Moffatt 2001; Jones et al. 2011; MacDonald et al. 1997; McDonough et al. 2005; Piiper 2000; Saitoh et al. 2009). Thus, it is possible that the shorter TD, the slower τ , and/or the decreased spatial heterogeneity of muscle deoxygenation kinetics (as the primary component τ) in the second bout reflected an improved distribution of local \dot{Q} and matching of muscle \dot{Q}_{O_2} to muscle \dot{V}_{O_2} in those fibers recruited early into exercise which may have delayed their fatigue, thereby reducing the recruitment of more type II fibers. Therefore, the

greater intracellular PO_2 (as reflected in the greater muscle $\dot{Q}_{O_2}/\dot{V}_{O_2}$, detected as lower [deoxy (Hb + Mb)]), during the primary component might be related to the reduction in the SC. Alternatively, the changes in muscle deoxygenation kinetics might have decreased the degree of metabolic perturbation, implicated in the $\dot{V}_{O_{2SC}}$, within already recruited fibers.

In conclusion, the relationship between priming-induced increase in baseline [total (Hb + Mb)] (Burnley et al. 2002; DeLorey et al. 2007; Jones et al. 2008) within and among muscles/regions and the reduced $\dot{V}_{O_{2SC}}$ suggests that resultant enhancements of D_mO_2 may be responsible for the overall faster \dot{V}_{O_2} kinetics. Augmented D_mO_2 would be expected to raise $P_{\text{intramyocyte } O_2}$ and thereby enhance mitochondrial oxidative regulation and reduce $\dot{V}_{O_{2SC}}$. The temporal displacement between increased [total (Hb + Mb)] at baseline and reduction in $\dot{V}_{O_{2SC}}$ implies that better oxygenation during the initial fast component (phase II) kinetics relates quantitatively to subsequently improved metabolic control. In contrast, neither the degree of reduction in baseline [deoxy (Hb + Mb)] nor the CVs of TD, and τ_p of [deoxy (Hb + Mb)] across the four sites (i.e., spatial variation) were related to the overall faster pulmonary \dot{V}_{O_2} kinetics. Indeed, our present results in combination with those of Saitoh et al. (2009) demonstrate that reduced heterogeneity of muscle deoxygenation is not requisite for priming exercise to speed overall \dot{V}_{O_2} kinetics. Further studies might usefully focus upon the extent to which increased or decreased heterogeneity is beneficial/detrimental under physiological and/or pathophysiological conditions.

Conflict of Interest

None declared.

References

- Bangsbo, J., P. Krstrup, J. González-Alonso, R. Boushel, and B. Saltin. 2000. Muscle oxygen kinetics at onset of intense dynamic exercise in humans. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 279:R899–R906.
- Bearden, S. E., and R. J. Moffatt. 2001. \dot{V}_{O_2} and heart rate kinetics in cycling transitions from an elevated baseline. *J. Appl. Physiol.* 90:2081–2087.
- Beaver, W. L., N. Lamarra, and K. Wasserman. 1981. Breath-by-breath measurement of true alveolar gas exchange. *J. Appl. Physiol.* 51:1662–1675.
- Beaver, W. L., K. Wasserman, and B. J. Whipp. 1986. A new method for detecting anaerobic threshold by gas exchange. *J. Appl. Physiol.* 60:2020–2027.
- Behnke, B. J., C. A. Kindig, T. I. Musch, W. L. Sexton, and D. C. Poole. 2002. Effects of prior contractions on muscle

- microvascular oxygen pressure at onset of subsequent contractions. *J. Physiol.* 539:927–934.
- Bhambhani, Y. N., S. M. Buckley, and T. Susaki. 1997. Detection of ventilatory threshold using near infrared spectroscopy in men and women. *Med. Sci. Sports Exerc.* 29:402–409.
- Bowen, T. S., H. B. Rossiter, A. P. Benson, T. Amano, N. Kondo, J. M. Kowalchuk, et al. 2013. Slowed oxygen uptake kinetics in hypoxia correlate with the transient peak and reduced spatial distribution of absolute skeletal muscle deoxygenation. *Exp. Physiol.* 98:1585–1596.
- Burnley, M., A. M. Jones, H. Carter, and J. H. Doust. 2000. Effects of prior heavy exercise on phase II pulmonary oxygen uptake kinetics during heavy exercise. *J. Appl. Physiol.* 89:1387–1396.
- Burnley, M., J. H. Doust, D. Ball, and A. M. Jones. 2002. Effects of prior heavy exercise on \dot{V}_{O_2} kinetics during heavy exercise are related to changes in muscle activity. *J. Appl. Physiol.* 93:167–174.
- Chin, L. M., J. M. Kowalchuk, T. J. Barstow, N. Kondo, T. Amano, T. Shiojiri, et al. 2011. The relationship between muscle deoxygenation and activation in different muscles of the quadriceps during cycle ramp exercise. *J. Appl. Physiol.* 111:1259–1265.
- Davis, M. L., and T. J. Barstow. 2013. Estimated contribution of hemoglobin and myoglobin to near infrared spectroscopy. *Respir. Physiol. Neurobiol.* 186:180–187.
- DeLorey, D. S., J. M. Kowalchuk, and D. H. Paterson. 2004. Effects of prior heavy intensity exercise on pulmonary O_2 uptake and muscle deoxygenation kinetics in young and older adult humans. *J. Appl. Physiol.* 97:998–1005.
- DeLorey, D. S., J. M. Kowalchuk, A. P. Heenan, G. R. duManoir, and D. H. Paterson. 2007. Prior exercise speeds pulmonary O_2 uptake kinetics by increases in both local muscle O_2 availability and O_2 utilization. *J. Appl. Physiol.* 103:771–778.
- Federspiel, W. J., and A. S. Popel. 1986. A theoretical analysis of the effect of the particulate nature of blood on oxygen release in capillaries. *Microvasc. Res.* 32:164–189.
- Ferreira, L. F., D. K. Townsend, B. J. Lutjemeier, and T. J. Barstow. 2005. Muscle capillary blood flow kinetics estimated from pulmonary O_2 uptake and near-infrared spectroscopy. *J. Appl. Physiol.* 98:1820–1828.
- Ferreira, L. F., D. J. Padilla, T. I. Musch, and D. C. Poole. 2006. Temporal profile of rat skeletal muscle capillary haemodynamics during recovery from contractions. *J. Physiol.* 573:787–797.
- Ferreira, L. F., S. Koga, and T. J. Barstow. 2007. Dynamics of noninvasively estimated microvascular O_2 extraction during ramp exercise. *J. Appl. Physiol.* 103:1999–2004.
- Fukuba, Y., N. Hayashi, S. Koga, and T. Yoshida. 2002. \dot{V}_{O_2} kinetics in heavy exercise is not altered by prior exercise with a different muscle group. *J. Appl. Physiol.* 92:2467–2474.
- Gerbino, A., S. A. Ward, and B. J. Whipp. 1996. Effects of prior exercise on pulmonary gas-exchange kinetics during high-intensity exercise in humans. *J. Appl. Physiol.* 80:99–107.
- Grassi, B., S. Pogliaghi, S. Rampichini, V. Quaresima, M. Ferrari, C. Marconi, et al. 2003. Muscle oxygenation and pulmonary gas exchange kinetics during cycling exercise on-transitions in humans. *J. Appl. Physiol.* 95:149–158.
- Grassi, B., H. B. Rossiter, and J. A. Zoladz. 2015. Skeletal muscle fatigue and decreased efficiency: two sides of the same coin? *Exerc. Sport Sci. Rev.* 43:75–83.
- Groebe, K., and G. Thews. 1990. Calculated intra- and extracellular PO_2 gradients in heavily working red muscle. *Am. J. Physiol. Heart Circ. Physiol.* 259:H84–H92.
- Hamaoka, T., T. Katsumura, N. Murase, S. Nishio, T. Osada, T. Sako, et al. 2000. Quantification of ischemic muscle deoxygenation by near infrared time-resolved spectroscopy. *J. Biomed. Opt.* 5:102–105.
- Heinonen, I., S. V. Nesterov, J. Kempainen, P. Nuutila, J. Knuuti, R. Laitio, et al. 2007. Role of adenosine in regulating the heterogeneity of skeletal muscle blood flow during exercise in humans. *J. Appl. Physiol.* 103:2042–2048.
- Hernández, A., J. R. McDonald, N. Lai, and L. B. Gladden. 2010. A prior bout of contractions speeds \dot{V}_{O_2} and blood flow on-kinetics and reduces the \dot{V}_{O_2} slow-component amplitude in canine skeletal muscle contracting in situ. *J. Appl. Physiol.* 108:1169–1176.
- Hogan, M. C., P. G. Arthur, D. E. Bebout, P. W. Hochachka, and P. D. Wagner. 1992. Role of O_2 in regulating tissue respiration in dog muscle working in situ. *J. Appl. Physiol.* 73:728–736.
- Johnson, M. A., J. Polgar, D. Weightman, and D. Appleton. 1973. Data on the distribution of fibre types in thirty-six human muscles. An autopsy study. *J. Neurol. Sci.* 18:111–129.
- Jones, A. M., K. Koppo, and M. Burnley. 2003. Effects of prior exercise on metabolic and gas exchange responses to exercise. *Sports Med.* 33:949–971.
- Jones, A. M., N. J. A. Berger, D. P. Wilkerson, and C. L. Roberts. 2006. Effects of “priming” exercise on pulmonary O_2 uptake and muscle deoxygenation kinetics during heavy-intensity cycle exercise in the supine and upright positions. *J. Appl. Physiol.* 101:1432–1441.
- Jones, A. M., J. Fulford, and D. P. Wilkerson. 2008. Influence of prior exercise on muscle [phosphorylcreatine] and deoxygenation kinetics during high-intensity exercise in men. *Exp. Physiol.* 93:468–478.
- Jones, A. M., B. Grassi, P. M. Christensen, P. Krstrup, J. Bangsbo, and D. C. Poole. 2011. Slow component of \dot{V}_{O_2} kinetics: mechanistic bases and practical applications. *Med. Sci. Sports Exerc.* 43:2046–2062.
- Kalliokoski, K. K., C. Scheede-Bergdahl, M. Kjaer, and R. Boushel. 2006. Muscle perfusion and metabolic heterogeneity: insights from noninvasive imaging techniques. *Exerc. Sport Sci. Rev.* 34:164–170.

- Koga, S., D. C. Poole, L. F. Ferreira, B. J. Whipp, N. Kondo, T. Saitoh, et al. 2007. Spatial heterogeneity of quadriceps muscle deoxygenation kinetics during cycle exercise. *J. Appl. Physiol.* 103:2049–2056.
- Koga, S., D. C. Poole, Y. Fukuoka, L. F. Ferreira, N. Kondo, E. Ohmae, et al. 2011. Methodological validation of the dynamic heterogeneity of muscle deoxygenation within the quadriceps during cycle exercise. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 301:R534–R541.
- Koga, S., Y. Kano, T. J. Barstow, L. F. Ferreira, E. Ohmae, M. Sudo, et al. 2012. Kinetics of muscle deoxygenation and microvascular PO₂ during contractions in rat: comparison of optical spectroscopy and phosphorescence-quenching techniques. *J. Appl. Physiol.* 112:26–32.
- Koga, S., R. C. I. Wüst, B. Walsh, C. A. Kindig, H. B. Rossiter, and M. C. Hogan. 2013. Increasing temperature speeds intracellular PO₂ kinetics during contractions in single *Xenopus* skeletal muscle fibers. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 304:R59–R66.
- Koga, S., H. B. Rossiter, I. Heinonen, T. I. Musch, and D. C. Poole. 2014. Dynamic heterogeneity of exercising muscle blood flow and O₂ utilization. *Med. Sci. Sports Exerc.* 46:860–876.
- Koppo, K., and J. Bouckaert. 2000. In humans the oxygen uptake slow component is reduced by prior exercise of high as well as low intensity. *Eur. J. Appl. Physiol.* 83:559–565.
- Ma, S., H. B. Rossiter, T. J. Barstow, R. Casaburi, and J. Porszasz. 2010. Clarifying the equation for modeling of \dot{V}_{O_2} kinetics above the lactate threshold. *J. Appl. Physiol.* 109:1283–1284.
- MacDonald, M., P. K. Pedersen, and R. L. Hughson. 1997. Acceleration of \dot{V}_{O_2} kinetics in heavy submaximal exercise by hyperoxia and prior high-intensity exercise. *J. Appl. Physiol.* 83:1318–1325.
- McDonough, P., B. J. Behnke, D. J. Padilla, T. I. Musch, and D. C. Poole. 2005. Control of microvascular oxygen pressures in rat muscles comprised of different fibre types. *J. Physiol.* 563:903–913.
- Murgatroyd, S. R., and L. A. Wylde. 2011. The power-duration relationship of high-intensity exercise: from mathematical parameters to physiological mechanisms. *J. Physiol.* 589:2443–2445.
- Murias, J. M., M. D. Spencer, D. S. Delorey, B. J. Gurd, J. M. Kowalchuk, and D. H. Paterson. 2011. Speeding of \dot{V}_{O_2} kinetics during moderate-intensity exercise subsequent to heavy-intensity exercise is associated with improved local O₂ distribution. *J. Appl. Physiol.* 111:1410–1415.
- Niwayama, M., L. Lin, J. Shao, N. Kudo, and K. Yamamoto. 2000. Quantitative measurement of muscle hemoglobin oxygenation using near-infrared spectroscopy with correction for the influence of a subcutaneous fat layer. *Rev. Sci. Instrum.* 71:4571–4575.
- Oda, M., Y. Yamashita, T. Nakano, A. Suzuki, K. Shimizu, I. Hirano, et al. 1999. Near-infrared time resolved spectroscopy system for tissue oxygenation monitor. *Proc SPIE* 3579:611–617.
- Ohmae, E., Y. Ouchi, M. Oda, T. Suzuki, S. Nobesawa, T. Kanno, et al. 2006. Cerebral hemodynamics evaluation by near-infrared time-resolved spectroscopy: correlation with simultaneous positron emission tomography measurements. *NeuroImage* 29:697–705.
- Paterson, N. D., J. M. Kowalchuk, and D. H. Paterson. 2005. Effects of prior heavy-intensity exercise during single-leg knee extension on \dot{V}_{O_2} kinetics and limb blood flow. *J. Appl. Physiol.* 99:1462–1470.
- Perrey, S., J. Scott, L. Mourot, and J. D. Rouillon. 2003. Cardiovascular and oxygen uptake kinetics during sequential heavy cycling exercises. *Can. J. Appl. Physiol.* 28:283–298.
- Piiper, J. 2000. Perfusion, diffusion and their heterogeneities limiting blood tissue O₂ transfer in muscle. *Acta Physiol. Scand.* 168:603–607.
- Poole, D. C., and A. M. Jones. 2005. Towards an understanding of the mechanistic bases of \dot{V}_{O_2} kinetics. Pp. 294–328 in A. M. Jones and D. C. Poole, eds. *Oxygen Uptake Kinetics in Sport, Exercise and Medicine*. Routledge, London.
- Poole, D. C., and A. M. Jones. 2012. Oxygen uptake kinetics. *Compr. Physiol.* 2:933–996.
- Poole, D. C., W. Schaffartzik, D. R. Knight, T. Derion, B. Kennedy, H. J. Guy, et al. 1991. Contribution of exercising legs to the slow component of oxygen uptake kinetics in humans. *J. Appl. Physiol.* 71:1245–1260.
- Richardson, R. S., E. A. Noyszewski, J. S. Leigh, and P. D. Wagner. 1998. Lactate efflux from exercising human skeletal muscle: role of intracellular PO₂. *J. Appl. Physiol.* 85:627–634.
- Rossiter, H. B. 2011. Exercise: Kinetic considerations for gas exchange. *Compr. Physiol.* 1:203–244.
- Rossiter, H. B., S. A. Ward, J. M. Kowalchuk, F. A. Howe, J. R. Griffiths, and B. J. Whipp. 2001. Effects of prior exercise on oxygen uptake and phosphocreatine kinetics during high-intensity knee-extension exercise in humans. *J. Physiol.* 537:291–303.
- Sahlin, K., J. B. Sørensen, L. B. Gladden, H. B. Rossiter, and P. K. Pedersen. 2005. Prior heavy exercise eliminates \dot{V}_{O_2} slow component and reduces efficiency during submaximal exercise in humans. *J. Physiol.* 564:765–773.
- Saitoh, T., L. F. Ferreira, T. J. Barstow, D. C. Poole, A. Ooue, N. Kondo, et al. 2009. Effects of prior heavy exercise on heterogeneity of muscle deoxygenation kinetics during subsequent heavy exercise. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 297:R615–R621.
- Scheuermann, B. W., B. D. Hoelting, M. L. Noble, and T. J. Barstow. 2001. The slow component of O₂ uptake is not accompanied by changes in muscle EMG during repeated bouts of heavy exercise in humans. *J. Physiol.* 531:245–256.
- Spencer, M. D., J. M. Murias, T. M. Grey, and D. H. Paterson. 2012. Regulation of \dot{V}_{O_2} kinetics by O₂ delivery: insights from acute hypoxia and heavy-intensity priming exercise in young men. *J. Appl. Physiol.* 112:1023–1032.

- Tordi, N., S. Perrey, A. Harvey, and R. L. Hughson. 2003. Oxygen uptake kinetics during two bouts of heavy cycling separated by fatiguing sprint exercise in humans. *J. Appl. Physiol.* 94:533–541.
- Whipp, B. J., N. Lamarra, and S. A. Ward. 1995. Obligatory anaerobiosis resulting from oxygen uptake-to-blood flow ratio dispersion in skeletal muscle: a model. *Eur. J. Appl. Physiol.* 71:147–152.
- Wilkerson, D. P., K. Koppo, T. J. Barstow, and A. M. Jones. 2004. Effect of prior multiple-sprint exercise on pulmonary O₂ uptake kinetics following the onset of perimaximal exercise. *J. Appl. Physiol.* 97:1227–1236.