



Corrigendum: Mequindox Induced Genotoxicity and Carcinogenicity in Mice

Qianying Liu^{1,2}, Zhixin Lei¹, Qin Wu¹, Deyu Huang¹, Shuyu Xie^{1,3}, Xu Wang^{1,3}, Yuanhu Pan^{1,3*} and Zonghui Yuan^{1,2,3,4*}

OPEN ACCESS

Edited and reviewed by:

Eleonore Fröhlich, Medical University of Graz, Austria

*Correspondence:

Yuanhu Pan panyuanhu@hotmail.com Zonghui Yuan yuan5802@mail.hzau.edu.cn

Specialty section:

This article was submitted to Predictive Toxicology, a section of the journal Frontiers in Pharmacology

Received: 25 October 2018 Accepted: 12 November 2018 Published: 29 November 2018

Citation:

Liu Q, Lei Z, Wu Q, Huang D, Xie S, Wang X, Pan Y and Yuan Z (2018) Corrigendum: Mequindox Induced Genotoxicity and Carcinogenicity in Mice. Front. Pharmacol. 9:1387. doi: 10.3389/fphar.2018.01387 ¹ National Reference Laboratory of Veterinary Drug Residues (HZAU) and MAO Key Laboratory for Detection of Veterinary Drug Residues, Huazhong Agricultural University, Wuhan, China, ² Key Laboratory of Preventive Veterinary Medicine in Hubei Province, Huazhong Agricultural University, Wuhan, China, ³ MOA Laboratory for Risk Assessment of Quality and Safety of Livestock and Poultry Products, Huazhong Agricultural University, Wuhan, China, ⁴ Hubei Collaborative Innovation Center for Animal Nutrition and Feed Safety, Wuhan, China

Keywords: mequindox, quinoxaline, carcinogenicity, genotoxicity, KM mice

A Corrigendum on

Mequindox Induced Genotoxicity and Carcinogenicity in Mice

by Liu, Q., Lei, Z., Wu, Q., Huang, D., Xie, S., Wang, X., et al. (2018). Front. Pharmacol. 9:361. doi: 10.3389/fphar.2018.00361

In the original article, there was a mistake in **Figures 5B,F** as published. The descriptions of **Figures 5B,F** did not match the figures displayed. The corrected **Figure 5** appears below. The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2018 Liu, Lei, Wu, Huang, Xie, Wang, Pan and Yuan. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

1




FIGURE 5 | Selected microphotographs of kidney, liver and testis following dietary exposure to MEQ in the carcinogenicity tests (200×, 400×). (A) Kidney from control group (200×); (B) Kidney from the M25 mg/kg group showing kidney interstitial small blood vessels congestion, glomerular congestion (400×); (C) Kidney from the M55 mg/kg group showing aggregation of lymphocyte into a group around the central veins (200×); (D) Kidney from the M110 mg/kg group showing degeneration and necrosis of renal tubular epithelial cells (400×); (E) Liver from control group (200×); (F) Liver from the M25 mg/kg group showing degeneration and necrosis of hepatic cells (400×); (G) Liver from the M55 showing neutrophilic infiltrate within and around bile duct (200×); (H) Liver from the M110 mg/kg group showing a broadened testicular interstitium (400×); (K) Testis at the M55 mg/kg group showing an irregular arrangement as well as a decreased number of spermatogenic cells (200×); (L) Testis at the M110 mg/kg group showing neutrophilic in the lumen (200×).