



ARTICLE

Isolation and Characterization of Halophilic *Kocuria salsicia* Strains from Cheese Brine

Hye-Young Youn and Kun-Ho Seo*

Center for One Health, College of Veterinary Medicine, Department of Veterinary Public Health, Konkuk University, Seoul 05029, Korea

OPEN ACCESS

Received November 19, 2021

Revised January 3, 2022

Accepted January 5, 2022

*Corresponding author : Kun-Ho Seo
Center for One Health, College of Veterinary
Medicine, Department of Veterinary Public
Health, Konkuk University, Seoul 05029,
Korea
Tel: +82-2-450-4121
Fax: +82-2-3436-4128
E-mail: bracstu3@konkuk.ac.kr

*ORCID
Hye-Young Youn
<https://orcid.org/0000-0003-4626-5859>
Kun-Ho Seo
<https://orcid.org/0000-0001-5720-0538>

Abstract *Kocuria salsicia* can survive in extreme environments and cause infections, including catheter-related bacteremia, in humans. Here, we investigated and evaluated the characteristics of nine *K. salsicia* strains (KS1–KS9) isolated from cheese brine from a farmstead cheese-manufacturing plant in Korea from June to December, 2020. *Staphylococcus aureus* American Type Culture Collection (ATCC) 29213 was used as a positive control in the growth curve analysis and biofilm-formation assays. All *K. salsicia* isolates showed growth at 15% salt concentration and temperatures of 15°C, 25°C, 30°C, 37°C, and 42°C. KS6 and KS8 showed growth at 5°C, suggesting that they are potential psychrotrophs. In the biofilm-formation analysis via crystal violet staining, KS6 exhibited the highest biofilm-forming ability at various temperatures and media [phosphate buffered saline, nutrient broth (NB), and NB containing 15% sodium chloride]. At 25°C and 30°C, KS3, KS6, and KS8 showed higher biofilm-forming ability than *S. aureus* ATCC 29213. The antimicrobial resistance of the isolates was evaluated using the VITEK® 2 system; most isolates were resistant to marbofloxacin and nitrofurantoin (both 9/9, 100%), followed by enrofloxacin (7/9, 77.8%). Five of the nine isolates (5/9, 55.6%) showed multidrug resistance. Our study reports the abilities of *K. salsicia* to grow in the presence of high salt concentrations and at relatively low temperatures, along with its multidrug resistance and tendency to form biofilms.

Keywords *Kocuria salsicia*, cheese brine, growth curve, biofilm, antimicrobial resistance

Introduction

Kocuria salsicia is a coccoid, gram-positive, and facultative anaerobic bacterium (Savini et al., 2010). It has been isolated from various animal hosts, soil, dairy products, the skin or oropharynx mucosa of humans, and high-salt and high-temperature environments (Basaglia et al., 2002). *Kocuria* spp., such as *K. kristinae*, *K. varians*, *K. rhizophila*, *K. rosea*, and *K. marina*, have been demonstrated to cause infections in humans (Lee et al., 2009). Sepsis and increased platelet and leukocyte counts are signs of *Kocuria* spp. infection (Dunn et al., 2011). Although few studies have evaluated the mechanisms of the infections and toxicity of *Kocuria* spp., biofilm has been suggested to be involved for adhesion and colonization (Meletis et al., 2012).

Infections associated with *Kocuria* spp. include urinary tract infections, cholecystitis, and catheter-related bacteremia (Kandi et al., 2016). Among *Kocuria* spp., *K. salsicia* is the causative agent of the first case of catheter-related bacteremia in Korea (Sohn et al., 2015).

The bacterial growth curve has been demonstrated to be useful in identifying the physiological characteristics of microorganisms; it indicates the phases of bacterial growth and tolerance in certain environments, such as various temperature and salt conditions (Zwietering et al., 1990). Biofilms are formed when microorganisms grow at a particular spot and reach a specific density, which can be predicted via a growth curve. Therefore, it is important to evaluate the growth curve for determining the biofilm-formation ability of the bacteria (Welch et al., 2012). Microbial attachment and biofilm formation on food contact surfaces in processing plants are major concerns in terms of survival of these microorganisms under extreme conditions, such as high osmotic pressure, and the risk of cross-contamination (Ryu and Beuchat, 2005). Recently, *Kocuria* spp. have gained prominence owing to a rise in the number of reports of human infections, signifying their pathogenic potential (Kandi et al., 2016). The infections caused by *Kocuria* spp. include urinary tract infections, catheter-associated bacteremia, and endocarditis, which might be associated with their biofilm-forming ability (Moreira et al., 2015; Sohn et al., 2015).

Currently, in farmstead manufacturing, which involves direct handling of livestock, antimicrobials are used to reduce biofilm-forming bacterial contamination; moreover, antibiotic resistance has increased over the past few decades (Mehli et al., 2017). The use of antimicrobials in clinical settings and food production is inefficient; nevertheless, the overuse of antimicrobials has resulted in the development of antimicrobial resistance in bacteria present in livestock products (European Food Safety Authority, 2019). Particularly, microbial contamination during cheese-making may act as an intermediate for transferring antimicrobial resistance genes to various bacteria, including non-pathogenic bacteria, and may lead to multidrug resistance development (Locatelli et al., 2016). Furthermore, these multidrug-resistant (MDR) bacteria act as a reservoir of antimicrobial resistance genes, facilitating their transmission to humans via food (Golob et al., 2019).

Many studies have focused on foodborne pathogens present in raw milk, cheese products, and environments in which farmstead cheese is produced (D'Amico and Donnelly, 2008; Fox et al., 2011; Kang et al., 2018; Mehli et al., 2017). However, no studies have focused on the microbial contamination of cheese brine, possibly because of its high salt concentration. To the best of our knowledge, this study is the first to report the isolation of *K. salsicia* from cheese brine from a farmstead cheese-manufacturing plant in Korea. The aims of the present study were to evaluate the (i) growth curve of *K. salsicia* strains isolated from cheese brine at different temperatures (5°C, 15°C, 25°C, 30°C, 37°C, and 42°C) and 15% salt concentration, (ii) biofilm-forming ability of *K. salsicia* strains at the different temperatures and media [phosphate buffered saline (PBS), nutrient broth (NB), and NB containing 15% sodium chloride (NaCl)], and (iii) antimicrobial susceptibility of *K. salsicia* strains as well as assess the potential risk posed by farmstead cheese (contaminated by organisms in the brine) with respect to the transfer of antimicrobial-resistant pathogens to humans.

Materials and Methods

Sample collection and *K. salsicia* isolation from cheese brine

From June to December, every month in 2020, a sterile bottle was used to collect two 500 mL bottles of string cheese brine per month (totaling 14 bottles) during cheese-making at a farmstead cheese house located in the Gyeong-gi province, Korea.

Salt concentration and pH of the brine were analyzed using a glass salimeter (Daedong, Seoul, Korea) and Orion Star™ A211 pH Benchtop Meter (Thermo Fisher Scientific, Waltham, MA, USA), respectively. The sample was transported to a laboratory refrigerator and analyzed within 4 h.

As there was no established selective medium for *K. salsicia* isolation, a loop of each bottle of cheese brine solution was streaked onto nutrient agar (Sigma-Aldrich, St. Louis, MO, USA) and tryptone soya agar (Oxoid, Basingstoke, UK), and incubated for 24 h at 37°C, in triplicate. Additionally, the colonies were cultured on Columbia agar containing 5% sheep blood (bioMérieux, Marcy l'Etoile, France) after a 48 h incubation to detect non-hemolytic and lemon-yellow colonies—cultural characteristics of *K. salsicia* (Kandi et al., 2016; Sohn et al., 2015). Typical colonies (non-hemolytic and lemon-yellow colonies) were sub-cultured on nutrient agar. Because of the lack of proper guidelines for *Kocuria* spp., we used the positive control (PC) for *Staphylococcus* spp. mentioned by Sohn et al. (2015). *Staphylococcus aureus* American Type Culture Collection (ATCC) 29213 was purchased from the ATCC (Manassas, VA, USA) and used as a PC strain (Bruins et al., 2007). *S. aureus* ATCC 29213 is a clinical isolate that has been applied for enteric and infectious disease research (Soni et al., 2015). Furthermore, *Escherichia coli* ATCC 8739, enterohemorrhagic *E. coli* (EHEC) ATCC 43894, and *Listeria monocytogenes* ATCC 51776 were purchased from the ATCC.

DNA extraction

DNA extraction was performed using the NucliSENS easyMAG system (bioMérieux). Briefly, each bacterial colony was added to lysis buffer (1.0 mL) and left to stand for 20 min at 25°C. The mixture was centrifuged at 15,770×g for 3 min using MIKRO 200 centrifuge (Hettich, Tuttlingen, Germany), and 1 mL of the supernatant was transferred to a well of a plastic vessel with 50 µL of magnetic silica and subjected to automated magnetic bead separation. DNA was then resuspended in 75 µL of elution buffer.

Identification and sequencing of *K. salsicia*

K. salsicia strains were isolated from cheese brine and identified via 16S rRNA sequencing. The 27F and 1492R primers were used for polymerase chain reaction (PCR). PCR products were sequenced using the same primers and ABI BigDye Terminator v3.1 Cycle Sequencing Kit (Applied Biosystems, Carlsbad, CA, USA). Sequencing was performed using an Applied Biosystems 3730XL DNA Analyzer obtained from Bionics (Seoul, Korea). Each 16S rRNA sequence was analyzed via the basic local alignment search tool (BLAST; <https://blast.ncbi.nlm.nih.gov/Blast.cgi>) using the National Center for Biotechnology Information 16S rRNA database to identify *K. salsicia*.

Growth curve analysis of *K. salsicia* and *S. aureus* at different temperatures

To analyze the growth curve at different temperatures (5°C, 15°C, 25°C, 30°C, 37°C, and 42°C), the cheese brine isolates and *S. aureus* ATCC 29213 were cultured in nutrient agar (Sigma-Aldrich) for 24 h at 37°C. NB (200 µL; Sigma-Aldrich) was filter-sterilized using a 0.2-µm syringe filter (Millipore, Bedford, MA, USA) and then mixed with each isolate at 0.5 McFarland turbidity. After mixing, 200 µL of each isolate was transferred to a 96-well plate (SPL Life Sciences, Pocheon, Korea). The growth curves were generated at 24 h intervals over 288 h at 5°C; 8 h intervals over 96 h at 15°C, 25°C, and 30°C; and 2 h intervals over 24 h at 37°C and 42°C by measuring the optical density (OD) at 595 nm using a Multiskan FC microplate reader (Thermo Fisher Scientific). The experiment was repeated in triplicate for each isolate.

Growth curve analysis of *K. salsicia* at 15% salt concentration and different temperatures compared with that of other pathogens

To analyze the growth of *K. salsicia* in a halophilic environment, three isolates (KS3, KS6, and KS8) among nine *K. salsicia* isolates were selected based on their relatively high biofilm production. KS3, KS6, and KS8 were compared with *S. aureus* ATCC 29213, *E. coli* ATCC 8739, EHEC ATCC 43894, and *L. monocytogenes* ATCC 51776 at a salt concentration of approximately 15% (w/v) based on the salt concentration of cheese brine in farmstead cheese manufacturing and different temperatures (Bintsis and Papademas, 2002; Bruins et al., 2007; Haastrup et al., 2018; Ingham et al., 2000; Larson et al., 1999). These four foodborne pathogens were used as quality control strains for analyzing the growth curve at 15% salt concentration. NB (200 μ L; Sigma-Aldrich) containing 15% NaCl (Sigma-Aldrich) was filter-sterilized using a 0.2- μ m syringe filter (Millipore) and then mixed with each strain at 0.5 McFarland turbidity. After mixing, 200 μ L of each strain was transferred to a 96-well plate (SPL Life Sciences). The growth curves were generated at 24 h intervals over 288 h at 5°C; 8 h intervals over 96 h at 15°C, 25°C, and 30°C; and 2 h intervals over 24 h at 37°C and 42°C by measuring the OD at 595 nm using a Multiskan FC microplate reader (Thermo Fisher Scientific). The experiment was repeated in triplicate for each strain.

Biofilm formation of *K. salsicia* at different temperatures

The biofilm-forming ability of the *K. salsicia* isolates was evaluated as previously described (Jeong et al., 2018). In brief, each colony of the isolates was added to 200 μ L of PBS (Sigma-Aldrich), NB (Sigma-Aldrich), and NB containing 15% NaCl and set to 0.5–0.6 McFarland turbidity. To assess the extent of biofilm formation in each microplate, 200 μ L of each sample was transferred to a 96-well polystyrene culture plate and incubated at 5°C, 15°C, 25°C, 30°C, 37°C, and 42°C for 24 h. The culture medium was discarded, and the microplate was washed twice with 200 μ L of PBS (Sigma-Aldrich). Cells from adherent biofilms were stained with 0.1% (w/v) crystal violet (100 μ L; Sigma-Aldrich) for 15 min at room temperature (20°C–25°C) and rinsed twice with PBS (Sigma-Aldrich). After removing the dye from stained cells using 99% ethanol (200 μ L), the amount of biofilm was quantified by measuring the absorbance of the solution at 595 nm using a Multiskan FC microplate reader (Thermo Fisher Scientific). The experiment was performed in triplicate.

Antimicrobial susceptibility testing of *K. salsicia*

Antimicrobial susceptibility tests were performed using the VITEK[®] 2 instrument (bioMérieux) with gram-positive susceptibility (AST-GP) cards (bioMérieux). The AST-GP cards contained amikacin (AMK), chloramphenicol (CHL), clindamycin (CLI), gentamicin (GEN), cefpodoxime (POD), enrofloxacin (ENO), erythromycin (ERY), marbofloxacin (MAR), minocycline (MIN), nitrofurantoin (NIT), pradofloxacin (PRA), and trimethoprim/sulfamethoxazole (SXT), and the AMK, CHL, CLI, GEN, ENO, ERY, MAR, MIN, NIT, PRA, and SXT results were interpreted in accordance with the Performance Standards for Antimicrobial Disk and Dilution Susceptibility Tests for Bacteria Isolated from Animals in the Clinical and Laboratory Standards Institute guidelines (CLSI VET01S; CLSI, 2020a). However, owing to the lack of POD minimum inhibitory concentration criteria in CLSI VET01S, Performance Standards for Antimicrobial Susceptibility Testing in the CLSI was used to interpret the result (CLSI, 2020b). As breakpoints for *K. salsicia* have not been established, the antimicrobial resistance test was performed with reference to criteria used for *S. aureus* ATCC 29213 (Sohn et al., 2015).

Statistical analysis

Data for biofilm formation and growth curve analysis are presented as the mean \pm SD. GraphPad Prism 7.00 (GraphPad

Software, San Diego, CA, USA) was used for data analyses. Biofilm formation data were analyzed using ANOVA followed by the Tukey method. $p < 0.05$ was considered significant.

Results and Discussion

Isolation and identification of *K. salsicia* strains in cheese brine

We identified nine *K. salsicia* isolates (KS1, KS2, KS3, KS4, KS5, KS6, KS7, KS8, and KS9) from a farmstead cheese house in Korea. The salt concentration and pH of the brine were 15%–18% (w/v) and 5.3, respectively. All *K. salsicia* strains were from different agar plates. Hemolytic *K. salsicia* colonies were not observed on Columbia agar containing 5% sheep blood but hemolytic *S. aureus* ATCC 29213 colonies were recorded. Consistently, 16S rRNA sequencing confirmed taxa of the cheese brine isolates as *K. salsicia* at the species level and their sequences were submitted to GenBank under accession numbers MW301599 for *K. salsicia* 1 (KS1), MW301601 for *K. salsicia* 2 (KS2), MW301600 for *K. salsicia* 3 (KS3), MW301603 for *K. salsicia* 4 (KS4), MW301604 for *K. salsicia* 5 (KS5), MW301605 for *K. salsicia* 6 (KS6), MW301606 for *K. salsicia* 7 (KS7), MW301607 for *K. salsicia* 8 (KS8), and MW301608 for *K. salsicia* 9 (KS9).

Growth curve analysis

The growth curves of KS1–KS9 were compared with that of *S. aureus* ATCC 29213 by measuring the OD of the cultures at 595 nm (Fig. 1). Growth of all *K. salsicia* isolates was observed at all temperatures (5°C, 15°C, 25°C, 30°C, 37°C, and 42°C), suggesting that it is a psychotropic bacterium. The growth of *K. salsicia* strains was the highest during the exponential phase at 5°C and 15°C, which lasted for at least 96 and 288 h, respectively (Figs. 1A, B). OD values of *K. salsicia* isolates (0.12–0.71) were higher than *S. aureus* ATCC 29213 (0.12–0.29) at 5°C (Fig. 1A). At 25°C and 30°C, *K. salsicia* isolates were in the stationary phase for approximately 60 h. All *K. salsicia* isolates showed OD values higher than those of *S. aureus* ATCC 29213 (PC; Figs. 1C, D).

Growth curves of KS3, KS6, and KS8 were compared with those of *S. aureus* ATCC 29213, *E. coli* ATCC 8739, EHEC ATCC 43894, and *L. monocytogenes* ATCC 51776 by measuring the OD of the cultures at 595 nm (Fig. 2). All microorganisms grew at 15°C, 25°C, 30°C, 37°C, and 42°C in NB containing 15% NaCl. *S. aureus* ATCC 29213 showed higher growth than the other strains tested. Particularly, *S. aureus* ATCC 29213 showed the highest OD value of approximately 0.40 at 15°C (Fig. 2B). The growth of KS3, KS6, and KS8 increased gradually at 15°C, 25°C, 30°C, 37°C, and 42°C; KS6 and KS8 showed slow growth at 5°C (Fig. 2A). *E. coli* ATCC 8739, EHEC ATCC 43894, and *L. monocytogenes* ATCC 51776 showed lower OD values (approximately 0.1 to 0.2) than *S. aureus* ATCC 29213 and *K. salsicia* isolates at all temperatures. In general, salt processing promotes the syneresis of whey from the curd, thereby reducing the moisture content of the cheese (McMahon et al., 2009). Although cheese salting is believed to decrease the population of undesirable contaminants, brine can also serve as a reservoir for certain salt-tolerant pathogens (Bintsis and Papademas, 2002). The psychrotroph *L. monocytogenes* survives for longer periods in brines stored at 4°C than in those stored at 12°C (Larson et al., 1999). *E. coli* O157:H7 and *Salmonella* Typhimurium can survive for several weeks in brine (Ingham et al., 2000). *Kocuria* spp. grow at a temperature range of 4°C–43°C and tolerate up to 15% NaCl concentration (Kim et al., 2004). Consistent with these studies, our results indicated that *K. salsicia* strains grew at a temperature range of 5°C–42°C and tolerated 15% NaCl concentration (Fig. 2). Although cheese brine does not provide a favorable condition for microorganisms to grow owing to its high salt concentration, it can cause cross-contamination because of the whey derived from cheese and improper temperature control (Mehli et al., 2017).




Fig. 1. Growth curves of *Staphylococcus aureus* ATCC 29213 and *Kocuria salsicia* isolates (KS1–KS9) cultured in nutrient broth (NB). Growth curves of cultures grown at 5°C, 15°C, 25°C, 30°C, 37°C, and 42°C were generated by measuring the optical density at 595 nm using a microplate reader. Figure legend denotes the order of the final values of negative control (NC), positive control (PC), and *K. salsicia* isolates (KS1–KS9). Values in parentheses are the average optical densities of NC, PC, and *K. salsicia* isolates (KS1–KS9). NC, NB; PC, *S. aureus* ATCC 29213; KS1–KS9, *K. salsicia* KS1–KS9; ATCC, American Type Culture Collection.

Biofilm-formation activity

The biofilm-forming ability of *K. salsicia* isolates (KS1–KS9) was evaluated in diverse temperatures and media. Although *K. salsicia* isolates did not grow well in PBS, the OD₅₉₅ value of KS9 (0.065) was significantly higher than that of *S. aureus* ATCC 29213 (0.052) at 37°C ($p < 0.05$; Table 1). Moreover, the biofilm-forming ability of KS6 was significantly different from that of *S. aureus* ATCC 29213 at the temperatures tested in NB ($p < 0.05$; Table 2). The highest OD₅₉₅ value for biofilms was observed for KS8 (0.371) followed by KS3 (0.361) at 30°C ($p < 0.05$). Notably, KS3, KS6, and KS8 showed better biofilm production abilities than *S. aureus* ATCC 29213 at four of the tested temperatures (25°C, 30°C, 37°C, and 42°C; $p < 0.05$). KS6 showed a significant difference in biofilm formation compared with *S. aureus* ATCC 29213, even at 5°C ($p < 0.05$). In NB containing 15% NaCl, all microorganisms grew at 5°C, 15°C, 25°C, 30°C, 37°C, and 42°C and formed biofilm. Among temperatures, all *K. salsicia* isolates (KS1–KS9) showed higher OD values (range 0.054 to 0.066) than *S.*




Fig. 2. Growth curves of *Kocuria salsicia* isolates (KS3, KS6, and KS8) compared with those of *Staphylococcus aureus* ATCC 29213, *Escherichia coli* ATCC 8739, Enterohemorrhagic *Escherichia coli* ATCC 43894, and *Listeria monocytogenes* ATCC 51776 cultured in nutrient broth (NB) containing 15% sodium chloride (NaCl). Growth curves of cultures grown at 5°C, 15°C, 25°C, 30°C, 37°C, and 42°C were generated using the optical density values at 595 nm measured using a microplate reader. Figure legend denotes the order of the final values of negative control (NC), positive control (PC), and *K. salsicia* isolates (KS3, KS6, and KS8). Values in parentheses are the average optical densities of NC, PC, and *K. salsicia* isolates (KS3, KS6, and KS8). NC, NB containing 15% NaCl; PC, *S. aureus* ATCC 29213+NB containing 15% NaCl; KS3, *K. salsicia* KS3+NB containing 15% NaCl; KS6, *K. salsicia* KS6+NB containing 15% NaCl; KS8, *K. salsicia* KS8+NB containing 15% NaCl; E, *E. coli* ATCC 8739+NB containing 15% NaCl; EHEC, Enterohemorrhagic *E. coli* ATCC 43894+NB containing 15% NaCl; LM, *Listeria monocytogenes* ATCC 51776+NB containing 15% NaCl; ATCC, American Type Culture Collection.

aureus ATCC 29213 at 15°C ($p < 0.05$; Table 3).

Most cases of *Kocuria* spp. infection have been hypothesized to be associated with catheter-associated bloodstream infections; however, this association was not established until recently (Barnes et al., 1999; Sohn et al., 2015). Only a few studies have reported the biofilm-forming ability of *Kocuria* spp. Therefore, studying the biofilm-forming ability of *K. salsicia* may provide valuable insights into the biofilm-forming potential of other members of this genus (Purty et al., 2013). Furthermore, temperature is a key regulator of bacterial biofilm formation, and the temperature changes occurring in food, as well as hospital conditions, influence biofilm formation by microorganisms (Di Ciccio et al., 2015; Nilsson et al., 2011).

Table 1. Biofilm-formation ability of *Kocuria salsicia* isolates (KS1–KS9) in phosphate buffered saline (PBS) compared with that of *Staphylococcus aureus* ATCC 29213 at 5°C, 15°C, 25°C, 30°C, 37°C, and 42°C

Strains	Optical density of isolates ¹⁾					
	Temperature (°C) ²⁾					
	5	15	25	30	37	42
NC	0.047±0.001	0.048±0.001	0.046±0.005	0.043±0.005	0.046±0.002	0.047±0.003
PC	0.051±0.006 ^a	0.050±0.005 ^a	0.051±0.001 ^a	0.060±0.011 ^a	0.052±0.002 ^a	0.053±0.003 ^a
KS1	0.048±0.003 ^a	0.052±0.005 ^a	0.053±0.002 ^a	0.059±0.002 ^a	0.056±0.002 ^a	0.053±0.001 ^a
KS2	0.053±0.004 ^a	0.053±0.000 ^a	0.056±0.002 ^a	0.062±0.003 ^a	0.057±0.003 ^a	0.053±0.001 ^a
KS3	0.053±0.004 ^a	0.053±0.007 ^a	0.054±0.000 ^a	0.064±0.003 ^a	0.051±0.001 ^a	0.055±0.003 ^a
KS4	0.048±0.001 ^a	0.050±0.003 ^a	0.052±0.003 ^a	0.055±0.002 ^a	0.053±0.003 ^a	0.055±0.003 ^a
KS5	0.053±0.005 ^a	0.054±0.002 ^a	0.053±0.001 ^a	0.054±0.004 ^a	0.057±0.001 ^a	0.054±0.002 ^a
KS6	0.049±0.003 ^a	0.053±0.003 ^a	0.059±0.005 ^a	0.061±0.001 ^a	0.059±0.001 ^a	0.056±0.002 ^a
KS7	0.050±0.004 ^a	0.053±0.003 ^a	0.055±0.003 ^a	0.054±0.002 ^a	0.052±0.002 ^a	0.053±0.001 ^a
KS8	0.056±0.006 ^a	0.051±0.006 ^a	0.055±0.002 ^a	0.059±0.005 ^a	0.055±0.000 ^a	0.053±0.000 ^a
KS9	0.057±0.005 ^a	0.059±0.008 ^a	0.055±0.004 ^a	0.064±0.004 ^a	0.065±0.005 ^b	0.057±0.001 ^a

¹⁾ Optical density of isolates is expressed as mean±SD.

²⁾ Biofilm formation by temperature (°C).

^{a,b} Different letters indicate statistical difference at p<0.05 compared to *S. aureus* ATCC 29213 (Tukey method).

ATCC, American Type Culture Collection; NC, negative control (PBS); PC, positive control (*S. aureus* ATCC 29213+PBS); KS1–KS9, *K. salsicia* KS1–KS9+PBS.

Table 2. Biofilm-formation ability of *Kocuria salsicia* isolates (KS1–KS9) in nutrient broth (NB) compared with that of *Staphylococcus aureus* ATCC 29213 at 5°C, 15°C, 25°C, 30°C, 37°C, and 42°C

Strains	Optical density of isolates ¹⁾					
	Temperature (°C) ²⁾					
	5	15	25	30	37	42
NC	0.069±0.007	0.064±0.003	0.068±0.005	0.075±0.007	0.062±0.004	0.048±0.001
PC	0.093±0.013 ^a	0.131±0.002 ^a	0.089±0.007 ^a	0.128±0.005 ^a	0.066±0.002 ^a	0.080±0.003 ^a
KS1	0.116±0.007 ^b	0.139±0.002 ^b	0.082±0.004 ^a	0.237±0.025 ^b	0.121±0.008 ^b	0.097±0.002 ^a
KS2	0.106±0.002 ^a	0.136±0.002 ^a	0.073±0.001 ^a	0.079±0.001 ^c	0.066±0.002 ^a	0.072±0.001 ^a
KS3	0.088±0.005 ^a	0.147±0.001 ^b	0.293±0.050 ^b	0.361±0.003 ^b	0.133±0.003 ^b	0.148±0.010 ^b
KS4	0.109±0.008 ^a	0.132±0.001 ^a	0.093±0.002 ^a	0.102±0.008 ^a	0.083±0.002 ^a	0.070±0.001 ^a
KS5	0.120±0.010 ^b	0.142±0.003 ^b	0.080±0.003 ^a	0.089±0.005 ^c	0.065±0.002 ^a	0.072±0.001 ^a
KS6	0.113±0.001 ^b	0.187±0.004 ^b	0.250±0.001 ^b	0.315±0.011 ^b	0.132±0.007 ^b	0.146±0.023 ^b
KS7	0.102±0.003 ^a	0.129±0.001 ^a	0.118±0.003 ^a	0.100±0.005 ^a	0.078±0.003 ^a	0.069±0.001 ^a
KS8	0.090±0.002 ^a	0.137±0.002 ^a	0.260±0.027 ^b	0.371±0.003 ^b	0.183±0.011 ^b	0.118±0.008 ^b
KS9	0.082±0.001 ^a	0.137±0.001 ^a	0.075±0.002 ^a	0.079±0.003 ^c	0.136±0.011 ^b	0.102±0.005 ^a

¹⁾ Optical density of isolates is expressed as mean±SD.

²⁾ Biofilm formation by temperature (°C).

^{a-c} Different letters indicate statistical difference at p<0.05 compared to *S. aureus* ATCC 29213 (Tukey method).

ATCC, American Type Culture Collection; NC, negative control (NB); PC, positive control (*S. aureus* ATCC 29213+NB); KS1–KS9, *K. salsicia* KS1–KS9+NB.

Table 3. Biofilm-formation ability of *Kocuria salsicia* isolates (KS1–KS9) in nutrient broth (NB) containing 15% sodium chloride (NaCl) compared with that of *Staphylococcus aureus* ATCC 29213 at 5°C, 15°C, 25°C, 30°C, 37°C, and 42°C

Strains	Optical density of isolates ¹⁾					
	Temperature (°C) ²⁾					
	5	15	25	30	37	42
NC	0.040±0.000	0.042±0.001	0.040±0.000	0.045±0.002	0.045±0.000	0.045±0.000
PC	0.047±0.001 ^a	0.049±0.000 ^a	0.053±0.001 ^a	0.051±0.001 ^a	0.066±0.000 ^a	0.066±0.000 ^a
KS1	0.056±0.003 ^b	0.062±0.001 ^b	0.079±0.010 ^b	0.076±0.006 ^b	0.071±0.014 ^a	0.064±0.006 ^a
KS2	0.051±0.000 ^a	0.054±0.001 ^b	0.054±0.000 ^a	0.084±0.002 ^b	0.066±0.007 ^a	0.076±0.009 ^a
KS3	0.057±0.002 ^b	0.063±0.002 ^b	0.067±0.001 ^b	0.096±0.004 ^b	0.077±0.004 ^a	0.071±0.004 ^a
KS4	0.052±0.001 ^a	0.055±0.001 ^b	0.056±0.002 ^a	0.067±0.000 ^b	0.056±0.004 ^a	0.054±0.005 ^a
KS5	0.055±0.002 ^b	0.055±0.001 ^b	0.054±0.001 ^a	0.065±0.003 ^a	0.059±0.002 ^a	0.058±0.004 ^a
KS6	0.061±0.007 ^b	0.057±0.000 ^b	0.059±0.001 ^a	0.069±0.005 ^b	0.065±0.007 ^a	0.064±0.007 ^a
KS7	0.054±0.001 ^a	0.054±0.001 ^b	0.054±0.001 ^a	0.062±0.008 ^a	0.055±0.006 ^a	0.063±0.006 ^a
KS8	0.053±0.000 ^a	0.066±0.001 ^b	0.059±0.001 ^a	0.080±0.000 ^b	0.070±0.004 ^a	0.068±0.002 ^a
KS9	0.054±0.001 ^a	0.059±0.002 ^b	0.053±0.001 ^a	0.083±0.011 ^b	0.076±0.001 ^a	0.066±0.005 ^a

¹⁾ Optical density of isolates is expressed as mean±SD.

²⁾ Biofilm formation by temperature (°C).

^{a,b} Different letters indicate statistical difference at $p < 0.05$ compared to *S. aureus* ATCC 29213 (Tukey method).

ATCC, American Type Culture Collection; NC, negative control (NB containing 15% NaCl); PC, positive control (*S. aureus* ATCC 29213+NB containing 15% NaCl); KS1–KS9, *K. salsicia* KS1–KS9+NB containing 15% NaCl.

Consistent with the findings of the present study, another study reported that the effect of the growth temperature on the formation of *S. aureus* biofilm is influenced by several environmental factors such as nutrient availability and growth medium (Banks and Bryers, 1991; de Jesus Pimentel-Filho et al., 2014). In the present study, the OD values of *K. salsicia* KS1–KS9 and *S. aureus* ATCC 29213 were different; however, the patterns of OD values in all temperatures and media were similar. As brine containing cheese whey may have abundant nutrients for the growth of microorganisms, it is important to evaluate the biofilm formation ability of *K. salsicia* isolates in different nutrient sources. KS3, KS6, and KS8 showed higher growth rates than *S. aureus* ATCC 29213; this difference in growth rates was larger at 25°C and 30°C than that at 37°C and 42°C. Further, low temperatures increase the hydrophilic properties of cells and alter the ability of the bacteria to adhere to hydrophobic materials such as polystyrene and the effect of temperature on biofilm formation also depends on the presence or absence of NaCl (Rode et al., 2007). Auto-aggregation of microorganisms increases with increased NaCl concentration; this auto-aggregation is highly correlated with the biofilm formation by foodborne pathogens (Xu et al., 2010). To the best of our knowledge, biofilm formation under various temperatures, nutrient conditions, and environmental conditions has not been extensively investigated for *Kocuria* spp. Our results suggest that *K. salsicia* can grow at a wide range of temperatures and survive in the cheese brine tank. Altuntas et al. (2004) reported that *K. rosea*, a catheter-associated bacterium, is vancomycin sensitive; however, antimicrobial treatment is ineffective until the catheter is removed, indicating that the biofilm formation on the surface of the catheter can protect the bacterial community from antimicrobial action (Savini et al., 2010). Thus, the ability to form biofilms can facilitate the co-existence of halophilic bacteria in the biofilm in the brine, leading to the contamination of the final products and damage to equipment. It could also lead to the development of resistance to antibacterial agents or disinfectants, resulting in serious hygiene problems and economic losses (Barnes et al., 1999).

Antimicrobial resistance of *K. salsicia* strains

The antimicrobial susceptibility of the *K. salsicia* isolates obtained from cheese brine was investigated (Table 4). Five of the nine isolates (55.6%) were MDR, showing resistance to at least three different classes of antimicrobials. Moreover, the isolates showed higher resistance to the fluoroquinolone class of antimicrobials, including MAR, ENO, and PRA, than to other antimicrobials. *Kocuria* spp. are sensitive to ampicillin, CLI, ERY, GEN, SXT, and cotrimoxazole antimicrobials (Savini et al., 2010; Sohn et al., 2015). Becker et al. (2008) reported that quinolone antimicrobials are effective against *Kocuria* spp. However, in our results, the *K. salsicia* isolates showed high minimal inhibitory concentration values against fluoroquinolone antimicrobials ENO and MAR (Table 4). These results suggest that *Kocuria* spp. have developed antimicrobial resistance to quinolones over the past decade. The antimicrobials sold for use in Korean livestock farms have increased by more than 70%, from 57 tons in 2010 to 97 tons in 2019 (APQA, 2019). The sales of ENO account for more than 70%–80% of quinolone antimicrobials and more than 2 tons of MAR have been sold since 2014 (APQA, 2019). In the present study, most antimicrobials were effective against the *K. salsicia* isolates (KS1–KS9); however, five of the nine isolates showed resistance to at least three antimicrobials. A previous study reported that MDR bacteria isolated from the cheese-making environment mainly acquired resistance genes from the environment and animal facilities (Kang et al., 2018). As antimicrobial resistance genes can be transferred between bacteria, which may occur during food production, it is necessary to perform antimicrobial stewardship at the farming stage (Jang et al., 2020).

Conclusion

Nine *K. salsicia* strains (KS1–KS9) were for the first time isolated from cheese brine in this study. They grew at a wide

Table 4. Antimicrobial susceptibility testing of the *Kocuria salsicia* isolates (KS1–KS9)

Sample (no. of isolates)	Isolate ID	MIC value (interpretation)												MDR
		Antimicrobial agent												
		AMK	CHL	CLI	GEN	POD	ENO	ERY	MAR	MIN	NIT	PRA	SXT	
Cheese brine	PC	≤2 (S)	8 (S)	≤0.12 (S)	≤0.5 (S)	2 (S)	0.12 (S)	≤0.25 (S)	0.25 (S)	0.25 (S)	16 (S)	0.12 (S)	≤10 (S)	– ¹⁾
	KS1	≤2 (S)	≤4 (S)	0.25 (S)	≤0.5 (S)	1 (S)	2 (I)	≤0.25 (S)	≥4 (R)	1 (I)	256 (R)	1 (I)	≤10 (S)	–
	KS2	4 (S)	≤4 (S)	≤0.12 (S)	1 (S)	2 (S)	≥4 (R)	≤0.25 (S)	≥4 (R)	≥16 (R)	256 (R)	2 (R)	≤10 (S)	+ ²⁾
	KS3	≤2 (S)	≤4 (S)	≤0.12 (S)	≤0.5 (S)	1 (S)	2 (I)	≤0.25 (S)	≥4 (R)	8 (R)	≥512 (R)	0.25 (S)	≤10 (S)	+
	KS4	≤2 (S)	8 (S)	0.5 (S)	≤0.5 (S)	2 (S)	≥4 (R)	≤0.25 (S)	≥4 (R)	1 (I)	256 (R)	1 (I)	≤10 (S)	–
	KS5	4 (S)	≤4 (S)	≤0.12 (S)	1 (S)	1 (S)	≥4 (R)	≤0.25 (S)	≥4 (R)	8 (R)	256 (R)	1 (I)	≤10 (S)	+
	KS6	≤2 (S)	≤4 (S)	0.25 (S)	≤0.5 (S)	1 (S)	≥4 (R)	≤0.25 (S)	≥4 (R)	1 (I)	256 (R)	0.5 (I)	≤10 (S)	–
	KS7	4 (S)	≤4 (S)	0.25 (S)	1 (S)	4 (I)	≥4 (R)	≤0.25 (S)	≥4 (R)	≥16 (R)	≥512 (R)	2 (R)	≤10 (S)	+
	KS8	≤2 (S)	≤4 (S)	0.25 (S)	≤0.5 (S)	0.5 (S)	≥4 (R)	≤0.25 (S)	≥4 (R)	≤0.5 (S)	256 (R)	1 (I)	≤10 (S)	–
KS9	4 (S)	≤4 (S)	0.25 (S)	1 (S)	0.5 (S)	≥4 (R)	≤0.25 (S)	≥4 (R)	4 (R)	256 (R)	2 (R)	≤10 (S)	+	

¹⁾ – indicates a negative result for antimicrobial resistance against more than three antimicrobial categories.

²⁾ + indicates a positive result for antimicrobial resistance against more than three antimicrobial categories.

MIC, minimum inhibitory concentration; AMK, amikacin; CHL, chloramphenicol; CLI, clindamycin; GEN, gentamicin; POD, cefpodoxime; ENO, enrofloxacin; ERY, erythromycin; MAR, marbofloxacin; MIN, minocycline; NIT, nitrofurantoin; PRA, pradofloxacin; SXT, trimethoprim/sulfamethoxazole; (R), resistant; (I), intermediate; (S), susceptible; MDR, multidrug-resistant; PC, positive control (*Staphylococcus aureus* ATCC 29213); KS1–KS9, *K. salsicia* KS1–KS9; ATCC, American Type Culture Collection.

range of temperatures, had potential biofilm-forming ability, and showed antimicrobial resistance. The results of the current study showed that brine could serve as an important reservoir for various halotolerant or halophilic microorganisms. Therefore, careful monitoring and hygienic handling of cheese brine are needed to prevent microbial contamination of the final product during cheese production in farmstead dairy plants. A major limitation of this study is that we did not evaluate the halophile-related gene clusters using whole genome sequencing nor perform a co-incubation growth analysis using cheese starter strains. Thus, further investigation is warranted to assess the halophilic characteristics of *K. salsicia* isolated from cheese brine and fate of *K. salsicia* during the ripening process.

Conflicts of Interest

The authors declare no potential conflict of interest.

Acknowledgments

The authors thank Yong-Seok Jang and Hyeon-Jin Kim for technical assistance and critical review of the manuscript and Kwang-Young Song for reviewing the manuscript. This research was supported by a grant (20162MFDS027) from the Ministry of Food and Drug Safety in 2021. The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Author Contributions

Conceptualization: Youn HY, Seo KH. Data curation: Youn HY. Formal analysis: Youn HY. Methodology: Youn HY. Investigation: Youn HY. Writing - original draft: Youn HY. Writing - review & editing: Youn HY, Seo KH.

Data Availability

The 16S rRNA sequences of the following *K. salsicia* isolates were deposited in the GenBank database: KS1 (accession number: MW301599), KS2 (accession number: MW301601), KS3 (accession number: MW301600), KS4 (accession number: MW301603), KS5 (accession number: MW301604), KS6 (accession number: MW301605), KS7 (accession number: MW301606), KS8 (accession number: MW301607), and KS9 (accession number: MW301608).

Ethics Approval

This article does not require IRB/IACUC approval because there are no human and animal participants.

References

- Altuntas F, Yildiz O, Eser B, Gündogan K, Sumerkan B, Çetin M. 2004. Catheter-related bacteremia due to *Kocuria rosea* in a patient undergoing peripheral blood stem cell transplantation. *BMC Infect Dis* 4:1-3.
- Animal and Plant Quarantine Agency [APQA]. 2019. National antibiotic use and resistance monitoring. Available from:

- https://www.mfds.go.kr/brd/m_231/view.do?seq=33047&srchFr=&srchTo=&srchWord=&srchTp=&itm_seq_1=0&itm_seq_2=0&multi_itm_seq=0&company_cd=&company_nm=&page=1. Accessed at Oct 10, 2021.
- Banks MK, Bryers JD. 1991. Bacterial species dominance within a binary culture biofilm. *Appl Environ Microbiol* 57:1974-1979.
- Barnes LM, Lo MF, Adams MR, Chamberlain AHL. 1999. Effect of milk proteins on adhesion of bacteria to stainless steel surfaces. *Appl Environ Microbiol* 65:4543-4548.
- Basaglia G, Carretto E, Barbarini D, Moras L, Scalone S, Marone P, De Paoli P. 2002. Catheter-related bacteremia due to *Kocuria kristinae* in a patient with ovarian cancer. *J Clin Microbiol* 40:311-313.
- Becker K, Rutsch F, Uekötter A, Kipp F, König J, Marquardt T, Peters G, von Eiff C. 2008. *Kocuria rhizophila* adds to the emerging spectrum of micrococcal species involved in human infections. *J Clin Microbiol* 46:3537-3539.
- Bintsis T, Papademas P. 2002. Microbiological quality of white-brined cheeses: A review. *Int J Dairy Technol* 55:113-120.
- Bruins MJ, Juffer P, Wolfhagen MJ, Ruijs GJ. 2007. Salt tolerance of methicillin-resistant and methicillin-susceptible *Staphylococcus aureus*. *J Clin Microbiol* 45:682-683.
- Clinical and Laboratory Standards Institute [CLSI]. 2020a. Performance standards for antimicrobial disk and dilution susceptibility tests for bacteria isolated from animals. 5th ed. CLSI, Wayne, PA, USA.
- Clinical and Laboratory Standards Institute [CLSI]. 2020b. Performance standards for antimicrobial susceptibility testing. 30th ed. CLSI, Wayne, PA, USA.
- D'Amico DJ, Donnelly CW. 2008. Enhanced detection of *Listeria* spp. in farmstead cheese processing environments through dual primary enrichment, PCR, and molecular subtyping. *J Food Prot* 71:2239-2248.
- de Jesus Pimentel-Filho N, de Freitas Martins MC, Nogueira GB, Mantovani HC, Vanetti MCD. 2014. Bovicin HC5 and nisin reduce *Staphylococcus aureus* adhesion to polystyrene and change the hydrophobicity profile and Gibbs free energy of adhesion. *Int J Food Microbiol* 190:1-8.
- Di Ciccio P, Vergara A, Festino AR, Paludi D, Zanardi E, Ghidini S, Ianieri A. 2015. Biofilm formation by *Staphylococcus aureus* on food contact surfaces: Relationship with temperature and cell surface hydrophobicity. *Food Control* 50:930-936.
- Dunn R, Bares S, David MZ. 2011. Central venous catheter-related bacteremia caused by *Kocuria kristinae*: Case report and review of the literature. *Ann Clin Microbiol Antimicrob* 10:31.
- European Food Safety Authority, European Centre for Disease Prevention and Control. 2019. The European Union summary report on antimicrobial resistance in zoonotic and indicator bacteria from humans, animals and food in 2017. *EFSA J* 17:e05598.
- Fox E, Hunt K, O'Brien M, Jordan K. 2011. *Listeria monocytogenes* in Irish Farmhouse cheese processing environments. *Int J Food Microbiol* 145:S39-S45.
- Golob M, Pate M, Kušar D, Dermota U, Avberšek J, Papič B, Zdovc I. 2019. Antimicrobial resistance and virulence genes in *Enterococcus faecium* and *Enterococcus faecalis* from humans and retail red meat. *BioMed Res Int* 2019:2815279.
- Haastrup MK, Johansen P, Malskær AH, Castro-Mejía JL, Kot W, Krych L, Arneborg N, Jespersen L. 2018. Cheese brines from Danish dairies reveal a complex microbiota comprising several halotolerant bacteria and yeasts. *Int J Food Microbiol* 285:173-187.
- Ingham SC, Su YC, Spangenberg DS. 2000. Survival of *Salmonella typhimurium* and *Escherichia coli* O157:H7 in cheese brines. *Int J Food Microbiol* 61:73-79.

- Jang YS, Kim DH, Bae D, Kim SH, Kim H, Moon JS, Song KY, Chon JW, Seo KH. 2020. Prevalence, toxin-typing, and antimicrobial susceptibility of *Clostridium perfringens* from retail meats in Seoul, Korea. *Anaerobe* 64:102235.
- Jeong D, Kim DH, Song KY, Seo KH. 2018. Antimicrobial and anti-biofilm activities of *Lactobacillus kefiranofaciens* DD2 against oral pathogens. *J Oral Microbiol* 10:1472985.
- Kandi V, Palange P, Vaish R, Bhatti AB, Kale V, Kandi MR, Bhoomagiri MR. 2016. Emerging bacterial infection: Identification and clinical significance of *Kocuria* species. *Cureus* 8:e731.
- Kang IB, Kim DH, Chon JW, Seo KH. 2018. Effect of microbial control measures on farmstead cheesemaking and antimicrobial resistance of *Staphylococcus aureus* and *Enterococcus* spp. isolates. *J Food Saf* 38:e12432.
- Kim SB, Nedashkovskaya OI, Mikhailov VV, Han SK, Kim KO, Rhee MS, Bae KS. 2004. *Kocuria marina* sp. nov., a novel actinobacterium isolated from marine sediment. *Int J Syst Evol Microbiol* 54:1617-1620.
- Larson AE, Johnson EA, Nelson JH. 1999. Survival of *Listeria monocytogenes* in commercial cheese brines. *J Dairy Sci* 82:1860-1868.
- Lee JY, Kim SH, Jeong HS, Oh SH, Kim HR, Kim YH, Lee JN, Kook JK, Kho WG, Bae IK, Shin JH. 2009. Two cases of peritonitis caused by *Kocuria marina* in patients undergoing continuous ambulatory peritoneal dialysis. *J Clin Microbiol* 47:3376-3378.
- Locatelli C, Cremonesi P, Bertocchi L, Zanoni MG, Barberio A, Drigo I, Varisco G, Castiglioni B, Bronzo V, Moroni P. 2016. Methicillin-resistant *Staphylococcus aureus* in bulk tank milk of dairy cows and effect of swine population density. *J Dairy Sci* 99:2151-2156.
- McMahon DJ, Motawee MM, McManus WR. 2009. Influence of brine concentration and temperature on composition, microstructure, and yield of feta cheese. *J Dairy Sci* 92:4169-4179.
- Mehli L, Hoel S, Thomassen GMB, Jakobsen AN, Karlsen H. 2017. The prevalence, genetic diversity and antibiotic resistance of *Staphylococcus aureus* in milk, whey, and cheese from artisan farm dairies. *Int Dairy J* 65:20-27.
- Meletis G, Gogou V, Palamouti M, Spiropoulos P, Xanthopoulou K, Tantou P, Thomoglou V, Rizou A, Thomoglou V. 2012. Catheter-related relapsing peritonitis due to *Kocuria varians* in a patient undergoing continuous ambulatory peritoneal dialysis. *Nefrología (Madrid)* 32:541-542.
- Moreira JS, Riccetto AGL, da Silva MTN, dos Santos Vilela MM. 2015. Endocarditis by *Kocuria rosea* in an immunocompetent child. *Braz J Infect Dis* 19:82-84.
- Nilsson RE, Ross T, Bowman JP. 2011. Variability in biofilm production by *Listeria monocytogenes* correlated to strain origin and growth conditions. *Int J Food Microbiol* 150:14-24.
- Purty S, Saranathan R, Prashanth K, Narayanan K, Asir J, Sheela Devi C, Kumar Amarnath S. 2013. The expanding spectrum of human infections caused by *Kocuria* species: A case report and literature review. *Emerg Microbes Infect* 2:1-8.
- Rode TM, Langsrud S, Holck A, Møretør T. 2007. Different patterns of biofilm formation in *Staphylococcus aureus* under food-related stress conditions. *Int J Food Microbiol* 116:372-383.
- Ryu JH, Beuchat LR. 2005. Biofilm formation by *Escherichia coli* O157:H7 on stainless steel: Effect of exopolysaccharide and Curli production on its resistance to chlorine. *Appl Environ Microbiol* 71:247-254.
- Savini V, Catavittello C, Masciarelli G, Astolfi D, Balbinot A, Bianco A, Febbo F, D'Amario C, D'Antonio D. 2010. Drug sensitivity and clinical impact of members of the genus *Kocuria*. *J Med Microbiol* 59:1395-1402.
- Sohn KM, Baek JY, Kim SH, Cheon S, Kim YS. 2015. Catheter-related bacteremia caused by *Kocuria salsicia*: The first case. *J Infect Chemother* 21:305-307.

- Soni I, Chakrapani H, Chopra S. 2015. Draft genome sequence of methicillin-sensitive *Staphylococcus aureus* ATCC 29213. *Genome Announc* 3:e01095-15.
- Welch K, Cai Y, Strømme M. 2012. A method for quantitative determination of biofilm viability. *J Funct Biomater* 3:418-431.
- Xu H, Zou Y, Lee HY, Ahn J. 2010. Effect of NaCl on the biofilm formation by foodborne pathogens. *J Food Sci* 75:M580-M585.
- Zwietering MH, Jongenburger I, Rombouts FM, Van't Riet K. 1990. Modeling of the bacterial growth curve. *Appl Environ Microbiol* 56:1875-1881.