

Outcome of Parkinson's Disease Patients Affected by COVID-19

There is extensive debate on the neurological consequences of corona virus disease 2019 (COVID-19) and the impact this might have for patients with neurodegenerative conditions, including Parkinson's disease (PD). Older advanced PD patients may represent a particularly vulnerable population, as respiratory muscle rigidity as well as impairment of cough reflex alongside preexisting dyspnoea may lead to increased severity of COVID-19.¹ In addition, there are indirect possible effects, such as the impact of stress, self-isolation, and anxiety, as well as the consequences of prolonged immobility because of the lockdown.^{2,3}

Several observations make the link between COVID-19 and PD particularly intriguing. Antibodies against coronavirus were found in the cerebrospinal fluid of PD patients more than 2 decades ago, suggesting a possible role for viral infections in neurodegeneration.⁴ Angiotensin-converting enzyme 2 (ACE2) receptors are highly expressed in dopamine neurons, and they are reduced in PD because of the degenerative process; therefore, severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2)-related brain penetration may cause additional harm and worsen symptoms and may increase the requirement of dopamine replacement therapy, as evident in 5 of our cases.^{5,6} Interestingly, the ability of coronaviruses to enter the brain through the nasal cavity determines anosmia/hyposmia and ageusia in many infected subjects, a symptomatology that closely resembles one of the most prominent premotor symptom of PD.⁷ Finally, the dopamine synthetic pathway is possibly involved in the pathophysiology of COVID-19, as ACE2 and dopamine decarboxylase coexpress and coregulate in nonneuronal cell types, which may indicate dopamine depletion and the need for considering levodopa as treatment.⁸

Outcomes of PD patients infected by SARS-CoV-2 are unknown. We present here the outcome of 10 clinical cases (Table 1) collected from the experience at the Parkinson and Movement Disorders Unit in Padua, Italy, and the Parkinson's Foundation Centre of Excellence at King's College Hospital in London, UK, from the beginning of March to the current period. The PD center in Padua has a catchment of 1022 patients, mainly in the province of Padua, which had 3407 cases of COVID-19, 2 of whom were advanced PD patients. Both were women residing in nursing homes with severe motor manifestations, and both were treated with levodopa therapy (Table 1). One remained asymptomatic, whereas the other, who in the last months had been suffering from

deteriorating cognition and hallucinations, developed respiratory problems and died. The King's center has 4000 PD patients in the catchment and is currently following more than 600 patients. In all, thus far, 8 cases have been identified with COVID-19, and clinical details are presented here. The King's COVID-19 PD group consists of 6 men and 2 women, all older than 60 years of age with severe motor dysfunction, comorbidities, and most requiring additional levodopa dosing following infection (Table 1). Anxiety and other nonmotor features, such as fatigue, orthostatic hypotension, cognitive impairment, and psychosis, also worsened during the infection. Fatigue was a dominant symptom during the SARS-CoV-2 infection in all cases on advanced therapies. Three patients died from COVID-19 pneumonia.

These findings suggest that PD patients of older age (mean, 78.3 years) with longer disease duration (mean, 12.7 years) are particularly susceptible to COVID-19 with a substantially high mortality rate (40%). Those on advanced therapies, such as deep brain stimulation or levodopa infusion therapy, seem especially vulnerable, and a mortality rate of 50% among our 4 such cases is of concern. Although 2 recent articles have addressed the issue of COVID-19 in PD, none specifically showed cases directly affected by COVID-19, and we believe ours is the first report of this nature.^{2,3} The report of Prasad et al focuses on the perceptions of SARS-CoV-2 infection in 100 PD patients and some symptoms listed tally with our observations in a real-life population with COVID-19.² Many national and charity guidelines do not list PD or specifically older subjects on advanced therapies as a susceptible group, and this information needs to be amended in light of these new data.⁹ ■

Acknowledgments: The views expressed are those of the authors and not necessarily those of the NHS, NIHR, or Department of Health. There are no funding sources to disclose. The authors also thank all clinicians and front-line staff engaged in the care of COVID-19 patients at King's College and Princess Royal Hospitals, London, UK, as well as at the Parkinson and Movement Disorders Unit in Padua, Italy.

Angelo Antonini, PhD,¹  Valentina Leta, MD,^{2,3}
James Teo, PhD,^{2,4} and K. Ray Chaudhuri, DSc^{2,3*} 
¹Parkinson and Movement Disorders Unit, University of Padua, Padua, Italy
²King's College London, Institute of Psychiatry, Psychology & Neuroscience, London, United Kingdom
³Parkinson's Foundation Centre of Excellence, King's College Hospital, London, United Kingdom
⁴King's College Hospital NHS Foundation Trust, London, United Kingdom

© 2020 International Parkinson and Movement Disorder Society

Key Words: COVID-19, Parkinson's disease, SARS-CoV-2

***Correspondence to:** Professor K. Ray Chaudhuri, Department of Basic and Clinical Neuroscience, The Maurice Wohl Clinical Neuroscience Institute, King's College London, Cutcombe Road, London, SE5 9RT, United Kingdom; E-mail: ray.chaudhuri@kcl.ac.uk

Relevant conflicts of interest/financial disclosures: None.

Funding agencies: None.

Received: 23 April 2020; **Accepted:** 24 April 2020

Published online 28 May 2020 in Wiley Online Library (wileyonlinelibrary.com). DOI: 10.1002/mds.28104

TABLE 1. Clinical features and outcomes of Parkinson's disease patients with corona virus disease 2019.

Patients	Age	Sex	PD duration (years)	PD therapy	Comorbidities	Clinical picture requiring SARS-CoV-2 testing	Therapeutic interventions (antibiotics and intensive care)	Outcome
1	76	F	28	<ul style="list-style-type: none"> Carbidopa-levodopa 25/100 mg 1 table t 3 times daily and 1/2 tablet once daily Rotigotine 4 mg 1 patch once daily Safinamide 50 mg 1 tablet once daily 	<ul style="list-style-type: none"> Dementia Dysphagia Severe joint deformities 	<ul style="list-style-type: none"> Fever 	<ul style="list-style-type: none"> No intensive care required 	Spontaneous recovery
2	79	F	12	<ul style="list-style-type: none"> Carbidopa-levodopa 25/100 mg 1 1/2 tablets 4 times daily 	<ul style="list-style-type: none"> Dementia Hallucinations 	<ul style="list-style-type: none"> Fever Cough Shortness of breath Confusion 	<ul style="list-style-type: none"> CPAP required (no resuscitation was advised) 	Died 14 days after onset of respiratory symptoms
3	81	M	10	<ul style="list-style-type: none"> Carbidopa-Levodopa CR 25/100 mg 1 tablet once daily Carbidopa-Levodopa 25/100 mg 2 tab 3 times daily 	<ul style="list-style-type: none"> Hypertension Ischemic heart disease Chronic kidney disease Dementia Angina pectoris Osteoporosis 	<ul style="list-style-type: none"> Fever Dry cough Shortness of breath 	<ul style="list-style-type: none"> Antibiotics (piperacillin/tazobactam + clarithromycin) No intensive care or CPAP required Required increased levodopa dosing 	Transferred to rehabilitation ward after 11 days of hospitalization
4	94	M	2	<ul style="list-style-type: none"> Carbidopa-Levodopa 25/100 mg 1 tablet 3 times daily 	<ul style="list-style-type: none"> Dementia Angina pectoris Osteoporosis 	<ul style="list-style-type: none"> Multiple falls and head injury Cough Delirium Arrhythmia 	<ul style="list-style-type: none"> Antibiotics (amoxicillin/clavulanic acid + clarithromycin) No intensive care or CPAP required 	Transferred to rehabilitation ward for respite care; 10 days of hospitalisation
5	87	M	6	<ul style="list-style-type: none"> Carbidopa-Levodopa 25/100 mg 1 tablet twice daily Carbidopa-Levodopa 25/100 mg 1 1/2 tablets twice daily 	<ul style="list-style-type: none"> Congestive Cardiac failure Chronic obstructive pulmonary disease Atrial fibrillation Orthostatic hypotension Asthma Anxiety disorder 	<ul style="list-style-type: none"> Syncopal episodes Hallucinations Fall with facial trauma Cough 	<ul style="list-style-type: none"> Antibiotics (amoxicillin/clavulanic acid) No intensive care. CPAP not possible because of facial trauma (no resuscitation was advised) Required increased levodopa dosing 	Died after 10 days of hospitalization
6	83	F	3	<ul style="list-style-type: none"> Carbidopa-Levodopa 25/100 mg 2 tablets twice daily and 1 1/2 tablets twice daily 	<ul style="list-style-type: none"> Anxiety disorder 	<ul style="list-style-type: none"> Worsening of mobility with fall and hyperCKemia Dry cough Fatigue Worsening of anxiety 	<ul style="list-style-type: none"> No intensive care or CPAP required Required increase in levodopa dosing 	Transferred to neurorehabilitation after 25 days of hospitalization

(Continues)

TABLE 1. Continued

Patients	Age	Sex	PD duration (years)	PD therapy	Comorbidities	Clinical picture requiring SARS-CoV-2 testing	Therapeutic interventions(antibiotics and intensive care)	Outcome
7	61	M	17	<ul style="list-style-type: none"> • I/LJ 1684 mg daily • Carbidopa-levodopa 25/100 mg 1 tablet once daily • Rotigotine 4 mg 1 patch once daily • Opicapone 50 mg 1 tablet once daily 	<ul style="list-style-type: none"> • Traumatic fractures secondary to falls • Orthostatic hypotension 	<ul style="list-style-type: none"> • Fever • Cough • Fatigue 	<ul style="list-style-type: none"> • Antibiotics • CPAP required 	Died during hospitalization
8	75	M	10	<ul style="list-style-type: none"> • I/LJ 1262 mg daily • Opicapone 50 mg 1 tablet once daily • Carbidopa-Levodopa CR 50/200 mg 1 tablet once daily • Carbidopa-Levodopa CR 25/100 mg 1 tablet once daily • Levodopa- Benserazide 100/50 mg 1 tablet 3 times daily 	<ul style="list-style-type: none"> • Orthostatic hypotension • Osteoporosis • Benign prostatic hyperplasia • Depression • Neuropathic pain 	<ul style="list-style-type: none"> • Fever • Cough • Fatigue 	<ul style="list-style-type: none"> • Antibiotics • CPAP required 	Died during hospitalization
9	72	M	15	<ul style="list-style-type: none"> • DBS bilateral STN • Subcutaneous Apomorphine infusion 46 mg daily • Rasagiline 1 mg 1 tablet once daily • Ropirinole CR 2 mg 1 tablet once daily • Levodopa-benserazide 50/12.5 mg 1 tablet 4 times daily • Carbidopa-Levodopa CR 25/100 mg 1 tablet 3 times daily 	<ul style="list-style-type: none"> • Asthma • Diabetes mellitus type II 	<ul style="list-style-type: none"> • Fever • Fatigue • Worsening of motor symptoms 	<ul style="list-style-type: none"> • Required increase in levodopa dosing • No intensive care or CPAP required 	Spontaneous recovery, waiting for transfer for rehabilitation
10	75	F	24	<ul style="list-style-type: none"> • I/LJ stopped due to recent PEG-J tube malfunctioning • Levodopa-Benserazide 100/25 mg six times daily • Carbidopa-Levodopa CR 25/100 mg once daily 	<ul style="list-style-type: none"> • Atrial fibrillation • Cognitive impairment 	<ul style="list-style-type: none"> • Fatigue • Cough 	<ul style="list-style-type: none"> • Required increase in levodopa dosing • No intensive care or CPAP required 	Recovering

CPAP, continuous positive airway pressure; CR, controlled release; DBS, deep brain stimulation; I/LJ, intrajejunal levodopa infusion; F, female; M, male; PD, Parkinson's disease; PEG-J, percutaneous endoscopic transgastric jejunostomy; SARS-CoV-2, severe acute respiratory syndrome coronavirus 2; STN, subthalamic nucleus.

References

1. van Wamelen DJ, Leta V, Johnson J, et al. Drooling in Parkinson's disease: prevalence and progression from the Non-motor International Longitudinal Study. *Dysphagia* 2020; <https://doi.org/10.1007/s00455-020-10102-5>
2. Prasad S, Holla VV, Neeraja K, et al. Parkinson's disease and COVID-19: perceptions and implications in patients and caregivers. *Mov Disord* 2020; <https://doi.org/10.1002/mds.28088>.
3. Helmich RC, Bloem BR. The impact of the COVID-19 pandemic on Parkinson's disease: hidden sorrows and emerging opportunities. *J Parkinsons Dis* 2020;10(2):351–354.
4. Fazzini E, Fleming J, Fahn S. Cerebrospinal fluid antibodies to coronavirus in patients with Parkinson's disease. *Mov Disord* 1992;7(2): 153–158.
5. Rodriguez-Perez AI, Garrido-Gil P, Pedrosa MA, et al. Angiotensin type 2 receptors: role in aging and neuroinflammation in the substantia nigra. *Brain Behav Immun* 2019;S0889-1591(19)31128-6.
6. Joglar B, Rodriguez-Pallares J, Rodriguez-Perez AI, Rey P, Guerra MJ, Labandeira-Garcia JL. The inflammatory response in the MPTP model of Parkinson's disease is mediated by brain angiotensin: relevance to progression of the disease. *J Neurochem* 2009;109(2):656–669.
7. Schaeffer E, Postuma RB, Berg D. Prodromal PD: a new nosological entity. *Prog Brain Res* 2020;252:331–356.
8. Nataf S. An alteration of the dopamine synthetic pathway is possibly involved in the pathophysiology of COVID-19. *J Med Virol* 2020; doi: <https://doi.org/10.1002/jmv.25826>.
9. Fasano A, Antonini A, Katzenschlager R, et al. Management of advanced therapies in Parkinson's disease patients in times of humanitarian crisis: the COVID-19 experience *Mov Disord Clin Pract* 2020;7(4): 361–372.