



Corrigendum

Korean J Anesthesiol 2020;73(3):267
<https://doi.org/10.4097/kjae.2016.69.4.368.e1>
pISSN 2005-6419 • eISSN 2005-7563

An ultrasound-guided fascia iliaca catheter technique does not impair ambulatory ability within a clinical pathway for total hip arthroplasty

Seshadri C. Mudumbai^{1,3}, T. Edward Kim^{1,3},
Steven K. Howard^{1,3}, Nicholas J. Giori^{2,3}, Steven Woolson^{2,3},
Toni Ganaway¹, Alex Kou¹, Robert King³,
Edward R. Mariano^{1,3}

Departments of ¹Anesthesiology, Perioperative and Pain Medicine, ²Orthopaedic Surgery, Stanford University School of Medicine, Stanford, ³Veterans Affairs Palo Alto Health Care System, Palo Alto, CA, USA

Korean Journal of Anesthesiology 2016;69(4):368-375.
<https://ekja.org/journal/view.php?doi=10.4097/kjae.2016.69.4.368>

The article by Mudumbai SC, et al. entitled, “An ultrasound-guided fascia iliaca catheter technique does not impair ambulatory ability within a clinical pathway for total hip arthroplasty.” contained an error in the result.

Before correction:

Page 371, The primary outcome, total ambulation distance in meters (median [Q1-Q3]), did not differ between the two groups (FIC 63 [30–120] vs. ITM (83 [48–1114]; P = 0.08).

The correct information is found below:

Page 371, The primary outcome, total ambulation distance in meters (median [Q1-Q3]), did not differ between the two groups (FIC 63 [30–120] vs. ITM (83 [48–114]; P = 0.08).

The authors apologize for any inconvenience this mistake may have caused.

© The Korean Society of Anesthesiologists, 2020

© This is an open-access article distributed under the terms of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.