

CORRECTION

Open Access

Correction: Comparative genomic profiling of Dutch clinical *Bordetella pertussis* isolates using DNA microarrays: identification of genes absent from epidemic strains

Audrey J King^{1*}, Tamara van Gorkom¹, Jeroen LA Pennings², Han GJ van der Heide¹, Qiushui He³, Dimitri Diavatopoulos⁴, Kees Heuvelman¹, Marjolein van Gent¹, Karin van Leeuwen¹, Frits R Mooi¹

Correction

After the publication of this work [1] we found out that Table 1 was not filled in correctly in the original article. The total number of strains analyzed in the *ptxP1* or the *ptxP3* lineage was not shown between the brackets, leading to incompleteness of the data. The revised Table 1 is shown below.

We regret any inconvenience that the incompleteness of Table 1 in the original article might have caused.

Author details

¹Laboratory for Infectious Diseases and Screening (LIS) Centre for Infectious Disease Control, National Institute for Public Health and the Environment (RIVM), Bilthoven, The Netherlands. ²Laboratory for Health Protection Research, National Institute for Public Health and the Environment (RIVM), Bilthoven, The Netherlands. ³Pertussis Reference Laboratory, National Public Health Institute, Finland. ⁴Department of Microbiology and Immunology, University of Melbourne, Victoria, Australia.

Received: 19 March 2010 Accepted: 24 March 2010

Published: 24 March 2010

References

1. King AJ, van Gorkom T, Pennings JL, van der Heide HG, He Q, Diavatopoulos D, et al: Comparative genomic profiling of Dutch clinical *Bordetella pertussis* isolates using DNA microarrays: identification of genes absent from epidemic strains. *BMC Genomics* 2008, **9**:311.

doi:10.1186/1471-2164-11-196

Cite this article as: King et al.: Correction: Comparative genomic profiling of Dutch clinical *Bordetella pertussis* isolates using DNA microarrays: identification of genes absent from epidemic strains. *BMC Genomics* 2010 **11**:196.

* Correspondence: Audrey.King@rivm.nl

¹Laboratory for Infectious Diseases and Screening (LIS) Centre for Infectious Disease Control, National Institute for Public Health and the Environment (RIVM), Bilthoven, The Netherlands

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



Table 1 Presence of regions of difference in Dutch clinical isolates from 1993 to 2004

Region of difference	BP-number	No. of genes	Size	<i>ptxP1</i> lineage	<i>ptxP3</i> lineage
Locus:			(kb)		
<i>RD-3</i>	BP0910A-BP0937	23	24.9	0% (17)	0% (26)
<i>RD-5</i>	BP1135-BP1141	7	8.6	0% (17)	0% (26)
<i>RD-6</i>	BP1158-BP1176	19	18.7	94% (17)	100% (26)
<i>RD-27</i>	BP1553	1	0.8	82% (17)	100% (26)
<i>RD-10</i>	BP1948-BP1966	18	22.7	100% (27)	0% (53)
<i>RD-28</i>	BP2122-BP2123	2	1.7	94% (17)	100% (26)
<i>RD-29</i>	BP2822-BP2839	17	16.9	94% (17)	100% (26)
<i>RD-18</i>	BP3314-BP3322	9	9.4	94% (17)	92% (26)
<i>RD-1*</i>	BP0515-BP0516	2	1	0% (17)	13% (32)

Regions of difference (RDs) are numbered according to Brinig *et al* [23]. *RD-1** is part of *RD-1* as was found by Brinig *et al* [23]. The percentage of strains harbouring the region and the number of strains analyzed are indicated. Size, refers to the size of the deletion.