




RESEARCH ARTICLE

Determination of the minimum protective dose of a glycoprotein-G-deficient infectious laryngotracheitis virus vaccine delivered via eye-drop to week-old chickens

Mesula G. Korsa, Joanne M. Devlin , Carol A. Hartley, Glenn F. Browning, Mauricio J. C. Coppo, José A. Quinteros , Carlos A. Loncoman , Adepeju E. Onasanya, Dulari Thilakarathne, Andrés Diaz-Méndez*

Asia-Pacific Centre for Animal Health, Melbourne Veterinary School, Faculty of Veterinary and Agricultural Sciences, The University of Melbourne, Parkville, Victoria, Australia

* andres.diaz@unimelb.edu.au



 OPEN ACCESS

Citation: Korsa MG, Devlin JM, Hartley CA, Browning GF, Coppo MJC, Quinteros JA, et al. (2018) Determination of the minimum protective dose of a glycoprotein-G-deficient infectious laryngotracheitis virus vaccine delivered via eye-drop to week-old chickens. PLoS ONE 13(12): e0207611. <https://doi.org/10.1371/journal.pone.0207611>

Editor: Yongchang Cao, Sun Yat-Sen University, CHINA

Received: April 4, 2018

Accepted: November 2, 2018

Published: December 6, 2018

Copyright: © 2018 Korsa et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper.

Funding: This work was funded by the Poultry Cooperative Research Centre through Sub-Project 1.1.10 awarded to project leader Joanne M Devlin and was funded by the Asia Pacific Centre for Animal Health. Mesula G. Korsa was supported by Australian Government Research Training Program (RTP) scholarship. There was no additional

Abstract

Infectious laryngotracheitis (ILT) is an upper respiratory tract disease of chickens that is caused by infectious laryngotracheitis virus (ILTV), an alphaherpesvirus. This disease causes significant economic losses in poultry industries worldwide. Despite widespread use of commercial live attenuated vaccines, many poultry industries continue to experience outbreaks of disease caused by ILTV. Efforts to improve the control of this disease have resulted in the generation of new vaccine candidates, including ILTV mutants deficient in virulence factors. A glycoprotein G deletion mutant vaccine strain of ILTV (Δ G ILTV), recently licenced as Vaxsafe ILT (Bioproperties Pty Ltd), has been extensively characterised *in vitro* and *in vivo*, but the minimum effective dose required to protect inoculated animals has not been determined. This study performed a vaccination and challenge experiment to determine the minimum dose of Δ G ILTV that, when delivered by eye-drop to seven-day-old specific pathogen-free chickens, would protect the birds from a robust challenge with a virulent field strain of virus (class 9 ILTV). A dose of $10^{3.8}$ plaque forming units was the lowest dose capable of providing a high level of protection against challenge, as measured by clinical signs of disease, tracheal pathology and virus replication after challenge. This study has shown that the Δ G ILTV vaccine strain is capable of inducing a high level of protection against a virulent field virus at a commercially feasible dose. These results lay the foundations upon which a commercial vaccine can be developed, thereby offering the potential to provide producers with another important tool to help control ILTV.

1. Introduction

Infectious laryngotracheitis virus (ILTV) is an alphaherpesvirus that causes acute upper respiratory tract disease in chickens and has significant economic importance for poultry industries throughout the world [1]. Commercially available attenuated vaccines are commonly used to

external funding received for this study. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

Abbreviations: ILTV, infectious laryngotracheitis virus; ILT, infectious laryngotracheitis; Δ gG ILTV, glycoprotein G-deficient ILTV; dpc, day post challenge; PFU, plaque forming unit; UnVx-Uch, unvaccinated and unchallenged; UnVx-Ch, unvaccinated and challenged; SPF, specific pathogen free.

help control the disease [2, 3]. Despite the use of these vaccines, many poultry industries continue to experience outbreaks of disease caused by ILTV. In Australia, natural recombination between two distinct live attenuated vaccine strains resulted in the generation of virulent recombinant field strains of ILTV, including the class 8 and 9 viruses [4]. These recombinant viruses have spread and have caused disease in major poultry-producing areas of Australia. Another recombinant field virus, class 10 ILTV, emerged in 2013 in Australian poultry and has also spread and caused outbreaks of disease [5]. Taken together, available data suggest that improvements to control strategies for ILTV are needed, including tools to control the recently emerged, virulent recombinant viruses.

Recently, efforts to improve the use of ILTV vaccines, and to develop new ILTV vaccines, have resulted in the availability of more options for control of the disease by poultry producers. In some countries, vectored ILTV vaccines are in widespread use. These vaccines use a virus vector, such as fowlpox virus or herpesvirus of turkeys, to deliver specific ILTV antigens [6–11]. The generation of ILTV mutants deficient in virulence factors (deletion mutant vaccines) has also been investigated and these have the potential to offer additional tools for disease control [2, 3]. Previous *in vivo* and *in vitro* studies have shown that ILTV deficient in the virulence factor glycoprotein G (gG, a chemokine binding protein) has characteristics that would make it suitable for use as an attenuated vaccine (Δ gG ILTV) [12–20]

The Δ gG ILTV vaccine strain has been extensively studied *in vivo*. The first study to investigate the potential efficacy of the Δ gG ILTV vaccine strain delivered the virus at a dose of 10^3 plaque forming units (PFU), via intratracheal inoculation to four-week-old specific pathogen free (SPF) chickens, followed by challenge of the birds with the CSW-1 field strain (class 4) of ILTV at seven weeks of age. Birds that received the Δ gG ILTV vaccine had significantly fewer clinical signs than unvaccinated birds [13]. The suitability of the Δ gG ILTV vaccine strain was then investigated using delivery methods suitable for mass vaccination programs, including delivery via drinking water, eye-drop or *in ovo* inoculation [12, 15–17]. The results from these studies demonstrated that the Δ gG ILTV vaccine strain has desirable safety and efficacy characteristics. Although this vaccine strain has shown promise as a potential alternative to conventionally attenuated live ILTV vaccines, the minimum effective dose required to protect inoculated birds has not been determined. The aim of this study was therefore to determine the minimum dose of Δ gG ILTV that, when delivered by eye-drop to seven-day-old SPF chickens, would protect the birds from a robust challenge with virulent, recombinant class 9 virus.

2. Materials and methods

2.1. Experimental design, animals and virus strains used in this study

The study was approved by the Animal Ethics Committee of the Faculty of Veterinary and Agricultural Sciences, The University of Melbourne (Animal Ethics ID -1413097) in accordance with institutional and national guidelines. After hatching, 121 SPF chicks were divided into six groups (five groups with 22 birds and one group with 11 birds) that were each housed in separate isolator units. Vaccine (Vaxsafe ILT) was obtained from the manufacturer (Bioproperties Pty Ltd) as freeze-dried product, five passages beyond the master seed. Inoculum containing the Δ gG ILTV vaccine strain was then prepared by re-suspending the freeze-dried product in commercial sterile diluent (Merial Select Inc.) according to the manufacturer's directions. At seven days of age, four groups, each of which contained 22 birds, were vaccinated with $10^{5.0}$ PFU, $10^{3.8}$ PFU, $10^{3.5}$ PFU, or $10^{3.2}$ PFU of the Δ gG ILTV vaccine strain via eye-drop in a 30 μ L volume. All ILTV titres were determined using PFU assays performed in LMH cells, as previously described [12]. The remaining two groups of birds were mock vaccinated by administering the same volume of sterile diluent. After mock-vaccination, one bird

in the group of 11 birds developed a leg injury (unrelated to mock-vaccination) and was euthanized for animal welfare reasons, consistent with Animal Ethics ID -1413097.

At 28 days of age the vaccinated groups of birds were challenged with class 9 ILTV, a virulent recombinant field strain ($10^{3.0}$ PFU/bird), resuspended in vaccine diluent, with approximately half of the dose ($10^{2.7}$ PFU) delivered onto the conjunctiva (40 μ L each eye) and the other half ($10^{2.7}$ PFU) into the trachea (150 μ L). Similar procedures were used to inoculate one of the unvaccinated groups of 22 birds with the same virulent field strain, as a positive control group (unvaccinated-challenged). The remaining unvaccinated group of 11 birds was kept as a negative control group (unvaccinated-unchallenged). Birds were scored for clinical signs of disease from three to six days post-challenge (dpc). Typical clinical signs due to challenge with wild type virus are expected to peak during this period [12]. Any birds that showed severe signs of disease were humanely euthanized by anaesthetic (halothane) overdose and these birds were included in the recorded mortalities. Any recorded severe clinical signs in these birds prior to being euthanised were also included in the cumulative clinical scores. All remaining birds were euthanized at 7 dpc. At necropsy, the severity of ILTV induced tracheal pathology was recorded [12], and conjunctival and tracheal swabs were collected and placed in 1 mL of sterile viral transport medium consisting of Dulbecco's minimal essential medium (DMEM, Sigma-Aldrich) supplemented with 10% v/v foetal bovine serum (Sigma-Aldrich), 10 mM HEPES, pH 7.6, 50 μ g/ml ampicillin (Sigma-Aldrich) and 2.5 μ g/ml amphotericin B (Astral Scientific). The samples were transported on ice and then immediately stored at -80°C until processing. Clinical scores, gross pathological lesions, virus detection and replication in the conjunctival and tracheal mucosa were used to assess the level of protection induced in response to the different doses of Δ gG ILTV and are described in further detail below.

2.2. Clinical scoring

Clinical signs of demeanour, dyspnoea, and conjunctivitis were scored as previously described [13]. Briefly, demeanour was scored as 0 (normal demeanour), 1 (depressed demeanour) or 2 (severely depressed demeanour). Similarly, conjunctivitis was scored as 0 (conjunctival mucosa normal), 1 (partial eye closure) or 2 (complete eye closure, marked conjunctivitis), and dyspnoea was scored as 0 (normal breathing), 1 (mild dyspnoea), 2 (moderate dyspnoea), 3 (marked dyspnoea), or 4 (severe gasping).

2.3 Gross tracheal pathology scoring

Gross tracheal pathology was scored as previously described [13]. Briefly, at necropsy, the trachea was removed and then cut lengthways for gross lesion scoring. The severity of the lesions was scored as 0 (normal), 1 (mild amount of mucus present), 2 (moderate amount of mucous present), 3 (large amount of mucus present, some blood also present, or diphtheritic material present but not appearing to block the trachea) or 4 (large amount of mucus present, significant blood also presents, or a diphtheritic plug present and blocking the trachea).

2.4 Virus detection and quantification

DNA was extracted from the medium containing the conjunctival and tracheal swabs that were collected during necropsy using the KingFisher Flex Purification System (ThermoFisher Scientific) following the manufacturer's instructions. Positive extraction control samples (diluted stocks of the SA2 ILTV vaccine strain; Zoetis) and negative extraction control samples (distilled water) were included in each extraction plate. Ninety microliter samples of eluted DNA were sealed and stored at -20°C until the extracts were tested for the presence of ILTV DNA by real-time quantitative PCR using oligonucleotide primer pairs that amplify a 113 bp

region of the UL15 gene of ILTV, as described previously [21]. To generate a standard curve, a 10-fold dilution series of the UL15 sequence cloned into pGEM-T (Promega) was prepared using a QIAgility robot (Qiagen) and included in triplicate in each run to enable quantitation of the ILTV genome concentration in each of the extracted samples. Only samples that produced amplicons with a melt curve that matched those of the standard curve samples were regarded as positive for the presence of ILTV DNA. Calculations of the concentration of the ILTV genome in the extracted samples were performed using Rotorgene Q version 2.1.0 (Qiagen) and the concentration in viral genome copy numbers per reaction were log₁₀ transformed for statistical analysis, with the lower limit of detection for the assay defined as 100 genome copies per reaction.

2.5 Statistical analyses

GraphPad Prism 6 (GraphPad Software) and Excel 2016 (Microsoft) were used for data analyses. Mann-Whitney tests were used to compare differences between the groups in the scores for clinical signs and gross tracheal pathology. A one-way analysis of variance, in conjunction with Dunnett's Multiple Comparisons test, was used to compare differences between the groups in viral genome concentrations. Fisher's Exact Test was used to compare differences between the groups in the proportions of birds positive for ILTV by qPCR and in mortality rates.

3. Results

Mortality rates, the severity of virus-induced clinical signs of disease (demeanour, conjunctivitis and dyspnoea), the severity of gross tracheal pathology, and the level of viral replication in tracheal and conjunctival mucosa were assessed after challenge. The vaccinated and control groups were compared in order to identify a suitable minimum effective dose of the vaccine.

3.1 Clinical signs of disease

Scores for demeanour, conjunctivitis and dyspnoea are summarised in Tables 1, 2 and 3, respectively. For all these parameters, the highest scores were seen in the positive control (unvaccinated-challenged) group, and the scores in the negative control (unvaccinated-unchallenged) group were all zero. The differences between the negative control and the positive control group were significant for all parameters. There were no significant differences between the group that received the highest dose of vaccine ($10^{5.0}$ PFU) and the negative control group at any time point for any of these parameters. This was also true for the group that received the second highest dose of vaccine ($10^{3.8}$ PFU), although when the scores were summed to yield cumulative scores, this group had a cumulative conjunctivitis score that was significantly higher than the cumulative score for the negative control group (Table 2). The scores for the groups of birds that received the lower doses of vaccine ($10^{3.2}$ or $10^{3.5}$ PFU) were significantly higher than those for the negative control group for all disease parameters on at least two of the four time points. In these groups, the cumulative scores for all parameters were significantly higher than those of the negative control group (Tables 1–3).

3.2 Viral detection and quantification in tracheal and conjunctival swabs

The results for virus quantification in tracheal and conjunctival swabs after challenge are presented in Fig 1. The groups vaccinated with the two higher doses of the vaccine ($10^{3.8}$ PFU or $10^{5.0}$ PFU) had concentrations of virus at both sites that were significantly lower than those of birds in the positive control group, but that were not significantly different from those of the

Table 1. Scores for demeanour on days 3 to 6 after challenge in unvaccinated birds and in birds vaccinated with different doses of the ΔgG ILTV vaccine administered via eye-drop.

Vaccine dose (PFU)	Challenge ‡	Median demeanour score (range)*				
		3 dpc [^]	4 dpc	5 dpc	6 dpc	Cumulative [†]
None	None	0 (0–0) ^A	0 (0–0) ^A	0 (0–0) ^A	0 (0–0) ^A	0 (0–0) ^A
None	Class 9 [‡]	1 (0–2) ^B	1 (1–2) ^B	1 (1–2) ^B	1 (1–1) ^B	3 (0–5) ^B
ΔgG ILTV (10 ^{5.0})	Class 9	0 (0–0) ^A	0 (0–0) ^A	0 (0–0) ^A	0 (0–0) ^A	0 (0–0) ^A
ΔgG ILTV (10 ^{3.8})	Class 9	0 (0–1) ^A	0 (0–1) ^A	0 (0–1) ^A	0 (0–1) ^A	0 (0–3) ^A
ΔgG ILTV (10 ^{3.5})	Class 9	0 (0–1) ^A	0 (0–2) ^C	1 (0–2) ^C	0 (0–1) ^A	1 (0–5) ^C
ΔgG ILTV (10 ^{3.2})	Class 9	0 (0–1) ^A	0 (0–1) ^{A, C}	1 (0–2) ^C	0 (0–1) ^A	1 (0–4) ^C

*Values labelled with the same uppercase superscript letter in the same column were not significantly different (P > 0.05, Mann-Whitney test)

‡ Class 9 challenge = 10³ PFU of class 9 ILTV delivered via eye-drop and intra-tracheal inoculation

[^] dpc = days post-challenge

[†] Cumulative = sum of scores on all days for each individual bird

<https://doi.org/10.1371/journal.pone.0207611.t001>

negative control (unvaccinated-unchallenged) birds. The groups vaccinated with the two lower doses of vaccine had concentrations of virus at both sites that were significantly higher than those of negative control birds. There was no significant difference in virus concentrations in the trachea of the birds that received the two lower doses of vaccine and that of birds in the positive control group, but the two lower doses of the vaccine were partially protective in the conjunctiva, resulting in significantly lower concentrations of virus at this site than in the positive control birds (Fig 1).

The proportions of birds that were positive after challenge for ILTV by qPCR are shown in Fig 2. A smaller proportion of birds that received the higher doses of the vaccine (10^{3.8} PFU or 10^{5.0} PFU) yielded ILTV positive conjunctival swabs (1/22 and 2/22 birds, respectively). The proportion of birds positive for ILTV DNA in the trachea after challenge was not significantly

Table 2. Scores for conjunctivitis on days 3 to 6 after challenge in unvaccinated birds and in birds vaccinated with different doses of ΔgG ILTV vaccine administered via eye-drop.

Vaccine dose (PFU)	Challenge ‡	Median conjunctivitis score (range)*				
		3 dpc [^]	4 dpc	5 dpc	6 dpc	Cumulative [†]
None	None	0 (0–0) ^A	0 (0–0) ^A	0 (0–0) ^A	0 (0–0) ^A	0 (0–0) ^A
None	Class 9 [‡]	0 (0–1) ^B	1 (0–2) ^B	1 (0–2) ^B	1 (0–1) ^B	1 (0–6) ^B
ΔgG ILTV (10 ^{5.0})	Class 9	0 (0–0) ^A	0 (0–0) ^A	0 (0–0) ^A	0 (0–0) ^A	0 (0–0) ^{A, C}
ΔgG ILTV (10 ^{3.8})	Class 9	0 (0–0) ^A	0 (0–0) ^A	0 (0–2) ^A	0 (0–0) ^A	0 (0–2) ^C
ΔgG ILTV (10 ^{3.5})	Class 9	0 (0–1) ^A	0 (0–1) ^{A, C}	0 (0–2) ^A	0 (0–2) ^C	0 (0–5) ^D
ΔgG ILTV (10 ^{3.2})	Class 9	0 (0–1) ^B	0 (0–2) ^C	0 (0–1) ^A	0 (0–1) ^{B, C}	0 (0–4) ^D

*Values labelled with the same uppercase superscript letter in the same column were not significantly different (P > 0.05, Mann-Whitney test)

‡ Class 9 challenge = 10³ PFU of class 9 ILTV delivered via eye-drop and intra-tracheal inoculation

[^] dpc = days post-challenge

[†] Cumulative = sum of scores on all days for each individual bird

<https://doi.org/10.1371/journal.pone.0207611.t002>

Table 3. Scores for dyspnoea on days 3 to 6 after challenge in unvaccinated birds and in birds vaccinated with different doses of ΔgG ILTV vaccine administered via eye-drop.

Vaccine dose (PFU)	Challenge [‡]	Median dyspnoea score (range)*				
		3 d.p.c [^]	4 d.p.c	5 d.p.c	6 d.p.c	Cumulative [†]
None	None	0 (0–0) ^A	0 (0–0) ^A	0 (0–0) ^A	0 (0–0) ^A	0 (0–0) ^A
None	Class 9 [‡]	1 (0–3) ^B	1 (0–3) ^B	1 (0–2) ^B	0 (0–1) ^B	2 (1–6) ^B
ΔgG ILTV (10 ^{5.0})	Class 9	0 (0–0) ^A	0 (0–0) ^A	0 (0–0) ^A	0(0–0) ^A	0 (0–0) ^A
ΔgG ILTV (10 ^{3.8})	Class 9	0 (0–1) ^A	0 (0–1) ^A	0 (0–2) ^A	0(0–1) ^A	0 (0–4) ^{A, C}
ΔgG ILTV (10 ^{3.5})	Class 9	0 (0–1) ^A	0 (0–2) ^A	0 (0–3) ^B	0(0–1) ^B	1 (0–4) ^C
ΔgG ILTV (10 ^{3.2})	Class 9	0 (0–1) ^B	1 (0–4) ^B	1 (0–2) ^B	0(0–1) ^B	2 (0–5) ^B

*Values labelled with the same uppercase superscript letter in the same column were not significantly different (P > 0.05, Mann Whitney test)

[‡] Class 9 challenge = 10³ PFU of class 9 ILTV delivered via eye-drop and intra-tracheal inoculation

[^] dpc = days post-challenge

[†] Cumulative = sum of scores on all days for each individual bird

<https://doi.org/10.1371/journal.pone.0207611.t003>

different between the negative control group and the groups that received the higher dose of the vaccine (10^{3.8} PFU or 10^{5.0} PFU), indicating a good level of protection in these vaccinated birds. In the group that received the lowest dose of vaccine, the proportion of ILTV positive birds (13/22) did not differ significantly from that of the positive control group (18/22). Fewer birds were positive for ILTV in the trachea compared to the conjunctiva (Fig 2).




Fig 1. Scatter plot of concentrations of ILTV as determined by qPCR on extracts from conjunctival or tracheal swabs of unvaccinated birds and birds vaccinated with different doses of ΔgG ILTV at day 7 after challenge with class 9 ILTV. Values labelled with the same upper-case letter (A, B, C) were not significantly different (P > 0.05, one-way analyses of variance in conjunction with Dunnett’s Multiple Comparisons test) in each panel (conjunctiva or trachea). GCN = genome copy number, UnVx-Uch = unvaccinated and unchallenged, UnVx-Ch = unvaccinated and challenged, PFU = plaque forming units. Means (horizontal solid lines) and standard deviations (error bars) are also shown. The cut off value for positivity was 100 genome copies per reaction and this is indicated by a horizontal dashed line.

<https://doi.org/10.1371/journal.pone.0207611.g001>




Fig 2. Bar graphs showing proportions of ILTV positive birds per group, as determined by qPCR on extracts from conjunctival or tracheal swabs of unvaccinated birds and birds vaccinated with different doses of Δ gG ILTV at day 7 after challenge with class 9 ILTV. Positivity was determined using the cut-off value of 100 genome copies number per reaction. UnVx-Uch = unvaccinated and unchallenged, UnVx-Ch = unvaccinated and challenged, PFU = plaque forming units. Values labelled with the same upper-case letter (A, B, C, D) were not significantly different ($P > 0.05$, Fisher's exact test) in each panel (conjunctiva or trachea).

<https://doi.org/10.1371/journal.pone.0207611.g002>

3.3 Gross tracheal pathology and mortalities

Mortalities and gross tracheal pathology scores after challenge are shown in Table 4. No mortalities were seen in the birds vaccinated with the higher doses of vaccine ($10^{3.8}$ PFU or $10^{5.0}$ PFU). There were four mortalities after challenge in each of the groups of birds that received the lower doses of vaccine ($10^{3.2}$ PFU or $10^{3.5}$ PFU). These mortality rates were not significantly different from that seen in the positive control group (10/22).

None of the birds in the negative control group had any tracheal pathology consistent with infection with ILTV. There was no significant difference in the tracheal pathology scores of the birds in the negative control group and those of the birds in the groups that received the two higher doses of vaccine ($10^{3.8}$ PFU or $10^{5.0}$ PFU). The tracheal pathology scores of the groups of birds that received the lower doses of vaccine ($10^{3.2}$ PFU or $10^{3.5}$ PFU) were significantly higher than the scores of the birds in the negative control group, but significantly lower than the scores of the birds in the positive control group.

4. Discussion

A dose of $10^{3.8}$ PFU was the lowest dose capable of providing a high level of protection against challenge with the class 9 ILTV. In this study, birds vaccinated with a dose of $10^{3.8}$ PFU were protected and did not develop clinical signs or tracheal pathology consistent with ILT. Replication of wild type virus was also greatly reduced in the conjunctival and tracheal mucosa of these birds but was not completely prevented. This is common for ILTV vaccines, which

Table 4. Mortality rates and gross pathological scores in unvaccinated birds and in birds vaccinated with different doses of ΔgG ILTV vaccine via eye-drop at day 7 after challenge.

Vaccine dose (PFU)	Challenge [‡]	N*	No. mortalities ^{†^} (%)	Median pathology score (range)
None	None	10	0 (0) ^{A, C}	0 (0–0) ^A
None	Class 9 [‡]	22	10 (45.5) ^B	1.5 (0–3) ^B
ΔgG ILTV (10 ^{5.0})	Class 9	22	0 (0) ^A	0 (0–1) ^A
ΔgG ILTV (10 ^{3.8})	Class 9	22	0 (0) ^A	0 (0–1) ^A
ΔgG ILTV (10 ^{3.5})	Class 9	22	4 (18.2) ^{B, C}	1 (0–3) ^C
ΔgG ILTV (10 ^{3.2})	Class 9	22	4 (18.2) ^{B, C}	1 (0–3) ^C

[‡] Class 9 challenge = 10³ PFU of class 9 ILTV delivered via eye-drop and intra-tracheal inoculation

*N = number of birds in group

[†] Values labelled with the same uppercase superscript letter in the same column were not significantly different (P > 0.05, Fishers exact test or Mann-Whitney test).

[^] Birds that died or were euthanized due to severe clinical signs of disease following challenge

<https://doi.org/10.1371/journal.pone.0207611.t004>

typically induce non-sterile immunity [12, 15, 17, 22, 23]. Increasing the vaccine dose to 10^{5.0} PFU did not significantly improve the level of protection compared to that induced by the minimum effective dose, whereas lower doses (10^{3.2} PFU or 10^{3.5} PFU) failed to fully protect the birds from clinical disease after challenge. Unprotected or incompletely protected birds may serve as reservoir of infection from which susceptible animals may become infected [5]. Thus, vaccines should not be used at a dose below the minimum effective dose. Using the lowest possible protective dose of vaccine (rather than a higher dose) is important for keeping vaccine production costs as low as possible, and thus the costs of vaccination as low as possible for producers.

Vaccination and challenge experiments are commonly performed to assess the level of protection induced by ILTV vaccines. Such studies are necessary because there are currently no easily measured correlates of protection (such as serum antibody levels) for ILTV infection, as cell mediated immunity [24, 25], rather than neutralising antibodies [26], are thought to protect against disease. Most ILTV vaccination and challenge studies assess levels of clinical protection by scoring clinical signs, tracheal pathology, and mortalities [2, 3, 12, 27]. Many also assess weight gain [15, 17, 27]. In this study, weight gain was not measured, as layer type birds do not typically have significant weight gains in short periods of time. Importantly, our study also assessed virus replication after challenge by measuring the viral genome concentration, as have some other vaccine efficacy studies [7, 12, 15, 17]. Vaccines that prevent clinical signs of disease but do not prevent replication of pathogens after challenge have been called ‘leaky’ or ‘imperfect’ vaccines and this is a common feature of ILTV vaccines [27–29]. Such vaccines may select for more virulent pathogens over time [30, 31] as has recently been demonstrated experimentally for Marek’s disease vaccines [30]. In our study, the concentrations of virus genomes in the conjunctival and tracheal mucosa were reduced to very low levels in birds that received high doses of the vaccine (10^{3.8} PFU or 10^{5.0} PFU) and only a small number of birds had detectable concentrations of ILTV DNA in the trachea or conjunctiva. These results suggest this vaccine induces significant protection against viral replication, although it should be noted that viral detection and quantification was assessed at one time point. The inclusion of additional time points would further clarify this observation, particularly earlier time points when viral titres are likely to be higher.

The results from this study demonstrate that Vaxsafe ILT induces a high level of protection when administered by eye-drop at a dose of $10^{3.8}$ PFU. This adds to the growing number of studies that have demonstrated the high level of safety and efficacy of the Δ gG ILTV strain when administered via different routes, using different challenge models [12, 13, 15, 16]. Glycoprotein G (gG) is a viral chemokine binding protein. Deletion of gG from the ILTV genome alters the immune response to ILTV infection [32], resulting in a more protective, cell-mediated immune response [20]. This vaccine also offers the potential to discriminate between vaccinated birds and birds infected with wildtype viruses using differential PCR [33] or ELISAs [19] in DIVA (differentiation of infected and vaccinated animals) control programmes. Although a number of gene-deleted ILTV strains have been shown to have *in vivo* phenotypes that may make them suitable for use as vaccines [12–16, 34–41], the Δ gG ILTV vaccine strain is the most extensively investigated of these deletion mutants, providing a high level of confidence in the performance of the vaccine under different conditions. This current study shows that the Δ gG ILTV vaccine strain is able to induce a high level of protection against a virulent recombinant field virus at a commercially feasible dose. These results lay the foundations for the application of a commercial vaccine product, offering poultry producers a new tool to help control ILTV.

Acknowledgments

Mesula G. Korska was supported by Australian Government Research Training Program (RTP) scholarship. This work was partially funded by the Australian Poultry Cooperative Research Centre. We thank Dr. Pollob Shil, June Daly and Janece Wheeler for providing care for the animals. We thank Bioproperties Pty Ltd, and particularly Dr Greg Underwood, for support, advice and valuable discussions.

Author Contributions

Conceptualization: Joanne M. Devlin, Carol A. Hartley, Glenn F. Browning, Andrés Diaz-Méndez.

Data curation: Mesula G. Korska, Andrés Diaz-Méndez.

Formal analysis: Mesula G. Korska, Mauricio J. C. Coppo.

Funding acquisition: Glenn F. Browning.

Investigation: Mesula G. Korska, Joanne M. Devlin, Carol A. Hartley, Mauricio J. C. Coppo, José A. Quinteros, Carlos A. Loncoman, Adepeju E. Onasanya, Dulari Thilakarathne, Andrés Diaz-Méndez.

Methodology: Mesula G. Korska, Joanne M. Devlin, Carol A. Hartley, Mauricio J. C. Coppo, José A. Quinteros, Carlos A. Loncoman, Adepeju E. Onasanya, Dulari Thilakarathne, Andrés Diaz-Méndez.

Project administration: Joanne M. Devlin, Carol A. Hartley, Glenn F. Browning, Andrés Diaz-Méndez.

Resources: Joanne M. Devlin, Glenn F. Browning, Andrés Diaz-Méndez.

Supervision: Joanne M. Devlin, Carol A. Hartley, Glenn F. Browning, Andrés Diaz-Méndez.

Writing – original draft: Mesula G. Korska, Joanne M. Devlin.

Writing – review & editing: Mesula G. Korsa, Joanne M. Devlin, Carol A. Hartley, Glenn F. Browning, Mauricio J. C. Coppo, José A. Quinteros, Carlos A. Loncoman, Adepeju E. Onasanya, Dulari Thilakarathne, Andrés Diaz-Méndez.

References

- Garcia M, Spatz S, Guy J. Infectious laryngotracheitis. In: Swayne DE, Glisson JR, McDougald LM, Nolan LK, Suarez DL, and Nair V, editors. *Diseases of Poultry*. 13th ed, Ames IA: Blackwell Publishing; 2013, p 161–79
- Coppo MJ, Noormohammadi AH, Browning GF, Devlin JM. Challenges and recent advancements in infectious laryngotracheitis virus vaccines. *Avian Pathol*. 2013; 42: 195–205. <https://doi.org/10.1080/03079457.2013.800634> PMID: 23718807
- García M. Current and future vaccines and vaccination strategies against infectious laryngotracheitis (ILT) respiratory disease of poultry. *Vet Microbiol*. 2016.
- Lee SW, Markham PF, Coppo MJ, Legione AR, Markham JF, Noormohammadi AH, et al. Attenuated vaccines can recombine to form virulent field viruses. *Science (New York, NY)*. 2012; 337: 188. <https://doi.org/10.1126/science.1217134> PMID: 22798607
- Agnew-Crumpton R, Vaz PK, Devlin JM, O'Rourke D, Blacker-Smith HP, Konsak-Ilievski B, et al. Spread of the newly emerging infectious laryngotracheitis viruses in Australia. *Infect, Genet and Evol*. 2016; 43: 67–73.
- Davison S, Gingerich EN, Casavant S, Eckroade RJ. Evaluation of the efficacy of a live fowlpox-vectored infectious laryngotracheitis/avian encephalomyelitis vaccine against ILT viral challenge. *Avian Dis*. 2006; 50: 50–4. <https://doi.org/10.1637/7398-062105R.1> PMID: 16617981
- Johnson DI, Vagnozzi A, Dorea F, Riblet SM, Mundt A, Zavala G, et al. Protection against infectious laryngotracheitis by in ovo vaccination with commercially available viral vector recombinant vaccines. *Avian Dis*. 2010; 54: 1251–9. <https://doi.org/10.1637/9401-052310-Reg.1> PMID: 21313847
- Vagnozzi A, Zavala G, Riblet SM, Mundt A, Garcia M. Protection induced by commercially available live-attenuated and recombinant viral vector vaccines against infectious laryngotracheitis virus in broiler chickens. *Avian Pathol*. 2012; 41: 21–31. <https://doi.org/10.1080/03079457.2011.631983> PMID: 22845318
- Godoy A, Icard A, Martinez M, Mashchenko A, Garcia M, El-Attrachea J. Detection of infectious laryngotracheitis virus antibodies by glycoprotein-specific ELISAs in chickens vaccinated with viral vector vaccines. *Avian Dis*. 2013; 57: 432–6. <https://doi.org/10.1637/10345-090312-Reg.1> PMID: 23901757
- Gimeno IM, Cortes AL, Guy JS, Turpin E, Williams C. Replication of recombinant herpesvirus of turkey expressing genes of infectious laryngotracheitis virus in specific pathogen free and broiler chickens following in ovo and subcutaneous vaccination. *Avian Pathol*. 2011; 40: 395–403. <https://doi.org/10.1080/03079457.2011.588196> PMID: 21812719
- Esaki M, Noland L, Eddins T, Godoy A, Saeki S, Saitoh S, et al. Safety and efficacy of a turkey herpesvirus vector laryngotracheitis vaccine for chickens. *Avian Dis*. 2013; 57: 192–8. <https://doi.org/10.1637/10383-092412-Reg.1> PMID: 24689173
- Devlin JM, Browning GF, Gilkerson JR, Fenton SP, Hartley CA. Comparison of the safety and protective efficacy of vaccination with glycoprotein-G-deficient infectious laryngotracheitis virus delivered via eye-drop, drinking water or aerosol. *Avian Pathol*. 2008; 37: 83–8. <https://doi.org/10.1080/03079450701802214> PMID: 18202954
- Devlin JM, Browning GF, Hartley CA, Gilkerson JR. Glycoprotein G deficient infectious laryngotracheitis virus is a candidate attenuated vaccine. *Vaccine*. 2007; 25: 3561–6. <https://doi.org/10.1016/j.vaccine.2007.01.080> PMID: 17316926
- Devlin J, Browning G, Hartley C, Kirkpatrick N, Mahmoudian A, Noormohammadi A, et al. Glycoprotein G is a virulence factor in infectious laryngotracheitis virus. *J Gen Virol*. 2006; 87: 2839–47. <https://doi.org/10.1099/vir.0.82194-0> PMID: 16963741
- Coppo MJ, Noormohammadi AH, Hartley CA, Gilkerson JR, Browning GF, Devlin JM. Comparative in vivo safety and efficacy of a glycoprotein G-deficient candidate vaccine strain of infectious laryngotracheitis virus delivered via eye drop. *Avian Pathol*. 2011; 40: 411–7. <https://doi.org/10.1080/03079457.2011.588686> PMID: 21812721
- Legione AR, Coppo MJ, Lee SW, Noormohammadi AH, Hartley CA, Browning GF, et al. Safety and vaccine efficacy of a glycoprotein G deficient strain of infectious laryngotracheitis virus delivered in ovo. *Vaccine*. 2012; 30: 7193–8. <https://doi.org/10.1016/j.vaccine.2012.10.023> PMID: 23084851
- Korsa MG, Browning GF, Coppo MJ, Legione AR, Gilkerson JR, Noormohammadi AH, et al. Protection induced in broiler chickens following drinking-water delivery of live infectious laryngotracheitis vaccines

- against subsequent challenge with recombinant field virus. *PLoS One*. 2015; 10: e0137719. <https://doi.org/10.1371/journal.pone.0137719> PMID: 26366738
18. Devlin JM, Hartley CA, Gilkerson JR, Coppo MJC, Vaz P, Noormohammadi AH, et al. Horizontal transmission dynamics of a glycoprotein G deficient candidate vaccine strain of infectious laryngotracheitis virus and the effect of vaccination on transmission of virulent virus. *Vaccine*. 2011; 29: 5699–704. <https://doi.org/10.1016/j.vaccine.2011.06.002> PMID: 21689710
 19. Shil NK, Markham PF, Noormohammadi AH, O'Rourke D, Devlin JM. Development of an enzyme-linked immunosorbent assay to detect chicken serum antibody to glycoprotein G of infectious laryngotracheitis virus. *Avian Dis*. 2012; 56: 509–15. <https://doi.org/10.1637/10054-010912-Reg.1> PMID: 23050467
 20. Coppo MJ, Devlin JM, Legione AR, Vaz PK, Lee S-W, Quinteros JA, et al. Infectious laryngotracheitis virus viral chemokine-binding protein glycoprotein G alters transcription of key inflammatory mediators in vitro and in vivo. *J Virol*. 2018; 92: e01534–17. <https://doi.org/10.1128/JVI.01534-17> PMID: 29070686
 21. Mahmoudian A, Kirkpatrick NC, Coppo M, Lee S-W, Devlin JM, Markham PF, et al. Development of a SYBR green quantitative polymerase chain reaction assay for rapid detection and quantification of infectious laryngotracheitis virus. *Avian Pathol*. 2011; 40: 237–42. <https://doi.org/10.1080/03079457.2011.553582> PMID: 21711182
 22. Guy JS, Barnes HJ, Morgan LM. Virulence of infectious laryngotracheitis viruses: comparison of modified-live vaccine viruses and North Carolina field isolates. *Avian Dis*. 1990; 106–13. PMID: 2157386
 23. Bagust TJ, Johnson MA. Avian infectious laryngotracheitis: virus-host interactions in relation to prospects for eradication. *Avian Pathol*. 1995; 24: 373–91. <https://doi.org/10.1080/03079459508419079> PMID: 18645796
 24. Honda T, Okamura H, Taneno A, Yamada S, Takahashi E. The role of cell-mediated immunity in chickens inoculated with the cell-associated vaccine of attenuated infectious laryngotracheitis virus. *J Vet Med Sci*. 1994; 56: 1051–5. PMID: 7696391
 25. Fahey KJ, York JJ, Bagust TJ. Laryngotracheitis herpesvirus infection in the chicken. II. The adoptive transfer of resistance with immune spleen cells. *Avian Pathol*. 1984; 13: 265–75. <https://doi.org/10.1080/03079458408418530> PMID: 18766843
 26. Fahey KJ, York JJ. The role of mucosal antibody in immunity to infectious laryngotracheitis virus in chickens. *J Gen Virol*. 1990; 71: 2401–5. <https://doi.org/10.1099/0022-1317-71-10-2401> PMID: 2172453
 27. Rodríguez-Avila A, Oldoni I, Riblet S, García M. Evaluation of the protection elicited by direct and indirect exposure to live attenuated infectious laryngotracheitis virus vaccines against a recent challenge strain from the United States. *Avian Pathol*. 2008; 37:287–92. <https://doi.org/10.1080/03079450802043742> PMID: 18568655
 28. Rodríguez-Avila A, Oldoni I, Riblet S, García M. Replication and transmission of live attenuated infectious laryngotracheitis virus (ILTV) vaccines. *Avian Dis*. 2007; 51:905–11. <https://doi.org/10.1637/8011-041907-REGR.1> PMID: 18251401
 29. Guy JS, Barnes HJ, Smith L. Increased virulence of modified-live infectious laryngotracheitis vaccine virus following bird-to-bird passage. *Avian Dis*. 1991:348–55. PMID: 1649591
 30. Read AF, Baigent SJ, Powers C, Kgosana LB, Blackwell L, Smith LP, et al. Imperfect vaccination can enhance the transmission of highly virulent pathogens. *PLoS Biol*. 2015; 13:e1002198. <https://doi.org/10.1371/journal.pbio.1002198> PMID: 26214839
 31. Devlin JM, Vaz PK, Coppo MJ, Browning GF. Impacts of poultry vaccination on viruses of wild bird. *Curr Opin Virol*. 2016; 19:23–9. <https://doi.org/10.1016/j.coviro.2016.06.007> PMID: 27359320
 32. Devlin JM, Viejo-Borbolla A, Browning GF, Noormohammadi AH, Gilkerson JR, Alcamí A, et al. Evaluation of immunological responses to a glycoprotein G deficient candidate vaccine strain of infectious laryngotracheitis virus. *Vaccine*. 2010; 28:1325–32. <https://doi.org/10.1016/j.vaccine.2009.11.013> PMID: 19932672
 33. Shil NK, Legione AR, Markham PF, Noormohammadi AH, Devlin JM. Development and validation of TaqMan real-time polymerase chain reaction assays for the quantitative and differential detection of wild-type infectious laryngotracheitis viruses from a glycoprotein G-deficient candidate vaccine strain. *Avian Dis*. 2015; 59: 7–13 PMID: 26292527
 34. García M, Spatz S, Cheng Y, Riblet S, Volkening J, Schneiders G. Attenuation and protection efficacy of open reading frame C (ORF C) gene deleted recombinant of infectious laryngotracheitis virus (ILTV). *J Gen Virol*. 2016.
 35. Veits J, Lüschoff D, Kindermann K, Werner O, Teifke JP, Mettenleiter TC, et al. Deletion of the non-essential UL0 gene of infectious laryngotracheitis (ILT) virus leads to attenuation in chickens, and UL0

- mutants expressing influenza virus haemagglutinin (H7) protect against ILT and fowl plague. *J Gen Virol.* 2003; 84: 3343–52. <https://doi.org/10.1099/vir.0.19570-0> PMID: 14645915
36. Fuchs W, Mettenleiter TC. The nonessential UL49. 5 gene of infectious laryngotracheitis virus encodes an O-glycosylated protein which forms a complex with the non-glycosylated UL10 gene product. *Virus Res.* 2005; 112: 108–14. <https://doi.org/10.1016/j.virusres.2005.03.002> PMID: 16022905
 37. Helferich D, Veits J, Teifke JP, Mettenleiter TC, Fuchs W. The UL47 gene of avian infectious laryngotracheitis virus is not essential for in vitro replication but is relevant for virulence in chickens. *J Gen Virol.* 2007; 88: 732–42. <https://doi.org/10.1099/vir.0.82533-0> PMID: 17325345
 38. Pavlova SP, Veits J, Blohm U, Maresch C, Mettenleiter TC, Fuchs W. In vitro and in vivo characterization of glycoprotein C-deleted infectious laryngotracheitis virus. *J Gen Virol.* 2010; 91: 847–57. <https://doi.org/10.1099/vir.0.016634-0> PMID: 19940061
 39. Mashchenko A, Riblet SM, Zavala G, Garcia M. In ovo vaccination of commercial broilers with a glycoprotein J gene-deleted strain of infectious laryngotracheitis virus. *Avian Dis.* 2013; 57: 523–31. <https://doi.org/10.1637/10413-100512-Reg.1> PMID: 23901771
 40. Schnitzlein WM, Winans R, Ellsworth S, Tripathy DN. Generation of thymidine kinase-deficient mutants of infectious laryngotracheitis virus. *Virology.* 1995; 209: 304–14. <https://doi.org/10.1006/viro.1995.1262> PMID: 7778265
 41. Han M, Kweon C, Mo I, Kim S. Pathogenicity and vaccine efficacy of a thymidine kinase gene deleted infectious laryngotracheitis virus expressing the green fluorescent protein gene. *Arch Virol* 2002; 147: 1017–31. <https://doi.org/10.1007/s00705-001-0794-y> PMID: 12021870