

Editorial

Prevention of alcohol-related health harm in the workplace

Prevention of alcohol-related health problems is one of the most central issue worldwide. In Global Burden of Disease (GBD) Study 2010, alcohol use was the third and the eighth leading risk factor for the disability-adjusted life years (DALYs) in men and women, respectively¹. World Health Organization has adopted the global strategy to reduce the harmful use of alcohol in 2010²) to reduce serious harmful health problems caused by alcohol abuse. The global action plan 2013–2020 has set the goal of at least 10% relative reduction in the harmful use of alcohol³).

In Japanese men, the proportion of the habitual alcohol drinkers who drink more than or equal to three days per week and drink one unit of alcohol (about 23g) each day they drink were more than 50% in 1995, while that decreased to about 35% in 2010⁴). During the same period from 1995 to 2010, the proportion of habitual drinkers were around 7–8% and no significant changes have been observed in Japanese women. The proportion of those with alcohol dependence diagnosed by ICD-10 or with alcohol use disorders assessed by AUDIT has not changed during the period between 2003 and 2013 in both sexes in the periodical nationwide survey⁵). Thus, harmful use of alcohol remains to be one of serious problems in Japan.

East Asian people have two important genetic polymorphisms in alcohol-metabolizing enzymes; *ADH1B* and *ALDH2*⁶). These two polymorphisms result in considerable differences in alcohol sensitivity and alcohol drinking behavior⁶). Those with high alcohol sensitivity have increased risk for esophageal cancer when they habitually drink heavily⁷). Asthmatic patients with high alcohol sensitivity has been associated with experience of alcohol-induced asthma⁸). On the other hand, those with low alcohol sensitivity have increased risk for alcoholism⁹).

Job stressors have been associated with heavy drinking in male Japanese workers^{10, 11}). A stressful psychosocial work environment has been found to be a risk factor for alcohol dependence in workers¹²). Long-term heavy exposure to ethanol would lead to many harmful consequences including alcoholism, hypertension, chronic liver diseases, pancreatitis, and alcohol-related cancers. Thus, it is of

importance for occupational physicians and health staffs to find not only alcohol-related health problems but also presence of job stressors which might be related to alcohol abuse.

Previously we reported that both daily hassles and the genetic polymorphism in the *ALDH2* gene were associated with problem drinking in male Japanese workers¹³). In east Asia, if public genetic testing service for the *ADH1B* and *ALDH2* gene is provided with enough ethical consideration, it would be of help to promote personalized prevention of harmful alcohol-related diseases, especially for workers who tend to drink heavily and have job stressors.

In Japan, Basic Act on Measures against Alcohol-related Health Harm was put into force in 2014¹⁴). Under this law, the national and local governments are going to make the basic plan for the promotion of measures against alcohol-related health harm. The act referred to responsibilities of the National and Local Government, business operators, citizens, physicians, and health promotion service providers.

There are many areas where occupational physicians and/or occupational health staffs can contribute including education and promotion of learning and public relation activities concerning alcohol related-problems, discovery of alcohol-related health harm, and health guidance and/or brief intervention¹⁵) for preventing occurrence, progression, and relapsing of alcohol-related health harm. They can also provide appropriate advice for workers who have driven under the influence of alcohol. The most important thing is that they can play important roles in rehabilitation of those with alcoholism into the worksite by understanding and supporting them.

Accumulating evidence shows that occasional or light/moderate drinking might be beneficial to health^{16–18}). On the other hand, even light/moderate drinking would increase risk for alcohol-related cancers^{19, 20}). Thus, more evidence is needed to determine truly beneficial amounts of alcohol consumption.

Occupational studies and practice aiming at promotion of healthy alcohol drinking and prevention of various alcohol-related health problems described above should be further encouraged worldwide in the coming decades.

References

- 1) Lim SS, Vos T, Flaxman AD, Danaei G, Shibuya K, Adair-Rohani H, Amann M, Anderson HR, Andrews KG, Aryee M, Atkinson C, Bacchus LJ, Bahalim AN, Balakrishnan K, Balmes J, Barker-Collo S, Baxter A, Bell ML, Blore JD, Blyth F, Bonner C, Borges G, Bourne R, Boussinesq M, Brauer M, Brooks P, Bruce NG, Brunekreef B, Bryan-Hancock C, Bucello C, Buchbinder R, Bull F, Burnett RT, Byers TE, Calabria B, Carapetis J, Carnahan E, Chafe Z, Charlson F, Chen H, Chen JS, Cheng AT, Child JC, Cohen A, Colson KE, Cowie BC, Darby S, Darling S, Davis A, Degenhardt L, Dentener F, Des Jarlais DC, Devries K, Dherani M, Ding EL, Dorsey ER, Driscoll T, Edmond K, Ali SE, Engell RE, Erwin PJ, Fahimi S, Falder G, Farzadfar F, Ferrari A, Finucane MM, Flaxman S, Fowkes FG, Freedman G, Freeman MK, Gakidou E, Ghosh S, Giovannucci E, Gmel G, Graham K, Grainger R, Grant B, Gunnell D, Gutierrez HR, Hall W, Hoek HW, Hogan A, Hosgood HD 3rd, Hoy D, Hu H, Hubbell BJ, Hutchings SJ, Ibeanusi SE, Jacklyn GL, Jasrasaria R, Jonas JB, Kan H, Kanis JA, Kassebaum N, Kawakami N, Khang YH, Khatibzadeh S, Khoo JP, Kok C, Laden F, Lalloo R, Lan Q, Lathlean T, Leasher JL, Leigh J, Li Y, Lin JK, Lipshultz SE, London S, Lozano R, Lu Y, Mak J, Malekzadeh R, Mallinger L, Marcenes W, March L, Marks R, Martin R, McGale P, McGrath J, Mehta S, Mensah GA, Merriman TR, Micha R, Michaud C, Mishra V, Mohd Hanafiah K, Mokdad AA, Morawska L, Mozaffarian D, Murphy T, Naghavi M, Neal B, Nelson PK, Nolla JM, Norman R, Olives C, Omer SB, Orchard J, Osborne R, Ostro B, Page A, Pandey KD, Parry CD, Passmore E, Patra J, Pearce N, Pelizzari PM, Petzold M, Phillips MR, Pope D, Pope CA 3rd, Powles J, Rao M, Razavi H, Rehfues EA, Rehm JT, Ritz B, Rivara FP, Roberts T, Robinson C, Rodriguez-Portales JA, Romieu I, Room R, Rosenfeld LC, Roy A, Rushton L, Salomon JA, Sampson U, Sanchez-Riera L, Sanman E, Sapkota A, Seedat S, Shi P, Shield K, Shivakoti R, Singh GM, Sleet DA, Smith E, Smith KR, Stapelberg NJ, Steenland K, Stöckl H, Stovner LJ, Straif K, Straney L, Thurston GD, Tran JH, Van Dingenen R, van Donkelaar A, Veerman JL, Vijayakumar L, Weintraub R, Weissman MM, White RA, Whiteford H, Wiersma ST, Wilkinson JD, Williams HC, Williams W, Wilson N, Woolf AD, Yip P, Zielinski JM, Lopez AD, Murray CJ, Ezzati M, AlMazroa MA, Memish ZA (2012) A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* **380**, 2224–60. [[Medline](#)] [[CrossRef](#)]
- 2) World Health Organization. Global strategy to reduce the harmful use of alcohol. http://www.who.int/substance_abuse/publications/global_strategy_reduce_harmful_use_alcohol/en/. Accessed August 3, 2017.
- 3) World Health Organization. Global Action Plan for the prevention and control of noncommunicable diseases 2013–2020. http://apps.who.int/iris/bitstream/10665/94384/1/9789241506236_eng.pdf#search=%27global+action+plan%27. Accessed August 3, 2017.
- 4) Ministry of Health, Labor and Welfare. The National Health and Nutrition Survey Japan. http://www.e-stat.go.jp/SG1/estat/GL08020101.do?_toGL08020101_&statCode=000001041744&requestSender=dsearch and http://www.nibiohn.go.jp/eiken/chosa/kokumin_eiyou/. Accessed August 3, 2017.
- 5) Osaki Y, Kinjo A, Higuchi S, Matsumoto H, Yuzuriha T, Horie Y, Kimura M, Kanda H, Yoshimoto H (2016) Prevalence and trends in alcohol dependence and alcohol use disorders in Japanese adults; Results from periodical nationwide surveys. *Alcohol Alcohol* **51**, 465–73. [[Medline](#)] [[CrossRef](#)]
- 6) Takeshita T, Morimoto K (1996) Effects of genetic polymorphisms in alcohol-metabolizing enzymes on alcohol hypersensitivity and alcohol-related health problems in orientals. *Environ Health Prev Med* **1**, 1–8. [[Medline](#)] [[CrossRef](#)]
- 7) Yang SJ, Yokoyama A, Yokoyama T, Huang YC, Wu SY, Shao Y, Niu J, Wang J, Liu Y, Zhou XQ, Yang CX (2010) Relationship between genetic polymorphisms of ALDH2 and ADH1B and esophageal cancer risk: a meta-analysis. *World J Gastroenterol* **16**, 4210–20. [[Medline](#)] [[CrossRef](#)]
- 8) Matsuse H, Shimoda T, Fukushima C, Mitsuta K, Kawano T, Tomari S, Saeki S, Kondoh Y, Machida I, Obase Y, Asai S, Kohno S (2001) Screening for acetaldehyde dehydrogenase 2 genotype in alcohol-induced asthma by using the ethanol patch test. *J Allergy Clin Immunol* **108**, 715–9. [[Medline](#)] [[CrossRef](#)]
- 9) Higuchi S, Matsushita S, Murayama M, Takagi S, Hayashida M (1995) Alcohol and aldehyde dehydrogenase polymorphisms and the risk for alcoholism. *Am J Psychiatry* **152**, 1219–21. [[Medline](#)] [[CrossRef](#)]
- 10) Hiro H, Kawakami N, Tanaka K, Nakamura K; Japan Work Stress and Health Cohort Study Group (2007) Association between job stressors and heavy drinking: age differences in male Japanese workers. *Ind Health* **45**, 415–25. [[Medline](#)] [[CrossRef](#)]
- 11) Morikawa Y, Nakamura K, Sakurai M, Nagasawa SY, Ishizaki M, Nakashima M, Kido T, Naruse Y, Nakagawa H (2014) The effect of age on the relationships between work-related factors and heavy drinking. *J Occup Health* **56**, 141–9. [[Medline](#)] [[CrossRef](#)]
- 12) Head J, Stansfeld SA, Siegrist J (2004) The psychosocial work environment and alcohol dependence: a prospective study. *Occup Environ Med* **61**, 219–24. [[Medline](#)] [[CrossRef](#)]
- 13) Takeshita T, Maruyama S, Morimoto K (1998) Relevance of both daily hassles and the ALDH2 genotype to problem drinking among Japanese male workers. *Alcohol Clin Exp Res* **22**, 115–20. [[Medline](#)] [[CrossRef](#)]
- 14) Measures against Alcohol-related Health Harm. Cabinet

- Office, Government of Japan. (in Japanese). <http://www8.cao.go.jp/alcohol/>. Accessed August 3, 2017.
- 15) Fleming MF, Mundt MP, French MT, Manwell LB, Stauffacher EA, Barry KL (2002) Brief physician advice for problem drinkers: long-term efficacy and benefit-cost analysis. *Alcohol Clin Exp Res* **26**, 36–43. [[Medline](#)] [[CrossRef](#)]
 - 16) Di Castelnuovo A, Costanzo S, Bagnardi V, Donati MB, Iacoviello L, de Gaetano G (2006) Alcohol dosing and total mortality in men and women: an updated meta-analysis of 34 prospective studies. *Arch Intern Med* **166**, 2437–45. [[Medline](#)] [[CrossRef](#)]
 - 17) Inoue M, Nagata C, Tsuji I, Sugawara Y, Wakai K, Tamakoshi A, Matsuo K, Mizoue T, Tanaka K, Sasazuki S, Tsugane S; Research Group for the Development and Evaluation of Cancer Prevention Strategies in Japan (2012) Impact of alcohol intake on total mortality and mortality from major causes in Japan: a pooled analysis of six large-scale cohort studies. *J Epidemiol Community Health* **66**, 448–56. [[Medline](#)] [[CrossRef](#)]
 - 18) Roerecke M, Rehm J (2012) The cardioprotective association of average alcohol consumption and ischaemic heart disease: a systematic review and meta-analysis. *Addiction* **107**, 1246–60. [[Medline](#)] [[CrossRef](#)]
 - 19) Bagnardi V, Blangiardo M, La Vecchia C, Corrao G (2001) A meta-analysis of alcohol drinking and cancer risk. *Br J Cancer* **85**, 1700–5. [[Medline](#)] [[CrossRef](#)]
 - 20) Bagnardi V, Rota M, Botteri E, Tramacere I, Islami F, Fedirko V, Scotti L, Jenab M, Turati F, Pasquali E, Pelucchi C, Bellocco R, Negri E, Corrao G, Rehm J, Boffetta P, La Vecchia C (2013) Light alcohol drinking and cancer: a meta-analysis. *Ann Oncol* **24**, 301–8. [[Medline](#)] [[CrossRef](#)]

Tatsuya TAKESHITA

Department of Public Health, School of Medicine,
Wakayama Medical University, Japan