

Identification of genes and pathways associated with pancreatic ductal adenocarcinoma by bioinformatics analyses

JIN LONG, ZHONGBO ZHANG, ZHE LIU, YUANHONG XU and CHUNLIN GE

Department of General Surgery, The First Hospital of China Medical University, Shenyang, Liaoning 110001, P.R. China

Received December 11, 2014; Accepted November 12, 2015

DOI: 10.3892/ol.2015.4042

Abstract. This study aimed to explore the underlying genes and pathways associated with pancreatic ductal adenocarcinoma (PDAC) by bioinformatics analyses. Gene expression profile GSE43795 was downloaded from the Gene Expression Omnibus database. The differentially expressed genes (DEGs) between six PDAC and five non-neoplastic pancreatic tissue samples were analyzed using the limma package. Gene ontology (GO) and pathway enrichment analyses of DEGs were performed, followed by functional annotation and protein-protein interaction (PPI) network construction. Finally, the sub-network was identified and pathway enrichment analysis was performed on the contained DEGs. A total of 374 downregulated and 559 upregulated DEGs were identified. The downregulated DEGs were enriched in GO terms associated with digestion and transport and pathways related to metabolism, while the upregulated DEGs were enriched in GO terms associated with the cell cycle and mitosis and pathways associated with the occurrence of cancer including the cell cycle pathway. Following functional annotation, the oncogene pituitary tumor-transforming 1 (*PTTG1*) was upregulated. In the PPI network and sub-network, cell division cycle 20 (*CDC20*) and BUB1 mitotic checkpoint serine/threonine kinase B (*BUB1B*) were hub genes with high connectivity degrees. Additionally, DEGs in the sub-network including cyclin B1 (*CCNB1*) were mainly enriched in the cell cycle and p53 signaling pathways. In conclusion, the cell cycle and p53 signaling pathways may play significant roles in PDAC, and DEGs including *CDC20*, *BUB1B*, *CCNB1* and *PTTG1* may be potential targets for PDAC diagnosis and treatment.

Introduction

Pancreatic ductal adenocarcinoma (PDAC) ranks among the most malignant of human cancers (1). Currently, the annual number of associated mortalities is similar to the disease's annual incidence (2). The poor prognosis of PDAC is correlated with the nonspecificity of symptoms, advanced disease at presentation and lack of effective adjuvant and systemic therapy (3). Currently, surgical extirpation for localized disease offers the only chance of long-term survival (3). Therefore, understanding the pathological mechanisms to detect PDAC as early as possible is an urgent requirement to enable further advances in therapeutic modalities and agents.

Previous studies have identified that the development of PDAC may involve certain genetic factors including the overexpression of oncogenes, inactivation of tumor suppressor genes or the deregulation of various signaling pathways (4). For instance, kirsten rat sarcoma viral oncogene homolog mutations have been observed to occur with increasing frequency in progressively later stages of pancreatic adenocarcinoma (5). Iacobuzio-Donahue *et al* demonstrated that tumor antigen p97, cathepsin L2 and kallikrein 10 were differentially expressed among PDACs (6). Additionally, the phosphoinositide 3-kinase signaling pathway is known to be activated in pancreatic cancer, which is due to the aberrant expression of phosphatase and tensin homolog. Progress achieved in understanding the mechanism of PDAC is likely to contribute to the treatment of this disease. However, no breakthrough treatments have been identified, so the present knowledge would appear to be insufficient.

In the present study, we downloaded microarray data of GSE43795 and identified the differentially expressed genes (DEGs) between PDAC and non-neoplastic pancreatic tissue (NN) samples to explore the molecular mechanisms of PDAC. Park *et al* (7) used the dataset GSE43795 to study the characterization of gene expression and activated signaling pathways in solid pseudopapillary neoplasms of the pancreas. However, the functional annotation and protein-protein interaction (PPI) of DEGs are still far from being clear. In the present study, we performed functional enrichment analyses and functional annotation. Finally, PPI networks and sub-networks were constructed and analyzed to study and identify target genes for the diagnosis and treatment of PDAC. We aimed to explore the underlying genes and pathways associated with PDAC. The findings from this study are likely to play a significant

Correspondence to: Dr Jin Long, Department of General Surgery, The First Hospital of China Medical University, 155 Nanjing Bei Street, Shenyang, Liaoning 110001, P.R. China
E-mail: jinlongjld@163.com

Key words: pancreatic ductal adenocarcinoma, differentially expressed genes, pathway enrichment analysis, protein-protein interaction network

role in PDAC genesis and may potentially serve as biomarkers in the diagnosis and treatment of PDAC.

Materials and methods

Affymetrix microarray data. The microarray data of GSE43795 were downloaded from the Gene Expression Omnibus (<http://www.ncbi.nlm.nih.gov/geo/>) database based on the platform of GPL10558 Illumina HumanHT-12 V4.0 expression beadchip. A total of six PDAC and five NN samples were used in this study to develop the Affymetrix microarray data (Affymetrix, Inc., Santa Clara, CA, USA).

Data pre-processing and differential expression analysis. Background correction, quartile data normalization and probe summarization were performed for the original array data, then they were converted into expression measures by the robust multiarray average (8) algorithm in the R affy package (9) (<http://www.bioconductor.org>).

For the GSE43795 dataset, the limma eBayes (10) method in Bioconductor (<http://www.bioconductor.org>) was used to identify genes which were differentially expressed between PDAC and NN samples. The \log_2 -fold change (\log_2FC) was calculated. $|\log_2FC| \geq 3$ and false discovery rate (FDR) < 0.01 were considered as the cutoff values for DEG screening.

Gene ontology and pathway enrichment analyses. Gene ontology (GO) (11) is a tool used for collecting a large number of gene annotation terms. The Kyoto Encyclopedia of Genes and Genomes (KEGG) knowledge database (12) is a collection of online databases dealing with genomes, enzymatic pathways and biological chemicals. The Database for Annotation, Visualization and Integrated Discovery (DAVID) (13), as a comprehensive set of functional annotation tools, has been developed for relating the functional terms with gene lists using a clustering algorithm. In order to analyze the DEGs at the functional level, we performed GO and KEGG pathway enrichment analyses using the DAVID online tool to obtain the enriched biological processes (BPs) and pathways. $P < 0.01$ was set as the threshold value.

Functional annotation of DEGs. Based on the data information of transcription factors (TFs), the DEGs were selected and annotated to determine whether these genes had the function of transcriptional regulation. The tumor suppressor gene database (TSGene) (14) integrates TSGs with large-scale experimental evidence to provide a comprehensive resource for the further investigation of TSGs and their molecular mechanisms in cancer. The tumor-associated gene (TAG) database (15) is used to save new genes that play a role in carcinogenesis. In this study, we extracted all known oncogenes and TSGs from the TAG and TSG databases.

PPI network construction and sub-network identification. The Search Tool for the Retrieval of Interacting Genes (STRING) database (16) is a precomputed global resource which was designed to evaluate PPI information. In the PPI network, each node stands for a gene and the edges represent the interactions between nodes. The degree indicates the number of edges linked to a given node and the nodes with a high degree are defined as the hub genes that possess essential biological

functions. In this study, the STRING online tool was applied to analyze the PPI network of DEGs and only those experimentally validated interactions with a combined score > 0.9 were selected as significant.

The BioNet package (<http://www.bioconductor.org/packages/release/bioc/html/BioNet.html>) (17) provides a comprehensive set of methods for the integrated analysis of gene expression data and biological networks. In the current study, we used BioNet to identify the sub-network in the PPI network with FDR $< 1.0E-06$.

Based on the DEGs in the sub-network, we performed KEGG pathway enrichment analysis with $P < 0.01$.

Results

Identification of DEGs. For the dataset GSE43795, a total of 979 transcripts were obtained following data pre-processing. Among them, 393 were downregulated transcripts corresponding to 374 DEGs and 586 were upregulated transcripts which corresponded to 559 DEGs.

GO and pathway enrichment analyses. Following GO and pathway analyses for down- and upregulated DEGs, the top five GO BP terms were collected and are shown in Table I. The GO terms enriched by downregulated DEGs were mainly related to digestion, transport and signaling while the GO terms enriched by upregulated DEGs were mainly associated with the cell cycle and mitosis.

The pathways enriched by downregulated DEGs were mainly related to digestion, absorption and metabolism, and included protein digestion and absorption and metabolism of xenobiotics by cytochrome P450. The pathways enriched by upregulated DEGs were mainly related to the occurrence and spread of cancer, including the cell cycle and p53 signaling pathways (Table II).

Functional annotation of DEGs. After researching the expression of TFs and TAGs, 15 TFs were downregulated, including prospero homeobox 1 and PBX/knotted 1 homeobox 2, and 19 TFs were upregulated, including vitamin D receptor and upstream transcription factor 1. In addition, among the downregulated DEGs, 22 genes were TAGs. Of these, 3 were oncogenes, 16 were TSGs and the effect of other 3 genes was uncertain. In the upregulated DEGs, 12 were oncogenes, including pituitary tumor-transforming 1 (*PTTGI*), 32 were TSGs and the effect of other 11 genes was uncertain (Table III).

PPI network construction and sub-network pathway enrichment analysis. Based on data from the STRING database, the PPI network was constructed (Fig. 1). Ten nodes were selected as hub genes (degree ≥ 12), including cell division cycle 20 (*CDC20*, degree=18) and BUB1 mitotic checkpoint serine/threonine kinase B (*BUB1B*, degree=16).

Using the BioNet package, sub-networks with 34 nodes were detected. Fig. 2 shows that *CDC20* was a hub gene with degree=16.

Following KEGG pathway enrichment analyses of the DEGs in the sub-network, we observed that these DEGs including cyclin B1 (*CCNBI*) were mainly enriched in the cell cycle, p53 signaling and oocyte meiosis pathways.

Table I. Gene ontology functional enrichment analysis for down- and upregulated differentially expressed genes (top 5).

| Term | Pathway | Count | P-value |
|--------------------|---------------------------|-------|----------|
| Downregulated DEGs | | | |
| GO: 0007586 | Digestion | 18 | 1.16E-10 |
| GO: 0006811 | Ion transport | 53 | 1.73E-08 |
| GO: 0007267 | Cell-cell signaling | 53 | 2.21E-08 |
| GO: 0044765 | Single-organism transport | 104 | 3.34E-08 |
| GO: 0006810 | Transport | 114 | 7.52E-07 |
| Upregulated DEGs | | | |
| GO: 0000280 | Nuclear division | 44 | 6.22E-15 |
| GO: 0007067 | Mitosis | 44 | 6.22E-15 |
| GO: 0000278 | Mitotic cell cycle | 71 | 1.64E-14 |
| GO: 0048285 | Organelle fission | 44 | 9.46E-14 |
| GO: 0051301 | Cell division | 48 | 9.29E-12 |

Count signifies the number of enriched DEGs. DEG, differentially expressed gene.

Table II. Pathway functional enrichment analysis for down- and upregulated differentially expressed genes.

| Term | Pathway | Count | P-value |
|--------------------|---|-------|-----------|
| Downregulated DEGs | | | |
| 04972 | Pancreatic secretion | 19 | 4.79E-12 |
| 04974 | Protein digestion and absorption | 13 | 9.90E-08 |
| 00260 | Glycine, serine and threonine metabolism | 8 | 9.43E-07 |
| 04964 | Proximal tubule bicarbonate reclamation | 6 | 1.78E-05 |
| 04950 | Maturity onset diabetes of the young | 6 | 2.99E-05 |
| 00980 | Metabolism of xenobiotics by cytochrome P450 | 8 | 4.11E-04 |
| 00982 | Drug metabolism-cytochrome P450 | 8 | 4.97E-04 |
| 00480 | Glutathione metabolism | 6 | 1.58 E-03 |
| 04971 | Gastric acid secretion | 7 | 2.65 E-03 |
| 04973 | Carbohydrate digestion and absorption | 5 | 4.96 E-03 |
| 04020 | Calcium signaling pathway | 11 | 5.38 E-03 |
| 04610 | Complement and coagulation cascades | 6 | 8.03 E-03 |
| 00250 | Alanine, aspartate and glutamate metabolism | 4 | 8.44 E-03 |
| 00750 | Vitamin B6 metabolism | 2 | 9.09 E-03 |
| 04976 | Bile secretion | 6 | 9.21 E-03 |
| Upregulated DEGs | | | |
| 04110 | Cell cycle | 15 | 9.27E-06 |
| 04115 | p53 signaling pathway | 11 | 9.35E-06 |
| 04512 | ECM-receptor interaction | 10 | 3.77E-04 |
| 05200 | Pathways in cancer | 20 | 4.11E-03 |
| 05146 | Amoebiasis | 9 | 7.08E-03 |
| 05412 | Arrhythmogenic right ventricular cardiomyopathy | 7 | 9.61E-03 |

Count signifies the number of enriched DEGs. DEG, differentially expressed gene.

Discussion

The identification of DEGs in PDAC is critical to the development of novel strategies to detect and treat this highly malignant cancer. In the present study, a total of 933 DEGs

were identified between PDAC and NN samples through gene expression profiling of GSE43795. The downregulated DEGs were mainly enriched in the BP terms associated with digestion, transport and signaling, and pathways associated with digestion, absorption and metabolism. The upregulated DEGs

Table III. Results of functional annotation of differentially expressed genes.

| TF count | TF name | TAG count | TAG name |
|--------------------|--|-----------|---|
| Downregulated DEGs | | | |
| 15 | <i>PROX1, PKNOX2, PBX1, PAX6, ONECUT1, NR5A2, NR4A2, NKX2-5, NKX2-2, LMO3, KLF15, INSM1, GATA4, FOSB, ESRRG</i> | 22 | TAG oncogenes: <i>PBX1, GATA4, FGFR1</i> ; TSGs: <i>ZBTB16, WNK2, SFRP5, SFRP1, SERPINI2, PROX1, PLCE1, PAX6, ONECUT1, NRCAM, GNMT, DIRAS3, C2orf40, BTG2, BEX2, ARID3B</i> ; Others: <i>SLC43A1, NR4A2, CHRM3</i> |
| Upregulated DEGs | | | |
| 19 | <i>VDR, USF1, SPI1, RUNX2, RUNX1, PITX1, MYCBP, LEF1, HOXC4, HOXB8, HOXB7, HOXA13, HOXA10, FOXM1, FOXF2, FOXD2, FOXD1, FOXA1, E2F7</i> | 55 | TAG oncogenes: <i>WISP1, TNFRSF6B, SPI1, RUNX2, PTTG1, NRAS, LCN2, LAMC2, HOXA10, HMMR, CEP55, CCNA2</i> ; TSGs: <i>TES, SFN, SERPINB5, SEC14L2, RASAL1, RARRES3, RARRES1, PYCARD, PRODH, MMP11, MFHAS1, JUP, ISG15, INPP4B, IGFBP3, HTRA1, HPGD, HOPX, GPRC5A, GLIPR1, GJB2, FANCD2, EGLN3, CHEK1, CEACAM1, CDH11, CASP8, CAPG, BUB1B, BLM, BIK, ABLIM3</i> ; Others: <i>TFAP2A, TACC3, RUNX1, PTK6, OAS1, ITGB4, FHL2, DHDH, CCNE2, BUB1, BIRC5</i> |

TF, transcription factor; TAG, tumor-associated gene; DEG, differentially expressed gene; TSG, tumor suppressor gene.




Figure 1. Protein-protein interaction network of differentially expressed genes (DEGs). Green nodes represent downregulated DEGs; red nodes represent upregulated DEGs. The size of the node indicates the connectivity degree and larger circles indicate a higher degree.

were mainly enriched in the BP terms associated with cell cycle and mitosis, and in the cell cycle and p53 signaling pathways. The oncogene *PTTG1* was upregulated following functional annotation. In the PPI network, the hub genes *CDC20* and *BUB1B* had higher connectivity degrees. Additionally, *CCNB1*,

CDC20 and *BUB1B* were enriched in several pathways in the sub-networks. This result suggested that these genes and pathways may play significant roles in the progression of PDAC.

Cancer is characterized by uncontrolled cell proliferation, and tumor cells have typically acquired damage to genes




Figure 2. Sub-network of differentially expressed genes (DEGs). Color indicates \log_2 -fold change (from low to high: green, pink, red). Green nodes represent downregulated DEGs; red nodes represent upregulated DEGs. Circles indicate that the node is significant in the network while squares indicate less significance.

that directly regulate their cell cycles (18). Research has revealed that the loss of appropriate cell cycle regulation leads to genomic instability (19). The cell cycle is believed to play a role in the etiology of spontaneous cancers (20). In the present study, the cell cycle pathway was observed to be enriched by several upregulated DEGs, including *CDC20* and *BUB1B*, which were also hub genes in the PPI network and sub-network. *CDC20* is one of the regulators of spindle checkpoints, and appears to act as a regulatory protein interacting with several other proteins at multiple points in the cell cycle (21). In mammals, *CDC20* is involved in anaphase onset and late mitotic events (22) and its expression is essential for cell division (23). At present, *CDC20* is frequently reported to be upregulated in numerous types of malignancies including pancreatic cancer (23). Chang *et al* suggested that *CDC20* expression may play a role in facilitating PDAC cell mitosis. For the other DEG, *BUB1B*, its encoded protein is also a key protein in the mitotic spindle checkpoint (24). It has been reported that overexpression of spindle assembly checkpoint molecules may result in DNA aneuploidy and carcinogenesis in mice (25). The high expression of *BUB1B* has often been reported to be associated with chromosomal instability in several malignancies, including kidney carcinomas, breast cancer and bladder cancer (24,26,27). Our results further confirm that aberrant *CDC20* and *BUB1B* expression and cell cycle pathways in which the two DEGs participated may play key roles in PDAC tumorigenesis and progression and may thus be useful as therapeutic targets.

In this study, the p53 signaling pathway was also observed to be enriched by upregulated DEGs including *CCNB1*. The

p53 protein inhibits malignant transformation through direct and indirect regulation of transcription of the genes associated with the cell cycle and apoptosis (21). Presently, *TP53* is the most frequently mutated gene in human cancer and its mutation frequency is up to 96% in pancreatic adenocarcinoma (28,29). *CCNB1* is a regulatory protein involved in mitosis, and plays an essential role in cell proliferation (30). In normal tissues, the expression level of *CCNB1* is low; however, it was noted to be overexpressed in tumors with *TP53* mutation, including colorectal, cervical and pancreatic cancer (31-33). In addition, *TP53* has been demonstrated to regulate the promoter of *CCNB1* in opposing ways (34). Briefly, the p53 signaling pathway and its enriched DEGs including *CCNB1* were closely associated with PDAC; thus, this pathway and these genes may be used as potential targets for PDAC treatment.

The results of functional annotation of DEGs revealed that the *PTTG1* oncogene was upregulated; in addition, it was noted to participate in the PPI network and sub-network. Our findings also revealed that *PTTG1* was enriched in the BP terms associated with mitosis. *PTTG1* is a regulatory protein, and plays a central role in chromosome stability, cell transformation and gene regulation (35,36). In particular, *PTTG1* is a critical mitotic checkpoint protein that helps hold sister chromatids together before entering anaphase (37). Research has identified that *PTTG1* expression is highly activated in rapidly proliferating cells (38). To date, the overexpression of *PTTG1* has been identified in numerous cancers tissues as well as in cell lines, including colon, ovarian, breast and various other solid tumors (39-41). In our study, the overexpression of *PTTG1* is consistent with the observations above. Taken

together, these data support the hypothesis that *PTTG1* may be a candidate molecular marker associated with PDAC progression and prognosis.

Although bioinformatics technologies have the potential to identify and validate candidate agents for critical diseases, certain limitations remain in this study. Firstly, the sample size for microarray analysis was small, which may have caused a high rate of false positive results. Secondly, this study lacked experimental verification. Further genetic and experimental studies with a larger sample size are still required in the future to confirm the results.

However, our data provide a comprehensive bioinformatics analysis of the DEGs and pathways which may be involved in PDAC. The findings of the present study may contribute to our understanding of the underlying molecular mechanisms of PDAC. DEGs including *CDC20*, *BUB1B*, *CCNB1* and *PTTG1* as well as the cell cycle pathway and p53 signaling pathway have the potential to be used as targets for PDAC diagnosis and treatment.

Acknowledgements

The authors wish to express thanks to Fenghe (Shanghai) Information Technology Co., Ltd. Their ideas and assistance provided a valuable added dimension to this research.

References

- Aguirre AJ, Bardeesy N, Sinha M, Lopez L, Tuveson DA, Horner J, Redston MS and DePinho RA: Activated Kras and Ink4a/Arf deficiency cooperate to produce metastatic pancreatic ductal adenocarcinoma. *Genes Dev* 17: 3112-3126, 2003.
- Garcia M, Jemal A, Ward EM, Center MM, Hao Y, Siegel RL and Thun MJ: Global Cancer Facts & Figures 2007. American Cancer Society, Atlanta, GA, 2007.
- Conlon KC, Klimstra DS and Brennan MF: Long-term survival after curative resection for pancreatic ductal adenocarcinoma. Clinicopathologic analysis of 5-year survivors. *Ann Surg* 223: 273-279, 1996.
- Sarkar FH, Banerjee S and Li Y: Pancreatic cancer: pathogenesis, prevention and treatment. *Toxicol Appl Pharmacol* 224: 326-336, 2007.
- Berthelemy P, Bouisson M, Escourrou J, Vaysse N, Rumeau JL and Pradayrol L: Identification of K-ras mutations in pancreatic juice in the early diagnosis of pancreatic cancer. *Ann Intern Med* 123: 188-191, 1995.
- Iacobuzio-Donahue CA, Ashfaq R, Maitra A, Adsay NV, Shen-Ong GL, Berg K, Hollingsworth MA, Cameron JL, Yeo CJ, Kern SE, *et al*: Highly expressed genes in pancreatic ductal adenocarcinomas: a comprehensive characterization and comparison of the transcription profiles obtained from three major technologies. *Cancer Res* 63: 8614-8622, 2003.
- Park M, Kim M, Hwang D, Park M, Kim WK, Kim SK, Shin J, Park ES, Kang CM, Paik YK and Kim H: Characterization of gene expression and activated signaling pathways in solid-pseudopapillary neoplasm of pancreas. *Mod Pathol* 27: 580-593, 2014.
- Irizarry RA, Hobbs B, Collin F, Beazer-Barclay YD, Antonellis KJ, Scherf U and Speed TP: Exploration, normalization and summaries of high density oligonucleotide array probe level data. *Biostatistics* 4: 249-264, 2003.
- Gautier L, Cope L, Bolstad BM and Irizarry RA: Affy-analysis of Affymetrix GeneChip data at the probe level. *Bioinformatics* 20: 307-315, 2004.
- Smyth GK: Limma: linear models for microarray data. In: *Bioinformatics and Computational Biology Solutions using R and Bioconductor*. Gentleman R, Carey V, Dudoit S, Irizarry R and Huber W (eds). Springer, New York, Springer, pp397-420, 2005.
- The Gene Ontology Consortium; Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP, Dolinski K, Dwight SS, Eppig JT, *et al*: Gene Ontology: tool for the unification of biology. *Nat Genet* 25: 25-29, 2000.
- Kanehisa M and Goto S: KEGG: Kyoto encyclopedia of genes and genomes. *Nucleic Acids Res* 28: 27-30, 2000.
- Huang DW, Sherman BT, Tan Q, Collins JR, Alvord WG, Roayaei J, Stephens R, Baseler MW, Lane HC and Lempicki RA: The DAVID Gene Functional Classification Tool: a novel biological module-centric algorithm to functionally analyze large gene lists. *Genome Biol* 8: R183, 2007.
- Zhao M, Sun J and Zhao Z: TSGene: a web resource for tumor suppressor genes. *Nucleic Acids Res* 41 (Database issue): D970-D976, 2013.
- Chen JS, Hung WS, Chan HH, Tsai SJ and Sun HS: In silico identification of oncogenic potential of fyn-related kinase in hepatocellular carcinoma. *Bioinformatics* 29: 420-427, 2013.
- Franceschini A, Szklarczyk D, Frankild S, Kuhn M, Simonovic M, Roth A, Lin J, Minguez P, Bork P, von Mering C and Jensen LJ: STRING v9.1: protein-protein interaction networks, with increased coverage and integration. *Nucleic Acids Res* 41 (Database issue): D808-D815, 2013.
- Beisser D, Klau GW, Dandekar T, Müller T and Dittrich MT: BioNet: an R-Package for the functional analysis of biological networks. *Bioinformatics* 26: 1129-1130, 2010.
- Sherr CJ: Cancer cell cycles. *Science* 274: 1672-1677, 1996.
- Hartwell LH and Kastan MB: Cell cycle control and cancer. *Science* 266: 1821-1828, 1994.
- Hunter T and Pines J: Cyclins and cancer II: cyclin D and CDK inhibitors come of age. *Cell* 79: 573-582, 1994.
- Kidokoro T, Tanikawa C, Furukawa Y, *et al*: CDC20, a potential cancer therapeutic target, is negatively regulated by p53. *Oncogene* 27: 1562-1571, 2008.
- Fung TK and Poon RY: A roller coaster ride with the mitotic cyclins. *Semin Cell Dev Biol* 16: 335-342, 2005.
- Chang DZ, Ma Y, Ji B, Liu Y, Hwu P, Abbruzzese JL, Logsdon C and Wang H: Increased CDC20 expression is associated with pancreatic ductal adenocarcinoma differentiation and progression. *J Hematol Oncol* 5: 15, 2012.
- Yamamoto Y, Matsuyama H, Chochi Y, Okuda M, Kawauchi S, Inoue R, Furuya T, Oga A, Naito K and Sasaki K: Overexpression of BUBR1 is associated with chromosomal instability in bladder cancer. *Cancer Genet Cytogenet* 174: 42-47, 2007.
- van Deursen JM: Rb loss causes cancer by driving mitosis mad. *Cancer cell* 11: 1-3, 2007.
- Pinto M, Vieira J, Ribeiro FR, Soares MJ, Henrique R, Oliveira J, Jerónimo C and Teixeira MR: Overexpression of the mitotic checkpoint genes BUB1 and BUBR1 is associated with genomic complexity in clear cell kidney carcinomas. *Cell Oncol* 30: 389-395, 2008.
- Scintu M, Vitale R, Prencipe M, Gallo AP, Bonghi L, Valori VM, Maiello E, Rinaldi M, Signori E, Rabitti C, *et al*: Genomic instability and increased expression of BUB1B and MAD2L1 genes in ductal breast carcinoma. *Cancer Lett* 254: 298-307, 2007.
- Petitjean A, Mathe E, Kato S, Ishioka C, Tavtigian SV, Hainaut P and Olivier M: Impact of mutant p53 functional properties on TP53 mutation patterns and tumor phenotype: lessons from recent developments in the IARC TP53 database. *Hum Mutat* 28: 622-629, 2007.
- Hingorani SR, Wang L, Multani AS, Combs C, Deramandt TB, Hruban RH, Rustgi AK, Chang S and Tuveson DA: Trp53R172H and KrasG12D cooperate to promote chromosomal instability and widely metastatic pancreatic ductal adenocarcinoma in mice. *Cancer Cell* 7: 469-483, 2005.
- Suzuki T, Urano T, Miki Y, Moriya T, Akahira J, Ishida T, Horie K, Inoue S and Sasano H: Nuclear cyclin B1 in human breast carcinoma as a potent prognostic factor. *Cancer Sci* 98: 644-651, 2007.
- Wang A, Yoshimi N, Ino N, Tanaka T and Mori H: Overexpression of cyclin B1 in human colorectal cancers. *J Cancer Res Clin Oncol* 123: 124-127, 1997.
- El-Ghobashy AA, Shaaban AM, Herod J, *et al*: Overexpression of cyclins A and B as markers of neoplastic glandular lesions of the cervix. *Gynecol Oncol* 92: 628-634, 2004.
- Weissmueller S, Machado E, Saborowski M, *et al*: Mutant p53 drives pancreatic cancer metastasis through cell-autonomous PDGF receptor β signaling. *Cell*, 157: 382-394, 2014. .
- Yin XY, Grove L, Datta NS, Katula K, Long MW and Prochownik EV: Inverse regulation of cyclin B1 by c-Myc and p53 and induction of tetraploidy by cyclin B1 overexpression. *Cancer Res* 61: 6487-6493, 2001.
- Zhang X, Horwitz GA, Prezant TR, *et al*: Structure, expression and function of human pituitary tumor-transforming gene (PTTG). *Mol Endocrinol* 13: 156-166, 1999.

36. Pei L and Melmed S: Isolation and characterization of a pituitary tumor-transforming gene (PTTG). *Mol Endocrinol* 11: 433-441, 1997.
37. Ying H, Furuya F, Zhao L, Araki O, West BL, Hanover JA, Willingham MC and Cheng SY: Aberrant accumulation of PTTG1 induced by a mutated thyroid hormone β receptor inhibits mitotic progression. *J Clin Invest* 116: 2972-2984, 2006.
38. Genkai N, Homma J, Sano M, Tanaka R and Yamanaka R: Increased expression of pituitary tumor-transforming gene (PTTG)-1 is correlated with poor prognosis in glioma patients. *Oncol Rep* 15: 1569-1574, 2006.
39. Heaney AP, Singson R, McCabe CJ, Nelson V, Nakashima M and Melmed S: Expression of pituitary-tumour transforming gene in colorectal tumours. *Lancet* 355: 716-719, 2000.
40. Solbach C, Roller M, Fellbaum C, Nicoletti M and Kaufmann M: PTTG mRNA expression in primary breast cancer: a prognostic marker for lymph node invasion and tumor recurrence. *Breast* 13: 80-81, 2004.
41. Puri R, Tousson A, Chen L and Kakar SS: Molecular cloning of pituitary tumor transforming gene 1 from ovarian tumors and its expression in tumors. *Cancer Lett* 163: 131-139, 2001.