



Cite this article: Nati JJH, Lindström J, Halsey LG, Killen SS. 2016 Is there a trade-off between peak performance and performance breadth across temperatures for aerobic scope in teleost fishes? *Biol. Lett.* **12**: 20160191. <http://dx.doi.org/10.1098/rsbl.2016.0191>

Received: 5 March 2016

Accepted: 6 September 2016

Subject Areas:

ecology

Keywords:

ecophysiology, metabolic rate, locomotion, thermal performance, environmental change

Author for correspondence:

Julie J. H. Nati

e-mail: j.nati.1@research.gla.ac.uk

Electronic supplementary material is available online at <https://dx.doi.org/10.6084/m9.fig-share.c.3469878>.

Is there a trade-off between peak performance and performance breadth across temperatures for aerobic scope in teleost fishes?

Julie J. H. Nati¹, Jan Lindström¹, Lewis G. Halsey² and Shaun S. Killen¹

¹Institute of Biodiversity, Animal Health and Comparative Medicine, University of Glasgow, Graham Kerr Building, Glasgow G12 8QQ, UK

²Department of Life Sciences, University of Roehampton, Holybourne Avenue, London SW15 4JD, UK

JJHN, 0000-0001-5613-8365; LGH, 0000-0002-0786-7585

The physiology and behaviour of ectotherms are strongly influenced by environmental temperature. A general hypothesis is that for performance traits, such as those related to growth, metabolism or locomotion, species face a trade-off between being a thermal specialist or a thermal generalist, implying a negative correlation between peak performance and performance breadth across a range of temperatures. Focusing on teleost fishes, we performed a phylogenetically informed comparative analysis of the relationship between performance peak and breadth for aerobic scope (AS), which represents whole-animal capacity available to carry out simultaneous oxygen-demanding processes (e.g. growth, locomotion, reproduction) above maintenance. Literature data for 28 species indicate that peak aerobic capacity is not linked to thermal performance breadth and that other physiological factors affecting thermal tolerance may prevent such a trade-off from emerging. The results therefore suggest that functional links between peak and thermal breadth for AS may not constrain evolutionary responses to environmental changes such as climate warming.

1. Introduction

For ectotherms, performance traits related to growth, reproduction and locomotion are often depicted using thermal performance curves [1,2] that illustrate how a trait responds to variation in environmental temperature (figure 1). Evolutionary thermal adaptation [2] may result in thermal specialists or thermal generalists, performing better over a narrow versus a broad range of temperatures, respectively (figure 1; [3]). Thermal and biochemical constraints on enzyme structure and function and membrane fluidity suggest that adaptations for increased performance at one temperature may cause decreased performance at other temperatures, resulting in a trade-off between peak performance (P_{\max}) at a thermal optimum and thermal performance breadth (T_{breadth}). Owing to these potential compromises, previous researchers have suggested that a 'jack of all temperatures is a master of none' [4].

While a trade-off between P_{\max} and T_{breadth} is predicted by theory [3,5], several studies have documented that an increased performance capacity at one temperature does not necessarily lead to reduced performance at other temperatures [4,6–9]. Notably, however, most studies have examined the differences in performance among populations of the same species, with few attempts to examine whether a trade-off exists across species. Therefore, it remains unknown whether any trade-off between P_{\max} and T_{breadth} generate interspecific constraints on thermal




Figure 1. Theoretical thermal performance curves illustrating thermal specialists (higher peak performance, in blue) and thermal generalists (higher performance breadth, in orange). Specialists have a higher peak performance (P_{\max}) at their optimum temperature (T_{opt}). Generalists have a lower P_{\max} but a wider breadth of temperatures over which they perform normally (T_{breadth} , here defined as the range of temperatures allowing 80% of P_{\max}). Critical minimum and maximum thermal limits (CT_{\min} and CT_{\max} , respectively) occur where performance equals zero. Inset: predicted negative correlation between P_{\max} and T_{breadth} if there is a trade-off between being a thermal specialist and a thermal generalist.

adaptation. Furthermore, most attempts to examine trade-offs between P_{\max} and T_{breadth} have focused on isolated components of locomotory performance (e.g. maximum speed, endurance). This approach may, however, fail to detect broader-scale trade-offs in organismal functioning. Aerobic scope (AS), in contrast, is an integrative trait, representing whole-animal cardiovascular and respiratory capacity to provide oxygen above maintenance requirements, for aerobic activities including growth, locomotion and reproduction [10,11]. In ectotherms, AS generally increases with temperature until T_{opt} and then usually decreases with further warming ([12], but see [13] for exceptions), potentially providing a composite measure to examine thermal sensitivity of whole-animal aerobic performance. AS is also ecologically relevant and has been related to geographical distribution [14], the capacity to cope with environmental stressors [15] and competitive ability [16]. Additionally, species with a higher AS tend to be more active and athletic [17], presumably because increased locomotion requires a greater allocation of oxygen to skeletal muscles [12]. If there is indeed a trade-off between P_{\max} and T_{breadth} , then ectothermic species that experience selection for increased peak AS to facilitate foraging, predator avoidance or migration may conversely have a reduced T_{breadth} [12].

We investigated the relationship between P_{\max} and T_{breadth} for AS across 28 species of teleost fish, to determine the extent to which adaptation for performance at a particular optimal temperature may impose constraints on performance across a range of temperatures. We focus on fishes because they are a diverse taxon that experiences shifts in thermal regimes over varying timescales. It has also been proposed that AS may influence the ability of fishes to respond to climate change [18].

2. Material and methods

Data for the AS of fish, calculated as the difference between standard and maximum metabolic rates (SMR and MMR), were

extracted from the literature where they were available at three or more temperatures for a species within a single study ($n = 28$ species; electronic supplementary material, table S1). A Gaussian model was fitted to the relationship between AS and temperature for each species to produce a thermal performance curve for each species [19]. Peak AS (P_{\max}) was defined as the highest value for AS along the thermal performance curve. The optimum temperature (T_{opt}) was the temperature corresponding to P_{\max} [1,11]. We used one dataset per species; when more than one dataset was available, we used the dataset that measured AS at the highest number of temperatures. If multiple datasets were available for a species within a given study, then we used that which gave the highest value for P_{\max} . Performance breadth (T_{breadth}) was the range of temperatures over which a species maintained at least 80% of peak AS [20]. As the 80% of P_{\max} designation for T_{breadth} range is somewhat arbitrary, we also examined 60%, 70% and 90% thresholds for T_{breadth} . The modelled values for T_{opt} and P_{\max} versus the highest values for measurement temperature and AS, respectively, which were observed in each source study are given in electronic supplementary material, figure S1. Data for an additional 15 species were not included because either: (i) AS did not increase or decrease appreciably over the temperatures tested (i.e. the performance 'curve' was flat; two species); (ii) AS decreased with no obvious peak across temperatures tested in the study, perhaps, because the lowest temperature in the study was above T_{opt} (three cases); (iii) AS increased with no obvious peak across temperatures tested in the study, perhaps, because the highest temperature in the study was below T_{opt} (nine cases) or (iv) AS continued to increase with temperature until CT_{\max} (one case).

Data were analysed using the phylogenetic generalized least-squares (PGLS) method applying a phylogeny generated from the comprehensive tree of life [21] (electronic supplementary material, figure S2 for more details on statistical analysis, including phylogenetic tree). $\log P_{\max}$ ($\text{mg O}_2 \text{ h}^{-1}$) was the response variable with T_{breadth} , T_{opt} and \log body mass (g) as explanatory variables. Six studies measured AS in fish acutely exposed to each temperature; we therefore constructed a separate model using only studies in which fish were thermally acclimated ($n = 20$ species, electronic supplementary material, table S2). Model residuals were checked for normality




Figure 2. (a) Relationship between log-transformed peak aerobic scope (P_{\max}) and thermal performance breadth (T_{breadth}). For visual representation, data were standardized for body mass and T_{opt} using residuals from a PGLS multiple regression of $\log P_{\max}$ versus \log body mass and T_{opt} ($\log P_{\max} = 1.0013(\log \text{mass}) + 0.0285(T_{\text{opt}}) - 0.9778$; $p < 0.001$, $r^2 = 0.924$), added to the fitted model value for body mass = 300 g and $T_{\text{opt}} = 20^\circ\text{C}$ (the mean body mass and T_{opt} for species used in this study, respectively). (b) Relationship between P_{\max} and optimum temperature (T_{opt}). For this panel, P_{\max} was standardized to a body mass of 300 g using residuals of a PGLS linear regression of $\log P_{\max}$ versus \log body mass ($\log P_{\max} = 0.9409(\log \text{mass}) - 0.3227$; $p < 0.001$; $r^2 = 0.864$). In both panels, each point represents one species ($n = 28$).

and homogeneity of variance. The significance level of all tests was $\alpha = 0.05$.

3. Results

PGLS analysis revealed no relationship between P_{\max} and T_{breadth} , regardless of the threshold used to define T_{breadth} (figure 2a, model details for 80% T_{breadth} threshold in table 1; for 60%, $p = 0.188$; 70%, $p = 0.368$; 90%, $p = 0.200$; electronic supplementary material, tables S3–S5). P_{\max} increased with T_{opt} (figure 2b; PGLS, effect of T_{opt} , $t = 4.240$, $p < 0.001$). Explanatory variables explained 92.2% of variation in P_{\max} (R^2). Trends were identical when PGLS models were performed using studies with acclimated animals only (electronic supplementary material, table S2, PGLS, effect of $T_{\text{breadth80\%}}$, $t = -0.752$, $p = 0.466$).

4. Discussion

We found no evidence of a trade-off between T_{breadth} and P_{\max} for AS across teleost fishes. Evolutionary and plastic changes to peak performance for AS may not necessarily lead to reduced performance over a broader range of

Table 1. Summary of the PGLS model testing for the effects of aerobic scope breadth ($T_{\text{breadth80\%}}$, °C), optimal temperature (T_{opt} , °C) and body mass (\log g) on P_{\max} (log mg O₂ h⁻¹). $r^2 = 0.937$, $F_{5,21} = 62.59$, $p < 0.001$, $n = 28$ species, $\lambda = 0.70$. For lifestyle categorization, the reference category is ‘benthic’.

term	estimate	s.e.m.	t	p
intercept	-1.000	0.252	-3.971	<0.001
$T_{\text{breadth80\%}}$	-0.01	0.008	-1.249	0.226
T_{opt}	0.03	0.007	4.240	<0.001
log mass	1.01	0.066	15.387	<0.001
lifestyle				
benthopelagic	0.072	0.110	0.660	0.516
pelagic	0.246	0.117	2.095	0.049

temperatures and so there may not be differentiation along a thermal specialist/generalist continuum with respect to AS in teleost fishes. Investment in factors such as gill surface area, heart pumping capacity, tissue vascularization and mitochondrial density should all act to increase P_{\max} [17,22]. The results here suggest that these features can also increase AS at other temperatures, particularly if compensatory mechanisms allow for plasticity in each in response to temperature (e.g. change in the concentration or isoforms of aerobic enzymes [23]). It has been proposed that AS may constrain geographical distributions of aquatic ectothermic species in the face of a warming climate and other aspects of environmental changes [18]. The findings here suggest that the evolutionary potential of P_{\max} is not constrained by prior adaptation for a wider T_{breadth} , or vice versa, depending on the degree of genetic correlation between P_{\max} and T_{breadth} . For example, stenothermal species, which may have experienced relaxed selection for T_{breadth} , may be able to readily evolve a broader performance breadth or shift P_{\max} in response to changing thermal regimes, at least for AS. There are also other factors influencing plastic and evolutionary responses to thermal regimes that perhaps override or obscure links between P_{\max} and T_{breadth} for AS. For example, a decreased T_{breadth} for AS may not compromise fitness to the same extent as a reduced ability to adjust cellular membrane fluidity in response to thermal variation [24]. This could preclude a negative correlation between P_{\max} and T_{breadth} for AS from arising across species through evolutionary processes. It should also be noted that there appears to be several fish species that may not show a decline in AS with increasing temperature before reaching CT_{max} [13]. In such species, P_{\max} must also not be constrained by performance breadth for AS.

Our phylogenetically informed analysis spans a variety of species with varying lifestyles. Still, complex interactions among thermal history, body size, lifestyle and evolutionary history could also mask a trade-off between P_{\max} and T_{breadth} for AS. Across fishes, AS differs among benthic, benthopelagic and pelagic species [17], and so it is also possible that selection on AS to support locomotory capacity may outweigh thermal effects on P_{\max} . This interpretation is supported by the observation that pelagic species had a higher P_{\max} in this study, as compared to benthic and benthopelagic species. (table 1). It must also be considered that

species may differ in the percentage of P_{\max} most relevant for defining T_{breadth} [12]. For example, species that perform long migrations might require a higher proportion of T_{breadth} to maximize lifetime fitness when compared with more sedentary species.

P_{\max} was higher in species with a higher T_{opt} providing support for the 'hotter is better' model [25]. AS in fishes generally increases with temperature until T_{opt} and then decreases as temperature increases further (though see [13]). It is unlikely that species would live in habitats with temperatures much higher than T_{opt} given that thermal performance curves can be asymmetrical, with performance dropping more steeply above T_{opt} [12]. To date, most studies have not measured AS at sufficient temperature points to permit complex asymmetrical modelling [19] and so we have little understanding of how curve asymmetry may be linked with thermal specialization for AS. This may be especially relevant for the many tropical fishes that have a decreased thermal window between CT_{\min} and CT_{\max} [26]. Future studies examining changes in AS with temperature in ectotherms should perform measurements across as many

temperatures as possible to permit complex modelling of asymmetrical performance curves.

Data accessibility. The data from this study are available as electronic supplementary material.

Ethics. All procedures comply with UK Home Office Project Licence 60/4461.

Authors' contributions. J.J.H.N., J.L. and S.S.K. conceived the study. J.J.H.N. and S.S.K. collected the data. J.J.H.N., J.L., L.G.H. and S.S.K. performed the analysis. J.J.H.N. drafted the manuscript with editorial assistance from J.L., L.G.H. and S.S.K. All authors give final approval of the version to be published and agree to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Competing interests. The authors declare no competing interests.

Funding. S.S.K. was supported by NERC Advanced Fellowship NE/J019100/1 and European Research Council Starting Grant 640004. J.J.H.N. was supported by an Aides à la Formation Recherche doctoral grant from the Fonds National de la Recherche Luxembourg (4005263).

Acknowledgements. We thank David McKenzie and four anonymous reviewers for comments on an earlier version of this manuscript.

References

- Huey RB, Stevenson R. 1979 Integrating thermal physiology and ecology of ectotherms: a discussion of approaches. *Am. Zool.* **19**, 357–366. (doi:10.1093/icb/19.1.357)
- Angilletta MJ. 2009 *Thermal adaptation: a theoretical and empirical synthesis*. Oxford, UK: Oxford University Press.
- Gilchrist GW. 1995 Specialists and generalists in changing environments. I. Fitness landscapes of thermal sensitivity. *Am. Nat.* **146**, 252–270. (doi:10.1086/285797)
- Huey RB, Hertz PE. 1984 Is a jack-of-all-temperatures a master of none? *Evolution* **38**, 441–444. (doi:10.2307/2408502)
- Levins R. 1962 Theory of fitness in a heterogeneous environment. I. The fitness set and adaptive function. *Am. Nat.* **96**, 361–373. (doi:10.1086/282245)
- Ojanguren AF, Reyes-Gavilán FG, Braña F. 2001 Thermal sensitivity of growth, food intake and activity of juvenile brown trout. *J. Therm. Biol.* **26**, 165–170. (doi:10.1016/S0306-4565(00)00038-3)
- Wilson RS, Franklin CE, Davison W, Kraft P. 2001 Stenotherms at sub-zero temperatures: thermal dependence of swimming performance in Antarctic fish. *J. Comp. Physiol. B* **171**, 263–269. (doi:10.1007/s003600000172)
- Purchase CF, Brown JA. 2000 Interpopulation differences in growth rates and food conversion efficiencies of young Grand Banks and Gulf of Maine Atlantic cod (*Gadus morhua*). *Can. J. Fish Aquat. Sci.* **57**, 2223–2229. (doi:10.1139/f00-204)
- Malloy KD, Targett TE. 1994 Effects of ration limitation and low temperature on growth, biochemical condition, and survival of juvenile summer flounder from two Atlantic coast nurseries. *Trans. Am. Fish. Soc.* **123**, 182–193. (doi:10.1577/1548-8659(1994)123<0182:EORLAL>2.3.CO;2)
- Clark TD, Sandblom E, Jutfelt F. 2013 Aerobic scope measurements of fishes in an era of climate change: respirometry, relevance and recommendations. *J. Exp. Biol.* **216**, 2771–2782. (doi:10.1242/jeb.084251)
- Fry FEJ. 1971 The effect of environmental factors on the physiology of fish. In *Fish physiology*, pp. 1–98. San Diego, CA: Academic Press.
- Farrell AP. 2016 Pragmatic perspective on aerobic scope: peaking, plummeting, pejus and apportioning. *J. Fish Biol.* **88**, 322–343. (doi:10.1111/jfb.12789)
- Lefevre S. 2016 Are global warming and ocean acidification conspiring against marine ectotherms? A meta-analysis of the respiratory effects of elevated temperature, high CO₂ and their interaction. *Conserv. Physiol.* **4**, cow009. (doi:10.1093/conphys/cow009)
- Payne NL *et al.* 2016 Temperature dependence of fish performance in the wild: links with species biogeography and physiological thermal tolerance. *Funct. Ecol.* **30**, 903–912. (doi:10.1111/1365-2435.12618)
- Claireaux G, Lefrancois C. 2007 Linking environmental variability and fish performance: integration through the concept of scope for activity. *Phil. Trans. R. Soc. B* **362**, 2031–2041. (doi:10.1098/rstb.2007.2099)
- Seth H, Gräns A, Sandblom E, Olsson C, Wiklander K, Johnsson JI, Axelsson M. 2013 Metabolic scope and interspecific competition in sculpins of Greenland are influenced by increased temperatures due to climate change. *PLoS ONE* **8**, e62859. (doi:10.1371/journal.pone.0062859)
- Killen SS, Glazier D, Rezende EL, Clark T, Atkinson D, Willener A, Halsey LG. 2016 Ecological influences and physiological correlates of metabolic rates in teleost fishes. *Am. Nat.* **187**, 592–606. (doi:10.1086/685893)
- Pörtner HO, Farrell AP. 2008 Physiology and climate change. *Science* **322**, 690–692. (doi:10.1126/science.1163156)
- Angilletta MJ. 2006 Estimating and comparing thermal performance curves. *J. Therm. Biol.* **31**, 541–545. (doi:10.1016/j.jtherbio.2006.06.002)
- Frisk M, Skov PV, Steffensen JF. 2012 Thermal optimum for pikeperch (*Sander lucioperca*) and the use of ventilation frequency as a predictor of metabolic rate. *Aquaculture* **324**, 151–157. (doi:10.1016/j.aquaculture.2011.10.024)
- Hinchliff CE *et al.* 2015 Synthesis of phylogeny and taxonomy into a comprehensive tree of life. *Proc. Natl Acad. Sci. USA* **112**, 12 764–12 769. (doi:10.1073/pnas.1423041112)
- Norin T, Malte H. 2012 Intraspecific variation in aerobic metabolic rate of fish: relations with organ size and enzyme activity in brown trout. *Physiol. Biochem. Zool.* **85**, 645–656. (doi:10.1086/665982)
- Prosser CL. 1991 *Comparative animal physiology, environmental and metabolic animal physiology*. Hoboken, NJ: John Wiley & Sons.
- Cossins A, Prosser C. 1978 Evolutionary adaptation of membranes to temperature. *Proc. Natl Acad. Sci. USA* **75**, 2040–2043. (doi:10.1073/pnas.75.4.2040)
- Angilletta MJ, Huey RB, Frazier MR. 2010 Thermodynamic effects on organismal performance: is hotter better? *Physiol. Biochem. Zool.* **83**, 197–206. (doi:10.1086/648567)
- Sunday JM, Bates AE, Dulvy NK. 2011 Global analysis of thermal tolerance and latitude in ectotherms. *Proc. R. Soc. B* **278**, 1823–1830. (doi:10.1098/rspb.2010.1295)