

SCIENTIFIC REPORTS



Correction: Publisher Correction

OPEN

Acoustic allometry revisited: morphological determinants of fundamental frequency in primate vocal production

Maxime Garcia^{1,2}, Christian T. Herbst¹, Daniel L. Bowling¹, Jacob C. Dunn^{3,4} & W. Tecumseh Fitch¹

A fundamental issue in the evolution of communication is the degree to which signals convey accurate (“honest”) information about the signaler. In bioacoustics, the assumption that fundamental frequency (f_0) should correlate with the body size of the caller is widespread, but this belief has been challenged by various studies, possibly because larynx size and body size can vary independently. In the present comparative study, we conducted excised larynx experiments to investigate this hypothesis rigorously and explore the determinants of f_0 . Using specimens from eleven primate species, we carried out an inter-specific investigation, examining correlations between the minimum f_0 produced by the sound source, body size and vocal fold length (VFL). We found that, across species, VFL predicted minimum f_0 much better than body size, clearly demonstrating the potential for decoupling between larynx size and body size in primates. These findings shed new light on the diversity of primate vocalizations and vocal morphology, highlighting the importance of vocal physiology in understanding the evolution of mammal vocal communication.

The question of when and why animal signals convey accurate information about the signaler or environment – the problem of “honest” communication – has a long history^{1–3}. In the domain of acoustic communication, important insights have recently come from applying a better understanding of the vocal production mechanism to this issue^{4–7}. These studies indicate that increased understanding of the signal production mechanism can play a central role in explaining what components of a signal convey honest information, and why (e.g. ref.⁸).

Contemporary understanding of vocal production in mammal communication has benefited greatly from adopting the source-filter theory of vocal production^{4,5,9,10}. According to this framework¹¹, originally developed for human speech and later applied to animal communication, a sound is produced by the vibrating vocal folds within the larynx (the sound source) and their vibration rate determines the fundamental frequency (hereafter f_0) of the acoustic signal. This source signal then propagates through the vocal tract, where airborne resonances (which vary with vocal tract length and shape) emphasize some frequencies, called formants.

The connection between the acoustic characteristics of vocalizations and the physical attributes of the signaler suggests that key aspects of sound production can be anatomically constrained, with much research focusing on the relationship between a caller’s body size and f_0 ^{7,12,13}. In particular, Morton¹² postulated that an animal’s body size should be negatively related to the frequency content of its voice (including f_0), although he did not specify the physical and/or physiological factors that would underlie this putative correlation.

The prediction of a correlation between body size and f_0 relies on two main assumptions⁵: 1) body size directly determines the size of the larynx and therefore, the length of the vocal folds (as vocal folds in mammalian larynges extend from the thyroid cartilage to the arytenoids¹⁴), and 2) that the resting (i.e., unstretched) vocal fold length (hereafter VFL) has a direct influence on f_0 . Biomechanical theory corroborates the latter condition, predicting that longer focal folds produce lower f_0 ¹⁵. However, the former assumption has been challenged, given that

¹Department of Cognitive Biology, University of Vienna, Althanstrasse 14, 1090, Vienna, Austria. ²ENES Lab, Université Lyon/Saint-Etienne, NEURO-PSI, CNRS UMR 9197, Saint-Etienne, France. ³Division of Biological Anthropology, University of Cambridge, Pembroke Street, Cambridge, CB2 3QG, UK. ⁴Animal and Environment Research Group, Anglia Ruskin University, East Road, Cambridge, CB1 1PT, UK. Correspondence and requests for materials should be addressed to M.G. (email: maxime.garcia@ymail.com) or W.T.F. (email: tecumseh.fitch@univie.ac.at)

larynx size is not necessarily constrained by body size^{5,8}. Indeed various intraspecific studies, in multiple species, have failed to reveal the expected size-frequency relationship, finding a weak or non-existent correlation between body size and f_0 within adults of a given species^{8,16–19}.

Research on primate vocal production has also followed this general line of thought regarding the body size – f_0 relationship. A literature-based analysis conducted by Hauser²⁰ concluded that ‘larger species produce relatively lower-pitched vocalizations than smaller species’, relying on an amalgam of various frequency measures determined by visual inspection of printed spectrograms. In Hauser’s study however, the methodology applied to designate ‘frequency’ pooled manual measurements of the dominant frequency (hereafter DF) and f_0 . The interpretation of Hauser’s results is difficult because f_0 and DF reflect different acoustic phenomena: while f_0 reflects the rate of vibration of the vocal folds, DF is defined as the frequency at which the radiated acoustic spectrum has its greatest amplitude (see e.g. ref.²¹). DF is influenced by both the spectral composition of the laryngeal sound source and the filtering characteristics of the vocal tract. Such a conflation of distinct causal factors could easily confound the quantitative estimation of the relationship between frequency and body size across species, as shown by a recent study conducted on a wide range of vocalizations from numerous primate and carnivore species⁷.

Another complicating factor is that f_0 can strongly depend on several parameters besides VFL. For example, an increase in subglottal pressure (hereafter Psub), determined by the air pressure from the lungs, typically leads to an increase in f_0 ^{22,23}. Likewise, an increase in the tension applied to the vocal folds has similar effects: stretching of the vocal folds by the action of the cricothyroid muscle¹⁵ increases tension and stiffness, leading to a higher f_0 ²⁴. Vocal fold mass may also affect f_0 , although this has recently been disputed²⁵. Finally, the vocal folds are multilayered structures²⁶ and layer composition varies across species^{27,28}, which could influence elasticity²⁹ and thus f_0 ³⁰.

The influence of these multiple factors means that for a given VFL and tissue composition, an animal can in principle greatly increase f_0 by increasing Psub²⁴ and vocal fold tension³¹. Analyses of vocalizations produced by free-moving animals, including the study conducted by Hauser²⁰ and the most recent large-scale analyses on the question^{6,7}, cannot account for these confounding factors. The use of experimentally-controlled *in vitro* phonation in an excised larynx setup offers a major advantage in this respect, providing accurate measurement and control of key factors, such as Psub and vocal fold tension³². Unlike *in vivo* conditions, excised larynx experiments also allow us to adjust and precisely document laryngeal geometry and vocal fold position.

In the present comparative study we use an automated excised larynx setup to investigate larynges from 11 primate species, phonated in a controlled laboratory setting, to examine the physical and physiological determinants of inter-specific variation of primate f_0 in detail. Our underlying physical model is given in equation (1)¹⁵, representing a simple string model of a vibrating vocal fold, where L is the VFL, σ is the tensile stress in the vocal fold and ρ the tissue density:

$$f_0 = \frac{1}{2L} \sqrt{\frac{\sigma}{2L}} \text{ [Hz]} \quad (1)$$

Equation (1) suggests that, given constant tissue density and VFL, the lowest f_0 is reached at minimal tensile stress. This condition can easily be met in an excised larynx preparation where vocal folds can be adducted without being elongated. At this stage (fixed tissue density and minimal tension), VFL should be the key determinant of f_0 . Because f_0 decreases with Psub^{23,24}, the lowest attainable f_0 should then occur at the lowest pressure inducing phonation, i.e. at phonation threshold pressure (hereafter PTP). Again, an excised larynx preparation allows this to be controlled by progressively adjusting pressure until reaching PTP, where f_0 should be at a minimum and mainly dependent on the resting VFL. Thus, for a fixed tissue density and with minimal tension and Psub, measuring the minimum f_0 (hereafter $\text{min}f_0$) that a given larynx can produce is predicted by theory to be the most appropriate standardized approach to determine to what degree f_0 provides an honest indicator of body size.

We investigated specifically how well $\text{min}f_0$ and other f_0 measures are predicted by both VFL and body size across species, using individual larynges from 11 different primate species. We carried out CT-scans of excised larynges from individuals of known body size, in order to obtain anatomical estimates of VFL for each specimen, and then phonated these same specimens in an excised larynx setup under controlled conditions of Psub and minimal vocal fold tension. Because larynx size and body size are not necessarily correlated, we predicted that VFL, rather than body size, should best predict the $\text{min}f_0$ of acoustic signals. Although the decoupling of larynx size and body size has been previously discussed in primate vocal production^{5,33}, the present study is the first empirical test of the physics underlying this prediction, using a controlled *in vitro* setup and matching anatomical and acoustical measurements from the same individuals. Based on the considerable diversity found both in primate vocal signals and vocal anatomy^{5,34–36} we discuss our results in the context of evolutionary pressures that may have influenced vocal production in primates and mammals more generally.

Results

Anatomical relationship between body size and VFL across species. Ordinary least squares (OLS) regression showed a significant positive relationship between log VFL and log body size ($r^2 = 0.35$, $\beta = 1.26$, $t = 2.51$, $P = 0.03$; Fig. 1a), which was confirmed by the phylogenetic generalized least squares (PGLS) regression ($r^2 = 0.52$, $\lambda = 1.00$, $t = 3.44$, $P = 0.007$). Excluding howler monkeys (which can be suspected to be outliers in this type of regression given their highly enlarged vocal apparatus³⁷) from the analysis (Fig. 1b) did not change the nature of the significant positive relationship ($\beta = 1.26$, $t = 6.76$, $P < 0.001$) but greatly improved the fit (OLS $r^2 = 0.85$); PGLS regression was equivalent to OLS ($\lambda = 0.00$; Fig. 1b). This supports both the outlier status of howler species and the potential for decoupling between larynx size and body size across primate species.




Figure 1. Decoupling of body size and vocal fold length. Bivariate plots illustrating the relationship between the base-10 logarithms of body size and VFL: **(a)** including howler species; **(b)** excluding howler species. Black lines depict OLS regressions and red lines depict PGLS regressions.

Acoustic allometry: Prediction of $\text{min}f_0$ from body size and VFL across species. Having found this potential for decoupling between larynx and body size, we then examined the inter-specific allometric relationship between these anatomical components and the acoustic production from the same specimens. OLS regressions indicated significant negative relationships for both log body size vs. log $\text{min}f_0$ ($\beta = -1.95$, $t = -2.82$, $P = 0.02$; Fig. 2a) and log VFL vs. log $\text{min}f_0$ ($\beta = -1.31$, $t = -6.52$, $P < 0.001$; Fig. 2b). Comparison of r-squared values suggest that log VFL was a much better predictor of $\text{min}f_0$ than log body size ($r^2 = 0.81$ vs. 0.41 , respectively) (see Supplementary Table S2). The PGLS regressions supported these results, again showing significant negative relationships and that log VFL was a better predictor of $\text{min}f_0$ than log body size ($r^2 = 0.81$, $\lambda = 0.00$, $t = -6.52$, $p < 0.001$ and $r^2 = 0.53$, $\lambda = 0.59$, $t = -3.48$, $p = 0.007$, respectively; Supplementary Table S2).

Repeating these analyses while excluding howler monkeys, similar results were obtained for both comparisons, with OLS regressions indicating significant negative relationships for both log body size vs. log $\text{min}f_0$ ($\beta = -1.95$, $t = -4.11$, $P = 0.004$; Fig. 2c) and log VFL vs. log $\text{min}f_0$ ($\beta = -1.48$, $t = -4.64$, $P = 0.002$; Fig. 2d). Once again prediction of $\text{min}f_0$ by log VFL was stronger than by log body size ($r^2 = 0.72$ vs. 0.67), although the difference was considerably reduced in comparison to the regression including howlers. This suggests that the inclusion of howlers is important for, but does not fully account for, the observed advantage of using VFL to predict $\text{min}f_0$ compared to body size. The PGLS regressions excluding the two howler species were similar to their OLS counterparts, changing neither the fits nor significance levels (Supplementary Table S2).

Acoustic allometry: Prediction of $\text{mean}f_0$ and $\text{max}f_0$ from body size and VFL across species.

Parallel analyses were run using $\text{max}f_0$ and $\text{mean}f_0$ instead of $\text{min}f_0$. As for log $\text{min}f_0$, the OLS and PGLS regressions showed that both log $\text{mean}f_0$ and log $\text{max}f_0$ were better predicted by log VFL than by log body size (with all regressions being significant – all P s ≤ 0.02). For the log VFL regressions, fits for log $\text{mean}f_0$ and log $\text{max}f_0$ ($r^2 = 0.7$ and $r^2 = 0.74$, respectively) were lower than for log $\text{min}f_0$ ($r^2 = 0.81$). PGLS regressions did not change these results. The same was not true for the body size regressions, where fits with log $\text{mean}f_0$ and log $\text{max}f_0$ ($r^2 = 0.46$ and $r^2 = 0.43$, respectively) were slightly higher than log $\text{min}f_0$ ($r^2 = 0.41$). PGLS regressions provided the same conclusions despite changing the fit of these models ($r^2 = 0.65$, $r^2 = 0.56$ and $r^2 = 0.53$ for $\text{mean}f_0$, log $\text{max}f_0$ and log $\text{min}f_0$, respectively).

Excluding howler species, the results remained much the same, with OLS regressions showing better predictions for log VFL ($\text{mean}f_0$ $r^2 = 0.67$, $\text{max}f_0$ $r^2 = 0.75$) compared to log body size ($\text{mean}f_0$ $r^2 = 0.6$, $\text{max}f_0$ $r^2 = 0.56$). For the log VFL regressions, the fit for log $\text{mean}f_0$ ($r^2 = 0.67$) was lower than that for log $\text{min}f_0$ ($r^2 = 0.72$), itself lower than that for log $\text{max}f_0$ ($r^2 = 0.75$). However for the body size regressions, fits with log $\text{mean}f_0$ and log $\text{max}f_0$ ($r^2 = 0.6$ and $r^2 = 0.56$, respectively) were both lower than log $\text{min}f_0$ ($r^2 = 0.67$). PGLS regressions did not change any of the results from the analyses excluding howler species. See Supplementary Table S2 for full statistics on all of the above regressions. Inspection of the residual errors from the model including howler species confirmed our motive for running it again without howlers, as both species (*Alouatta caraya* and *Alouatta sara*) showed the highest absolute residuals in our regression.

Driving pressure: Role of P_{sub} in determining $\text{min}f_0$. A Wilcoxon signed-rank test showed that the subglottal pressure at which $\text{min}f_0$ is obtained (mean \pm SE = 9.38 ± 0.94) was significantly lower than the pressure at which $\text{max}f_0$ was obtained (mean \pm SE = 18.73 ± 2) ($Z = -3.94$, $P < 0.001$; see Supplementary Table S3 for raw data). This corroborates the expectation that f_0 is positively correlated with P_{sub} ¹⁵, and supports the approach we used, i.e using minimal subglottal pressure in order to obtain a standardized comparison of f_0 (through $\text{min}f_0$) across species.




Figure 2. Acoustic allometry from primate laryngeal specimens. Bivariate plots illustrating relationships between the base-10 logarithms of (a) body size and $\text{min}f_0$ and (b) VFL and $\text{min}f_0$, for all 11 primate species considered here; (c,d) show the same relationships excluding the 2 howler monkey species. Black lines depict OLS regressions and red lines depict PGLS regressions.

Discussion

This study is the first empirical examination of the physical and physiological factors underlying size-frequency allometry across multiple primate species. Using a sample of 11 species for which the length of the laryngeal vocal folds (ranging from 7.46 to 64.4 mm) and size of the entire body (ranging from 30 to 98 cm) was known, we recorded *in vitro* phonation in a setup that allows vocal fold tension to be kept at a minimal level while maintaining precise control over subglottal pressure. While previous conclusions have typically been drawn from averages over a large number of species and/or vocalization samples (e.g. refs^{6,7}), our approach has the advantage of investigating acoustic allometry with matching anatomical and vocal production data. This provides an unprecedented opportunity to explore the causal determinants of f_0 with a constrained interpretation of the mechanisms at work in this process.

As predicted by Morton¹², and echoing more recent findings^{6,7}, we found that calls from larger species indeed have lower f_0 , as shown by the significant negative $\text{min}f_0$ - body size correlation in our data. In agreement with theoretical predictions¹⁵, we also found that calls produced with longer vocal folds have a lower $\text{min}f_0$. Additionally, our data show that VFL is the best predictor of the minimum fundamental frequency attainable by phonation of the specimens larynges (Fig. 2a and b). PGLS analyses (that controlled for non-independence of data points due to shared ancestry of species) confirmed these results, as VFL was still, by far, the stronger predictor of $\text{min}f_0$ in these analyses (Fig. 2a and b).

In addition to documenting the moderate strength of the VFL - body size regression (Fig. 1a), these results also illustrate the considerable variability of relative laryngeal size across primate species, independent of body size. This decoupling between larynx size and overall body size can occur because laryngeal growth is not tightly constrained by the rest of the body⁵. The soft cartilaginous structure of the larynx combined with its location,

Family	Species	Common name	Sex	Body length (cm)	VFL (mm)	Min f_0 (Hz)	Epiglottis position
Atelidae	<i>Alouatta caraya</i>	Black howler	F	57	57.76	35.61	Covering
Atelidae	<i>Alouatta sara</i>	Bolivian red howler	M	65.5	64.40	27.61	Retracted
Atelidae	<i>Ateles fusciceps</i>	Black-headed spider monkey	M	53.5	20.36	81.47	Covering
Hominidae	<i>Gorilla gorilla</i>	Western gorilla	F	94	35.40	27.44	Retracted
Cercopithecidae	<i>Macaca fuscata</i>	Japanese macaque	F	72.6	17.15	91.49	Covering
Cercopithecidae	<i>Macaca silenus</i>	Lion-tailed macaque	F	53	15.70	123.81	Retracted
Cercopithecidae	<i>Macaca sylvanus</i>	Barbary macaque	F	59	14.87	185.85	Covering
Hominidae	<i>Pan troglodytes</i>	Chimpanzee	F	98	38.25	88.32	Retracted
Cercopithecidae	<i>Papio hamadryas</i>	Hamadryas baboon	M	78	25.05	106.98	Covering
Cebidae	<i>Saimiri sciureus</i>	Common squirrel monkey	M	30	7.46	658.48	Covering
Lemuridae	<i>Varecia variegata</i>	Black-and-white ruffed lemur	F	50	17.86	161.37	Retracted

Table 1. Primate species used in the study, including specimen sex, body size (from anatomical measurements), vocal fold length (estimated from CT-scan measurements) and min f_0 values (from excised larynx experiments); epiglottis position when min f_0 was obtained is also indicated for each species.

connected loosely in primates via muscles and ligaments to the skull, jaw and sternum, allows its independent growth during development⁵, potentially influenced by hormone levels or other size-independent factors³⁸. This peculiar anatomical independence can allow larynx size to be sensitive to various selective pressures that may differ from those acting on body size.

Ecological factors are among the potential selective forces acting on vocalization frequency. For example, species-specific habitat could have fine-tuned laryngeal anatomy by favorably selecting vocalizations produced within a certain frequency range (the ‘acoustic adaptation hypothesis’³⁹). Hauser²⁰ suggested that the lower frequency range found for *Macaca silenus* could be one such example, as the tropical rainforest home to this species might impair the propagation of higher frequencies⁴⁰. Although our analyses focus on min f_0 rather than complete frequency range, this suggestion is supported by the fact that a species of relatively similar size (*Macaca sylvanus*), inhabiting more open habitats⁴¹, has a min f_0 50% higher than that of *M. silenus* (185.85 Hz and 123.81 Hz, respectively). While the propagation of these two frequencies *per se* might not differ much in tropical and open habitats, the apparent predisposition of the *M. sylvanus* larynx to produce higher-pitched vocalizations is worthy of further investigation and environmental propagation experiments.

Second, species-specific socio-ecology also has the potential to influence laryngeal anatomy independently from body size, so that it better suits the requirements of a given species’ vocal communication system⁴². The apes included in this study provide an illustrative case of this possibility: despite being very close in terms of measured body length (94 cm for the female gorilla, 98 cm for the female chimpanzee), and vocal fold length (38.25 mm vs. 35.4 mm, respectively), min f_0 in the chimpanzee was over 3 times higher than that of the gorilla (88.32 Hz and 27.44 Hz, respectively; Table 1). Structural aspects of vocal fold composition differ between these two species⁴³, and such histological differences may result from selection for different communicative needs and call usage inherent to these species’ social systems. Chimpanzees live in fission-fusion systems⁴⁴ and vocalize mostly in long-distance communicative contexts using loud, high-frequency pant-hoots⁴⁵. Gorillas, on the other hand, live in more cohesive social groups⁴⁶ and typically vocalize at closer range mostly using low frequency grunts^{45,46}. A vocal fold structure suitable for higher-frequency call production in chimpanzees and lower frequency call production in gorillas could thus contribute to explaining why the theoretically-predicted correlation between min f_0 and VFL does not lead to similar observations in these close relatives of humans. Additional histological data would be required to evaluate this hypothesis, focusing for instance on vocal fold elasticity as this parameter has been shown to affect f_0 ³⁰.

Finally, sexual selection is an evolutionary force for which there is already some evidence of an influence on laryngeal growth, leading to a decoupling of larynx size from overall body size⁶. Howler monkeys provide one of the most drastic example of hypertrophied vocal apparatus^{37,47} and thus have disproportionately low frequency vocalizations^{33,42}. However, males howlers’ larynges and hyoids are enlarged to a much greater extent than those of females⁴². As outlined in a recent study⁴⁸, mating systems appear to strongly influence f_0 dimorphism in anthropoid primates, including humans⁴⁹. Appropriate playback experiments⁵⁰ will be necessary to investigate the effect of acoustic traits in howlers vocalizations that are potentially relevant to sexual partners and/or competitors, as previously done in other species (e.g. refs^{51–53}). Size exaggeration often occurs via behavioral/anatomical adaptations affecting formants⁶. However, howler laryngeal hypertrophy affects both vocal fold length and vocal tract morphology (as air sacs fill the enlarged thyro-hyoid apparatus and may act as a resonance chamber^{37,54}). This explains the abnormally low f_0 and formants that characterize howler species vocalizations⁴², given that their vocal folds and vocal tracts are considerably larger than those of similarly sized primates (e.g., macaques). In this context, it appears evident that howler vocalizations do not provide honest signals about the size of the caller when making across taxa comparisons. However, similar to red deer vocal tract elongation during roaring⁵⁵, or koala descended larynx for bellowing¹⁹, howler vocalizations are likely to provide a case of honest signalling when considering vocal production within the species⁴²; again, answering this will require further research, combining anatomical investigation and playback of resynthesized signals.

Because our larynges were from dead animals, all oscillations observed were generated by passive airflow. It must be noted that such conditions do not necessarily reproduce the lowest possible f_o s: vocal fold oscillation may in a few cases be induced by active contraction of the vocal fold musculature (the so-called “active” theory of phonation, as apparently applicable to cat purring⁵⁶). Because of limits on the rate of muscular contraction, active phonation is only expected to be possible for f_o s below ~40 Hz^{34,57}. An exception is the superfat laryngeal muscles of bats, which are specially adapted to contract up to 180 times a second in some species⁵⁸. Aside from cat purring, however, mammalian phonation is typically generated without periodic muscular contraction at each f_o period, and instead produced by the airflow passing through the glottis (the myoelastic-aerodynamic theory (MEAD; refs^{28,32,59}). The excised larynx methodology applied here illustrates the generality of the MEAD principle, broadening the range of primate species to which it can be applied. For example, our experiments show that the MEAD principle is sufficient to explain the low f_o s of the two howler species we investigated, as we were able to get low frequencies for these species (*A. caraya*: 35.61 Hz; *A. sara*: 25.42 Hz), comparable to those observed in their natural vocalizations⁴², entirely with passive phonation. While not definitely ruling out the possibility of active phonation in our species, this suggests that no active contraction of the laryngeal muscles is required to produce the low-frequency calls typical of howler species.

Based on theoretical predictions and the reasoning outlined in the introduction, we used $\text{min}f_o$ as our standard frequency measure. With our setup, we had no means of controlling whether the higher end of our applied pressure range (on which $\text{max}f_o$, and thus $\text{mean}f_o$, theoretically depend) was physiologically relevant (i.e., matched what the living animal is capable of; pressure measurements, via tracheal catheter, would be necessary to determine this). $\text{Min}f_o$ is therefore the only frequency measure obtained with clear boundary conditions, and thus that can reasonably be assumed to have physiological relevance. Given this, it is intriguing that all f_o measures ($\text{min}f_o$, $\text{mean}f_o$, $\text{max}f_o$) were significantly negatively related with VFL, and that $\text{min}f_o$ had a weaker correlation with body size than $\text{mean}f_o$ and $\text{max}f_o$ (Supplementary Table S2). This highlights the importance of caution while conducting acoustic allometry research based on non-controlled acoustic data from living animals; it is crucial to avoid false positives by broadly sampling the vocalization types utilized in the analyses.

Two further points support the use of $\text{min}f_o$ and minimal P_{sub} in future analyses of this sort: 1) the fit of regressions between VFL and $\text{max}f_o$ (as well as VFL and $\text{mean}f_o$) were not as good as those between VFL and $\text{min}f_o$ (Supplementary Table S2) and 2) P_{sub} at $\text{min}f_o$ was significantly lower than P_{sub} at $\text{max}f_o$. In-depth investigation of the $P_{\text{sub}}-f_o$ relationship would be valuable, but is beyond the scope of this study. Such investigations require extreme caution, as preservation of the biomechanical properties of vocal fold tissue (e.g. viscoelasticity) may be affected by the time elapsed between death and specimen freezing, which in turn may have affected PTP⁶⁰ and potentially altered the resulting $\text{min}f_o$. While we acknowledge this potential limitation in our protocol, we point out (1) the difficulty of gathering such a collection of specimens: for many species it is rare to be able to acquire even a single specimen within a time span of a decade or more (e.g. apes or howler monkeys) and (2) that given the systematic variation observed in the relationship between VFL and $\text{min}f_o$, the data collected here offer unique insights into the physical determinant of vocal frequency and the decoupling between larynx and body size in a representative sample of primates. We would also like to call attention to the fact that, although our selection of larynges was based on specimen availability and thus included either male or female larynges, the large range of body sizes observed across the species considered here should limit the impact of any potential sexual dimorphism on our results. However, given the more limited size variation, along with the potential for size dimorphism, within a species, future studies investigating the decoupling between larynx and body size at an intra-specific level should ensure the selection of larynges from the same sex.

Ultimately, by adopting a novel allometric approach, our study both confirms the theoretical prediction that vocal fold length is a main determinant of f_o , and provides evidence that laryngeal growth is not tightly constrained by overall body size (at least in the primate species investigated here). Our results call attention to the considerable anatomical variation across species that can be observed in primate vocal production systems, most of which has barely been investigated. As illustrated by the hypotheses we raise, this variability offers great potential for future in-depth studies of how various selective pressures may have driven diversity in vocal production and anatomy in primates and other mammals. Further work using excised larynx systems like that described here are critical to improving our understanding of mammalian vocal production mechanisms, and thus of the functions of mammal vocal communication viewed from an evolutionary perspective.

Methods

Data collection. *Anatomical specimens.* As part of the specimen acquisition program at the Department of Natural Sciences, National Museums Scotland, the remains of deceased European zoo animals are regularly collected and processed. Our larynges came from these zoo specimens and all samples came from animals that had died of natural causes. For each individual, body size was measured as the distance between the ischium of the pelvis and the top of the skull (head-body length, without tail). Body length was preferred over body weight because: (1) obesity is a potential problem in zoo animals⁶¹; (2) bodies can dehydrate once deceased⁶² making post mortem weight dependent on measurement delay; and (3) weight data could not be obtained for some of our specimens. Larynges from the cadavers of 11 individuals, each of a different primate species, were excised, frozen and stored at -20° Celsius at the National Museums Scotland before being shipped to the Department of Cognitive Biology, University of Vienna (see Supplementary Text for additional information on the freezing method). Each larynx was then thawed, cleaned, inspected, photographed and measured in preparation for X-ray CT scanning, after which they were refrozen and stored at -20° C. The primates used for this study were chosen to represent a wide range of body sizes and phylogenetic diversity (Fig. 3 and Table 1).

CT scans. Two procedures were applied, depending on the size of the specimen: the larynx of the smallest species (squirrel monkey, *Saimiri sciureus*.) was scanned using micro CT, while ordinary CT was used for the




Figure 3. Consensus tree of phylogenetic distances among the species examined in this study. Tree based on a combination of 2 to 16 DNA sequences among 11 mitochondrial and 6 autosomal genes retrieved from mitochondrial and autosomal data available from *10kTrees*⁷¹, version 3 at <http://10ktrees.nunn-lab.org/project.html>; see Supplementary Table S1.

other 10 larynges. All CT scans were performed at the University of Veterinary Medicine Vienna. Macro CT scans were made using a Siemens SOMATOM Emotion helical CT-scanner (Siemens AG, Munich, Germany), and the micro-CT scan was made using an Xradia microXCT-400 (0.4x lens; Carl Zeiss X-ray Microscopy, Pleasanton, CA). For macro-CT scans, specimens were positioned in ventral recumbancy on X-ray-transparent styrofoam plates and scanned frozen. Scanning parameters were adjusted to specimen size, using 110–130 kV source voltage and 80–110 mA beam intensity. Reconstructed image slices measured 512×512 pixels. Depending on larynx size, the dimensions of reconstructed voxels varied between $238\text{--}340 \mu\text{m}^2$ in the xy plane and $200\text{--}500 \mu\text{m}$ in the z plane. Due to its small size and longer scanning time, the *Saimiri* specimen was thawed prior to micro-CT scanning and mounted vertically inside a sealed Falcon tube, the bottom of which was partially filled with phosphate-buffered saline to prevent dehydration. The specimen was scanned at 40 keV source voltage and $200 \mu\text{A}$ beam intensity. Reconstructed slices measured 512×512 pixels and the voxel resolution of reconstructed volumes was $35 \mu\text{m}^3$.

Excised larynx experiments. A detailed description of the setup used in this study has been given elsewhere⁶³. Before use in excised larynx experiments (Table 1) each specimen was thawed, then prepared by removing excess tissue and tracheal rings, before being mounted on a vertical subglottic tube. The tube diameter was adjusted to match specimen size such that an airtight seal was formed with the trachea. Larynx stability and support were ensured using a combination of adjustable plastic support structures (made of LEGO blocks, Billund, Denmark) and custom-made 3D-printed plastic mounts placed on the left lateral, right lateral and anteriorly sides of the larynx.

Phonation was obtained by passing a controlled flow of warm ($\sim 37^\circ\text{C}$) humid (100%) air through the mounted larynx. Vocal folds were adducted using 2 manually controlled micromanipulators (Warzhauser MM33, Tamm, Germany) mounted on a tilting platform. For standardization purposes, the degree of adduction was fixed when phonation could be reliably induced with minimal airflow and tension on the vocal folds, and attained a steady phonation (assessed by ear and via examination of the electroglottographic (EGG) signal during the experiment). Custom-made copper EGG electrodes were placed on both sides of the thyroid cartilage, at the level of the vocal folds, for an optimal recording of vocal fold vibrations. Psub was controlled using “ELLApp” software (created in Python by CTH). Acoustic, EGG and sound intensity were recorded using a DPA 4061 omnidirectional microphone (positioned at a variable but known distance from the vocal folds), a Glottal Enterprises EG 2-1000 two-channel electroglottograph (lower cutoff-frequency 2 Hz) and an NL-52 RION sound pressure level-meter (located 30 cm from the vocal folds; settings ‘fast acquisition’ and ‘dB C’ weighting), respectively. All signals were acquired, synchronized and stored within ELLApp.

Phonation and data acquisition followed an adjustable computer-controlled sequence. Pressure sweeps were applied to each excised larynx, consisting of a slow linear increase in Psub followed by a slow linear decrease of the same duration; the lowest Psub value was set just below the PTP, and the highest value varied with specimen size. Each larynx was exposed to 4–8 pressure sweeps, 2–4 with the epiglottis covering the airway and 2–4 with the epiglottis retracted. The aim of epiglottis manipulation was to evaluate whether a source-filter interaction (so-called “feedback” system, refs^{5,64}) exists between the vibrating vocal folds and what is left of the vocal tract in our setup, i.e. the space between the glottis and the epiglottis. The number of sweeps was chosen to allow us to evaluate repeatability of acoustic production while avoiding damage or drying of the sound source. Throughout the experiments, larynges were kept moist using a spray-bottle containing saline solution (0.9% NaCl).




Figure 4. Isosurface of large and small laryngeal specimens. Panel a (*Alouatta sara*) and panel b (*Macaca fuscata*) show the homologous landmarks used to establish the vocal fold length proxy. L1: Dorsal apical cricoid; L2: Ventral basal thyroid; L3: Ventral apical thyroid. VF: segment used as the skeletal proxy for vocal fold length (not to scale).

Data analysis. *Anatomical measurements: CT scans.* Both macro and micro CT data were analyzed using AMIRA software (version 5.6.0). Along their length, the vocal folds are composed of a membranous and a cartilaginous section⁶⁵. Soft tissue geometry is difficult to visualize in CT, and direct measurement of VFL would have required tissue destruction and perhaps histology to be accurately determined. Thus our aim was to obtain clear 3D visualization of the laryngeal cartilages (hyoid bone, thyroid, cricoid and arytenoid cartilages⁶⁶) to estimate the total vocal fold length (membranous + cartilaginous length). After creating an isosurface model of these cartilages, VFL was estimated based on measurement of homologous landmarks placed at the intersection of the mid-sagittal plane and the cricoid and thyroid cartilages (Fig. 4). The most dorsal point for our vocal fold estimate was placed at the apex of the cricoid cartilage on the midline. The most ventral possible attachment point of the vocal folds was defined as the midpoint between the apex and the base of the thyroid cartilage (as the thyroid attachment of the vocal fold could not be consistently visualized from the CT data).

Signal analysis. The analysis of f_0 from acquired signals was conducted using the autocorrelation function in Praat⁶⁷, and in ELLApp. After synchronization of the various input signals in ELLApp, EGG signals were annotated in Praat and f_0 was extracted with appropriately adjusted settings (Praat function ‘To Pitch (ac)...’, creating a Praat ‘PitchTier’ object; see Supplementary Text for details). Settings were adjusted both relying on visual inspection of the spectrograms (to identify and exclude non-periodic regimes; time step was automatically computed as $0.75/\text{pitch_floor}$, which varied between 20 Hz and 620 Hz depending on the pressure sweep and species analyzed) and of the waveform (to further identify and exclude ambiguous nonlinear phenomena like subharmonics^{68,69}). The raw f_0 data produced by Praat (termed “PitchTier” in Praat) was then edited manually in order to exclude any pitch-tracking errors made by the automated f_0 extraction algorithm. We excluded all parts of the signals characterized by non-periodic oscillation of the vocal folds (with periodicity defined as a minimum of 10 regular consecutive vocal fold oscillatory cycles), as well as those regions where Praat’s automated calculation did not correspond to the lowest partial visible in the spectrogram and/or the main oscillation in the waveform. The minimum, maximum and mean f_0 ($\text{min}f_0$, $\text{max}f_0$ and $\text{mean}f_0$, respectively) were queried from this corrected pitch object based on the remaining annotated sections, using the Praat ‘Get minimum...’, ‘Get maximum...’ and ‘Get mean...’ functions.

Using the calibrated data from ELLApp, we also extracted the P_{sub} values obtained at $\text{min}f_0$ and $\text{max}f_0$ to evaluate the effect of P_{sub} on f_0 , and verify our approach of using minimal tension and P_{sub} to attain $\text{min}f_0$.

Statistics. Following assessment of data normality using Shapiro-Wilk tests, body size, VFL and $\text{min}f_0$ were log-transformed (base 10; see raw data values Table 1) and the following OLS linear regressions (i.e. standard linear regression models) were computed: $\log \text{min}f_0$ vs. \log body size, $\log \text{min}f_0$ vs. \log VFL, and \log body size vs. \log VFL. Additionally, due to the potential influence of species relatedness, PGLS regressions (which accounts for the potential non-independence of data points due to shared phylogenetic history; see ref.⁷⁰) were also computed on the same set of variables, controlling for the potential effects of phylogenetic covariance⁷⁰, using the consensus phylogenetic tree shown in Fig. 3 (created using *10kTrees*⁷¹, version 3 (<http://10ktrees.nunn-lab.org/project.html>)).

In order to evaluate the validity of focusing on $\min f_0$, for comparative purposes we conducted the same set of analyses on $\max f_0$ and $\text{mean} f_0$. Because howler monkeys are clear anatomical outliers due to their greatly enlarged vocal apparatus^{37,42,47}, we also conducted the same set of analyses omitting the two howler specimens to evaluate whether our results were solely driven by these extreme cases. Finally, ‘Sex’ and ‘Epiglottis’ variables were included either alone or together in preliminary linear and PGLS models, but showed no significant effect in any of all the possible combinations. These two variables were thus omitted from subsequent analyses.

To evaluate the effect of subglottal pressure on f_0 , P_{sub} values at $\min f_0$ and $\max f_0$ were compared using a Wilcoxon signed-rank test.

All statistics and computations were done in R⁷² using the ‘lm’ function for OLS regressions and the ‘ppls’ function (‘caper’ package) for PGLS regressions⁷³. Two-tailed P-values are reported and significance level is set at 0.05.

References

- Zahavi, A. The cost of honesty (further remarks on the handicap principle). *J. Theor. Biol.* **67**, 603–605 (1977).
- Krebs, J. R. & Dawkins, R. In *Behavioural Ecology: an evolutionary approach*. ((eds J. R. Krebs & N. B. Davies) 380–402 (Blackwell Scientific Publications, 1984).
- Johnstone, R. A. Sexual selection, honest advertisement and the handicap principle: reviewing the evidence. *Biol. Rev. (Camb.)* **70**, 1–65 (1995).
- Taylor, A. M. & Reby, D. The contribution of source-filter theory to mammal vocal communication research. *J. Zool.* **280**, 221–236, <https://doi.org/10.1111/j.1469-7998.2009.00661.x> (2010).
- Fitch, W. T. & Hauser, M. D. Vocal production in nonhuman primates: acoustics, physiology, and functional constraints on “honest” advertisement. *Am J Prim* **37**, <https://doi.org/10.1002/ajp.1350370303> (1995).
- Charlton, B. D. & Reby, D. The evolution of acoustic size exaggeration in terrestrial mammals. *Nat. Commun.* **7**, 12739, <https://doi.org/10.1038/ncomms12739> (2016).
- Bowling, D. *et al.* Body size and vocalization in primates and carnivores. *Scientific reports* **7**, 41070 (2017).
- Reby, D. & McComb, K. Anatomical constraints generate honesty: acoustic cues to age and weight in the roars of red deer stags. *Anim. Behav.* **65**, 519–530, <https://doi.org/10.1006/anbe.2003.2078> (2003).
- Koda, H. *et al.* Soprano singing in gibbons. *Am. J. Phys. Anthropol.* **149**, 347–355, <https://doi.org/10.1002/ajpa.22124> (2012).
- Koda, H., Tokuda, I. T., Wakita, M., Ito, T. & Nishimura, T. The source-filter theory of whistle-like calls in marmosets: Acoustic analysis and simulation of helium-modulated voices. *J. Acoust. Soc. Am.* **137**, 3068–3076, <https://doi.org/10.1121/1.4921607> (2015).
- Fant, G. Acoustic theory of speech production. (Mouton, 1960).
- Morton, E. S. On the occurrence and significance of motivation-structural rules in some bird and mammal sounds. *Am. Nat.* **111**, 855–869 (1977).
- Ohala, J. J. An ethological perspective on common cross-language utilization of F0 of voice. *Phonetica* **41**, 1–16 (1984).
- Harrison, D. F. N. *The anatomy and physiology of the mammalian larynx*. (Cambridge University Press, 1995).
- Titze, I. R. *Principles of voice production*. (National Center for Voice and Speech (2nd Edition), 2000).
- Masataka, N. Lack of correlation between body size and frequency of vocalizations in young female Japanese macaques (*Macaca fuscata*). *Folia Primatol. (Basel)* **63**, 115–118 (1994).
- Pfefferle, D., West, P. M., Grinnell, J., Packer, C. & Fischer, J. Do acoustic features of lion, *Panthera leo*, roars reflect sex and male condition? *J. Acoust. Soc. Am.* **121**, 3947–3953, <https://doi.org/10.1121/1.2722507> (2007).
- Pisanski, K. *et al.* Vocal indicators of body size in men and women: a meta-analysis. *Anim. Behav.* **95**, 89–99, <https://doi.org/10.1016/j.anbehav.2014.06.011> (2014).
- Charlton, B. D. *et al.* Cues to body size in the formant spacing of male koala (*Phascolarctos cinereus*) bellows: honesty in an exaggerated trait. *J. Exp. Biol.* **214**, 3414–3422, <https://doi.org/10.1242/jeb.061358> (2011).
- Hauser, M. D. The evolution of nonhuman primate vocalizations: effects of phylogeny, body weight, and social context. *Am. Nat.* **528**–542 (1993).
- Gingras, B., Boeckle, M., Herbst, C. T. & Fitch, W. T. Call acoustics reflect body size across four clades of anurans. *J. Zool.* **289**, 143–150, <https://doi.org/10.1111/j.1469-7998.2012.00973.x> (2013).
- Ladefoged, P. & McKinney, N. P. Loudness, sound pressure, and subglottal pressure in speech. *J. Acoust. Soc. Am.* **35**, 454–460, <https://doi.org/10.1121/1.1918503> (1963).
- Titze, I. R. On the relation between subglottal pressure and fundamental frequency in phonation. *J. Acoust. Soc. Am.* **85**, 901–906 (1989).
- Titze, I. R. Comments on the myoelastic - aerodynamic theory of phonation. *J. Speech Hear. Res.* **23**, 495–510 (1980).
- Titze, I. R. Vocal fold mass is not a useful quantity for describing F0 in vocalization. *J. Speech Lang. Hear. Res.* **54**, 520–522, [https://doi.org/10.1044/1092-4388\(2010/09-0284\)](https://doi.org/10.1044/1092-4388(2010/09-0284)) (2011).
- Hirano, M. Morphological structure of the vocal cord as a vibrator and its variations. *Folia Phoniatrica et Logopaedica* **26**, 89–94 (1974).
- Kurita, S., Nagata, K. & Hirano, M. In *Vocal fold physiology: Contemporary research and clinical issues* (eds D. M. Bless & J. H. Abbs) 3–21 (College Hill, 1983).
- Herbst, C. T. In *Vertebrate sound production and acoustic communication* (eds A. Roderick Suthers, Tecumseh W. Fitch, R. Richard Fay & N. Arthur Popper) 159–189 (Springer International Publishing, 2016).
- Alipour, F., Jaiswal, S. & Vigmstad, S. Vocal fold elasticity in the pig, sheep, and cow larynges. *J. Voice* **25**, 130–136, <https://doi.org/10.1016/j.jvoice.2009.09.002> (2011).
- Alipour-Haghighi, F. & Titze, I. R. Elastic models of vocal fold tissues. *J. Acoust. Soc. Am.* **90**, 1326–1331 (1991).
- Titze, I., Riede, T. & Mau, T. Predicting achievable fundamental frequency ranges in vocalization Across species. *PLoS Comput. Biol.* **12**, e1004907, <https://doi.org/10.1371/journal.pcbi.1004907> (2016).
- Titze, I. R. & Alipour, F. *The myoelastic aerodynamic theory of phonation*. (National Center for Voice and Speech, 2006).
- Fitch, W. T. & Hauser, M. D. In *Acoustic Communication* (ed. A. M. Ed. Simmons, Fay, R. R. & Popper, A. N.) 65–137 (Springer, 2002).
- Fitch, W. T. In *Primate audition: ethology and neurobiology*. Frontiers in Neuroscience. (CRC Press, 2002).
- Brown, C. H. In *Primate audition: ethology and neurobiology*. Frontiers in Neuroscience. (CRC Press, 2002).
- Hammerschmidt, K. & Fischer, J. In *The evolution of communicative creativity: From fixed signals to contextual flexibility*. 93–119 (The MIT Press, 2008).
- Schön, M. A. The anatomy of the resonating mechanism in howling monkeys. *Folia Primatol. (Basel)* **15**, 117–132 (1971).
- Fitch, W. T. & Giedd, J. Morphology and development of the human vocal tract: a study using magnetic resonance imaging. *J. Acoust. Soc. Am.* **106**, 1511–1522 (1999).
- Ey, E. & Fischer, J. The “Acoustic Adaptation Hypothesis”—a review of the evidence from birds, anurans and mammals. *Bioacoustics* **19**, 21–48, <https://doi.org/10.1080/09524622.2009.9753613> (2009).
- Ken, M., Douglas, Q. & Marler, P. Sound transmission and its significance for animal vocalization: II. Tropical forest habitats. *Behav. Ecol. Sociobiol.* **2**, 291–302 (1977).

41. Fa, J. E. Habitat distribution and habitat preference in Barbary macaques (*Macaca sylvanus*). *International Journal of Primatology* **5**, 273–286, <https://doi.org/10.1007/bf02735762> (1984).
42. Dunn, J. C. *et al.* Evolutionary trade-off between vocal tract and testes dimensions in Howler monkeys. *Curr. Biol.* **25**, 2839–2844, <https://doi.org/10.1016/j.cub.2015.09.029> (2015).
43. Kelemen, G. The anatomical basis of phonation in the chimpanzee. *J. Morphol.* **82**, 229–256, <https://doi.org/10.1002/jmor.1050820205> (1948).
44. Lehmann, J. & Boesch, C. To fission or to fusion: effects of community size on wild chimpanzee (*Pan troglodytes verus*) social organisation. *Behav. Ecol. Sociobiol.* **56**, 207–216, <https://doi.org/10.1007/s00265-004-0781-x> (2004).
45. Marler, P. In *Growing points in ethology* (eds P. P. G. Bateson & R. A. Hinde) (Cambridge U Press, 1976).
46. Harcourt, A. H., Stewart, K. J. & Hauser, M. Functions of wild Gorilla ‘close’ calls. I. repertoire, context, and interspecific comparison. *Behaviour* **124**, 89–122 (1993).
47. Schön Ybarra, M. Morphological adaptations for loud phonation in the vocal organ of howling monkeys. *Primate Rep* **22**, 19–24 (1988).
48. Puts, D. A. *et al.* Sexual selection on male vocal fundamental frequency in humans and other anthropoids. *Proc. Biol. Sci.* **283**, <https://doi.org/10.1098/rspb.2015.2830> (2016).
49. Pisanski, K. *et al.* Volitional exaggeration of body size through fundamental and formant frequency modulation in humans. *Scientific Reports* **6**, 34389, <https://doi.org/10.1038/srep34389> (2016).
50. Fischer, J., Noser, R. & Hammerschmidt, K. Bioacoustic field research: A primer to acoustic analyses and playback experiments with primates. *Am. J. Primatol.* **75**, 643–663, <https://doi.org/10.1002/ajp.22153> (2013).
51. Mitani, J. C. Sexual selection and adult male orangutan long calls. *Anim. Behav.* **33**, 272–283, [https://doi.org/10.1016/S0003-3472\(85\)80141-X](https://doi.org/10.1016/S0003-3472(85)80141-X) (1985).
52. Charlton, B. D., Ellis, W. A. H., Brumm, J., Nilsson, K. & Fitch, W. T. Female koalas prefer bellows in which lower formants indicate larger males. *Anim. Behav.* **84**, 1565–1571, <https://doi.org/10.1016/j.anbehav.2012.09.034> (2012).
53. Reby, D. *et al.* Red deer stags use formants as assessment cues during intrasexual agonistic interactions. *Proc. R. Soc. Lond. B* **272**, 941–947, <https://doi.org/10.1098/rspb.2004.2954> (2005).
54. de Boer, B. Acoustic analysis of primate air sacs and their effect on vocalization. *J. Acoust. Soc. Am.* **126**, 3329–3343, <https://doi.org/10.1121/1.3257544> (2009).
55. Fitch, W. T. & Reby, D. The descended larynx is not uniquely human. *Proc. Biol. Sci.* **268**, 1669–1675, <https://doi.org/10.1098/rspb.2001.1704> (2001).
56. Remmers, J. E. & Gautier, H. Neural and mechanical mechanisms of feline purring. *Respir. Physiol.* **16**, 351–361, [https://doi.org/10.1016/0034-5687\(72\)90064-3](https://doi.org/10.1016/0034-5687(72)90064-3) (1972).
57. Hirose, H., Ushijima, T., Kobayashi, T. & Sawashima, M. An experimental study of the contraction properties of the laryngeal muscles in the cat. *Ann. Otol. Rhinol. Laryngol.* **78**, 297–306 (1969).
58. Elemans, C. P. H., Mead, A. F., Jakobsen, L. & Ratcliffe, J. M. Superfast muscles set maximum call rate in echolocating bats. *Science* **333**, 1885–1888, <https://doi.org/10.1126/science.1207309> (2011).
59. van den Berg, J. Myoelastic-aerodynamic theory of voice production. *J. Speech Lang. Hear. Res.* **1**, 227–244, <https://doi.org/10.1044/jshr.0103.227> (1958).
60. Chan, R. W. & Titze, I. R. Dependence of phonation threshold pressure on vocal tract acoustics and vocal fold tissue mechanics. *J. Acoust. Soc. Am.* **119**, 2351–2362, <https://doi.org/10.1121/1.2173516> (2006).
61. Schwitzer, C. & Kaumanns, W. Body weights of ruffed lemurs (*Varecia variegata*) in European zoos with reference to the problem of obesity. *Zoo Biol.* **20**, 261–269, <https://doi.org/10.1002/zoo.1026> (2001).
62. Aturaliya, S. & Lukasewycz, A. Experimental forensic and bioanthropological aspects of soft tissue taphonomy: 1. Factors influencing postmortem tissue desiccation rate. *J. Forensic Sci.* **44**, 893–896 (1999).
63. Herbst, C. T. *et al.* Glottal opening and closing events investigated by electroglottography and super-high-speed video recordings. *J. Exp. Biol.* **217**, 955–963, <https://doi.org/10.1242/jeb.093203> (2014).
64. Flanagan, J. L. Source-system interaction in the vocal tract. *Ann. N. Y. Acad. Sci.* **155**, 9–17, <https://doi.org/10.1111/j.1749-6632.1968.tb56744.x> (1968).
65. Busuttil, A., Davis, B. C. & Maran, A. G. The soft tissue/cartilage relationship in the laryngeal glottis. *J. Laryngol. Otol.* **95**, 385–391 (1981).
66. Storck, C. *et al.* Developing a 3D model of the laryngeal cartilages using HRCT data and MIMICS’s segmentation software. *Logoped. Phoniater. Vocol.* **35**, 19–23, <https://doi.org/10.3109/14015430903552378> (2010).
67. Praat: doing phonetics by computer [Computer program]. v. Version 5.4.01 (University of Amsterdam, The Netherlands, Retrieved from <http://www.praat.org/> 2014).
68. Fitch, W. T., Neubauer, J. & Herzog, H. Calls out of chaos: the adaptive significance of nonlinear phenomena in mammalian vocal production. *Anim. Behav.* **63**, 407–418, <https://doi.org/10.1006/anbe.2001.1912> (2002).
69. Herzog, H., Berry, D., Titze, I. R. & Saleh, M. Analysis of vocal disorders with methods from nonlinear dynamics. *J. Speech Hear. Res.* **37**, 1008–1019 (1994).
70. Symonds, M. R. E. & Blomberg, S. P. In *Modern phylogenetic comparative methods and their application in evolutionary biology: concepts and practice*. (ed Zsolt László Garamszegi) 105–130 (Springer Berlin Heidelberg, 2014).
71. Arnold, C., Matthews, L. J. & Nunn, C. L. The 10kTrees website: A new online resource for primate phylogeny. *Evol. Anthropol.* **19**, 114–118, <https://doi.org/10.1002/evan.20251> (2010).
72. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. <http://www.R-project.org/> (2015).
73. Caper: Comparative Analyses of Phylogenetics and Evolution in R. v. R package version 0.5.2 (2013).

Acknowledgements

We thank Andrew Kitchener and Georg Hantke of the Department of Natural Sciences, National Museums Scotland for access to the specimens used in this study, and Riccardo Hofer for help creating the excised larynx system, and for designing and 3D-printing custom-made laryngeal holders. CTH’s contribution was funded by an APART grant awarded by the Austrian Academy of Sciences. DLB’s contribution was funded by a Lisa Meitner fellowship (M1773) from the Austrian Science Fund (FWF). WTF acknowledges the support of ERC Advanced Grant SOMACCA (AdG #230604) in the development of the excised larynx setup.

Author Contributions

M.G., C.T.H., D.L.B. and W.T.F. conceived and designed the study; M.G., D.L.B. and W.T.F. wrote the manuscript, C.T.H. and J.C.D. helped writing the manuscript; M.G., C.T.H., D.L.B. and W.T.F. collected the data; M.G. and J.C.D. conducted the analyses, D.L.B. helped conduct the analyses.

Additional Information

Supplementary information accompanies this paper at <https://doi.org/10.1038/s41598-017-11000-x>.

Competing Interests: The authors declare that they have no competing interests.

Change History: A correction to this article has been published and is linked from the HTML version of this paper. The error has not been fixed in the paper.

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2018