










Original Article
Pediatrics



Effects of Sedation Performed by an Anesthesiologist on Pediatric Endoscopy: a Single-Center Retrospective Study in Korea

Sung Min Yang ¹, Dae Yong Yi ^{1,2}, Geun Joo Choi ³, In Seok Lim ^{1,2},
Soo Ahn Chae ^{1,2}, Sin Weon Yun ^{1,2}, Na Mi Lee ^{1,2}, Su Yeong Kim ¹ and
Eung Sang Choi ⁴

¹Department of Pediatrics, Chung-Ang University Hospital, Seoul, Korea

²Department of Pediatrics, Chung-Ang University College of Medicine, Seoul, Korea

³Department of Anesthesiology, Chung-Ang University Hospital, Chung-Ang University College of Medicine, Seoul, Korea

⁴Department of Pediatrics, Hanil General Hospital, Seoul, Korea



Received: Nov 27, 2019

Accepted: Mar 24, 2020

Address for Correspondence:

Dae Yong Yi, MD, PhD

Division of Pediatric Gastroenterology and Hepatology, Department of Pediatrics, Chung-Ang University Hospital, Chung-Ang University College of Medicine, 102 Heukseok-ro, Dongjak-gu, Seoul 06973, Korea.
E-mail: meltemp2@hanmail.net

© 2020 The Korean Academy of Medical Sciences.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID iDs

Sung Min Yang 


<https://orcid.org/0000-0001-6057-3458>

Dae Yong Yi 


<https://orcid.org/0000-0002-4168-7131>

Geun Joo Choi 


<https://orcid.org/0000-0002-4653-4193>

In Seok Lim 


<https://orcid.org/0000-0002-7047-7627>

Soo Ahn Chae 

<https://orcid.org/0000-0002-9825-8194>

Sin Weon Yun 

<https://orcid.org/0000-0001-8947-703X>

Na Mi Lee 

<https://orcid.org/0000-0003-0580-8677>

<https://jkms.org>

ABSTRACT


Background: Endoscopy is used for diagnosing and treating various digestive diseases in children as well as in adults. However, in pediatric patients, it is recommended that sufficient sedation should be ensured before conducting endoscopy, since insufficient sedation may cause serious complications. However, in Korea, no studies have yet described the types of sedation drugs, effects of sedation, and efficiency of endoscopy with respect to the sedation instructor. Thus, we investigated the effectiveness of sedative procedures performed by anesthesiologists.

Methods: We retrospectively reviewed the medical records of patients aged < 18 years who underwent endoscopy during March 2014–July 2019. Data of sedation instructors, sedation drugs and their doses, complications, and the recovery after sedation were evaluated.

Results: Of 257 patients, 217 underwent esophagogastroduodenoscopy (EGD) and 40 underwent colonoscopies. Before EGD, 29 patients (13.4%) underwent sedation by the pediatric endoscopist and 188 (86.6%) were sedated by the anesthesiologist. The anesthesiologist performed the sedation for all 40 patients who underwent colonoscopy. Endoscopic examinations performed by the anesthesiologist were relatively more time-consuming (401.0 ± 135.1 seconds vs. 274.9 ± 106.1 seconds, $P < 0.001$). We observed that in patients who underwent EGD, there was a difference in the dose of midazolam administered ($P = 0.000$). When comparing EGD and colonoscopy in patients undergoing sedation by the anesthesiologist, there were no significant differences in the doses of midazolam and ketamine, but the dose of propofol increased for colonoscopy (2.50 ± 0.95 mg/kg vs. 4.71 ± 1.66 mg/kg, $P = 0.000$). The cognitive recovery time according to drug dose was associated with propofol only in EGD with a shorter endoscopy time. The longer cognitive recovery time in colonoscopy and the discharge time of EGD and colonoscopies were not associated with propofol use.

Conclusion: When sedation is performed by an anesthesiologist, various drugs are used with sufficient doses and complications are reduced, but the discharge time does not change. For performing pediatric endoscopy in Korea, anesthesiologists should be considered for inducing anesthesia.

Keywords: Endoscopy; Pediatric Patients; Anesthesia; Propofol

Su Yeong Kim 

<https://orcid.org/0000-0001-6463-0210>

Eung Sang Choi 

<https://orcid.org/0000-0002-0878-7333>

Disclosure

The authors have no potential conflicts of interest to disclose.

Authors Contributions

Conceptualization: Yun SW, Choi ES. Data curation: Choi GJ. Writing - original draft: Yang SM, Yi DY. Writing - review & editing: Yi DY, Lim IS, Chae SA, Lee NM, Kim SY.

INTRODUCTION

Endoscopic examination has undergone massive advancements to enable its use in pediatric patients, while being used for the last 30 years to diagnose and treat various gastrointestinal (GI) diseases. In addition to routine diagnostic esophagogastroduodenoscopy (EGD) or colonoscopy, procedures such as polypectomy, endoscopic hemostasis, therapeutic foreign body removal, endoscopic retrograde cholangiopancreatography, and endoscopic ultrasound are also conducted in pediatric patients.¹⁻³ The prevalence of using endoscopies for diagnoses and treatments in pediatric patients is gaining popularity particularly in Korea, as observed in other countries, and the number of trials is gradually increasing. However, unlike adults, pediatric patients may experience increased discomfort while undergoing invasive tests such as endoscopy, because of preconceived abstinence from the procedure or anxiety and adherence issues.⁴⁻⁶ Moreover, endoscopy is a painful procedure, unlike magnetic resonance imaging or electroencephalography, often requiring the use of analgesic medications. Especially, the risk of complications increases if the patient cannot cooperate with the procedure.^{6,7} Therefore, it is increasingly common to plan deep-level sedation for pediatric patients prior to undergoing endoscopy.^{8,9} However, in clinical practice in healthcare institutions in Korea, sedation is induced and monitored in most cases by the endoscopy operator who performs the endoscopic examination rather than a specialist in anesthesiology. This challenge is multi-factorial, and there is a dearth of data on the type and dose of the drugs used for sedation. Further, there is a considerable level of panic among pediatric patients during endoscopic procedures, thereby eliciting forcible suppression by assistants.

Most importantly, in Korea, no studies have yet described the types of sedation drugs, effects of sedation, and efficiency of endoscopy with respect to the sedation instructor. Hence, in this study, we investigated the effectiveness of sedative procedures performed by anesthesiologists.

METHODS

Patient selection

We retrospectively reviewed the medical records of patients aged < 18 years who underwent endoscopy at the Chung-Ang University Hospital during March 2014–July 2019. The endoscopy room in the hospital is used jointly for adult and pediatric patients. At our hospital, during adult and pediatric endoscopies, sedation is performed and monitored by anesthesiologists and nurses of the anesthesiology department since 2014; this is in contrast to the procedure followed in most other hospitals in Korea. The drugs used for sedation in endoscopy include intravenous midazolam (Midazolam Bukwang Inj. 5 mg/5 mL, 1 mg/mL; Bukwang Pharmaceutical Co., Ltd., Seoul, Korea), ketamine (Huons Ketamine HCl Inj. 10 mL, 50 mg/mL; HUONS, Seongnam, Korea), and propofol (50 mL [vial] of Fresofol Mct 1% Inj. 50 mL [vial] 1% strain, 10 mg/mL; Fresenius Kabi Austria GmbH, Graz, Austria). Subsequently, the postprocedural recovery time was checked by an anesthesiologist. This is a routine process performed for almost all endoscopic procedures; however, the endoscope operator may perform the sedation intermittently during the weekend or when an anesthesiologist is unavailable. During the study period, the endoscope operator was common for all pediatric endoscopies, and sedations in all cases were performed by the same operator or a single anesthesiologist.

Endoscopic examinations performed in the neonatal intensive care unit, operation room, and intensive care unit were excluded from the study. Patients with neurological problems or structural abnormalities in the GI tract, such as esophageal duplication, were excluded because these problems might affect the type and dose of sedation drugs and the recovery time. We also excluded the cases of therapeutic endoscopies, such as those involving bleeding control, polypectomy, or foreign body removal. Finally, we analyzed the records of all patients aged < 18 years who underwent upper or lower GI endoscopy for simple diagnosis or examination in general with no underlying illnesses. The final selection of patients was classified based on the sedation performed by the endoscope operator and sedation performed by the anesthesiologist.

We collected data of the patients' clinical background, the doctor who conducted sedation, type and dose of the drug, complications, and extent of recovery after sedation, through medical records. The examination time was measured since the first endoscopic image was recorded while the endoscope was inserted into the oral cavity or through the anus to the time point of recording the last image just before the tube was removed from the oral cavity or anus to complete the examination. The degree of recovery from anesthesia was determined by checking eye-opening, verbal response, and orientation recovery as cognitive parameters and correlated with the time to reach a ≥ 9 score per the standard of discharge in the Aldrete scoring system.^{10,11} To exclude variations according to the combination of drugs used, we analyzed only those for whom almost the same dose of midazolam was used first for inducing sedation and then for maintenance of sedation with increased propofol dosage. Patients for whom sedation was maintained through ketamine were excluded from the analysis of recovery. The recovery time was determined based on the drug dose per body weight according to the procedure time.

Statistical analysis

Statistical analysis was performed using SPSS 18.0 statistical software (SPSS Inc., Chicago, IL, USA). The χ^2 test and Student's *t*-test were used to compare the groups and the mean and standard deviations were presented. The Mann-Whitney test was used to compare the group that underwent colonoscopy and EGD with the group that underwent only colonoscopy. The linear regression test was used to compare the sedation time and recovery time according to the use of propofol.

Ethics statement

This study was conducted with approval from the Institutional Review Board (IRB) of the Chung-Ang University Hospital (IRB No. 1706-004-16074), and informed consent was waived due to the retrospective nature of the study.

RESULTS

Clinical characteristics of all pediatric patients undergoing endoscopy

Between March 2014 and July 2019, 257 patients aged under 18 years underwent diagnostic endoscopies at Chung-Ang University Hospital, which were conducted by the same pediatric endoscopist (Table 1). Of them, 217 patients (mean age, 11.8 ± 4.1 years; 40.1% male) underwent EGD and 40 patients (mean age, 15.1 ± 2.7 years; 45.0% male) underwent colonoscopy. Overall, 54 (24.9%) and 16 (40.0%) patients were diagnosed with organic diseases, such as GI invasion of Henoch-Schönlein purpura, *Helicobacter pylori* gastritis, and

Table 1. Clinical characteristics of all pediatric patients undergoing endoscopy

Variables	Esophagogastroduodenoscopy (n = 217)	Colonoscopy (n = 40)	P value
Age, yr	11.8 ± 4.1	15.1 ± 2.7	0.000 ^a
Gender ratio, male:female	87 (40.1):130 (59.9)	18 (45.0):22 (55.0)	0.562
Height, cm	149.0 ± 20.3	161.7 ± 10.6	0.000 ^a
Body weight, kg	44.0 ± 17.5	59.2 ± 18.6	0.000 ^a
Body mass index, kg/m ²	18.9 ± 4.1	22.4 ± 6.0	0.001 ^a
Final diagnosis			0.048 ^a
Functional problems	163 (75.1)	24 (60.0)	
Organic diseases	54 (24.9)	16 (40.0)	
Sedation instructing doctor			0.011 ^a
Pediatric endoscopist	29 (13.4)	0	
Anesthesiologist	188 (86.6)	40 (100)	

Data are presented as mean ± standard deviation or number (%).

^aSignificant findings at $P < 0.05$.

inflammatory bowel disease, based on EGD and colonoscopy findings. Of the patients who underwent EGD, 29 (13.4%) underwent sedation by the pediatric endoscopist and 188 (86.6%) underwent sedation by the anesthesiologist. All 40 patients who underwent colonoscopy were sedated by the anesthesiologist.

Table 2 displays the different clinical characteristics according to the sedation instructor for 217 patients who underwent EGD. There were no differences in age, sex, height, weight, or body mass index between both groups. When the anesthesiologist performed the sedation, organic diseases were less frequently diagnosed (22.9% vs. 37.9%); however, this difference was statistically nonsignificant. Endoscopic examinations consumed more time when the anesthesiologist induced anesthesia than when the sedation was performed by the pediatric endoscopist (401.0 ± 135.1 seconds vs. 274.9 ± 106.1 seconds, $P < 0.001$).

Comparison of drugs for sedation in pediatric patients undergoing endoscopy

Among the 217 pediatric patients who underwent EGD, sedation drugs were compared according to the sedation instructor (**Table 3**). Midazolam was used in all patients of both groups, but there was a significant difference in the dosage and the dose per weight (4.35 ± 0.91 mg vs. 0.89 ± 1.47 mg, $P = 0.000$; 0.10 ± 0.03 mg/kg vs. 0.03 ± 0.08 mg/kg, $P = 0.000$, respectively). Ketamine did not show any difference between both groups in terms of the total dosage ($P = 0.776$); however, higher doses per body weight were used by the anesthesiologist (0.50 ± 0.10 mg/kg vs. 0.65 ± 0.42 mg/kg, $P = 0.005$). The pediatric endoscopist did not use propofol, but the anesthesiologist used a 2.50 ± 0.95 mg/kg dose of propofol in 181 patients (96.3%).

Table 2. Clinical characteristics of esophagogastroduodenoscopy patients according to the sedation instructor

Variables	Pediatric endoscopist (n = 29)	Anesthesiologist (n = 188)	P value
Age, yr	12.4 ± 3.7	11.8 ± 4.2	0.427
Sex, male	11 (37.9)	76 (40.4)	0.799
Height, cm	152.6 ± 15.9	148.4 ± 20.9	0.219
Body weight, kg	47.0 ± 16.5	43.6 ± 17.6	0.321
Body mass index, kg/m ²	19.7 ± 5.2	18.8 ± 3.9	0.292
Final diagnosis			0.081
Functional diseases	18 (62.1)	145 (89.0)	
Organic diseases	11 (37.9)	43 (22.9)	
Total endoscopy time, sec	274.9 ± 106.1	401.0 ± 135.1	0.000 ^a

Data are presented as mean ± standard deviation or number (%).

^aSignificant findings at $P < 0.05$.

Table 3. The drugs used for anesthesia in esophagogastroduodenoscopy patients according to the sedation instructors

Variables	Pediatric endoscopist (n = 29)	Anesthesiologist (n = 188)	P value
Midazolam	29 (100)	188 (100)	N/C
Dose, mg	4.35 ± 0.91	0.89 ± 1.47	0.000 ^a
Dose/body weight, mg/kg	0.10 ± 0.03	0.03 ± 0.08	0.000 ^a
Ketamine	10 (34.5)	90 (47.9)	0.178
Dose, mg	23.50 ± 6.69	21.12 ± 26.21	0.776
Dose/body weight, mg/kg	0.50 ± 0.10	0.65 ± 0.42	0.005 ^a
Propofol	0 (0)	181 (96.3)	0.000 ^a
Dose, mg	-	111.00 ± 60.57	N/C
Dose/body weight, mg/kg	-	2.50 ± 0.95	N/C

Data are presented as mean ± standard deviation or number (%).

N/C = not checked.

^aSignificant findings at $P < 0.05$.

Of the 40 colonoscopy cases, 36 were those of colonoscopy with EGD and 4 underwent colonoscopy alone. No significant differences were observed in the drugs used or their doses between the patients who underwent colonoscopy with EGD and those who underwent colonoscopy only; the endoscopy duration was 35.0 (25–65) minutes and 26.5 (26–36) minutes for both sub-groups, respectively ($P = 0.028$).

We compared the doses of drugs used by the anesthesiologist for inducing anesthesia before EGD and colonoscopy (Table 4). The number of cases of midazolam administration, the dose, and dose per body weight used did not differ between the two groups ($P = 0.175$, $P = 0.539$, and $P = 0.344$, respectively). Ketamine was used in more patients who underwent EGD alone than in those who underwent EGD and colonoscopy (47.9% vs. 20.0%, $P = 0.001$); however, the dose in patients using ketamine did not differ between the two groups (dose and dose per body weight, $P = 0.54$ and $P = 0.165$, respectively). Propofol was used in both groups. Unlike the cases of administering midazolam and ketamine, when propofol was used, the total drug dose and dose per body weight increased since colonoscopy lasted longer than EGD (111.00 ± 60.57 mg vs. 268.08 ± 98.25 mg, $P = 0.000$; 2.50 ± 0.95 mg/kg vs. 4.71 ± 1.66 mg/kg, $P = 0.000$, respectively).

Comparison of recovery time after sedation and dose per weight of propofol according to endoscopy time

Tables 5 and 6 show the comparison of recovery time after sedation and the dose per weight of propofol according to endoscopy time. The recovery time was measured in 80 (midazolam dose, 0.015 ± 0.008 mg/kg) patients who underwent EGD alone and 25 (midazolam dose, 0.016 ± 0.003 mg/kg) patients who underwent colonoscopy. In EGD, which had a relatively short endoscopy time, the dose of propofol per endoscopy time was significantly correlated

Table 4. Comparison of drug doses in patients sedated by the anesthesiologist

Variables	EGD (n = 188)	Colonoscopy (n = 40)	P value
Midazolam	188 (100)	39 (97.5)	0.175
Dose, mg	0.89 ± 1.47	1.04 ± 0.30	0.539
Dose/body weight, mg/kg	0.03 ± 0.08	0.04 ± 0.16	0.344
Ketamine	90 (47.9)	8 (20.0)	0.001 ^a
Dose, mg	21.12 ± 26.21	26.88 ± 17.92	0.545
Dose/body weight, mg/kg	0.65 ± 0.42	0.44 ± 0.21	0.165
Propofol	181 (96.3)	40 (100)	0.609
Dose, mg	111.00 ± 60.57	268.08 ± 98.25	0.000 ^a
Dose/body weight, mg/kg	2.50 ± 0.95	4.71 ± 1.66	0.000 ^a

Data are presented as mean ± standard deviation or number (%).

EGD = esophagogastroduodenoscopy.

^aSignificant findings at $P < 0.05$.

Table 5. Comparison of recovery time according to the propofol dose per weight and endoscopic time: Recovery characteristics of patients who underwent EGD with maintenance doses of propofol (n = 80)

Variables	Time, min	P value	β
Eye opening	19.91 ± 11.10	0.024 ^a	0.253
Verbal	20.49 ± 10.74	0.018 ^a	0.264
Orientation	22.04 ± 10.03	0.019 ^a	0.263
Score 9	23.81 ± 9.97	1.000	0.185

EGD = esophagogastroduodenoscopy; Score 9 = the time required for accomplishing an Aldrete score \geq 9.

^aSignificant findings at $P < 0.05$.

Table 6. Comparison of recovery time according to the propofol dose per weight and endoscopic time: Recovery characteristics of patients who underwent colonoscopy with maintenance doses of propofol (n = 25)

Variables	Time, min	P value	β
Eye opening	23.80 ± 9.82	0.342	0.198
Verbal	24.88 ± 9.60	0.303	0.214
Orientation	26.52 ± 9.12	0.385	0.182
Score 9	28.40 ± 9.19	0.666	0.091

EGD = esophagogastroduodenoscopy; Score 9 = the time required for accomplishing an Aldrete score \geq 9.

with cognitive parameters such as eye-opening, verbal response, and time of orientation recovery ($P = 0.024$, $P = 0.018$, and $P = 0.019$, respectively). Conversely, the recovery time with an Aldrete score of 9 and above as standard of discharge was not correlated with the dose of propofol per the endoscopy time ($P = 1.000$). In contrast, colonoscopies had a relatively long endoscopy duration and the dosage of propofol according to the duration of endoscopy was nonsignificantly associated with any indicator of recovery.

Drug use for endoscopic sedation and subsequent complications

Some complications were reported in 1 patient (3.4%) and 23 patients (12.2%) in both groups ($P = 0.214$). Among the sedations performed by the endoscopist, we noted one case of ketamine overdose caused hypotension due to a dose measurement error that required oxygen administration. Among the cases of sedation performed by the anesthesiologist, one patient, who cried severely before the endoscopic procedure, required 5-litre O₂ administration because of $\leq 90\%$ desaturation with cough and secretion after the endoscopy. Additionally, there was 1 case of bradycardia and 1 case of dystonia. In other cases, complications, such as transient cough, hiccup, snoring, irritability, chilling, intravenous line leakage, dizziness, and diplopia, were noted; however, these cases recovered without special treatment. Unlike when sedation was performed by an endoscopic operator, in the case of sedation performed by the anesthesiologist, there were no cases of complaints of pain, memory related to the sleep process, or discomfort due to insufficient sedation.

DISCUSSION

As mentioned previously, the anesthesiologist determines the use of drugs for sedation and monitors the process during endoscopic procedures, which occurs only at the Chung-Ang University Hospital in Korea. Therefore, to the best of our knowledge, this is the first report in Korea on anesthesia conducted by an anesthesiologist in pediatric endoscopy. In the present study, we report the successful implementation of performing sedation by anesthesiologists as opposed to the conventional practice of inducing anesthesia by pediatric endoscopists.

Substantial immobilization of uncomfortable and emotionally disturbed pediatric patients is essential for the successful implementation of pediatric GI procedures, such

as endoscopy.^{12,13} In some cases, light sedation was used, but was often considered as insufficient. Therefore, currently, general anesthesia is regularly used.^{13,14} Nevertheless, there are limitations to this for reasons such as the lack of anesthesiologists and cost-related concerns. Additionally, it is not an easy procedure to perform, especially in Korea. With this background, our experience of sedation performed by anesthesiologists in the endoscopy room revealed important aspects. First, when sedation was performed by an anesthesiologist, various drugs could be used in sufficient doses. Antidote-free analgesic drugs such as ketamine were also safe to use, and propofol was also available in sufficient doses. Nonetheless, significant complications were infrequent and were mostly temporary. Secondly, the endoscopist could concentrate on the endoscopy itself and could sufficiently examine the patient since the patient's sedation need not be monitored. Sedation-related monitoring was conducted by a specialist, and, in the cases of incomplete sedation, endoscopies could be performed in a stable state due to the use of additional drugs. Moreover, patient satisfaction was also very high.

The guidelines of the American Society for Gastrointestinal Endoscopy recommend this sedation protocol to ensure the safety, comfort, and cooperation of pediatric patients undergoing endoscopy.¹⁴⁻¹⁶ Nearly all children who receive GI procedures are advised to opt for endoscopist-administered moderate sedation or anesthesiologist-administered deep sedation, and general anesthesia. Moreover, previous studies emphasized the advantages of anesthesiologist-administered sedation.^{5,9,15,16} In addition to the same advantages we experienced, previous studies reported that sedation can be adequately performed in patients with various risk factors, and the risks of complications were lower. In particular, the effects of sedation with propofol were emphasized. Propofol is an ultra-short acting sedative agent that rapidly crosses the blood-brain barrier, enabling early-onset action. Because of its very short distribution half-life and prompt recovery, it has recently become preferable in endoscopy.¹⁷ Previous studies confirmed that propofol can be used relatively safely without severe complications in pediatric endoscopy.^{12,18-21} However, the use of propofol alone does not rule out the possibility of cardiorespiratory complications, which may occur with a significantly higher incidence than with other regimens.²² There is still considerable debate about endoscopist-administered propofol.^{23,24} The use of propofol in pediatric patients requires further analysis.²⁵ Therefore, procedures such as pediatric endoscopy should be proactive and specialists in anesthesiology should be recruited such as, anesthesiologists, in Korea to induce sedation. In our experience, the safety, effectiveness, and patient satisfaction with propofol administration by anesthesiologists were excellent.

In our case, the effects of the use of midazolam or ketamine did not differ for colonoscopy according to the procedure time; only the dose of propofol was increased for prolonged procedures. This was because propofol was used to maintain sedation after its induction. Because of this characteristic, low doses of midazolam were used for induction before administering propofol. This combination is widely used to relieve patient tension, increase the effectiveness of other drugs, and reduce side effects.²⁶⁻²⁹ As shown in the review by Yoon et al.,¹⁷ propofol is used for inducing sedation in EGD, but in colonoscopy, an additional propofol dose is required to maintain sedation during prolonged endoscopies.

In pediatric patients, recovery after anesthesia is influenced by various factors. It may also be affected by the type of examination or the usual anxiety, with the variations depending on the drug type, combination of drugs, and the dose of each drug.³⁰⁻³⁴ In our study, recovery after sedation was expected to vary according to the drug type, such as midazolam, ketamine, and

propofol, and their doses. We also expected it to vary depending on dose per body weight, obesity, and procedure time. However, it was difficult to assess all these variations because of the limitations of a retrospective study design and the small number of patients in each group. In the case of sedation performed by an anesthesiologist, since the midazolam doses were low during induction in all patients and propofol was used in more patients, recovery time was confirmed only based on the dose of propofol per body weight. Prolonged colonoscopies with a higher dose of propofol showed no correlation with recovery time. However, in EGD with a short endoscopy time and low propofol dose, simple cognitive parameters such as eye-opening, verbal response, and orientation differed significantly depending on the dose of propofol. The Aldrete score, which reflects overall sleep time and standard of discharge after sedation, was not associated with the propofol dose in EGD or colonoscopy. Our results demonstrate that increasing the propofol dose does not negatively impact the time of discharge.

Meanwhile, extensive investigations of propofol-related recovery in pediatric endoscopy are necessary. In particular, the results of a randomized double-blinded study on recovery time after MRI conducted by Schmitz et al.³⁰ were similar to ours. Compared to that in the patient group that was sedated using relatively high doses of only propofol and that using ketamine with relatively low-dose propofol, the recovery time was shorter in the group sedated with low-dose propofol. In general, propofol affects the central nervous system via cerebral vasoconstriction, which is directly proportional to the administered dose.¹⁸ Cerebral blood flow and a decrease in metabolic demand for oxygen may be associated with the action of propofol.^{35,36} Propofol was reported to affect the normal neurological functions, but its clinical significance is still unclear.¹⁸ Increasing sedation time through propofol use in children has been associated with prolonged sedation in premature infants or motor function reduction and blindness in 23-month-old infants.^{37,38} These neurological effects and differences in recovery time are thought to be related, but further research is needed to confirm this finding. The effects and half-life of propofol differ according to age group and weight.¹⁸ Therefore, there is a risk of overdose in obese patients, and sedation may be insufficient if the dose is reduced.³⁹⁻⁴¹ In our study, recovery time was determined using dose per body weight according to the procedure time; however, additional prospective studies stratifying patients according to body weight or age group may provide further details on this issue.

This study had several limitations; first, it was not a prospective study that evaluated a specific drug, dose, or method of administration, but a retrospective study that analyzed medical records over the 5-year study period. Second, this study may reflect Korea's policies on the use of anesthetic drugs, internal hospital regulations, or personal preferences of anesthesiologists. Despite these few limitations, our study is the first report in Korea on sedation conducted by anesthesiologists in pediatric patients for endoscopy, which shows the various advantages of this approach. We suggest that during pediatric endoscopy, sedation should be performed by anesthesiologists in more centers and that drug combinations for sedation and recovery from sedation should be further investigated.

REFERENCES

1. Thomson M, Tringali A, Dumonceau JM, Tavares M, Tabbers MM, Furlano R, et al. Paediatric gastrointestinal endoscopy: European Society for Paediatric Gastroenterology Hepatology and Nutrition and European Society of Gastrointestinal Endoscopy Guidelines. *J Pediatr Gastroenterol Nutr* 2017;64(1):133-53.

[PUBMED](#) | [CROSSREF](#)

2. Wyllie R, Kay MH. Gastrointestinal endoscopy in infants and children. *Pediatr Rev* 1993;14(9):352-9.
[PUBMED](#) | [CROSSREF](#)
3. Pall H, Lerner D, Khlevner J, Reynolds C, Kurowski J, Troendle D, et al. Developing the pediatric gastrointestinal endoscopy unit: a clinical report by the endoscopy and procedures committee. *J Pediatr Gastroenterol Nutr* 2016;63(2):295-306.
[PUBMED](#) | [CROSSREF](#)
4. Sargin M, Uluer MS, Aydogan E, Hanedan B, Tepe Mİ, Eryilmaz MA, et al. Anxiety levels in patients undergoing sedation for elective upper gastrointestinal endoscopy and colonoscopy. *Med Arh* 2016;70(2):112-5.
[PUBMED](#) | [CROSSREF](#)
5. Isik IA, Iyilikçi L, Ozturk Y, Adiyaman E. sedation practice outside the operating room for pediatric gastrointestinal endoscopy. *Indian Pediatr* 2015;52(11):989-90.
[PUBMED](#) | [CROSSREF](#)
6. Coté CJ, Wilson S; American Academy of Pediatrics; American Academy of Pediatric Dentistry. Guidelines for monitoring and management of pediatric patients before, during, and after sedation for diagnostic and therapeutic procedures. *Pediatrics* 2019;143(6):e20191000.
[PUBMED](#) | [CROSSREF](#)
7. Lightdale JR, Liu QY, Sahn B, Troendle DM, Thomson M, Fishman DS, et al. Pediatric endoscopy and high-risk patients: a clinical report from the NASPGHAN Endoscopy Committee. *J Pediatr Gastroenterol Nutr* 2019;68(4):595-606.
[PUBMED](#) | [CROSSREF](#)
8. Lee MC. Sedation for pediatric endoscopy. *Pediatr Gastroenterol Hepatol Nutr* 2014;17(1):6-12.
[PUBMED](#) | [CROSSREF](#)
9. Chung HK, Lightdale JR. Sedation and monitoring in the pediatric patient during gastrointestinal endoscopy. *Gastrointest Endosc Clin N Am* 2016;26(3):507-25.
[PUBMED](#) | [CROSSREF](#)
10. McGrath B, Chung F. Postoperative recovery and discharge. *Anesthesiol Clin North America* 2003;21(2):367-86.
[PUBMED](#) | [CROSSREF](#)
11. Ead H. From Aldrete to PADSS: reviewing discharge criteria after ambulatory surgery. *J Perianesth Nurs* 2006;21(4):259-67.
[PUBMED](#) | [CROSSREF](#)
12. Khalila A, Shavit I, Shaoul R. Propofol sedation by pediatric gastroenterologists for endoscopic procedures: a retrospective analysis. *Front Pediatr* 2019;7:98.
[PUBMED](#) | [CROSSREF](#)
13. Green SM, Klooster M, Harris T, Lynch EL, Rothrock SG. Ketamine sedation for pediatric gastroenterology procedures. *J Pediatr Gastroenterol Nutr* 2001;32(1):26-33.
[PUBMED](#) | [CROSSREF](#)
14. Fredette ME, Lightdale JR. Endoscopic sedation in pediatric practice. *Gastrointest Endosc Clin N Am* 2008;18(4):739-51.
[PUBMED](#) | [CROSSREF](#)
15. Lightdale JR, Acosta R, Shergill AK, Chandrasekhara V, Chathadi K, Early D, et al. Modifications in endoscopic practice for pediatric patients. *Gastrointest Endosc* 2014;79(5):699-710.
[PUBMED](#) | [CROSSREF](#)
16. van Beek EJ, Leroy PL. Safe and effective procedural sedation for gastrointestinal endoscopy in children. *J Pediatr Gastroenterol Nutr* 2012;54(2):171-85.
[PUBMED](#) | [CROSSREF](#)
17. Yoon SW, Choi GJ, Lee OH, Yoon IJ, Kang H, Baek CW, et al. Comparison of propofol monotherapy and propofol combination therapy for sedation during gastrointestinal endoscopy: a systematic review and meta-analysis. *Dig Endosc* 2018;30(5):580-91.
[PUBMED](#) | [CROSSREF](#)
18. Chidambaran V, Costandi A, D'Mello A. Propofol: a review of its role in pediatric anesthesia and sedation. *CNS Drugs* 2015;29(7):543-63.
[PUBMED](#) | [CROSSREF](#)
19. Kamat PP, McCracken CE, Gillespie SE, Fortenberry JD, Stockwell JA, Cravero JP, et al. Pediatric critical care physician-administered procedural sedation using propofol: a report from the Pediatric Sedation Research Consortium Database. *Pediatr Crit Care Med* 2015;16(1):11-20.
[PUBMED](#) | [CROSSREF](#)

20. Rajasekaran S, Hackbarth RM, Davis AT, Kopec JS, Cloney DL, Fitzgerald RK, et al. The safety of propofol sedation for elective nonintubated esophagogastroduodenoscopy in pediatric patients. *Pediatr Crit Care Med* 2014;15(6):e261-9.
[PUBMED](#) | [CROSSREF](#)
21. Kamat PP, McCracken CE, Gillespie SE, Fortenberry JD, Stockwell JA, Cravero JP, et al. Pediatric critical care physician-administered procedural sedation using propofol: a report from the Pediatric Sedation Research Consortium Database. *Pediatr Crit Care Med* 2015;16(1):11-20.
[PUBMED](#) | [CROSSREF](#)
22. Narula N, Masood S, Shojaee S, McGuinness B, Sabeti S, Buchan A. Safety of propofol versus nonpropofol-based sedation in children undergoing gastrointestinal endoscopy: a systematic review and meta-analysis. *Gastroenterol Res Pract* 2018;2018:6501215.
[PUBMED](#) | [CROSSREF](#)
23. Kim EH, Lee SK. Endoscopist-directed propofol: pros and cons. *Clin Endosc* 2014;47(2):129-34.
[PUBMED](#) | [CROSSREF](#)
24. Rex DK. Endoscopist-directed propofol. *Gastrointest Endosc Clin N Am* 2016;26(3):485-92.
[PUBMED](#) | [CROSSREF](#)
25. Akbulut UE, Kartal S, Dogan U, Akcali GE, Kalayci S, Kirci H. Propofol with and without midazolam for diagnostic upper gastrointestinal endoscopies in children. *Pediatr Gastroenterol Hepatol Nutr* 2019;22(3):217-24.
[PUBMED](#) | [CROSSREF](#)
26. Amini A, Arhami Dolatabadi A, Kariman H, Hatamabadi H, Memary E, Salimi S, et al. Low-dose fentanyl, propofol, midazolam, ketamine and lidocaine combination vs. regular dose propofol and fentanyl combination for deep sedation induction; a randomized clinical trial. *Emerg (Tehran)* 2018;6(1):e57.
[PUBMED](#)
27. Loh G, Dalen D. Low-dose ketamine in addition to propofol for procedural sedation and analgesia in the emergency department. *Ann Pharmacother* 2007;41(3):485-92.
[PUBMED](#) | [CROSSREF](#)
28. Phillips W, Anderson A, Rosengreen M, Johnson J, Halpin J. Propofol versus propofol/ketamine for brief painful procedures in the emergency department: clinical and bispectral index scale comparison. *J Pain Palliat Care Pharmacother* 2010;24(4):349-55.
[PUBMED](#) | [CROSSREF](#)
29. Ghojzadeh M, Sanaie S, Paknezhad SP, Faghiih SS, Soleimanpour H. Using ketamine and propofol for procedural sedation of adults in the emergency department: a systematic review and meta-analysis. *Adv Pharm Bull* 2019;9(1):5-11.
[PUBMED](#) | [CROSSREF](#)
30. Schmitz A, Weiss M, Kellenberger C, O'Gorman Tuura R, Klaghofer R, Scheer I, et al. Sedation for magnetic resonance imaging using propofol with or without ketamine at induction in pediatrics-A prospective randomized double-blinded study. *Paediatr Anaesth* 2018;28(3):264-74.
[PUBMED](#) | [CROSSREF](#)
31. Sargin M, Uluer MS, Aydogan E, Hanedan B, Tepe Mİ, Eryılmaz MA, et al. Anxiety levels in patients undergoing sedation for elective upper gastrointestinal endoscopy and colonoscopy. *Med Arh* 2016;70(2):112-5.
[PUBMED](#) | [CROSSREF](#)
32. Hayhoe S, Pallett S, Zani J, Trott J. Reduction of postanesthetic pediatric distress: a coordinated approach. *J Perianesth Nurs* 2018;33(3):312-318.e1.
[PUBMED](#) | [CROSSREF](#)
33. Moncel JB, Nardi N, Wodey E, Pouvreau A, Ecoffey C. Evaluation of the pediatric post anesthesia discharge scoring system in an ambulatory surgery unit. *Paediatr Anaesth* 2015;25(6):636-41.
[PUBMED](#) | [CROSSREF](#)
34. Jain A, Gombar S, Ahuja V. Recovery profile after general anaesthesia in paediatric ambulatory surgeries: desflurane versus propofol. *Turk J Anaesthesiol Reanim* 2018;46(1):21-7.
[PUBMED](#) | [CROSSREF](#)
35. Szabó EZ, Luginbuehl I, Bissonnette B. Impact of anesthetic agents on cerebrovascular physiology in children. *Paediatr Anaesth* 2009;19(2):108-18.
[PUBMED](#) | [CROSSREF](#)
36. Karsli C, Luginbuehl I, Farrar M, Bissonnette B. Propofol decreases cerebral blood flow velocity in anesthetized children. *Can J Anaesth* 2002;49(8):830-4.
[PUBMED](#) | [CROSSREF](#)
37. Bacon RC, Razis PA. The effect of propofol sedation in pregnancy on neonatal condition. *Anaesthesia* 1994;49(12):1058-60.
[PUBMED](#) | [CROSSREF](#)

38. Lanigan C, Sury M, Bingham R, Howard R, Mackersie A. Neurological sequelae in children after prolonged propofol infusion. *Anaesthesia* 1992;47(9):810-1.
[PUBMED](#) | [CROSSREF](#)
39. Chidambaran V, Sadhasivam S, Diepstraten J, Esslinger H, Cox S, Schnell BM, et al. Evaluation of propofol anesthesia in morbidly obese children and adolescents. *BMC Anesthesiol* 2013;13(1):8.
[PUBMED](#) | [CROSSREF](#)
40. Igarashi T, Nagata O, Iwakiri H, Ikeda M, Uezono S, Ozaki M. Two cases of intraoperative awareness during intravenous anesthesia with propofol in morbidly obese patients. *Masui* 2002.51(11):1243-7.
[PUBMED](#)
41. Olutoye OA, Yu X, Govindan K, Tjia IM, East DL, Spearman R, et al. The effect of obesity on the ED(95) of propofol for loss of consciousness in children and adolescents. *Anesth Analg* 2012;115(1):147-53.
[PUBMED](#) | [CROSSREF](#)