

Research

Open Access

## Raised houses reduce mosquito bites

J Derek Charlwood\*<sup>1</sup>, Joao Pinto<sup>2</sup>, Patrica R Ferrara<sup>3</sup>, Carla A Sousa<sup>4</sup>,  
Conceicao Ferreira<sup>5</sup>, Vilfrido Gil<sup>5</sup> and Virgillo E do Rosário<sup>2</sup>

Address: <sup>1</sup>Danish Bilharziasis Laboratories, Jaegersborg Allé 1-D, Charlottenlund, DK 2920, Denmark, <sup>2</sup>Centro de Malária e Outras Doenças Tropicais, Instituto de Higiene e Medicina Tropical, Universidade Nova de Lisboa, Rua da Junqueira 96, 1349-008 Lisbon, Portugal, <sup>3</sup>SIMA Project, c/o Technoserve, Rua Bas N'Tchinga 509, Bairro da Coop, Maputo, Mozambique, <sup>4</sup>Unidade de Entomologia Médica Instituto de Higiene e Medicina Tropical, Universidade Nova de Lisboa, Rua da Junqueira 96, 1349-008 Lisbon, Portugal and <sup>5</sup>Centro Nacional de Endemias, Caixa Postal 218, São Tomé, Democratic Republic of São Tomé and Príncipe

Email: J Derek Charlwood\* - dc@bilharziasis.dk; Joao Pinto - jpinto@ihmt.unl.pt; Patrica R Ferrara - patricia.ferrara@tvcabo.co.mz; Carla A Sousa - casousa@ihmt.unl.pt; Conceicao Ferreira - malarst@cstome.net; Vilfrido Gil - malarst@cstome.net; Virgillo E do Rosário - cmdt@ihmt.unl.pt

\* Corresponding author

Published: 10 December 2003

Received: 19 June 2003

*Malaria Journal* 2003, 2:45

Accepted: 10 December 2003

This article is available from: <http://www.malariajournal.com/content/2/1/45>

© 2003 Charlwood et al; licensee BioMed Central Ltd. This is an Open Access article: verbatim copying and redistribution of this article are permitted in all media for any purpose, provided this notice is preserved along with the article's original URL.

### Abstract

**Background:** In many parts of continental Africa house construction does not appear to impede entry of malaria vectors and, given their generally late biting cycle, the great majority of transmission takes place indoors. In contrast, many houses in São Tomé, 140 km off the coast of Gabon, are raised on stilts and built of wooden planks. Building on stilts is a time-honoured, but largely untested, way of avoiding mosquito bites. Exposure may also be affected by mosquito activity times and age composition of host-seeking females. A study was therefore undertaken on the island of São Tomé to determine if exposure to *Anopheles gambiae*, the only vector on the island, varied with house construction or time of the night.

**Methods:** A series of all-night landing collections were undertaken out of doors at ground level, inside houses at ground level, on the verandas of, and inside houses built on stilts. The gonotrophic age of an unselected sample of insects from the first three hours of landing collection (18:00–21:00) was determined by dissection. In addition, 1,149 miniature light-trap collections were obtained from 125 houses in the study area. Numbers collected were related to house construction.

**Results:** Biting of *An. gambiae* took place primarily outside at ground level. Less than one third of biting occurred inside houses. Houses built on stilts had half the number of *An. gambiae* in them compared to those built at ground level. Conversely houses with an eaves gap had more *An. gambiae* in them than houses without such a gap. Gonotrophic age did not affect house entry rates in *An. gambiae*. House construction affected *Culex quinquefasciatus* less than *An. gambiae*. Mean density per house, derived from a series of 1,490 randomly assigned light-trap collections, was over-dispersed with 18% of houses having 70% of the vectors.

**Conclusion:** House construction plays an important role in determining exposure to malaria vectors in São Tomé. Neighbours can have very different exposure levels. Recommendations for improvement in control are given.

## Background

In most areas of Africa the great bulk of feeding by the principal malaria vectors, *Anopheles funestus* and *Anopheles gambiae*, takes place in the latter part of the night, indoors [1]. Entry rates and hence transmission are affected by house construction. Thus, houses built of brick and with zinc roofs are associated with lower levels of malaria-associated anaemia in Tanzania compared to poorly built mud walled houses [2]. Similarly, in Sri Lanka house construction appears to be an important determinant of exposure to malaria [3]. Improvements in house construction, in particular screening against mosquito entry, have, therefore, long been recommended as a way of reducing exposure [4].

One way to do this is to sleep on a raised structure or in a house built on stilts [4]. The practise was described by Herodotus in Egypt 2450-2420 B.P. [5] and has been independently adopted by a number of different groups of people. For example, in Northern Vietnam the hill people build their houses on stilts to avoid *Anopheles minimus*, which seldom fly more than 2–3 m above the ground [6]. The anophelines feed on the animals kept under the houses and avoid the people. When people from the lowlands move into the hills they bring their culture with them. Their houses are built on the ground and animals are stabled alongside. As a result, *An. minimus*, which will feed on cattle but prefers humans, becomes an active vector of malaria [6].

Malaria was introduced in the archipelago of São Tomé and Príncipe shortly after colonization in the early 1500's [7]. Sporozoite rates in *An. gambiae*, the only vector present on the islands, are generally less than 1% [8,9] and, despite high vector densities, the disease reaches only meso-hyperendemic levels [10,11]. A reason for this may be the propensity for the vector to feed on non-human hosts, particularly dogs and pigs [12]. These feeds are all taken out of doors. Given that *An. gambiae* is generally thought to prefer human blood and is considered to be 'indifferent' to the location of the host [1], the observed blood feeding patterns suggest that entry into houses in São Tomé for the mosquito is difficult. According to an early visitor, many of the free African inhabitants on the islands built their houses of sawn timber planks on stilts, to avoid ground-level humidity and mosquito bites [7]. This type of construction still predominates on both islands. However, in a recent survey, people living in houses raised on stilts in Príncipe, the smaller of the two islands that make up the two main islands in the archipelago, did not have a lower prevalence of malaria than people living at ground level [13].

Present vector control measures are all based against mosquitoes biting indoors. Should a high proportion of infec-

tions be acquired outdoors, alternative strategies of vector control need to be considered. In this study we examine the influence of house construction on biting rates of the local *An. gambiae*. The gonotrophic age of insects was also determined in order to assess whether the protective effect, if any, of building a house on stilts is reduced because older insects bite at higher levels [14]. Results are discussed with respect to novel vector control strategies that might be adopted on the islands.

## Methods

Studies were undertaken in Riboque, a peri-urban area close to the capital city centre. Riboque has numerous houses with sawn timber walls and zinc-roofed houses; many built on stilts of varying heights, in close proximity to each other. Banana plants and a variety of fruit trees grow throughout the area and a small stream (Lukumi) runs through the middle. During the rainy season the stream is liable to flooding and there are extensive swamp areas. Climate is equatorial with annual mean temperatures around 25°C and relative humidity of 80%. There are two short dry seasons, June-August and December-January.

A series of all-night landing collections were undertaken out of doors at ground level, inside houses at ground level, on the verandas of, and inside houses built on stilts in the wet season of 1997. Mosquitoes were collected with an aspirator as they landed on the exposed lower legs of adult men working in six-hour shifts, all of who were offered prophylaxis according to the protocols of the Centro Nacional de Endemias de São Tomé (CNE). Following the initial all-night collections a series two-hour landing collections (from 2100–2300 h) were undertaken each month outside, at a ground level sentinel site, and numbers were compared to those caught in sentinel light-traps.

Each hour's collection was kept in individual containers and the gonotrophic age of an unselected sample of insects from the first three hours of landing collection (18:00–21:00) was determined by dissection of the female's ovaries and spermathecae following the schema described by Charlwood *et al.* [8].

Host-seeking mosquitoes were also caught with Centre for Disease Control (CDC) miniature light-traps run all night close to the bed of sleepers protected by an untreated mosquito net [8]. In 1997 and 1998, houses for light-trap collections were selected at random, from a list of 25 houses. Five traps were run per night. An additional trap was run in a sentinel house. From May 1997 the number of houses in the study increased to 157. New sentinel houses were employed in the event that houses in use became unavailable.

**Table 1: Mean numbers of *Anopheles gambiae* and *Culex quinquefasciatus* collected per hour in landing catches, according to location of collection site, Riboque, São Tomé.**

Mean number collected per man-hour (95% CI)	Outdoor ground level	Veranda	Indoor ground level	Indoor upper level
<i>An. gambiae</i>	16.51 (12.8–21.2)	8.10 (5.5–11.7)	3.58 (2.9–4.4)	2.38 (1.7–3.3)
<i>Cx. quinquefasciatus</i>	5.11 (3.9–6.6)	5.14 (3.5–7.4)	6.18 (5.1–7.4)	4.23 (3.2–5.5)

**Table 2: Physiological age of *Anopheles gambiae* collected in landing collections, according to location of collection site, Riboque, São Tomé.**

Physiological age ( <i>An. gambiae</i> )	Location				$\chi^2$	
	Outdoor ground level	Veranda	Indoor ground level	Indoor upper level		
Virgin	167	91	49	14	0.30	n.s.
Plug positive	103	44	26	3	0.84	n.s.
Nulliparous II	38	21	12	3	0.57	n.s.
Parous with Sac	193	100	58	6	0.04	n.s.
Parous No-sac	90	55	39	8	0.39	n.s.
Proportion parous	0.48	0.50	0.52	0.41		

The data from the all-night landing collections were balanced by eliminating measurements outside the time-frame over which indoor catches in elevated houses were obtained (18 March – 8 April 1997) and also houses over-sampled during this timeframe. In all 93 house-hours were systematically selected for each classification (inside/outside and ground level/elevated). Mosquito counts were log-transformed and a linear model, using S-PLUS 4.5 [15], was fitted to the log-transformed counts for *An. gambiae* and *Cx. quinquefasciatus*


## Results

Sixteen all night landing collections were performed out of doors at ground level, 22 inside houses at ground level, 15 on the verandas of, and eight inside houses built on stilts. A total of 8,926 *An. gambiae*, 6,539 *Culex quinquefasciatus*, 15 *Aedes aegypti* and a small number of unidentified culicines were collected. Differences in mosquito density were observed according to location of collection site and time of collection (Fig 1). More *An. gambiae* were collected outdoor biting to indoor biting and biting at ground level to biting at elevated levels. Table 1 shows linear model results for each species. *Cx. quinquefasciatus* biting densities were lower with statistical significance only for elevation. The fit for *An. gambiae* was much stronger than for *Cx. quinquefasciatus* (F statistic of 74.14 on 3 and 68 degrees of freedom and p-value less than 0.0001 for *An. gambiae* vs. 2.622 on 3 and 68 degrees of freedom p-value of 0.05 for *Cx. quinquefasciatus*). The effect of elevation


was weak for *Cx. quinquefasciatus* ( $p = 0.05$ , with increased ground level biting, whereas there was no statistically significant difference between indoor and outdoor biting. Results for *An. gambiae* on the other hand were quite different, with both house elevation and indoor/outdoor effects being statistically significant ( $p$ -values less than 0.0001). There was a similar interaction strength in the *An. gambiae* data compared to that of *Cx. quinquefasciatus* (both having  $p = 0.04$ ).

Gonotrophic age distribution of the 2, 818 *An. gambiae* dissected from the first three hours of collection was similar among different collection sites (Table 2).


From March 1997 to October 1999, 1,149 light-trap collections were obtained from 125 bedrooms in the study area. The number of *An. gambiae* and *Cx. quinquefasciatus* in light-traps were not significantly correlated (Spearman rank correlation coefficient on log-transformed ( $x+1$ ) numbers,  $\rho = 0.166$ ,  $n = 125$ ,  $p = 0.065$ ). The distribution of *An. gambiae* was over-dispersed with 70% of the mosquitoes being collected from 18% of the houses (Fig 2a). Densities varied greatly between neighbouring houses according to construction. For example, eighteen-fold differences in mean densities of *An. gambiae* per night were recorded between houses five meters apart one of which was built at ground level whilst the other was built on stilts two metres high (mean per night = 68.7, s. d. =



**Figure 1**  
Biting cycles of *Anopheles gambiae* according to house type and location in Riboque, São Tomé.



**Figure 3**  
Scatter plot of mean number of *An. gambiae* collected per hour from outdoor ground level sentinel collections and mean number collected from indoor sentinel light-trap collections, Riboque, São Tomé.



**Figure 2**  
Rank distribution of houses according to mean number of mosquitoes collected from light-traps. a) *An. gambiae* b) *Cx. quinquefasciatus*.

85.5, n = 69 compared to mean per night = 3.7, s. d. = 6.8, n = 23).

The concentration in numbers of *Cx. quinquefasciatus* was not so apparent (fig 2b). The number of both *An. gambiae* and *Cx. quinquefasciatus* in light-traps were negatively correlated with the house being built on stilts (Spearman rank correlation coefficient on log transformed numbers (x+1), rho = -0.71, n = 25, p < 0.0001 and rho = -0.54, n = 25, p = 0.006 respectively). The number of *An. gambiae* in light-traps, but not the number of *Cx. quinquefasciatus*, was positively associated with the presence of an eave gap (respective Spearman rank correlations rho = 0.567, n = 25, p = 0.0032 compared to rho = 0.177, n = 25, p = 0.4). There was, however, no association for either species between the number of windows, doors or inhabitants that a house had and the number of mosquitoes caught (data not shown).

A mean of 16.2 (s.d. 5.8) light-trap collections were run in sentinel houses per month and a mean of 7 (s.d. 14) two hour landing collections undertaken per month. The relationship between mean numbers of *An. gambiae* caught per month in sentinel collections and per hour in routine outdoor landing catches is shown in Fig 3. Mean monthly number of mosquitoes caught by both methods were highly correlated ( $y = 0.196x + 15.8, R^2 = 0.776$ ).

## Discussion

The ancient Egyptians were right. People in houses raised on stilts suffer fewer mosquito attacks than people living on the ground. House construction affects indoor exposure to malaria vectors in suburban São Tomé just as it does in rural Vietnam [6]. Most *An. gambiae* were caught outside, at ground level and the least were caught inside houses built on stilts. Houses built at ground level had more mosquitoes in them than houses built on stilts. This can give rise to an 18-fold difference in exposure to malaria vectors between houses less than 10 m apart. Not unexpectedly higher mosquito numbers in light-traps were associated with houses in which there was an eaves gap [16]. In Príncipe, people living in houses with an eaves gap were also associated with a higher risk of infection with malaria parasites than those from other houses [14]. House construction or location of the host (inside or outside) affected *Cx. quinquefasciatus* less than it did *An. gambiae*. While *Cx. quinquefasciatus* did show a weak preference for ground level biting to elevated levels there was no preference for indoor over outdoor biting and the poor overall fit suggested that other factors, not accounted for in the analysis, were important. At the time of the study biting rates for *Cx. quinquefasciatus* were, on the whole, lower than for *An. gambiae*, which could have been the result of the peak season for *Cx. quinquefasciatus* occurring at a different time of the year. The wide variation of biting rates for *An. gambiae* among individual houses suggests that even in this species more parameters could be added to the model.

Light-trap collections undertaken inside houses and outdoor landing catches undertaken in high density areas were well correlated. Light trap collections may therefore be a suitable alternative to landing collections in estimating seasonal/longitudinal fluctuations of *An. gambiae*, thus avoiding unnecessary exposure of collectors while performing landing captures. The age structure, at least in the first three hours of the night, of *An. gambiae* caught inside or outside or in ground-level or elevated locations were similar. Virgin female *An. gambiae*, which have to fly at least two meters above the ground in order to mate [17], appear to seek hosts by flying close to the ground in much the same way as other age groups. Given that the age structure of the insects at ground level outside was similar throughout the night [18] it is likely that the age structure of insects caught in the other locations also remains constant throughout the night. The constancy of parous rates implies that potentially infective insects will bite at all times of the night, regardless of location.

At the time of the study, many people in São Tomé watched communal television out of doors. Similarly, potential purchasers of the catch will wait on the beach for the return of the fishermen at night and, depending on the

tides, can spend many hours outside. The availability of cheap repellents [19] would reduce risk in such situations.

The concentration of mosquitoes in a small number of houses also implies that, if high risk houses can be easily identified and if it is socially acceptable, some sort of focal control might be feasible. Spraying high density houses (with an insecticide differing from the one used on nets) would expose much of the house-entering fraction of the mosquito population to two different insecticide groups and reduce the likelihood of resistance developing, much as combined therapy might reduce resistance developing in parasites.

Spraying does not, however, provide personal protection and cannot be easily implemented in a house-by-house way. Bednets have been widely available in São Tomé since 1995. Poorer households are still more likely to be inadequately protected and at low coverage rates unimpregnated bednets may even increase transmission in children [20]. Covering the eaves of houses with insecticide impregnated netting or installing netting ceilings has been shown to reduce mosquito entry rates and malaria transmission in continental Africa [21,22]. Such systems could be installed in houses with eaves in São Tomé. This would also provide a measure of protection to those who cannot afford bed nets.

## Abbreviations

CDC – Centres for Disease Control, CI – Confidence Interval CNE – Centro Nacional de Endemias, s. d. – standard deviation

## Authors' contributions

Charlwood – Designed the study, supervised collections, dissected the mosquitoes and wrote the manuscript

Pinto – Dissected the mosquitoes and wrote the manuscript

Ferrara-Undertook the statistical analyses

Sousa – Dissected the mosquitoes and wrote the manuscript

Ferreira – Provided logistical support in São Tomé

Gil – Provided logistical support in São Tomé

Rosario – Provided logistical support in Portugal

## Acknowledgements

We would like to thank the collectors from Riboque and the entomology technicians of the CNE for their assistance during this work. Thanks to Tom Smith for assistance with data analysis and for his comments on an earlier draft of the manuscript. Financial support from INCODC/EU project

"Studies on mosquito vectors in West Africa aimed at malaria control and epidemiology" (IC18CT960030). J. Pinto is supported by Fundação para a Ciência e Tecnologia, Portugal (SFRH/BPD/10086/2002). We would like to thank Maria do Céu (Univ. Coimbra) for showing us the reference to house construction in São Tomé.

## References

- Gillies MT, De Meillon B: **The Anophelinae of Africa South of the Sahara**. Publication No. 54. South African Institute for Medical Research, Johannesburg 1968.
- Kahigwa E, Schellenberg D, Sanz S, Aponte JJ, Wigayi J, Mshinda H, Alonso P, Menendez C: **Risk factors for presentation to hospital with severe anaemia in Tanzanian children: a case-control study**. *Trop Med Int Health* 2002, **7**:823-830.
- Gamage-Mendis AC, Carter R, Mendis C, de Zoysa APK, Herath PRJ, Mendis KN: **Clustering of malaria infections within an endemic population: risk of malaria associated with the type of housing construction**. *Am J Trop Med Hyg* 1991, **45**:77-85.
- Lindsay SW, Emerson PM, Charlwood JD: **Reducing malaria by mosquito-proofing houses**. *Trends Parasitol* 2002, **18**:510-514.
- Herodotus: **The histories**. Edited by: A de Sélincourt. Penguin Books, London; 1966.
- Laderman C: **Malaria and Progress: some historical and ecological considerations**. *Soc Sci Med* 1975, **9**:587-594.
- Caldeira AM: **Viagens de um piloto português do século XVI à costa de África e a São Tomé**. Comissão Nacional para as Comemorações dos Descobrimentos Portugueses, Lisbon 12000.
- Charlwood JD, Pinto J, Sousa CA, Ferreira C, Petrarca V, do Rosario VE: **'A mate or a meal' – Pre-gravid behaviour of female Anopheles gambiae from the islands of São Tomé and Príncipe, West Africa**. *Malaria J* 2003, **2**:9.
- Pinto J, Sousa CA, Gil V, Ferreira C, Gonçalves L, Lopes D, Petrarca V, Charlwood JD, do Rosário VE: **Malaria in São Tomé and Príncipe: Parasite prevalences and vector densities**. *Acta Trop* 2000, **76**:185-193.
- Martet G, Conceição S, Cordoliani G, Delolme H, Raphenon G, Hovette P, Doury JC, Baudon D, Touze JE, Lecamus JL: **Le paludisme en République de São Tomé e Príncipe. Evaluation épidémiologique et chimiorésistance de P. falciparum**. *Bull Soc Path Exot* 1991, **84**:273-280.
- Loureiro LF, Cesário AM, Franco AS, Rosário VE, Benito A, Ferreira C, Eggelte TE: **Malaria in São Tomé and Príncipe: prevalence and drug-susceptibility**. *Ann Trop Med Parasitol* 1994, **90**:223-224.
- Sousa CA, Pinto J, Almeida APG, Ferreira C, do Rosário VE, Charlwood JD: **Dogs as a favoured host choice of Anopheles gambiae sensu stricto (Diptera:Culicidae) of São Tomé, West Africa**. *J Med Entomol* 2001, **38**:122-125.
- Hagmann R, Charlwood JD, Gil V, Ferreira C, do Rosario V, Smith T: **Malaria and its possible control on the island of Príncipe**. *Malaria J* 2003, **2**:15.
- Snow WF, Wilkes TJ: **Age composition and vertical distribution of mosquito populations in The Gambia, West Africa**. *J Med Entomol* 1977, **13**:507-513.
- S-PLUS 4.5 Professional**. Release 2 MathSoft, Seattle 1997.
- Lindsay SW, Snow RW: **The trouble with eaves; house entry by vectors of malaria**. *Trans R Soc Trop Med Hyg* 1988, **82**:645-646.
- Charlwood JD, Pinto J, Sousa CA, Ferreira C, do Rosário VE: **The swarming and mating behaviour of Anopheles gambiae (Diptera: Culicidae) from São Tomé Island**. *J Vector Ecol* 2002, **27**:178-183.
- Charlwood JD, Pinto J, Sousa CA, Ferreira C, Gil V, de Rosario V: **Mating does not affect the biting behaviour of Anopheles gambiae from the islands of São Tomé and Príncipe, West Africa**. *Ann Trop Med Parasitol* 2003, **97**:751-756.
- Charlwood JD: **Repellent 'soap' for malaria vector control in Papua New Guinea**. *PNG Med J* 1987, **30**:99-103.
- Charlwood JD: **Getting a measure of bednets in Africa**. *Trends Parasitol* 2002, **18**:386.
- Cuzin-Ouattara N, Van den Brock AHA, Habluetzel A, Diabaté A, Sanogo-Ilboudo E, Diallo DA, Cousens SN, Esposito F: **Wide-scale installation of insecticide-treated curtains confers high levels of protection against malaria transmission in a hyperendemic area of Burkina Faso**. *Trans R Soc Trop Med Hyg* 1999, **93**:473-479.
- Lindsay SW, Jawara M, Paine K, Pinder M, Walraven GE, Emerson PM: **Changes in house design reduce exposure to malaria mosquitoes**. *Trop Med Int Health* 2003, **8**:512-517.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:  
http://www.biomedcentral.com/info/publishing\_adv.asp

