

CORRECTION


Correction: High fibroblast growth factor 23 levels are associated with decreased ferritin levels and increased intravenous iron doses in hemodialysis patients

The *PLOS ONE* Staff

The affiliation for the last author is incorrect. Takanori Shibata is not affiliated with #6 but with #2 Showa University, School of Medicine, Division of Nephrology, Department of Medicine, Tokyo, Japan. The publisher apologizes for the error.

Reference

1. Honda H, Michihata T, Shishido K, Takahashi K, Takahashi G, Hosaka N, et al. (2017) High fibroblast growth factor 23 levels are associated with decreased ferritin levels and increased intravenous iron doses in hemodialysis patients. *PLoS ONE* 12(5): e0176984. doi:[10.1371/journal.pone.0176984](https://doi.org/10.1371/journal.pone.0176984) PMID: [28475601](https://pubmed.ncbi.nlm.nih.gov/28475601/)



OPEN ACCESS

Citation: The *PLOS ONE* Staff (2017) Correction: High fibroblast growth factor 23 levels are associated with decreased ferritin levels and increased intravenous iron doses in hemodialysis patients. *PLoS ONE* 12(6): e0179139. <https://doi.org/10.1371/journal.pone.0179139>

Published: June 1, 2017

Copyright: © 2017 The PLOS ONE Staff. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.