

MITOGENOME ANNOUNCEMENT

 OPEN ACCESS 

The complete mitochondrial genome of *Tanacetipathes thamnea* Warner, 1981 (Antipatharia: Myriopathidae)

Diego Francisco Figueroa , David Hicks and Nicole Jewel Figueroa

School of Earth, Environmental, and Marine Sciences, University of Texas Rio Grande Valley, One West University Boulevard, Brownsville, TX, USA

ABSTRACT

Specimens of the black coral *Tanacetipathes thamnea* were collected from the Northwestern Gulf of Mexico. The complete mitochondrial genome of one of these specimens was obtained from genomic DNA by next-generation sequencing technology on the Illumina HiSeq 2500. Only three species of black corals have a completely sequenced mitochondrial genome. These were used to reconstruct the phylogeny for the order Antipatharia. The mitochondrial genome of *T. thamnea* is 17,712 base pairs and contains 13 protein-coding genes, 2 ribosomal RNAs, and 2 transfer RNAs in the following order: 16s RNA, COX3, COX1 (with intron), ND4L, COX2, ND4, ND6, ATP8, ATP6, and ND5 (with intron and copies of ND1 and ND3), tRNA-Trp, ND2, 12s RNA, CYTB, tRNA-Met. The gene arrangement is the same as that for *Myriopathes japonica* with a nearly identical sequence (99.35% identical). These results show that the mitochondrial genome within the family Myriopathidae is highly conserved.

ARTICLE HISTORY

Received 2 October 2019

Accepted 9 November 2019

KEYWORDS

Black coral; mitogenome; phylogeny; Gulf of Mexico

There are only three species of black corals (order Antipatharia) with completely sequenced mitochondrial genomes, *Myriopathes japonica*, *Chrysopathes formosa*, and *Stichopathes lutkeni* (Brugler and France 2007; Kayal et al. 2013). We present the first complete mitochondrial genome for the genus *Tanacetipathes* (GenBank: MN265369). The genus *Tanacetipathes* has 11 species (WoRMS Editorial Board 2019). *Tanacetipathes thamnea* is distributed throughout the western Atlantic, including the Gulf of Mexico (WoRMS Editorial Board 2019).

Specimens were collected at Aransas Bank (latitude 27° 35' N and longitude 96° 27' W) off the coast of Texas in the Gulf of Mexico and preserved in 95% ethanol. An individual polyp was picked from one specimen (UTRGV Coastal Studies Laboratory accession #D08-28B) and genomic DNA was extracted using the PureLink Genomic DNA Kit (Thermo Fisher Scientific Co., Waltham, MA). An indexed DNA library was prepared with the Nextera X2 kit (Illumina). This library was multiplexed with 95 other indexed libraries and sequenced on a 100 bp paired-end lane of Illumina HiSeq 2500 (Illumina, San Diego, CA) at Harvard's Biopolymers facility. Reads were assembled de novo using CLC Genomics Workbench version 11 (Qiagen). The contigs from the assembly included the full mitochondrial genome with an average coverage of 54, which was then annotated using the mitochondrial genome for *M. japonica* (GenBank: JX456459) as a reference.

The mitochondrial genome of *T. thamnea* is 17,712 base pairs. It contains 13 protein-coding genes, 2 ribosomal RNAs, and 2 transfer RNAs, all in the major strand in the following order: 16s RNA, COX3, COX1 (with intron), ND4L, COX2, ND4, ND6, ATP8, ATP6, ND5 (with intron and copies of ND1 and ND3), tRNA-Trp, ND2, 12s RNA, CYTB, and tRNA-Met. The nucleotide composition is 26.5% A, 18.2% C, 22.0% G, and 33.4% T (40.2% C + G and 59.8% A + T).

A concatenated alignment was created with the mitochondrial genomes of *M. japonica* (GenBank: JX456459), *Chrysopathes formosa* (GenBank: DQ304771), *Stichopathes lutkeni* (GenBank: JX023266), and *Metridium senile* (GenBank: HG423143) by extracting and aligning individual genes and RNAs using MUSCLE version 3.8 (Edgar 2004). Phylogenetic analyses were performed by maximum likelihood using Mega-X (Kumar et al. 2018) with bootstrap values from 1000 replicates and evolutionary model GTR + G+I. The resulting phylogeny shows that *T. thamnea* is most closely related to *M. japonica* (Figure 1). Both of these species belong to the family Myriopathidae. Sister to this clade is *Chrysopathes formosa* from the family Cladopathidae while *Stichopathes lutkeni*, from the family Antipathidae, is in the basal position. The mitochondrial genome within the family Myriopathidae is highly conserved. The mitochondrial genomes of *T. thamnea* and *M. japonica* are 99.35% identical.

The rate of evolution of the mitochondrial genome in this family is very low and it rivals the slow rate found within Octocorals, where mitogenomes of congeners only vary by

CONTACT Diego Francisco Figueroa  diego.figueroa@utrgv.edu  School of Earth, Environmental, and Marine Sciences, University of Texas Rio Grande Valley, One West University Boulevard, Brownsville, TX 78520, USA

© 2019 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.
This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.




Figure 1. Black coral (Antipatharia) maximum-likelihood tree based on complete mitochondrial genomes. The anemone *Metridium senile* is used as the outgroup. Branch labels show bootstrap support.

0.3–3.0% (e.g. Figueroa and Baco 2014). In Octocorals, this can be explained by a mitochondrial repair gene (*MutS*) found within their mitogenome (e.g. McFadden et al. 2006). A similar repair gene has not been found in Antipatharians, therefore, the reason for the slow rate of evolution of the mitochondrial genome within the family Myriopathidae remains to be explained.

Disclosure statement

The authors report no conflicts of interest. The authors alone are responsible for the content and writing of the manuscript.

Funding

Specimen collections of the south Texas coast were supported by TPWD-ARP [Grant No. 475342, 2016–2018] to David Hicks. Research reported in this publication was supported in part by an Institutional Grant [NA14OAR4170102 to Diego Figueroa and David Hicks] to the Texas Sea Grant College Program from the National Sea Grant Office, National Oceanic and Atmospheric Administration, U.S. Department of Commerce and by the Gulf Research Program of the National Academies of Sciences, Engineering, and Medicine under the Grant Agreement number [2000007266] to Diego Figueroa. The content is solely the responsibility of the authors and does not necessarily represent the official views of the Gulf Research Program of the National Academies of Sciences, Engineering, and Medicine.

ORCID

Diego Francisco Figueroa <http://orcid.org/0000-0002-0220-2912>

References

- Brugler MR, France SC. 2007. The complete mitochondrial genome of the black coral *Chrysopathes formosa* (Cnidaria:Anthozoa:Antipatharia) supports classification of antipatharians within the subclass Hexacorallia. Mol Phylogenet Evol. 42(3):776–788.
- Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput. Nucleic Acids Res. 32(5):1792–1797.
- Figueroa DF, Baco AR. 2014. Complete mitochondrial genomes elucidate phylogenetic relationships of the deep-sea octocoral families *Coralliidae* and *Paragorgiidae*. Deep Sea Res Part II Top Stud Oceanogr. 99:83–91.
- Kayal E, Roure B, Philippe H, Collins A, Lavrov D. 2013. Cnidarian phylogenetic relationships as revealed by mitogenomics. BMC Evol Biol. 13(1):5.
- Kumar S, Stecher G, Li M, Knyaz C, Tamura K. 2018. MEGA X: molecular evolutionary genetics analysis across computing platforms. Mol Biol Evol. 35(6):1547–1549.
- McFadden CS, France SC, Sánchez JA, Alderslade P. 2006. A molecular phylogenetic analysis of the *Octocorallia* (Cnidaria: Anthozoa) based on mitochondrial protein-coding sequences. Mol Phylogenet Evol. 41(3):513–527.
- WoRMS Editorial Board. 2019. World register of marine species. [accessed 2019 Aug 06]. <http://www.marinespecies.org> at VLIZ.