

the interactions between cannabis and chronic conditions, multiple comorbidities, polypharmacy, and mental health. The goal of this scoping review was to analyze the literature that addresses mental health outcomes of seniors who use cannabis, in order to answer the main research question: What is the relationship between older adults' use of cannabis and mental health? Following Arksey and O'Malley's five-stage framework, 10 electronic databases were searched along with a hand search of references. The search revealed 7000+ peer-reviewed and grey literature sources. 233 full-text sources were assessed for eligibility, with a total of 25 articles included. Thematic content analysis produced four major themes which addressed: (1) Usage characteristics; (2) User characteristics; (3) Outcomes; and (4) Physical and mental health considerations. Findings from this scoping review are positioned in terms of their implications for research, practice, and policy. While more in-depth, qualitative methods are required to develop further research, several harm-reduction strategies may be immediately utilized by both users and healthcare practitioners. It is critical that older adults and their physicians are able to make cannabis-related decisions with evidence-informed guidance to prevent problematic cannabis use and ensure positive mental health outcomes.

THE INFLUENCE OF COGNITIVE FUNCTION ON ADHERENCE TO BREAST CANCER SCREENING AMONG OLDER AMERICAN WOMEN

Chien-Ching Li,¹ Fritzi Flores,¹ Alicia K. Matthews,² Bryan James,³ and Raj Shah,⁴ 1. *Rush University, Rush University, Illinois, United States*, 2. *University of Illinois at UIC, Chicago, Illinois, United States*, 3. *Rush University, Chicago, Illinois, United States*, 4. *Rush, Rush University Medical Center, Illinois, United States*

Cognitive decline and impairment among older adults have become an important public health issue. Previous research shows older women have a greater prevalence of Alzheimer's disease than Men. Among women, breast cancer is one of the most common types of cancer. Over half of breast cancer deaths occur in women aged 65 and older. Therefore, early detection of breast cancer through mammogram screening is important among older women. This study aimed to examine the influence of cognitive function on adherence to mammogram breast cancer screening among older American women aged 65 and older. Data from the Health and Retirement Study (2012-2016) was obtained and analyzed. The independent variable of the study was cognitive function (normal, not normal). Adherence to mammogram (low, moderate, high) was the dependent variable. Multinomial regression was performed to examine the association between cognitive function and adherence to mammogram after controlling for demographic covariates. In the study, 33.3% of respondents had impaired cognitive function and 21.7% showed low adherence to mammogram screening. Regression results found that older women with impaired cognitive function were more likely to be in low adherence group (OR=1.30, p=0.01) or moderate adherence group (OR=1.47, p<0.001) relatively to be in high adherence group compared to older women with normal cognitive function. The development and implementation of interventions are needed for reducing barriers to accessing cancer screening services in high-risk vulnerable

populations. This submission is considered late-breaking research because study findings were obtained from a recently completed student's master's project.

THE LINK BETWEEN ENVIRONMENT, AGE, AND HEALTH IN A LARGE COHORT OF COMPANION DOGS FROM THE DOG AGING PROJECT

Brianah McCoy,¹ Layla Brassington,¹ Greer Dolby,¹ Kelly Jin,² Devin Collins,³ Matthew Dunbar,⁴ and Noah Snyder-Mackler,¹ 1. *Arizona State University, Tempe, Arizona, United States*, 2. *Allen Institute for Brain Science, Allen Institute for Brain Science, Washington, United States*, 3. *University of Washington, University of Washington, Washington, United States*, 4. *University of Washington, UNiversity of Washington, Washington, United States*

Exposure to social environmental adversity strongly predicts health and survival in many species such as non-human primates, wild mammals, and humans. However, little is known about how the health and mortality effects of these social determinants vary across the lifespan. Using the companion dog, which serves as a powerful comparative model for human health and aging due to our shared biology and environment, we examined which components of the social environment impact health, and how the effects vary with age, in dogs. We first drew on detailed survey data from owners of 27,547 dogs from the Dog Aging Project and identified six factors that together explained 35% of the variation in dog's social environment. These factors all predicted measures of health, disease, and mobility, when controlling for dog age and weight. Factors capturing measures of financial and household adversity were linked to poorer companion dog health, while factors associated with the social companions, like dogs and adults, were linked to better health. Interestingly, some of these effects differed across a dog's lifespan: for instance, the effect of neighborhood disadvantage on disease instances was strongest in older dogs. Together, our findings point to similar links between adversity and health in companion dogs, and set up future work on the molecular and biological changes associated with environmental variation in order to identify ways to mitigate or even reverse the negative environmental effects.

THE MODIFIED MINI MENTAL STATE (3MS) TEST - DEVELOPMENT OF AN ONLINE TRAINING PROGRAM

Tara Rose,¹ Evelyn Teng,¹ Chia Ying Chen,¹ Elyse Manzo,² Katherine Erickson,³ and Helena Chui,¹ 1. *University of Southern California, Los Angeles, California, United States*, 2. *University of Southern California, Huntington Beach, California, United States*, 3. *University of Southern California, Moraga, California, United States*

The Modified Mini Mental State (3MS), a screening test for cognitive abilities, can be administered in approximately 10 minutes and has a score range of 0-100. Early detection of cognitive impairment is important for clinical care. An individual's score change over time can also help assess disease progression and treatment effects. The Web of Science, an online database, reports 1,864 publications using the 3MS, including 412 in the last 5 years. Articles came from 52 countries in 7 continents, with languages in English (98.7%), French (0.5%), Spanish (0.5%), German