



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Short Communication

Risk of hospitalisation or death in households with a case of COVID-19 in England: an analysis using the HOSTED data set



J.A. Hall, R.J. Harris*, A. Zaidi, G. Dabrera, J.K. Dunbar

UK Health Security Agency, London, UK

ARTICLE INFO

Article history:

Received 11 April 2022

Received in revised form

16 July 2022

Accepted 19 July 2022

Available online 8 August 2022

Keywords:

COVID-19

SARS-CoV-2

Household contacts

Hospitalisation

Mortality

Household transmission

England

Surveillance

Secondary user service

HOSTED

ABSTRACT

Objective: To determine whether household contacts of confirmed cases of COVID-19 have an increased risk of hospitalisation or death.

Methods: We used the HOSTED data set of index cases of COVID-19 in England between June and November 2020, linked to Secondary Uses Service data on hospital episodes and Office for National Statistics' mortality data. Multivariable logistic regression models of the odds of household contacts being hospitalised or dying within six weeks of an index case, adjusted for case type, age, sex and calendar month were calculated. Excess risk was determined by comparing the first six weeks after the index case with 6–12 weeks after the index case in a survival analysis framework.

Results: Index cases were more likely to be hospitalised or die than either secondary cases or non-cases, having adjusted for age and sex. There was an increased risk of hospitalisation for non-cases (adjusted hazard ratio (aHR) 1.10; 95% confidence interval (CI) 1.04, 1.16) and of death (aHR 1.57; 95% CI 1.14, 2.16) in the first six weeks after an index case, compared to 6–12 weeks after.

Conclusion: Risks of hospitalisation and mortality are predictably higher in cases compared to non-cases. The short-term increase in risks for non-case contacts following diagnosis of the index case may suggest incomplete case ascertainment among contacts, although this was relatively small.

Crown Copyright © 2022 Published by Elsevier Ltd on behalf of The Royal Society for Public Health. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Introduction

The Household Transmission Evaluation Dataset (HOSTED)¹ provides a unique opportunity to explore the risk of hospitalisation and death in household contacts of confirmed cases of COVID-19 to determine if there is any excess risk to them, regardless of whether they are diagnosed with COVID-19 themselves. This is particularly important in the scenario of limited testing where cases may be missed, as was the case in England early in the pandemic.

Methods

The HOSTED methodology has been described elsewhere¹; in brief, it is an ongoing surveillance system that has identified the residential household contacts of laboratory-confirmed cases in England since 20th April 2020, including both Pillar 1 (testing of

persons for a clinical need in healthcare or as part of a public health investigation) and Pillar 2 (community-based testing accessible to members of the public). Linkages with hospital episodes from Secondary Uses Service (SUS) data (a national administrative dataset based on healthcare providers' clinical activities) and ONS (Office for National Statistics) mortality data for all anonymised cases and contacts within HOSTED enable us to investigate whether there is any increased risk in hospitalisation or death for household contacts of confirmed cases of COVID-19.

There is very limited evidence on hospitalisation and deaths of household contacts of confirmed COVID-19 cases. Data from a cohort study in Scotland have shown that between 1st March and 6th June 2020, household contacts of cases of COVID-19 in the general public had a risk of admission with COVID-19 of 0.05%; it was higher for healthcare workers and their household contacts.²

We considered individuals in households where the index case occurred between 1st June 2020 and 8th November 2020, extracted on the 31st January 2021, to allow for complete follow-up of hospitalisations within six weeks of the index case, and a further six weeks buffer in case of reporting delays in the SUS

* Corresponding author. UKHSA, Nobel House, 17 Smith Square, London, SW1P 3JR, UK. Tel.: +442083276428.

E-mail address: feedback.c19epi@ukhsa.gov.uk (R.J. Harris).

data. Hospitalised individuals were grouped using ICD-10 codes into 'COVID' (U07 and derivatives), 'pneumonia (B97 and J12) and possible interest', and 'other' (all other ICD-10 codes). ICD-10 codes of 'possible interest' were comorbidities thought to be risk factors for adverse outcomes of COVID-19 on discussion with clinicians. This predominantly included diseases of the cardiovascular, cerebrovascular, respiratory and renal systems, cancer, diseases that cause immunosuppression or require treatment with immunosuppressives, diabetes and obesity. The pneumonia codes were selected in case of misdiagnosis of a COVID-19 case as another viral pneumonia. This grouping together captured people in whom a diagnosis of COVID-19 could have been missed. We had data on whether the person had died, but not their cause of death.

Statistical analysis

The proportion hospitalised or dying within 6 weeks of the index case testing positive (starting from the specimen date of the positive test) was modelled using logistic regression; covariates included case category (index case, secondary case, contact without positive test), age group (0–34, 35–54, 55–69 and 70+ years) and sex. Time trends were considered by including calendar month, and age-specific trends. To ascertain whether there was any excess risk of hospitalisation in non-cases, we examined hospitalisation rates in the first six weeks after the index case compared to 6–12 weeks in a survival analysis framework. Hazards were assumed constant within each time interval and estimated hazard ratios (HRs) adjusted for age, sex and calendar time. Analysis was restricted to patients with 18 weeks observable follow-up.

Ethics approval

The HOSTED surveillance system was reviewed and approved by the PHE Research Ethics Governance Group. The data were collected and linked by NHS Digital. The data were processed lawfully under GDPR Article 6(1)e and 9(2)i and shared under Regulation 3(4) of the Health Service (Control of Patient Information) Regulations 2002.

Results

In England, there were 1.68 million individuals living in a household in which a confirmed case occurred between 1st June and 8th November 2020. The median household size was 4 (interquartile range (IQR): 3–5); the median age of individuals in the data set was 32 years (IQR 19–50 years), with 326,606 children younger than 16 years (19.4%) and 63,994 older than 70 years (3.8%). In 74.5% of the data for this period, the index case occurred in October/November, when cases were increasing rapidly in England, but before the emergence of the Alpha variant.

Hospitalisation

A total of 49,516 individuals (2.95%) were hospitalised within 42 days of the index case date. 28,843 of 477,034 (6.05%) index cases were hospitalised, and 4685 of 92,243 (5.08%) secondary cases were hospitalised. In comparison, among household contacts without laboratory-diagnosed COVID-19, 13,876 were hospitalised out of 1.05 million (1.32%).

In logistic regression of all persons in the data set, index cases were most likely to be hospitalised (aOR 4.49; 95% CI 4.40, 4.59)

compared to non-cases, after adjusting for age and sex, as were secondary cases (aOR 3.54; 95% CI 3.40, 3.65). Rates of hospitalisation increased with age, as did the risk associated with being a case compared to a contact, regardless of reason for admission. Of those aged over 70 years, 27.9% were admitted with COVID-19 within six weeks of laboratory confirmation and 4.14% were admitted with pneumonia or other potentially relevant ICD-10 code, compared to 1.26% and 0.35%, respectively, in index cases aged 0–34 years. Among household contacts who were not laboratory-confirmed COVID-19 cases, 0.35% of those aged over 70 years were admitted with COVID-19, 3.18% with pneumonia or other relevant condition and 0.58% for other reasons, according to the SUS data. For every age group, the percentages of non-COVID-19 admissions are higher in COVID-19 cases than in those not diagnosed with COVID-19.

Hospitalisations due to any cause in index cases of COVID-19 fell in all age groups from June to October 2020, though the risks decreased most for those younger than 35 years. Hospitalisation in non-cases also fell slightly in every age group between June- and October 2020, and younger age groups were consistently more likely to be hospitalised for reasons other than COVID-19/pneumonia.

Examining only the non-COVID-19 admissions, both index and secondary cases were more likely to be admitted to hospital in the six weeks following a positive test for SARS-CoV-2, particularly if they were older than 70 years, both for pneumonia/other relevant conditions or for another reason.

Mortality

A total of 6414 individuals (0.38%) died within 42 days of the index case date. 5230 of 477,034 index cases died (1.10%), 596 of 92,243 secondary cases died (0.65%), and 419 of 1.05 million non-cases died (0.04%).

Death was considerably more likely in index cases (aOR 22.9; 95% CI 20.7, 25.3) and secondary cases (aOR 13.1; 95% CI 11.5, 14.8) than in individuals not diagnosed with COVID-19, having adjusted for age and sex. In terms of trends, the risk of death in index cases reduced over time with a similar pattern to hospitalisation, with the greatest reduction over time in the younger groups. Trends are more stable for secondary cases, although data are sparse and confidence intervals wide. For those not diagnosed, there are declining trends in the youngest and oldest age groups, similar to hospitalisations, but confidence intervals are wide and the results are not significant.

Excess risk

The adjusted HR for hospitalisation within six weeks vs. 6–12 weeks was 1.10 (95% CI 1.04, 1.16), indicating a modest increase in hospitalisation rates in non-cases around the time of the index case. For mortality, the adjusted HR was 1.57 (95% CI 1.14, 2.16) for the first six weeks after the index case compared to 7–12 weeks after. As shown in Fig. 1, this was driven by increased hospitalisation and deaths in those older than 55 years.

Discussion

Index cases had the highest risk of hospitalisation and death followed by secondary cases compared to household contacts who did not become laboratory-confirmed cases (non-cases). The higher




Fig. 1. Proportion of non-diagnosed contacts hospitalised (any cause) or dying per week since date of index case positive, by age. Y-axis scales vary between panels.

risk in index cases could be due to testing being biased towards more severe index cases, whereas case ascertainment may be less dependent on severity for secondary cases.

We found some evidence of a modestly increased risk of hospitalisation in household contacts of laboratory-confirmed cases of COVID-19 who did not become laboratory-confirmed cases themselves. This suggests that, over the timeframe considered, case ascertainment has been good, if not complete. We found a higher risk of admission than the Scottish study, which is likely because we included all hospitalisations, not only those for COVID-19(2).

Mortality may be increased by around 50% in non-cases immediately following the index case, although absolute mortality rates remain low and confidence intervals for any excess risk were relatively wide. This may indicate incomplete case ascertainment if a person died before being tested. Routine postmortem testing for SARS-CoV-2 could reveal the true burden of the disease.³

Strengths and limitations

The HOSTED data set is large, covering all laboratory-confirmed cases and their household contacts in England. However, as a passive surveillance system, the data are subject to several limitations, including incomplete case ascertainment and a lack of information on testing uptake which could introduce bias. Without genomics data, we cannot confirm household transmission versus secondary cases having acquired their infection elsewhere. However, self-isolation of households following COVID-19 symptoms even before confirmation reduces the likelihood of acquiring an infection outside of the household. Previous sensitivity analysis showed that secondary attack rates within the household were robust to changing the definition of a secondary case from 2–14 days after the index case to 4–14 days after the index case.

Author statements

Acknowledgements

The authors wish to thank Heather Pinches, Phillip Bowker and Paul Ellingham as well as the rest of the NHS Digital team whose hard work and dedication enabled the creation of these data during the UK COVID-19 pandemic response. The authors also wish to thank Sarah Woodhall for her integral role in developing the HOSTED project.

Data availability statement

The data underlying this article cannot be shared publicly due to the legal and policy controls placed on data used as part of the government’s response to the COVID-19 pandemic.

Funding

This work was undertaken as part of the core functions of Public Health England in relation to the surveillance of communicable diseases and outbreak response.

Competing interests

None declared.

References

- Hall JA, Harris RJ, Zaidi A, Woodhall SC, Dabrera G, Dunbar JK. HOSTED-England’s Household Transmission Evaluation Dataset: preliminary findings from a novel passive surveillance system of COVID-19. *Int J Epidemiol* 2021;**50**(3):743–52.
- Shah ASV, Wood R, Gribben C, Caldwell D, Bishop J, Weir A, et al. Risk of hospital admission with coronavirus disease 2019 in healthcare workers and their households: nationwide linkage cohort study. *BMJ* 2020;**371**:m3582.
- Hall JA, Harris RJ, Emmett HE, Lowe B, Singanayagam A, Twhig KA, et al. On the sensitivity and specificity of post-mortem upper respiratory tract testing for SARS-CoV-2. *J Infect Dis* 2021 Aug 2;**224**(3):389–94.