

RESEARCH ARTICLE

# Comparing Effects of Biologic Agents in Treating Patients with Rheumatoid Arthritis: A Multiple Treatment Comparison Regression Analysis

Ingunn Fride Tvete<sup>1\*</sup>, Bent Natvig<sup>2</sup>, Jørund Gåsemyr<sup>2</sup>, Nils Meland<sup>3</sup>, Marianne Røine<sup>3</sup>, Marianne Klemp<sup>4,5</sup>

**1** The Norwegian Computing Center, Oslo, Norway, **2** Department of Mathematics, University of Oslo, Oslo, Norway, **3** Smerud Medical Research International AS, Oslo, Norway, **4** Department of Pharmacology, University of Oslo, Oslo, Norway, **5** The Norwegian Knowledge Centre for the Health Services, Oslo, Norway

\* [ingunn.fride.tvete@nr.no](mailto:ingunn.fride.tvete@nr.no)



CrossMark  
click for updates

## OPEN ACCESS

**Citation:** Tvete IF, Natvig B, Gåsemyr J, Meland N, Røine M, Klemp M (2015) Comparing Effects of Biologic Agents in Treating Patients with Rheumatoid Arthritis: A Multiple Treatment Comparison Regression Analysis. *PLoS ONE* 10(9): e0137258. doi:10.1371/journal.pone.0137258

**Editor:** Masataka Kuwana, Nippon Medical School Graduate School of Medicine, JAPAN

**Received:** April 17, 2015

**Accepted:** August 13, 2015

**Published:** September 10, 2015

**Copyright:** © 2015 Tvete et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** All relevant data are within the paper and its Supporting Information files.

**Funding:** This work was financed by the Norwegian Center for Research-based Innovation 'Statistics for innovation', Project Number 970141669. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing Interests:** The authors have declared that no competing interests exist.

## Abstract

Rheumatoid arthritis patients have been treated with disease modifying anti-rheumatic drugs (DMARDs) and the newer biologic drugs. We sought to compare and rank the biologics with respect to efficacy. We performed a literature search identifying 54 publications encompassing 9 biologics. We conducted a multiple treatment comparison regression analysis letting the number experiencing a 50% improvement on the ACR score be dependent upon dose level and disease duration for assessing the comparable relative effect between biologics and placebo or DMARD. The analysis embraced all treatment and comparator arms over all publications. Hence, all measured effects of any biologic agent contributed to the comparison of all biologic agents relative to each other either given alone or combined with DMARD. We found the drug effect to be dependent on dose level, but not on disease duration, and the impact of a high versus low dose level was the same for all drugs (higher doses indicated a higher frequency of ACR50 scores). The ranking of the drugs when given without DMARD was certolizumab (ranked highest), etanercept, tocilizumab/ abatacept and adalimumab. The ranking of the drugs when given with DMARD was certolizumab (ranked highest), tocilizumab, anakinra, rituximab, golimumab/ infliximab/ abatacept, adalimumab/ etanercept. Still, all drugs were effective. All biologic agents were effective compared to placebo, with certolizumab the most effective and adalimumab (without DMARD treatment) and adalimumab/ etanercept (combined with DMARD treatment) the least effective. The drugs were in general more effective, except for etanercept, when given together with DMARDs.

## Introduction

Disease modifying treatment has traditionally been given to rheumatoid arthritis (RA) patients at an early stage of the disease [1–2]. Early intervention could reduce joint damage and repress the RA progress [3]. For RA patients, biologic therapy has been a newer line of treatment typically following the firstly given disease modifying anti-rheumatic drugs (DMARDs). Biologic agents block certain chemicals in the blood from activating the immune system and hence protect patients' joints [4]. A joint treatment regime of both biologic and DMARD therapy is recommended [5].

When assessing clinical efficacy of biologic agents most randomized controlled trials (RCTs) have compared the effect of one biologic agent versus placebo, either with or without additional DMARD treatment in both or one of the treatment arms. Few trials actively compared the effect of one biologic agent to another. Also, many trials contained multiple treatment arms which differ both with respect to dosing regime and endpoints measured. The great variety in comparisons and trial design has made it challenging to compare and rank biologic agents. We have taken a multiple treatment comparison (MTC) regression modelling approach. We have developed a model including all trials comparing the effect of a biologic drug against placebo or another biologic drug, with or without DMARD treatment. An advantage with MTCs is that one obtains treatment comparisons not directly observed. For example, with some trials comparing treatment one to treatment two and some trials comparing treatment two to treatment three there was an indirect comparison of treatment one to treatment three. Also, observed treatment comparisons could be strengthened from the indirect estimates. Hence, by taking this approach one could compare and rank all biologics with respect to their clinical effect.

Among the RA biologic agents on the market there have been five TNF inhibiting anti-inflammatory drugs (adalimumab, certolizumab, etanercept, golimumab, and infliximab), one interleukin-1 (IL-1) receptor antagonist (anakinra), one T-cell selective co-stimulation modulator (abatacept), one chimeric monoclonal CD20 antibody (rituximab), and one anti-IL-6 (tocilizumab).

Most systematic reviews compared some of these drugs, a few compared all of them, by including various trials reporting treatment effects [6]. A model approach to take into consideration whether patients have been given joint DMARD and biologic agent treatment or just a biologic agent alone has been called for [6]. This paper demonstrates such an approach by constructing a comprehensive statistical model, including both direct and indirect comparisons of the effect of all the nine biologic agents assessed in published trials. The objective of this paper was to compare and rank biologic agents' clinical efficacy, when given alone or in combination with DMARDs in RA patients. We have developed a novel MTC regression analysis approach where we also take the disease duration and dose level into account.

## Materials and Methods

### Literature search

To identify published trials, we searched MEDLINE, Centre for Reviews and Dissemination, The Cochrane Library and the Drug Effectiveness Review Project's website for systematic reviews assessing efficacy of biologic drugs in RA patients using included drugs (abatacept, adalimumab, anakinra, certolizumab, etanercept, golimumab, infliximab, rituximab and tocilizumab) and indication (rheumatoid arthritis) as search terms and mapped to MeSH (Medical Subject Headings) terminology. We attempted to identify additional trials through hand

searches of reference lists of included trials and reviews. The search was last updated December 2014.

Among the systematic reviews identified, we selected the most comprehensive and updated of high quality, which we used to identify 93 possible eligible publications. After closer inspection, we found that 39 publications were not relevant for our analysis and therefore excluded. We found that 18 lacked the ACR50 response, one focused only on children, nine were not randomized, one reported a trial which was not double blind, eight considered other already reported trials and two contained a comparison including NSAID treatment which we did not consider. The analysis was therefore based on 54 publications [7–60]. One of the 54 publications reported two different comparison groups [22], giving a total of 55 comparison groups. A list of the included publications with an overview of the different comparisons is presented in [Table 1](#). Also see details of the study selection process in the flowchart in [Fig 1](#) and Table A in [S1 File](#). When considering publications reporting on the same trial at various time points we chose publications who reported trial duration closest to one year, see Table A in [S1 File](#) for details.

## Study selection

Two people independently reviewed full text of all the publications for inclusion. If both reviewers agreed that the reported trial did not meet eligibility criteria it was excluded. Records were considered for exclusion if they did not meet pre-established eligibility criteria with respect to trial design or duration, patient population, interventions, outcomes, and comparisons to medications outside our scope of interest. We considered adult patients diagnosed with RA who participated in a randomized, double blind clinical trial evaluating efficacy of biologic agents. We wanted not to distinguish between mode of administration (intravenous or oral) and to follow the intention to treat principle. We focused on the drugs' effect and used the number of patients experiencing a 50% improvement according to the American College of Rheumatology (ACR) scale [61] (ACR50) as endpoint. Efficacy endpoints are the response ratios between biologic agent treated groups and placebo (with or without DMARD treatment). We take into consideration the influence that the disease duration at trial start and the dose level used can have on the treatment effect in scenario analyses.

**Data.** [Table 1](#) presents the trials entering the analysis, together with the drugs considered in each trial. For the number of participants and ACR50 responders in the trials, see the Table C in [S1 File](#). We did not distinguish between patients given DMARD or DMARD plus placebo, since both groups were therapeutically comparable. [Fig 2](#) displays the network of direct and indirect comparisons, where DM indicated either DMARD treatment alone or joint placebo and DMARD treatment. There were for example 11 group comparisons of ADA versus placebo treatment. In total there were 111 comparisons over 54 publications. All nine drugs were compared with placebo when given jointly with DMARD, but there were no comparisons of infliximab, anakinra, golimumab or rituximab without joint DMARD treatment.

The duration of RA was given as the average number of years of disease duration at trial start for each treatment arm, see Table D in [S1 File](#). The dose levels were defined as either low or high, see Tables E and F in [S1 File](#).

## Statistical analysis

We constructed a joint model for assessing the comparable relative effect between the various biologic drugs, placebo and DMARD for all trials, and performed a Bayesian statistical analysis based on this model inspired by Klemp et al. [62], now including regression terms taking the explanatory variables disease duration and dose level into consideration. The analysis

**Table 1. The publications and the different treatment arms.**

	Publication	Treat <sub>arm 1</sub> *	Treat <sub>arm 2</sub>	Treat <sub>arm 3</sub>	Treat <sub>arm 4</sub>	Treat <sub>arm 5</sub>	Treat <sub>arm 6</sub>	Treat <sub>arm 7</sub>
1	Abe 2006	P+DM	INF+DM	INF+DM				
2	Emery 2008a	P+DM	TOC+DM	TOC+DM				
3	Genovese 2008a	DM	TOC+DM					
4	Maini 2006	P+DM	TOC	TOC	TOC	TOC+DM	TOC+DM	TOC+DM
5	Smolen 2008	P+DM	TOC+DM	TOC+DM				
6	Chen 2009	DM	ADA+DM					
7	Cohen 2002	P+DM	ANA+DM	ANA+DM	ANA+DM	ANA+DM	ANA+DM	
8	Cohen 2004	P+DM	ANA+DM					
9	Edwards 2004	DM	RIT+DM					
10	Emery 2006b	P+DM	RIT+DM	RIT+DM				
11	Emery 2009	P+DM	GOL+DM	GOL+DM				
12	Emery 2010	P+DM	RIT+DM	RIT+DM				
13	Fleischmann 2009	P	CER					
14	Kremer 2006	P+DM	ABA+DM					
15	Lipsky 2000	P+DM	INF+DM	INF+DM	INF+DM	INF+DM		
16	Maini 1999	P+DM	INF+DM	INF+DM	INF+DM	INF+DM		
17	Miyasaka 2008	P	ADA	ADA	ADA			
18	Moreland 1999	P	ETA	ETA				
19	Putte 2003	P	ADA	ADA	ADA			
20	Putte 2004	P	ADA	ADA	ADA	ADA		
21	Quinn 2005	P+DM	INF+DM					
22	Schiff 2008	P+DM	ABA+DM	INF+DM				
23	Schiff 2008	INF+DM	ABA+DM					
24	Nishimoto 2004	P	TOC	TOC				
25	Nishimoto 2007	DM	TOC					
26	Nishimoto 2009	P+DM	TOC+DM					
27	Smolen 2009	P+DM	CER+DM	CER+DM				
28	StClair 2004	P+DM	INF+DM	INF+DM				
29	Weinblatt 1999	P+DM	ETA+DM					
30	Weinblatt 2003	P+DM	ADA+DM	ADA+DM	ADA+DM			
31	Westhovens 2006b	P+DM	INF+DM	INF+DM				
32	Zhang 2006	P+DM	INF+DM					
33	Furst 2003	P	ADA					
34	Genovese 2005a	P	ABA					
35	Kay 2008	P+DM	GOL+DM					
36	Keystone 2004	P+DM	ADA+DM	ADA+DM				
37	Keystone 2008a	P+DM	CER+DM	CER+DM				
38	Kim 2007	P+DM	ADA+DM					
39	Klareskog 2004	P+DM	ETA+DM					
40	Kremer 2005	P+DM	ABA+DM	ABA+DM				
41	Keystone 2009a	P+DM	GOL+DM					
42	Cohen 2006	P+DM	RIT+DM					
43	Emery 2008b	P+DM	TOC+DM	TOC+DM				
44	Detert 2013	P+DM	ADA+DM					
45	Kavanaugh 2013	P+DM	ADA+DM					
46	Tak 2011	P+DM	RIT+DM	RIT+DM				
47	Choy 2012	P+DM	CER+DM					

(Continued)

Table 1. (Continued)

	Publication	Treat <sub>arm 1</sub> *	Treat <sub>arm 2</sub>	Treat <sub>arm 3</sub>	Treat <sub>arm 4</sub>	Treat <sub>arm 5</sub>	Treat <sub>arm 6</sub>	Treat <sub>arm 7</sub>
48	Kremer 2010	P+DM	GOL+DM	GOL+DM				
49	Tanaka 2012	P+DM	GOL+DM	GOL+DM				
50	Kremer 2011	P+DM	TOC+DM	TOC+DM				
51	Jones 2010	P+DM	TOC					
52	Yazici 2012	P	TOC					
53	Breedveld 2006	DM	ADA+DM	ADA				
54	Westhovens 2009	P+DM	ABA+DM					
55	Weinblatt 2013	ADA+DM	ABA+DM					

\*P: placebo, DM: DMARD, ADA: adalimumab, CER: certolizumab, ETA: etanercept, GOL: golimumab, INF: infliximab, ANA: anakinra, ABA: abatacept, RIT: rituximab, TOC: tocilizumab.

doi:10.1371/journal.pone.0137258.t001

embraced all treatment and comparator arms over the 54 publications. This way all measured effects of any biologic agent contributed to the comparison of all biologic agents relative to each other either alone or combined with DMARD.

In the following we let P and DM denote placebo and DMARD treatment respectively. Similarly, ADA: adalimumab, CER: certolizumab, ETN: etanercept, GOL: golimumab, INF: infliximab, ANA: anakinra, TOC: tocilizumab, ABA: abatacept and RIT: rituximab. The studies = 1, . . . , S, S = 55, had various numbers of arms, ranging from two to seven.

We present for simplicity first a model without explanatory variables and then include them. Let  $r_{ij}$  denote the number who achieves an ACR50 score in study  $i$  in arm  $j$ ,  $i = 1, . . . , S$ ,





Fig 1. Flow chart. Study selection procedure.

doi:10.1371/journal.pone.0137258.g001



**Fig 2. Direct and indirect comparisons.** DM indicates either DMARD alone or placebo and DMARD treatment together.

doi:10.1371/journal.pone.0137258.g002

$j = 1, \dots, a_i$ , where  $a_i$  is the number of arms in study  $i$ , and  $a_i$  varies between two and seven arms. We let  $m_{ij}$  denote the number of patients in study  $i$  in arm  $j$ .

Let  $p_{ki}$  denote the probability in trial  $i$  to achieve an ACR50 score for treatment  $k$ , where  $k$  could be one of 16 different treatments: CER, ADA, ETN, TOC, ABA, DM/DM+P, INF+DM, ABA+DM, ANA+DM, CER+DM, GOL+DM, ADA+DM, TOC+DM, RIT+DM, ETN+DM and placebo (P).  $r_{ij}$  follows a binomial distribution. We assume that the multiplicative treatment effects relative to placebo are given by the effect-ratios  $\gamma^{kP} = p_{ki}/p_{Pi}$ , where  $k$  denotes treatment with CER, ADA, ETN, TOC, ABA, DM/DM+P, INF+DM, ABA+DM, ANA+DM, CER+DM, GOL+DM, ADA+DM, TOC+DM, RIT+DM and ETN+DM ( $k$  denotes now one of 15 different treatments, that is without placebo). We assume for now that the effect-ratios are constant over studies independent of explanatory variables.

Let  $A_i$  denote all treatment arms in study  $i$ . The likelihood is given by:

$$\prod_{i=1}^S \prod_{j \in A_i} (p_{pi} \gamma^{jP})^{r_{ij}} (1 - p_{pi} \gamma^{jP})^{m_{ij} - r_{ij}}.$$

This Bayesian approach is hence based on the construction of probability distributions for the parameters to be estimated ( $p_{pi}$ , the  $\gamma$ s). This does not mean that these parameters should be interpreted as random variables, but our knowledge of the parameters is uncertain, and we describe this uncertainty through probability distributions.

Probability distributions describing our initial uncertainty are called prior distributions (that is, before the data are collected). When the study results are taken into account, the prior distributions are updated by Bayes' formula to posterior distributions. With little information a priori we choose non-informative priors. Hence, we assume a priori that  $\gamma^{jP} \sim \text{normal}(0, \sigma_\phi^2)$ ,  $j = \text{CER}, \dots, \text{ETN+DM}$  and  $\sigma_\phi \sim \text{uniform}(0, 20)$ .

Parameters to be estimated in the model are the  $p_{pi}$ s,  $\gamma$ s and  $\sigma_\phi$ . We will assume a priori that  $p_{pi}$ ,  $i = 1, \dots, S$ , are uniformly distributed, with an upper limit dependent upon the  $\gamma$ s in study  $i$  in the following way:

$$p_{pi} \sim \text{uniform}(0, \min(1, \min_{j \in A_i} 1/\gamma^{j \cdot P})).$$

Hence, the upper bound of the placebo response probabilities and medication response ratios constrain each other; a larger prior support for the placebo response probabilities ( $p_{pi}$ ) induces a smaller prior support for the medication response ratios (the  $\gamma$ s). One must balance these two as shown in the equation above.

We now consider a model with explanatory variables. When we let the multiplicative treatment effects be dependent upon duration of disease ( $X^V$ ) and dose level ( $X^D$ ), where we let the impact of duration of disease and dose level be independent upon the given drug, we define

$$G_{ij} = \gamma^{jP} e^{\beta_D X_{ij}^D + \beta_V X_{ij}^V}.$$

The likelihood is now given by:

$$\prod_{i=1}^S \prod_{j \in A_i} (p_{pi} G_{ij})^{r_{ij}} (1 - p_{pi} G_{ij})^{m_{ij} - r_{ij}}.$$

Now we assume a priori that

$$p_{pi} \sim \text{uniform}(0, \min(1, \min_{j \in A_i} 1/G_{ij})).$$

The placebo-probabilities  $p_{pi}$  are hence assumed a priori to be uniformly distributed, in a range dependent upon disease duration and dose level in study  $i$ .

We let a priori  $\beta^V \sim \text{normal}(0, \sigma_V^2)$  and similarly  $\beta^D \sim \text{normal}(0, \sigma_D^2)$ . The  $\beta$ s could be independent upon which TNF-inhibitor drug was given (giving one estimated  $\beta$ ), as described above, or dependent giving one  $\beta$  for each of the TNF-inhibitor drugs. We let the a priori distributions for  $\sigma_V$  and  $\sigma_D$  both be uniformly distributed over (0,20).

This model was fitted in WinBUGS[63] run from R[64]. We did 500 000 burn-ins and thereafter 500 000 new updates where we took out every 200. This gave 2 500 samples from the full posterior conditional distribution for each of the parameters entering the model. Table A in S1 File contains the estimated parameters in the model.

We considered varieties of the model where the effect was dependent upon disease duration or dose level or both. The effect of disease duration and drug level could also be drug dependent or not. This resulted in eight model variations to be explored, in addition to one without any explanatory variables.

From the joint model, we estimated the relative effect of each biologic agent versus placebo (with or without DMARD treatment) and versus other biologic agents. As seen in Fig 2 biologic agents were given alone (without DMARD) for five drugs, while joint biologic and DMARD therapy was given for all the nine drugs included. There were  $4 \cdot 5/2 = 10$  unique drug comparisons when given alone and  $9 \cdot 8/2 = 36$  unique drug comparisons when given jointly with DMARD. Altogether this encompasses 46 comparisons of biologic agents against each other.

## Results

We included 54 publications for our MTC regression analysis, published between 1999 and 2013. Overall there were 19 798 patients given biologic treatment therapy, 1 165 given placebo and 8 037 given DMARD or joint DMARD and placebo treatment. The patient characteristic average disease duration for a treatment arm ranged from 0.13 to 13.1 years. Dose level was

**Table 2. Probabilities that one agent was better than another, given alone (top) and together with DMARD (bottom).**

Biological agents given alone								
	ETN	TOC	ABA	ADA				
CER	0.69	0.98	0.89	1				
ETN	-	0.72	0.75	0.98				
TOC	-	-	0.54	1				
ABA	-	-	-	0.87				
Joint biological and DMARD therapy								
	TOC+DM	ANA+DM	RIT+DM	GOL+DM	INF+DM	ABA+DM	ADA+DM	ETN+DM
CER+DM	0.97	0.98	1	1	1	1	1	1
TOC+DM	-	0.73	1	1	1	1	1	1
ANA+DM	-	-	0.88	0.96	0.98	0.99	1	1
RIT+DM	-	-	-	0.87	0.97	0.99	1	1
GOL+DM	-	-	-	-	0.62	0.72	0.9	0.94
INF+DM	-	-	-	-	-	0.7	0.94	0.96
ABA+DM	-	-	-	-	-	-	0.91	0.93
ADA+DM	-	-	-	-	-	-	-	0.76

doi:10.1371/journal.pone.0137258.t002

either low (in 51 arms) or high (in 48 arms). The trials lasted between 12 and 54 weeks, most of them lasting 24 weeks.

In the initial model fitting, the inclusion of explanatory variables did not alter the effect estimates of the agents much compared to not including them. Especially the ranking of the drugs relative to each other did not change. When we examined the impact of disease duration on treatment effect we found no significant effect. This was true both when the effect was drug dependent and when it was not. When we examined the impact of dose level on treatment effect we found that higher doses were associated with higher effect compared to lower doses (statistically significant), with a coefficient for dose level of 0.392, see Table B in [S1 File](#). This was true only when the dose effect was drug independent.


When assuming an impact of both disease duration and dose level on treatment effect we found the effect of disease duration not to be statistically significant and the effect of dose level to be statistically significant when specified as drug independent. Hence, we concluded in our analysis that the disease duration had no impact on the drug effect but that dose level did, and the impact of a high versus low dose level was the same for all biologic drugs examined (in models with a drug dependent dose effect all dose parameters except the one for joint tocilizumab and DMARD treatment versus placebo were not significant).

We obtained two sets of results, one when the drugs were given alone and one when the drugs were given jointly with DMARD.

[Table 2](#) displays the probability that one agent was better than another. The probability that certolizumab was better than etanercept when drugs were given alone was 0.69 while the probability that joint certolizumab and DMARD treatment was better than joint tocilizumab and DMARD treatment was 0.97 and so on. All drugs except etanercept had higher response ratios (drugs versus placebo) when taken jointly with DMARD.

Based on the samples from the posterior distribution the agents were ranked according to the relative effect of drug versus placebo, giving histograms displaying the ranking ([Fig 3](#) and [4](#)). The height of the bars gives the probability of being ranked from position one to last. The effect ratios were the estimated effect of drug versus placebo treatment ([Fig 3](#)) or of joint drug and DMARD versus placebo treatment ([Fig 4](#)), see Table B in [S1 File](#). Certolizumab was ranked






**Fig 3. Histograms and ranking I.** ACR50 effects as the response ratio versus placebo, drugs given alone.

doi:10.1371/journal.pone.0137258.g003

as number one 62.8% of the time when given alone and 95.3% of the time when given with DMARD. Adalimumab was ranked as the least effective 84.4% of the times when given alone. Etanercept was 73.1% of the times ranked as the least effective when given with DMARD. The



**Fig 4. Histograms and ranking II.** ACR50 effects as the response ratio versus placebo, drugs were given together with DMARD.

doi:10.1371/journal.pone.0137258.g004

effect ratio of abatacept versus placebo treatment was 4.42 and the corresponding effect of tocilizumab was close in value, 4.14. Also, the probability that tocilizumab was better than abatacept was close to one half (0.54), indicating that these two drugs were equally effective.

The ranking of the drugs when given alone was, from most to least effective, certolizumab, etanercept, tocilizumab/ abatacept and adalimumab. All drugs were however effective, with response-ratios ranging from 2.54 (credibility interval 2.02 to 3.21) for adalimumab to 7.55 (credibility interval 4.36 to 10.23) for certolizumab.

The ranking of the drugs when given together with DMARD was, from most to least effective, certolizumab, tocilizumab, anakinra, rituximab, golimumab/ infliximab/ abatacept and adalimumab/etanercept. The effect ratio of joint golimumab and DMARD versus placebo treatment was 4.61 and the corresponding effects of joint infliximab and DMARD and joint abatacept and DMARD versus placebo treatment were 4.43 and 4.28 respectively. The probabilities that joint golimumab and DMARD treatment was better than joint infliximab and DMARD and joint abatacept and DMARD treatment were 0.62 and 0.72 respectively. The probability that joint infliximab and DMARD treatment was better than joint abatacept and DMARD treatment was 0.7. Hence, one could argue that these three drugs were not that different with respect to effect.

The effect ratio of joint adalimumab and DMARD versus placebo treatment was 4.00 and the corresponding effect of joint etanercept and DMARD treatment was 3.71. Also, the probability that joint adalimumab and DMARD treatment was better than joint etanercept and DMARD treatment was 0.76. These two treatments also seem to be similar with respect to effect. All agents were effective when given together with DMARD, with response-ratios ranging from 3.71 (credibility interval 2.74 to 5.05) for etanercept to 10.99 (credibility interval 7.31 to 16.42) for certolizumab.

The ranking of drugs given jointly with DMARD was for certolizumab, abatacept and adalimumab the same as the ranking when drugs were given without DMARD treatment. Etanercept was on the other hand ranked higher and tocilizumab lower when given alone compared to given with DMARD treatment. Tocilizumab had an almost twice as high response ratio while the effect of etanercept was almost halved when given with DMARD. Hence, unlike the other drugs, etanercept had a much higher effect when given alone than with joint DMARD treatment, but the uncertainty concerning the effect of exclusive etanercept treatment was very large.

## Discussion

All biological agents were effective compared to placebo, with certolizumab the most effective. The drugs were in general more effective when given together with DMARDs which was in harmony with current guidelines [5]. The dose level influenced the treatment effect (statistically significant), with a higher dose indicating a larger effect. We found no significant difference between longer versus shorter disease duration on the treatment effect. Higher doses could of course also lead to more adverse effects, an issue not focused on in this paper, and our finding should be interpreted with this in mind. Still, as various studies and even treatment arms had different dosing regimens it was important to correct for it in the analysis, giving estimated relative drug effects not confounded by dose level.

The way patients have been included in the trials and treated throughout the trial period could have changed over the publication years 1999 through 2013, possibly biasing the results. Clearly this would be a problem if some drugs were studied in earlier years and others in later years. This was not the case. Also, we found no apparent time trends regarding disease duration. The variation in trial duration could also potentially bias the ranking of the treatments. However, the different biologic agents were reasonably similarly distributed with respect to trial duration, with trials involving anakinra as perhaps an exception, all lasting only 24 or 26 weeks.

Previous MTC analyses of clinical trials reporting effects of biologic drugs indicated different rankings of the drugs [6], and this analysis has been an additional contribution.

For example, Nixon et al. [65] compared the effect of etanercept, infliximab, adalimumab and anakinra in 13 publications, none with head-to-head comparisons. Different models gave varying degrees of efficacy. In a final model they found anakinra to be the least effective. Singh et al. [66] compared abatacept, adalimumab, etanercept, infliximab, rituximab and anakinra from 27 clinical trials, none contained head-to-head comparisons. They reported a low effect of anakinra based on the results from three trials only. Divine et al. [67] and Turkstra et al. [68] both compared all nine biologic drugs and also ranked anakinra as the least effective drug. However, in all the mentioned publications there was little evidence of statistically significant differences in effects between anakinra and the other biologic drugs. In our analysis, anakinra (together with DMARD treatment) was ranked as number three. Golimumab was also ranked as relatively low in effect in both Divine et al. [67] and Mandema et al. [69], while this study ranked it somewhat higher.

Thorlund et al. [6] stated: “*Since future RA MTCs will undoubtedly continue to include both monotherapy and combination therapy trials, it is important to establish whether the concomitant use of DMARDs does in fact yield an effect modification on the relative comparative treatment effects. This will require more refined analyses than what have currently been performed among the MTCs*”. Our approach answers this request by distinguishing between joint DMARD or exclusively biologic drug use, and by letting this difference be drug dependent. And we find that for especially etanercept treatment this distinction truly matters. Our comparison network (Fig 2) resembles the one suggested in Thorlund et al. [6].

An advantage of our approach was the capability to include results from many trials into one model simultaneously, whether the trials compared biologic drugs alone or with DMARD treatment. In previous MTC-analyses only Mandema et al. [69] reported many trials (50), but it was unclear how many went into the analysis of ranking the drugs with respect to the ACR50 score. Our analysis included both direct and indirect comparisons from 54 publications in the same network, and we interpreted the results by probabilities tied to comparisons and ranking of the drugs. This approach could be particularly useful in situations when new drugs are introduced to establish a possible therapeutic benefit compared to already existing drugs.

A possible weakness of any such MTC analysis is that the trials combined might differ with respect to patient characteristics, follow-up time or other trial procedures, potentially introducing heterogeneity in the treatment effects. An MTC regression analysis will try to capture some of the possible differences, and we have considered the impact of dose level and disease duration in this analysis. Other explanatory variables could also have been considered, as geographical information on trial location or more detailed information on previous patient history. For some trials this could be available information, but not always. Correcting for many factors could exclude studies due to lack of information, introducing selection bias.

The drugs were ranked with respect to efficacy. We did not consider adverse events. Of course, adverse events are also important. This was however a more complicated task as the publications reported adverse events differently. Also, some publications did not report adverse events at all.

With many MTC analyses within the RA field published over the last five years with slightly different results it is clear that ranking these drugs has been challenging. The final result would depend on the chosen model, which trials are included, how heterogeneity is handled, if and which explanatory variables are taken into consideration and if and how joint treatment with other drugs are considered. We believe that our unified approach of including all relevant comparisons, both direct and indirect comparisons that exist, while specifying whether the drugs

were given alone or in combination with DMARDs, is a sensible approach giving the possibility to rank all biologic drugs with respect to comparative effectiveness.

## Supporting Information

### S1 PRISMA Checklist.

(DOC)

### S1 File.

(DOCX)

## Author Contributions

Conceived and designed the experiments: IFT BN JG MK. Performed the experiments: IFT MK MR NM. Analyzed the data: IFT BN JG MK. Wrote the paper: IFT BN JG MK MR NM.

## References

1. Lard LR, Visser H, Spreyer I, et al. Early versus delayed treatment in patients with recent-onset rheumatoid arthritis: comparison of two cohorts who received different treatment strategies. *Am J Med* 2001; 111: 446–51. PMID: [11690569](#)
2. Olsen NJ, Stein M. New drugs for rheumatoid arthritis. *N Engl J Med* 2004; 350: 2167–79. *J Rheumatol Suppl.* 2002 Nov; 66:3–8. PMID: [15152062](#)
3. Emery P. Evidence supporting the benefit of early intervention in rheumatoid arthritis. *J Rheumatol Suppl.* 2002 Nov; 66: 3–8. PMID: [12435162](#)
4. <http://www.nhs.uk/Conditions/Rheumatoid-arthritis/Pages/Treatment.aspx>
5. Singh JA, Furst DE, Bharat A, Curtis JR, Kavanaugh AF, Kremer JM, et al. Update of the 2008 American College of Rheumatology recommendations for the use of disease-modifying antirheumatic drugs and biologic agents in the treatment of rheumatoid arthritis. *Arthritis Care & Research* May 2012; Vol. 64, No. 5, 625–639.
6. Thortlund K, Druyts E, Aviña-Zubieta JA, Wu P, Mills EJ. Why the findings of published multiple treatment comparison meta-analyses of biologic treatments for rheumatoid arthritis are different: an overview of recurrent methodological shortcomings. *Ann Rheum Dis* 2013; 72: 1524–1535 doi: [10.1136/annrheumdis-2012-201574](#) PMID: [23087184](#)
7. Abe T, Takeuchi T, Miyasaka N, Hashimoto H, Kondo H, Ichikawa Y, et al. A Multicenter, Double-Blind, Randomized, Placebo Controlled Trial of Infliximab Combined with Low Dose Methotrexate in Japanese Patients with Rheumatoid Arthritis. *J Rheumatol* 2006; 33; 37–44. PMID: [16395748](#)
8. Emery P, Keystone E, Tony HP, Cantagrel A, van Vollenhoven R, Sanchez A, et al. IL-6 receptor inhibition with tocilizumab improves treatment outcomes in patients with rheumatoid arthritis refractory to anti-tumour necrosis factor biologicals: results from a 24-week multicentre randomised placebo-controlled trial. *Ann Rheum Dis* 2008; 67:1516–1523. doi: [10.1136/ard.2008.092932](#) PMID: [18625622](#)
9. Genovese MC, McKay JD, Nasonov EL, Mysler EF, da Silva NA, Alecock E, et al. Interleukin-6 receptor inhibition with tocilizumab reduces disease activity in rheumatoid arthritis with inadequate response to disease-modifying antirheumatic drugs: the tocilizumab in combination with traditional disease-modifying antirheumatic drug therapy study. *Arthritis Rheum.* 2008; 58:2968–80. doi: [10.1002/art.23940](#) PMID: [18821691](#)
10. Maini RN, Taylor PC, Szechinski J, Pavelka K, Bröll J, Balint G, et al. Double-Blind Randomized Controlled Clinical Trial of the Interleukin-6 Receptor Antagonist, Tocilizumab, in European Patients With Rheumatoid Arthritis Who Had an Incomplete Response to Methotrexate. *ARTHRITIS & RHEUMATISM* Vol. 54, No. 9, September 2006, 2817–2829.
11. Smolen JS, Beaulieu A, Rubbert-Roth A, Ramos-Remus C, Rovinsky J, Alecock E, et al. Effect of interleukin-6 receptor inhibition with tocilizumab in patients with rheumatoid arthritis (OPTION study): a double-blind, placebo-controlled, randomised trial. *Lancet* 2008; 371: 987–97. doi: [10.1016/S0140-6736\(08\)60453-5](#) PMID: [18358926](#)
12. Chen DY, Chou SJ, Tsu-Yi Hsieh TY, Chen YH, Chen HH, Hsieh CW, et al. Randomized, Double-blind, Placebo-controlled, Comparative Study of Human Anti-TNF Antibody Adalimumab in Combination with Methotrexate and Methotrexate Alone in Taiwanese Patients with Active Rheumatoid Arthritis. *J Formos Med Assoc* 2009, Vol 108, No 4.

13. Cohen S, Hurd E, Cush J, Schiff M, Weinblatt ME, Moreland LW, et al. Treatment of Rheumatoid Arthritis With Anakinra, a Recombinant Human Interleukin-1 Receptor Antagonist, in Combination With Methotrexate. *ARTHRITIS & RHEUMATISM*. Vol. 46, No. 3, March 2002, 614–624.
14. Cohen SB, Moreland LW, Cush JJ, Greenwald MW, Block S, Shergy WJ, et al for the 990145 Study Group. A multicentre, double blind, randomised, placebo controlled trial of anakinra (Kineret), a recombinant interleukin 1 receptor antagonist, in patients with rheumatoid arthritis treated with background methotrexate. *Ann Rheum Dis* 2004; 63:1062–1068. PMID: [15082469](#)
15. Edwards JCW, Szczepanski L, Szechinski J, Filipowicz-Sosnowska A, Emery P, Close DR, et al. Efficacy of B-Cell–Targeted Therapy with Rituximab in Patients with Rheumatoid Arthritis. *N Engl J Med* 2004; 350:2572–81. PMID: [15201414](#)
16. Emery P, Fleischmann R, Filipowicz-Sosnowska A, Schechtman J, Szczepanski L, Kavanaugh A, et al for the DANCER Study Group. The Efficacy and Safety of Rituximab in Patients With Active Rheumatoid Arthritis Despite Methotrexate Treatment. *ARTHRITIS & RHEUMATISM* Vol. 54, No. 5, May 2006b, 1390–1400.
17. Emery P, Fleischmann RM, Moreland LW, Hsia EC, Strusberg I, Durez P, et al. Golimumab, a Human Anti–Tumor Necrosis Factor Monoclonal Antibody, Injected Subcutaneously Every Four Weeks in Methotrexate-Naive Patients With Active Rheumatoid Arthritis. *ARTHRITIS & RHEUMATISM* Vol. 60, No. 8, August 2009, 2272–2283.
18. Emery P, Deodhar A, Rigby WF, Isaacs JD, Combe B, Racewicz AJ, et al. Efficacy and safety of different doses and retreatment of rituximab: a randomised, placebo-controlled trial in patients who are biological naïve with active rheumatoid arthritis and an inadequate response to methotrexate (Study Evaluating Rituximab’s Efficacy in MTX in adequate responders (SERENE)). *Ann Rheum Dis* 2010; 69:1629–1635. doi: [10.1136/ard.2009.119933](#) PMID: [20488885](#)
19. Fleischmann R, Vencovsky J, van Vollenhoven RF, Borenstein D, Box J, Coteur G, et al. Efficacy and safety of certolizumab pegol monotherapy every 4 weeks in patients with rheumatoid arthritis failing previous diseasemodifying antirheumatic therapy: the FAST4WARD study. *Ann Rheum Dis* 2009; 68:805–811. doi: [10.1136/ard.2008.099291](#) PMID: [19015206](#)
20. Kremer JM, Genant HK, Moreland LW, Russell AS, Emery P, Abud-Mendoza C, et al. Effects of Abatacept in Patients with Methotrexate-Resistant Active Rheumatoid Arthritis. *Ann Intern Med*. 2006; 144:865–876. PMID: [16785475](#)
21. Lipsky PE, van der Heijde D, St. Clair W, Furst DE, Breedveld FC, Kalden JRK, et al for the Anti–Tumor Necrosis Factor T Infliximab and Methotrexate in the Treatment of Rheumatoid Arthritis trial in Rheumatoid Arthritis with Concomitant Therapy Study Group. *N Engl J Med* 2000; 343:1594–1602. PMID: [11096166](#)
22. Maini R, St Clair EW, Breedveld F, Furst D, Kalden J, Weisman M, et al for the ATTRACT Study Group. Infliximab (chimeric anti-tumour necrosis factor  $\alpha$  monoclonal antibody) versus placebo in rheumatoid arthritis patients receiving concomitant methotrexate: a randomised phase III trial. *Lancet* 1999; 354: 1932–39. PMID: [10622295](#)
23. Miyasaka N: The CHANGE Study Investigators. Clinical investigation in highly disease-affected rheumatoid arthritis patients in Japan with adalimumab applying standard and general evaluation: the CHANGE study. *Mod Rheumatol* (2008) 18: 252–262. doi: [10.1007/s10165-008-0045-0](#) PMID: [18330677](#)
24. Moreland LW, Schiff MH, Baumgartner SW, Tindall EA, Fleischmann RM, Bulpitt KJ, et al. Etanercept Therapy in Rheumatoid Arthritis: A Randomized, Controlled Trial. *Ann Intern Med*. 1999; 130: 478–486. PMID: [10075615](#)
25. van de Putte LBA, Rau R, Breedveld FC, Kalden JK, Malaise MG, van Riel PLCM, et al. Efficacy and safety of the fully human anti-tumour necrosis factor  $\alpha$  monoclonal antibody adalimumab (D2E7) in DMARD refractory patients with rheumatoid arthritis: a 12 week, phase II study. *Ann Rheum Dis* 2003; 62:12 1168–1177.
26. van de Putte LBA, Atkins C, Malaise M, Sany J, Russell AS, van Riel PLCM, et al. Efficacy and safety of adalimumab as monotherapy in patients with rheumatoid arthritis for whom previous disease modifying antirheumatic drug treatment has failed. *Ann Rheum Dis* 2004; 63:5 508–516.
27. Quinn MA, Conaghan PG, O’Connor PJ, Karim Z, Greenstein A, Brown A, et al. Very Early Treatment With Infliximab in Addition to Methotrexate in Early, Poor-Prognosis Rheumatoid Arthritis Reduces Magnetic Resonance Imaging Evidence of Synovitis and Damage, With Sustained Benefit After Infliximab Withdrawal. *ARTHRITIS & RHEUMATISM* Vol. 52, No. 1, January 2005, 27–35.
28. Schiff M, Keiserman M, Codding C, Songcharoen S, Berman A, Naylor S, et al. Efficacy and safety of abatacept or infliximab vs placebo in ATTEST: a phase III, multi-centre, randomised, double-blind, placebo-controlled study in patients with rheumatoid arthritis and an inadequate response to methotrexate. *Ann Rheum Dis*. 2008 Aug; 67(8): 1096–103. PMID: [18055472](#)

29. Nishimoto N, Yoshizaki K, Miyasaka N, Yamamoto K, Kawai S, Takeuchi T, et al. Treatment of rheumatoid arthritis with humanized anti-interleukin-6 receptor antibody: a multicenter, double-blind, placebo-controlled trial. *Arthritis Rheum.* 2004; 50: 1761–9. PMID: [15188351](#)
30. Nishimoto N, Hashimoto J, Miyasaka N, Yamamoto K, Kawai S, Takeuchi T, et al. Study of active controlled monotherapy used for rheumatoid arthritis, an IL-6 inhibitor (SAMURAI): evidence of clinical and radiographic benefit from an x ray reader-blinded randomised controlled trial of tocilizumab. *Ann Rheum Dis* 2007; 66: 1162–1167. PMID: [17485422](#)
31. Nishimoto N, Miyasaka N, Yamamoto K, Kawai S, Takeuchi T, Azuma J, et al. Long-term safety and efficacy of tocilizumab, an anti-IL-6 receptor monoclonal antibody, in monotherapy, in patients with rheumatoid arthritis (the STREAM study): evidence of safety and efficacy in a 5-year extension study. *Ann Rheum Dis* 2009; 68: 1580–4. doi: [10.1136/ard.2008.092866](#) PMID: [19019888](#)
32. Smolen J, Landewe RB, Mease P, Brzezicki J, Mason D, Luijckens K, et al. Efficacy and safety of certolizumab pegol plus methotrexate in active rheumatoid arthritis: the RAPID 2 study. A randomised controlled trial. *Ann Rheum Dis* 2009; 68: 797–804. doi: [10.1136/ard.2008.101659](#) PMID: [19015207](#)
33. St. Clair EW, van der Heijde DMFM, Smolen JS, Maini RN, Bathon JM, Emery P, et al for the Active-Controlled Study of Patients Receiving Infliximab for the Treatment of Rheumatoid Arthritis of Early Onset Study Group. Combination of infliximab and methotrexate therapy for early rheumatoid arthritis: A randomized, controlled trial. *Arthritis & Rheumatism*, 2004, 50: 3432–3443.
34. Weinblatt ME, Kremer JM, Bankhurst AD, Bulpitt KJ, Fleischmann RM, Fox RI, et al. A trial of etanercept, a recombinant tumor necrosis factor receptor: Fc fusion protein, in patients with rheumatoid arthritis receiving methotrexate. *N Engl J Med.* 1999 Jan 28; 340(4): 253–9. PMID: [9920948](#)
35. Weinblatt ME, Keystone EC, Furst DE, Moredland LW, Weisman MH, Birbara CA, et al. Adalimumab, a fully human anti-tumor necrosis factor alpha monoclonal antibody, for the treatment of rheumatoid arthritis in patients taking concomitant methotrexate: the ARMADA trial. *Arthritis Rheum* 2003; 48: 35–45. PMID: [12528101](#)
36. Westhovens R, Yocum D, Han J, Berman A, Strusberg I, Geusens P, et al for the START Study Group. The Safety of Infliximab, Combined With Background Treatments, Among Patients With Rheumatoid Arthritis and Various Comorbidities. *ARTHRITIS & RHEUMATISM* Vol. 54, No. 4, April 2006, 1075–1086.
37. Zhang FC, Hou Y, Huang F, Wu DH, Bao CD, Ni LQ, et al. Infliximab versus placebo in rheumatoid arthritis patients receiving concomitant methotrexate: a preliminary study from China. *APLAR J Rheumatol* 2006, 9(2): 127–130,
38. Furst DE, Schiff MH, Fleischmann RM, Strand V, Birbara CA, Compagnone D, et al. Adalimumab, a fully human anti tumor necrosis factor-alpha monoclonal antibody, and concomitant standard antirheumatic therapy for the treatment of rheumatoid arthritis: results of STAR (Safety Trial of Adalimumab in Rheumatoid Arthritis). *J Rheumatol* December 2003 30(12): 2563–2571.
39. Genovese M, Becker J-C, Schiff M, Luggen M, Sherrer Y, Kremer J, et al: Abatacept for rheumatoid arthritis refractory to tumor necrosis factor alpha inhibition. *N Engl J Med* 2005; 353: 1114–23. PMID: [16162882](#)
40. Kay J, Matteson EL, Dasgupta B, Nash P, Durez P, Hall S, et al. Golimumab in patients with active rheumatoid arthritis despite treatment with methotrexate: a randomized, double-blind, placebo controlled, dose-ranging study. *Arthritis Rheum* 2008a; 58: 964–75.
41. Keystone EC, Kavanaugh AF, Sharp JT, Tannenbaum H, Hua Y, Teoh LS, et al. Radiographic, Clinical, and Functional Outcomes of Treatment With Adalimumab (a Human Anti-Tumor Necrosis Factor Monoclonal Antibody) in Patients With Active Rheumatoid Arthritis Receiving Concomitant Methotrexate Therapy. *ARTHRITIS & RHEUMATISM* Vol. 50, No. 5, May 2004, 1400–1411.
42. Keystone E, van der Heijde D, Mason D Jr, Landewé R, Vollenhoven RV, Combe B, et al. Certolizumab pegol plus methotrexate is significantly more effective than placebo plus methotrexate in active rheumatoid arthritis: findings of a fifty-two-week phase III, multicenter, randomized, double-blind, placebo-controlled, parallel-group study. *Arthritis Rheum* 2008; 58: 3319–29. doi: [10.1002/art.23964](#) PMID: [18975346](#)
43. Kim HY, Lee SK, Song YW, Yoo DH, Koh EM, Bin Yoo B, et al. A randomized, double-blind, placebo-controlled, phase III study of the human anti-tumor necrosis factor antibody adalimumab administered as subcutaneous injections in Korean rheumatoid arthritis patients treated with Methotrexate. *APLAR Journal of Rheumatology* 2007; 10: 9–16.
44. Klareskog L, van der Heijde D, de Jager JP, Gough A, Kalden J, Malaise M, et al for TEMPO (Trial of Etanercept and Methotrexate with Radiographic Patient Outcomes) study investigators. Therapeutic effect of the combination of etanercept and methotrexate compared with each treatment alone in patients with rheumatoid arthritis: double-blind randomised controlled trial. *Lancet.* 2004 Feb 28; 363 (9410): 675–81. PMID: [15001324](#)

45. Kremer JM, Dougados M, Emery P, Durez P, Sibilia J, Shergy W, et al. Treatment of Rheumatoid Arthritis With the Selective Costimulation Modulator Abatacept. *ARTHRITIS & RHEUMATISM* Vol. 52, No. 8, August 2005, 2263–2271.
46. Keystone EC, Genovese MC, Klareskog L, Hsia EC, Hall ST, Miranda PC, et al. Golimumab, a human antibody to tumour necrosis factor a given by monthly subcutaneous injections, in active rheumatoid arthritis despite methotrexate therapy: the GO-FORWARD Study. *Ann Rheum Dis* 2009; 68:789–796. doi: [10.1136/ard.2008.099010](https://doi.org/10.1136/ard.2008.099010) PMID: [19066176](https://pubmed.ncbi.nlm.nih.gov/19066176/)
47. Cohen SB, Emery P, Greenwald MW, Dougados M, Furie RA, Genovese MC, et al for the REFLEX Trial Group. Rituximab for Rheumatoid Arthritis Refractory to Anti-Tumor Necrosis Factor Therapy: Results of a Multicenter, Randomized, Double-Blind, Placebo-Controlled, Phase III Trial Evaluating Primary Efficacy and Safety at Twenty-Four Weeks. *ARTHRITIS & RHEUMATISM*; Vol. 54, No. 9, September 2006, 2793–2806.
48. Emery P, Breedveld FC, Hall S, Durez P, Chang DJ, Robertson D, et al. Comparison of methotrexate monotherapy with a combination of methotrexate and etanercept in active, early, moderate to severe rheumatoid arthritis (COMET): a randomised, double-blind, parallel treatment trial. *Lancet* 2008b; 372: 375–82.
49. Detert J, Bastian H, Listing J, Weiß A, Wassenberg S, Liebhaber A, et al. Induction therapy with adalimumab plus methotrexate for 24 weeks followed by methotrexate monotherapy up to week 48 versus methotrexate therapy alone for DMARD-naïve patients with early rheumatoid arthritis: HIT HARD, an investigator-initiated study. *Ann Rheum Dis* 2013; 72: 844–850. doi: [10.1136/annrheumdis-2012-201612](https://doi.org/10.1136/annrheumdis-2012-201612) PMID: [22739990](https://pubmed.ncbi.nlm.nih.gov/22739990/)
50. Kavanaugh A, Fleischmann RM, Emery P, Kupper H, Redden L, Guette B, et al. Clinical, functional and radiographic consequences of achieving stable low disease activity and remission with adalimumab plus methotrexate or methotrexate alone in early rheumatoid arthritis: 26-week results from the randomised, controlled OPTIMA study. *Ann Rheum Dis* 2013; 72: 64–71. doi: [10.1136/annrheumdis-2011-201247](https://doi.org/10.1136/annrheumdis-2011-201247) PMID: [22562973](https://pubmed.ncbi.nlm.nih.gov/22562973/)
51. Tak PP, Rigby WF, Rubbert-Roth A, Peterfy CG, van Vollenhoven RF, Stohl W, et al. Inhibition of joint damage and improved clinical outcomes with rituximab plus methotrexate in early active rheumatoid arthritis: the IMAGE trial. *Ann Rheum Dis* 2011; 70: 39–46. doi: [10.1136/ard.2010.137703](https://doi.org/10.1136/ard.2010.137703) PMID: [20937671](https://pubmed.ncbi.nlm.nih.gov/20937671/)
52. Kremer JM, Blanco R, Brzosko M, Burgos-Vargas R, Halland AM, Vernon E, et al. Tocilizumab Inhibits Structural Joint Damage in Rheumatoid Arthritis Patients with an Inadequate Response to Methotrexate: results from the double-blind treatment phase of a randomized placebo-controlled trial of tocilizumab safety and prevention of structural joint damage at one year. *Arthritis Rheum* 2011; 63: 609–621. doi: [10.1002/art.30158](https://doi.org/10.1002/art.30158) PMID: [21360490](https://pubmed.ncbi.nlm.nih.gov/21360490/)
53. Choy E, McKenna F, Vencovsky J, Valente R, Goel N, VanLunen B, Davies O, et al. Certolizumab pegol plus MTX administered every 4 weeks is effective in patients with RA who are partial responders to MTX. *Rheumatology* 2012; 51: 1226–1234. doi: [10.1093/rheumatology/ker519](https://doi.org/10.1093/rheumatology/ker519) PMID: [22344576](https://pubmed.ncbi.nlm.nih.gov/22344576/)
54. Tanaka Y, Harigai M, Takeuchi T, Yamanaka H, Ishiguro N, Yamamoto K, et al. Golimumab in combination with methotrexate in Japanese patients with active rheumatoid arthritis: results of the GO-FORTH study. *Ann Rheum Dis* 2012; 71: 817–24. doi: [10.1136/ard.2011.200317](https://doi.org/10.1136/ard.2011.200317) PMID: [22121129](https://pubmed.ncbi.nlm.nih.gov/22121129/)
55. Kremer J, Ritchlin C, Mendelsohn A, Baker D, Kim L, Xu Z, et al. Golimumab, a New Human Anti-Tumor Necrosis Factor-Antibody, Administered Intravenously in Patients With Active Rheumatoid Arthritis: Forty-Eight-Week Efficacy and Safety Results of a Phase III Randomized, Double-Blind, Placebo-Controlled Study. *ARTHRITIS & RHEUMATISM* Vol. 62, No. 4, April 2010, 917–928.
56. Jones G, Sebba A, Gu J, Lowenstein MB, Calvo A, Gomez-Reino JJ, et al. Comparison of tocilizumab monotherapy versus methotrexate monotherapy in patients with moderate to severe rheumatoid arthritis: the AMBITION study. *Ann Rheum Dis* 2010; 69: 88–96. doi: [10.1136/ard.2008.105197](https://doi.org/10.1136/ard.2008.105197) PMID: [19297346](https://pubmed.ncbi.nlm.nih.gov/19297346/)
57. Yazici Y, Curtis JR, Ince A, Baraf H, Malamet RL, Teng LL, et al. Efficacy of tocilizumab in patients with moderate to severe active rheumatoid arthritis and a previous inadequate response to disease-modifying antirheumatic drugs: the ROSE study. *Ann Rheum Dis* 2012; 71: 198–205. doi: [10.1136/ard.2010.148700](https://doi.org/10.1136/ard.2010.148700) PMID: [21949007](https://pubmed.ncbi.nlm.nih.gov/21949007/)
58. Breedveld FC1, Weisman MH, Kavanaugh AF, Cohen SB, Pavelka K, van Vollenhoven R, et al. The PREMIER study: A multicenter, randomized, double-blind clinical trial of combination therapy with adalimumab plus methotrexate versus methotrexate alone or adalimumab alone in patients with early, aggressive rheumatoid arthritis who had not had previous methotrexate treatment. *Arthritis Rheum*. 2006 Jan; 54 (1): 26–37. PMID: [16385520](https://pubmed.ncbi.nlm.nih.gov/16385520/)
59. Westhovens R, Robles M, Ximenes AC, Nayiager S, Wollenhaupt J, Durez P, et al. Clinical efficacy and safety of abatacept in methotrexate-naïve patients with early rheumatoid arthritis and poor prognostic factors. *Ann Rheum Dis* 2009; 68: 1870–7. doi: [10.1136/ard.2008.101121](https://doi.org/10.1136/ard.2008.101121) PMID: [19124524](https://pubmed.ncbi.nlm.nih.gov/19124524/)



60. Weinblatt ME, Schiff M, Valente R, Heijde D, Citera G, Zhao C, et al. Head-to-head comparison of subcutaneous abatacept versus adalimumab for rheumatoid arthritis: Findings of a phase IIIb, multinational, prospective, randomized study. *Arthritis Rheum*. 2013 Jan; 65(1): 28–38. doi: [10.1002/art.37711](https://doi.org/10.1002/art.37711) PMID: [23169319](https://pubmed.ncbi.nlm.nih.gov/23169319/)
61. <http://www.rheumatology.org/>
62. Klemp M, Tvette IF, Skomedal T, Gaasemyr J, Natvig B, Aursnes I. A review and bayesian meta-analysis of clinical efficacy and adverse effects of 4 atypical neuroleptic drugs compared with haloperidol and placebo. *J Clin Psychopharmacol* 2011; 31, 6, 698–704.
63. Thomas A. BUGS: a statistical modelling package RTA/BCS Modular Languages Newsletter. 1994; Vol 2, 36–38.
64. R Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0, URL <http://www.R-project.org/>.
65. Nixon RM, Bansback N, Brennan A. Using mixed treatment comparisons and meta-regression to perform indirect comparisons to estimate the efficacy of biologic treatments in rheumatoid arthritis. *Statist. Med*. 2007; 26: 1237–1254.
66. Singh JA, Christensen R, Wells GA, Suarez-Almazor ME, Buchbinder R, Lopez-Olivo MA, et al. A network meta-analysis of randomized controlled trials of biologics for rheumatoid arthritis: a Cochrane overview *CMAJ*. 2009 Nov 24; 181(11):787–96. doi: [10.1503/cmaj.091391](https://doi.org/10.1503/cmaj.091391) PMID: [19884297](https://pubmed.ncbi.nlm.nih.gov/19884297/)
67. Divine EB, Alfonso-Cristancho R, Sullivan SD. Effectiveness of biologic therapies for rheumatoid arthritis: an indirect comparisons approach. *Pharmacotherapy* 2011; Vol 31, No 1.
68. Turkstra E, Ng SK, Scuffham PA. A mixed treatment comparison of the short-term efficacy of biologic disease modifying anti-rheumatic drugs in established rheumatoid arthritis. *Current Medical Research & Opinion* 2011; Vol. 27, No. 10, 1885–1897.
69. Mandema JW, Salinger DH, Gibbs MA. A dose–response meta-analysis for quantifying relative efficacy of biologics in rheumatoid arthritis. *Clin Pharmacol Ther*. 2011 Dec; 90(6):828–35. doi: [10.1038/clpt.2011.256](https://doi.org/10.1038/clpt.2011.256) PMID: [22048227](https://pubmed.ncbi.nlm.nih.gov/22048227/)