

# BMJ Open Quantifying audio visual alcohol imagery in popular Indian films: a content analysis

Rohith Bhagawath,<sup>1</sup> Muralidhar M Kulkarni,<sup>1</sup> John Britton,<sup>2</sup> Jo Cranwell,<sup>3</sup> Monika Arora,<sup>4,5</sup> Gaurang P Nazar,<sup>4,5</sup> Somya Mullapudi,<sup>1</sup> Veena G Kamath<sup>1</sup>

**To cite:** Bhagawath R, Kulkarni MM, Britton J, *et al*. Quantifying audio visual alcohol imagery in popular Indian films: a content analysis. *BMJ Open* 2021;**11**:e040630. doi:10.1136/bmjopen-2020-040630

► Prepublication history and additional material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2020-040630>).

Received 29 May 2020

Revised 04 January 2021

Accepted 23 January 2021



© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY. Published by BMJ.

<sup>1</sup>Community Medicine, Kasturba Medical College, Manipal Academy of Higher Education, Manipal, Karnataka, India

<sup>2</sup>Division of Epidemiology and Public Health, University of Nottingham, Nottingham, UK

<sup>3</sup>Tobacco Control Research Group, Department for Health, University of Bath, Bath, UK

<sup>4</sup>Health Related Information Dissemination Amongst Youth, HRIDAY, New Delhi, India

<sup>5</sup>Health Promotion Division, Public Health Foundation of India, New Delhi, India

## Correspondence to

Professor John Britton; [j.britton@outlook.com](mailto:j.britton@outlook.com)

## ABSTRACT

**Objectives** Though exposure to alcohol imagery in films is a significant determinant of uptake and severity of alcohol consumption among young people, there is poor evidence regarding the content of alcohol imagery in films in low-income and middle-income countries. We have measured alcohol imagery content and branding in popular Indian films, in total and in relation to language and age rating.

**Design** In this observational study we measured alcohol imagery semiquantitatively using 5-minute interval coding. We coded each interval according to whether it contained alcohol imagery or brand appearances.

**Setting** India.

**Participants** None. Content analysis of a total of 30 national box office hit films over a period of 3 years from 2015 to 2017.

**Primary and secondary outcome measures** To assess alcohol imagery in Indian films and its distribution in relation to age and language rating has been determined.

**Results** The 30 films included 22 (73%) Hindi films and 8 (27%) in regional languages. Seven (23%) were rated suitable for viewing by all ages (U), and 23 (77%) rated as suitable for viewing by children subject to parental guidance for those aged under 12 (UA). Any alcohol imagery was seen in 97% of the films, with 195 of a total of 923 5-minute intervals, and actual alcohol use in 25 (83%) films, in 90 (10%) intervals. The occurrence of these and other categories of alcohol imagery was similar in U-rated and UA-rated films, and in Hindi and local language films. Episodes of alcohol branding occurred in 10 intervals in five films.

**Conclusion** Almost all films popular in India contain alcohol imagery, irrespective of age rating and language. Measures need to be undertaken to limit alcohol imagery in Indian films to protect the health of young people, and to monitor alcohol imagery in other social media platforms in future.

## BACKGROUND

Consumption of alcohol is a major avoidable cause of morbidity and mortality.<sup>1</sup> According to the WHO, there are 2.3 billion current alcohol users around the world,<sup>2</sup> and in 2016 alcohol consumption led to 2.8 million deaths.<sup>3</sup> Global alcohol consumption has increased by 38% in the last decade,<sup>4</sup>

## Strengths and limitations of this study

- Our study, to the best of our knowledge is the first of its kind to measure alcohol imagery in popular films in India.
- The study demonstrates that almost all popular Indian films contain alcohol imagery.
- We included only top 10 national box office hit films in the years 2015, 2016 and 2017 since coding films is time-consuming.
- Films in only four Indian languages figured in the box office hits; while films in several other regional languages did not.
- Although our study did not include any A-rated (adult) or S-rated (restricted to specialised audience) films, our findings related entirely to films classified by the Central Board of Film Certification as suitable for young people.

indicating that the burden of morbidity and mortality caused by alcohol is likely to grow.

Alcohol consumption is a significant public health problem in India.<sup>5</sup> Although rare among women, nearly one in three men now consume alcohol,<sup>6</sup> with an average intake of 18.3 L per year.<sup>2</sup> Although the mean age at which people begin consuming alcohol is 21 years,<sup>7 8</sup> 1.3% of children aged 10–17 report alcohol consumption,<sup>9</sup> and studies reveal that early onset of use of alcohol in India correlates with chronic heavy drinking patterns later in adult life.<sup>7 8</sup>

Research among adolescents in developed countries indicates that in addition to paid-for advertising, exposure to media imagery of alcohol products and alcohol consumption, particularly in films, plays an important role in promoting the uptake of alcohol consumption among young people.<sup>10</sup> However, while national film age-rating or classification systems typically include restrictions on alcohol imagery allowed in films rated suitable for viewing by children or young people, analysis of alcohol imagery in popular UK films demonstrates that alcohol content

continues to occur frequently in films marketed to these groups.<sup>11</sup> To our knowledge, however there has to date been no evaluation of the extent to which popular films in India include alcohol imagery. This study was therefore carried out to quantify alcohol imagery, in relation to the age classification and language of the film, in a sample of the most popular films in India in the years 2015–2017.

## METHODS

We used Indian national box office ratings data to identify the top 10 Indian films by box office revenue in each of the years 2015, 2016 and 2017.<sup>12</sup> The Central Board of Film Certification (CBFC) age rating (comprising U (unrestricted public exhibition), UA (parental guidance for children below the age of 12 years), A (restricted to adults) and S (restricted to any special class of persons)) along with the genre and language of each film was noted. We used the validated 5-minute interval coding,<sup>13 14</sup> as described in previous studies, to code the presence of alcohol imagery in each interval as any use, actual use, implied use, other alcohol references and alcohol brand appearances. Actual use of alcohol was coded if an actor/actress in the film was shown consuming alcohol, while implied alcohol use typically involved verbal comments related to alcohol or non-verbal actions such as holding a glass or a bottle appearing to contain an alcoholic drink. Other alcohol references usually comprised the appearance of alcohol bottles or beer mugs. An occurrence in any of the above categories was coded as present or absent for each interval. Multiple appearances in the same category in the same interval were coded as a single event, and appearances in different categories as separate events. The same appearance transitioning into two or more intervals was coded as two or more events, as appropriate. Accuracy of results was ensured by two coders independently coding each of the movies with quality check done by two other coders. Any discrepancy was cleared after discussing with the investigators of the study. For alcohol branding we noted occurrences of clear, unambiguous alcohol branding on-screen, subgrouped as branding on a product used in a scene, branding on a product not used in a scene, branded merchandise, advertisements visible in scenes of alcohol content and any other alcohol-related advertisement. The total number of alcohol brands shown in were counted and listed.

## Patient and public involvement

No patients were involved in this study.

## Analysis

Frequencies and percentages were calculated for the language, age rating, alcohol usage and brand appearance. Mean and standard deviation was calculated for runtime of the film. Proportions of films and intervals were compared for alcohol content between films according to age classification and language using  $\chi^2$  test.

## RESULTS

The 30 films included seven (23%) that were U-rated and 23 (77%) UA-rated, and 22 (73%) in Hindi and 8 (27%) in regional languages. The regional language included four Tamil, three Telugu and one Malayalam films. The mean (SD) film runtime of the 30 films was 151.2 (12.8) min, and their cumulative duration 4535 min, which we coded in 923 5-minute intervals. Of these, 229 (25%) were in U and 694 (75%) in UA films, and 665 (72%) in Hindi and 258 (28%) in regional language films. The list of the films with their language, genre, age rating and the number of intervals is provided in the online supplemental file 1. Any alcohol appearance was observed in 195 (21%) 5-minute intervals occurring in 29 (97%) of the films. The proportion of intervals containing alcohol was slightly lower in U-rated (38 intervals, 17%) than UA-rated (157 intervals, 23%) films, though this difference was statistically not significant ( $p>0.05$ ); and the same in Hindi (140 intervals, 21%) and regional language (55 intervals, 21%) films ( $p>0.05$ ).

Actual alcohol use appeared in 25 (83%) films, in 90 (10%) intervals. All seven U-rated films included actual alcohol use, which occurred in 19 (50%) of the 38 intervals including any alcohol imagery, with implied use and other alcohol references occurring in 11 (29%) and 8 (11%) of alcohol intervals, respectively. Actual alcohol use occurred in 18 (78%) of the 23 UA rated films in 71 (45%) of the 157 intervals including alcohol, while implied use and other alcohol references occurred in 55 (35%) and 37 (24%) of alcohol intervals, respectively. Actual alcohol use occurred in 18 (82%) of the 22 Hindi films, in 64 (46%) of the 140 intervals containing alcohol, with implied use and other alcohol references occurring in 54 (39%) and 25 (17%) of alcohol intervals, respectively; and in 7 of the 8 regional language films in 26 (47%) of the 55 intervals containing alcohol, with implied use and other alcohol references each occurring in 12 (22%) and 20 (40%) alcohol intervals, respectively, as shown in table 1.

Alcohol brand appearances occurred in 10 intervals in five films, one of which was U-rated and one in a regional language. The brands, Signature, Royal Stag, Carlsberg, Sula Wine, Old Smuggler, Kingfisher Premium, Kisset, McDonald, Brown Horse and Tuborg appeared once each, and Vat 69 twice.

## DISCUSSION

This study, to our knowledge the first of its kind in India, demonstrates that alcohol imagery occurred in almost all of this sample of films popular in India, all of which were classified by the CBFC in India as suitable either for unrestricted viewing by children (U), or for by children under the age of 12 with parental guidance (UA).<sup>15</sup> Most of the alcohol imagery comprised actual or implied use. In contrast with our earlier analysis of tobacco imagery in these films,<sup>13</sup> we found no evidence of greater alcohol content in films made in local languages. However, the proportion of both





classification only if it involves glamorisation or glorification, but evidence on the effect of tobacco exposure in films suggests that these considerations are unimportant: all tobacco imagery promotes tobacco use, irrespective of who, how or in what other circumstances the product is consumed.<sup>31</sup> To date, evidence on exposure to alcohol imagery is less developed and further work is needed to establish the extent of this effect. However, the prudent approach to avoidable risks is to avoid them, and since the inclusion of alcohol imagery in films aimed at children is entirely avoidable, protecting children from future alcohol use and consequent problems justifies more rigorous controls on the alcohol content of films aimed at children and young people in India.

**Acknowledgements** The authors appreciate the assistance of the project staff of the Preventing smoking uptake study in procuring and coding films.

**Contributors** JB, MMK and RB conceptualised the study. The coding sheet was developed by RB, MMK, JC and GPN. Coding analysis was done by RB, MMK and VGK. Interpretation of data and drafting the manuscript was done by RB, MMK, VGK, SM and MA. Final approval of the version was read and approved by all authors.

**Funding** This work was supported by the Medical Research Council (grant number MR/P008933/1) of the UK under the Global Alliance for Chronic Lung diseases program.

**Competing interests** None declared.

**Patient consent for publication** Not required.

**Ethics approval** IEC: 893/2018

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** Data are available upon reasonable request. Additional data on the alcohol content found in each film are available on request from rohit.bhagawath@manipal.edu.

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

## REFERENCES

- Gowing LR, Ali RL, Allsop S, *et al*. Global statistics on addictive behaviours: 2014 status report. *Addiction* 2015;110:904–19.
- WHO. Global status report on alcohol and health. World Health organization, 2018. Available: [http://www.who.int/substance\\_abuse/publications/global\\_alcohol\\_report/msbgsruprofiles.pdf](http://www.who.int/substance_abuse/publications/global_alcohol_report/msbgsruprofiles.pdf)
- Griswold MG, Fullman N, Hawley C, *et al*. Alcohol use and burden for 195 countries and territories, 1990–2016: a systematic analysis for the global burden of disease study 2016. *The Lancet* 2018;392:1015–35. doi:10.1016/S0140-6736(18)31310-2
- Manthey J, Shield KD, Rylett M, *et al*. Global alcohol exposure between 1990 and 2017 and forecasts until 2030: a modelling study. *Lancet* 2019;393:2493–502.
- Solanki MS. Alcohol use disorder. National health portal, 2015. Available: <https://www.nhp.gov.in/healthyliving/alcohol-use-disorder>
- NFHS 4. National family health survey. Mumbai, 2016. Available: [http://rchiips.org/nfhs/factsheet\\_NFHS-4.shtml](http://rchiips.org/nfhs/factsheet_NFHS-4.shtml)
- Soundararajan S, Narayanan G, Agrawal A. Relation between age at first alcohol drink & adult life drinking patterns in alcohol-dependent patients. *Indian J Med Res* 2019;146:606–11.
- Johnson PR, Banu S, Ashok MV. Severity of alcoholism in Indian males: correlation with age of onset and family history of alcoholism. *Indian J Psychiatry* 2010;52:243–9.
- Ambekar A, Chadda RK, Khandelwal SK. Magnitude of substance use in India. Ministry of social justice and Empowerment government of India, 2019. Available: [http://socialjustice.nic.in/writereaddata/UploadFile/Magnitude\\_Substance\\_Use\\_India\\_REPORT.pdf](http://socialjustice.nic.in/writereaddata/UploadFile/Magnitude_Substance_Use_India_REPORT.pdf)
- Hanewinkel R, Sargent JD, Hunt K, *et al*. Portrayal of alcohol consumption in movies and drinking initiation in low-risk adolescents. *Pediatrics* 2014;133:973–82.
- Lyons A, McNeill A, Gilmore I, *et al*. Alcohol imagery and branding, and age classification of films popular in the UK. *Int J Epidemiol* 2011;40:1411–9.
- Box office India.. Available: <https://www.boxofficeindia.com/> [Accessed 20 Jan 2019].
- Kulkarni MM, Kamath VG, Cranwell J, *et al*. Assessment of tobacco imagery and compliance with tobacco-free rules in popular Indian films. *Tob Control* 2020;29:119–21.
- Lyons A, McNeill A, Britton J. Alcohol imagery on popularly viewed television in the UK. *J Public Health* 2014;36:426–34.
- Ministry of Information and Broadcasting. Central board of film certification. Available: <https://www.cbfcindia.gov.in/main/guidelines.html>
- UNESCO Institute for Statistics. Record number of films produced, 2016. Available: <http://uis.unesco.org/en/news/record-number-films-produced#slideoutmenu>
- Lyons A, McNeill A, Chen Y, *et al*. Tobacco and tobacco branding in films most popular in the UK from 1989 to 2008. *Thorax* 2010;65:417–22.
- Mekemson C, Glik D, Titus K, *et al*. Tobacco use in popular movies during the past decade. *Tob Control* 2004;13:400–2.
- Dalton MA, Tickle JJ, Sargent JD, *et al*. The incidence and context of tobacco use in popular movies from 1988 to 1997. *Prev Med* 2002;34:516–23.
- Worth KA, Dal Cin S, Sargent JD. Prevalence of smoking among major movie characters: 1996–2004. *Tob Control* 2006;15:442–6.
- Adachi-Mejia AM, Dalton MA, Gibson JJ. Tobacco brand appearances in movies before and after the master settlement agreement. *JAMA* 2005;293:2341e2.
- Sargent JD, Tickle JJ, Beach ML, *et al*. Brand appearances in contemporary cinema films and contribution to global marketing of cigarettes. *Lancet* 2001;357:29–32.
- Harrop A, Daniels M. Methods of time sampling: a reappraisal of momentary time sampling and partial interval recording. *J Appl Behav Anal* 1986;19:73–7.
- Everett SA, Schnuth RL, Tribble JL. Tobacco and alcohol use in top-grossing American films. *J Community Health* 1998;23:317–24.
- Escamilla G, Craddock AL, Kawachi I. Women and smoking in Hollywood movies: a content analysis. *Am J Public Health* 2000;90:412e4.
- Hanewinkel R, Tanski SE, Sargent JD. Exposure to alcohol use in motion pictures and teen drinking in Germany. *Int J Epidemiol* 2007;36:1068–77.
- Cranwell J, Murray R, Lewis S, *et al*. Adolescents' exposure to tobacco and alcohol content in YouTube music videos. *Addiction* 2015;110:703–11.
- Hendriks H, Van den Putte B, Gebhardt WA, *et al*. Social drinking on social media: content analysis of the social aspects of alcohol-related posts on Facebook and Instagram. *J Med Internet Res* 2018;20:e226.
- Barker AB, Smith J, Hunter A, *et al*. Quantifying tobacco and alcohol imagery in Netflix and Amazon prime instant video original programming accessed from the UK: a content analysis. *BMJ Open* 2019;9:e025807.
- What young India watches. BARC India Newsletter, 2018. Available: [https://www.barcindia.co.in/resources/What Young India Watches.pdf](https://www.barcindia.co.in/resources/What%20Young%20India%20Watches.pdf)
- National Center for Chronic Disease Prevention and Health Promotion (US) Office on Smoking and Health. Preventing Tobacco Use Among Youth and Young Adults: A Report of the Surgeon General [Internet]. Centers for Disease Control and Prevention (US), 2012. Available: <https://www.ncbi.nlm.nih.gov/books/NBK99237/>