

Transcriptional Regulation of Chemical Diversity in *Aspergillus fumigatus* by LaeA

Robyn M. Perrin¹, Natalie D. Fedorova², Jin Woo Bok¹, Robert A. Cramer Jr.³, Jennifer R. Wortman², H. Stanley Kim⁴, William C. Nierman^{2,5}, Nancy P. Keller^{1*}

1 Department of Plant Pathology, University of Wisconsin-Madison, Madison, Wisconsin, United States of America, **2** The Institute for Genomic Research, Rockville, Maryland, United States of America, **3** Department of Molecular Genetics and Microbiology, Duke University Medical Center, Durham, North Carolina, United States of America, **4** Department of Medicine, Korea University Medical College, Seoul, Korea, **5** Department of Biochemistry and Molecular Biology, The George Washington University School of Medicine, Washington, D. C., United States of America

Secondary metabolites, including toxins and melanins, have been implicated as virulence attributes in invasive aspergillosis. Although not definitively proved, this supposition is supported by the decreased virulence of an *Aspergillus fumigatus* strain, $\Delta laeA$, that is crippled in the production of numerous secondary metabolites. However, loss of a single LaeA-regulated toxin, gliotoxin, did not recapitulate the hypovirulent $\Delta laeA$ pathotype, thus implicating other toxins whose production is governed by LaeA. Toward this end, a whole-genome comparison of the transcriptional profile of wild-type, $\Delta laeA$, and complemented control strains showed that genes in 13 of 22 secondary metabolite gene clusters, including several *A. fumigatus*-specific mycotoxin clusters, were expressed at significantly lower levels in the $\Delta laeA$ mutant. LaeA influences the expression of at least 9.5% of the genome (943 of 9,626 genes in *A. fumigatus*) but positively controls expression of 20% to 40% of major classes of secondary metabolite biosynthesis genes such as nonribosomal peptide synthetases (NRPSs), polyketide synthases, and P450 monooxygenases. Tight regulation of NRPS-encoding genes was highlighted by quantitative real-time reverse-transcription PCR analysis. In addition, expression of a putative siderophore biosynthesis NRPS (NRPS2/*sidE*) was greatly reduced in the $\Delta laeA$ mutant in comparison to controls under inducing iron-deficient conditions. Comparative genomic analysis showed that *A. fumigatus* secondary metabolite gene clusters constitute evolutionarily diverse regions that may be important for niche adaptation and virulence attributes. Our findings suggest that LaeA is a novel target for comprehensive modification of chemical diversity and pathogenicity.

Citation: Perrin RM, Fedorova ND, Bok JW, Cramer RA Jr., Wortman JR, et al. (2007) Transcriptional regulation of chemical diversity in *Aspergillus fumigatus* by LaeA. PLoS Pathog 3(4): e50. doi:10.1371/journal.ppat.0030050

Introduction

Aspergillus fumigatus is a saprophytic filamentous fungus with no known sexual stage. Prolific production of asexual spores (conidia) and nearly ubiquitous distribution in the environment ensures constant host exposure to its spores, at a density of 1 to 100 conidia/m⁻³ [1]. The innate immune system enables spores to be eliminated from lung epithelial tissue with ease in immunocompetent vertebrates. However, immunocompromised individuals are at risk for pulmonary disease as a consequence of *A. fumigatus* infection. Of particular concern is invasive aspergillosis, which occurs when hyphal growth proliferates throughout pulmonary or other tissues. Invasive aspergillosis has an associated mortality rate ranging from 50% to 90% depending on the patient population [2]. As the number of immunocompromised patients has increased in recent decades due to immunosuppressive chemotherapy treatments, HIV/AIDS, and solid organ and bone marrow transplantation, the incidence of invasive aspergillosis has increased more than 4-fold in developed nations [2].

Several *A. fumigatus* secondary metabolites or natural products (e.g., conidial melanins and mycotoxins) have been implicated as affecting virulence [3–7]. However, the exact mechanisms by which many of these compounds might affect disease outcome are unknown, nor is it clear in most cases whether these factors play direct or indirect roles in pathogenicity. In contrast to most genes involved in primary

metabolism, genes encoding secondary metabolite biosynthetic enzymes exist in contiguous clusters within the genome [8,9]. LaeA was originally identified as a transcriptional regulator of secondary metabolite gene clusters in *Aspergillus nidulans* and *A. fumigatus* [10,11], including gliotoxin in the latter. Gliotoxin has long been suggested to be a major virulence attribute in invasive aspergillosis [12–14]. However, whereas a $\Delta laeA$ mutant shows reduced virulence in a mouse model of invasive aspergillosis [11], inactivation of gliotoxin biosynthesis alone does not [15–17]. Therefore, we reasoned that because LaeA is a transcriptional regulator, perhaps acting at a chromatin remodeling level [9,18], a microarray experiment comparing the transcriptomes of $\Delta laeA$, wild-type, and complemented $\Delta laeA$ control strains would yield further insight into LaeA-mediated *A. fumigatus* virulence

Editor: Scott G. Filler, David Geffen School of Medicine at the University of California Los Angeles, United States of America

Received: September 28, 2006; **Accepted:** February 15, 2007; **Published:** April 13, 2007

Copyright: © 2007 Perrin et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abbreviations: NRPS, nonribosomal peptide synthetase; QRT-PCR, quantitative reverse-transcription PCR; RT, reverse-transcription; SAM, significance analysis of microarrays

* To whom correspondence should be addressed. E-mail: npk@plantpath.wisc.edu

Author Summary

Patients with suppressed immune systems due to cancer treatments, HIV/AIDS, or organ transplantation are at high risk of infection from microbes. Some of the most deadly infections for such patients arise from a fungal pathogen, *Aspergillus fumigatus*. This species, like several of its close relatives, can produce an array of small chemical compounds that influences both the infection process and its environmental niche outside of the host. The genes dedicated to production of each compound are clustered adjacent to each other in the genome. One protein named LaeA is a master regulator of such clustered small molecule genes, and removal of the gene encoding LaeA cripples the organism's ability to infect. We conducted a genome-wide microarray experiment to identify small molecule gene clusters controlled by the presence of LaeA in *A. fumigatus*. In doing so, we identified actively expressed gene clusters critical for small molecule production and potentially involved in disease progression. These results also provide insight into evolutionary events shaping the organism's collection of chemical compounds.

attributes. We uncovered an unprecedented view of LaeA global regulation of mycotoxin islands, nearly all found in nonsyntenic regions of the *Aspergillus* genome. Because secondary metabolite gene cluster regions are evolutionarily diverse and may affect virulence attributes, LaeA is a novel target for comprehensive modification of chemical diversity.

Results/Discussion

Microarray Data Generation and Quality Assessment

Transcriptional profiles of the wild-type, $\Delta laeA$, and $\Delta laeA$ complemented strain were determined by comparisons of relative transcript levels between (1) $\Delta laeA$ versus wild-type and (2) wild-type versus complemented control strain. All strains were grown under identical conditions (25 °C, liquid shaking culture, glucose minimal media, 60 h) for three biological replicates. The condition and time point were chosen on the basis of optimal production of secondary metabolites [10,11]. The comparison of $\Delta laeA$ versus wild-type was used to determine gene expression patterns specific to the $\Delta laeA$ mutant, while the wild-type versus complemented strain comparison was conducted as a control, because the difference between these two strains is the presence of an ectopic copy of a selectable marker for hygromycin resistance. The processed signal intensity ratios for the three $\Delta laeA$ versus wild-type replicates were analyzed using the significance analysis of microarrays (SAM) method [19], as described in Materials and Methods. In total, 943 genes were significantly differentially expressed. Figure S1 shows a heat map of a subset of these loci, depicting normalized expression ratios for the three $\Delta laeA$ versus wild-type experiments and the three wild-type versus complemented control experiments. The high quality of the data is indicated by the consistency of color between the replications and the relative lack of color in the control lanes.

Patterns of Transcriptional Regulation

Of the 943 genes showing significant differences in expression between $\Delta laeA$ and wild-type by SAM analysis, 415 showed increased expression in $\Delta laeA$ and 528 showed decreased expression. Table 1 and Figure S2 indicate functional categories for these genes (defined as described by the

Gene Ontology Consortium, <http://www.geneontology.org>). The most remarkable discovery was the near-global suppression of secondary metabolite gene expression in the $\Delta laeA$ mutant. Nearly all (97%) of the secondary metabolite gene cluster loci showed decreased expression in $\Delta laeA$, with a mere three genes in this category showing increased expression in $\Delta laeA$. This was in contrast to all other functional categories, which showed substantial proportions of both increased and decreased expression in the mutant, possibly reflecting indirect effects due to loss of production of multiple metabolites. In addition to genes with unknown function (39%) and genes involved in secondary metabolism (11%), other major categories included genes encoding proteins involved in transmembrane transport (8%) and those involved in information processing (4%), and cell wall biogenesis (4%). Statistical analysis of the overrepresentation of different Gene Ontology categories and Pfam protein domains within the set of 943 differentially regulated genes is shown in Tables S1 and S2, respectively.

Interestingly, LaeA appeared to influence expression of a subset of species- and lineage-specific genes not strongly conserved with other fungal species. Only 18% and 44% of all genes significantly differentially expressed in the mutant have putative orthologs in *Saccharomyces cerevisiae* and *Neurospora crassa*, respectively, compared to an average of 33% and 58% of all *A. fumigatus* genes. Many, but not all, of these genes were classified as secondary metabolism genes. Moreover, there are about 120 differentially expressed genes; again, most, but not all, are present in secondary metabolism clusters (Table 2), which have no detectable orthologs in *Aspergillus oryzae* and *A. nidulans*. Considering this overwhelming tight and directed transcriptional control of secondary metabolite loci by LaeA, below we focus on such genes as possible members of the LaeA-regulated *A. fumigatus* pathogenicity arsenal.

Regulation of Chemical Diversity by LaeA

Although initial genome analysis suggested the presence of 26 secondary metabolite gene clusters [20], subsequent analysis (G. Turner, N. D. Fedorova, V. Joardar, J. R. Wortman, and W. C. Nierman, unpublished data) has provided support for only 22 clusters. Of the 13 secondary metabolite gene clusters whose expression was influenced by LaeA in the condition used for microarray analysis, ten are particularly strongly affected, with a majority of genes within these clusters being significantly down-regulated in $\Delta laeA$ as indicated by SAM. Three additional clusters have at least one gene encoding a critical enzyme such as a nonribosomal peptide synthetase (NRPS) or a polyketide synthase showing decreased expression in $\Delta laeA$. Additionally, 38% (23 of 71) of all P450 monooxygenases show differential expression in $\Delta laeA$, also associated with secondary metabolite biosynthesis and/or detoxification. Fifteen of these genes encoding P450 monooxygenases are found in secondary metabolite gene clusters. Table S3 gives normalized expression ratio values for all 22 gene clusters in *A. fumigatus*. Table 2 summarizes the current state of knowledge regarding function of LaeA-regulated secondary metabolite gene clusters. These include clusters dedicated to production of conidial melanins, fumitremorigens, gliotoxin, and ergot alkaloids such as festuclavine, elymoclavine, and fumigaclavines A, B, and C (Table 2) [4,17,21–27]. Figure 1 depicts the chromosomal landscape of those regions most strongly regulated by LaeA.

Table 1. Gene Ontology Categories for Loci Regulated by LaeA

Gene Ontology Biological Process	LaeA	Down-Regulated	Highly Down-Regulated	Moderately Down-Regulated	Up-Regulated	Highly Up-Regulated	Moderately Up-Regulated	0–300 kb	Telomere-Proximal LaeA	Genome
Transmembrane transport	76	22	9	13	54	26	28	135	18	583
Cell wall biogenesis	37	20	12	8	17	7	10	39	8	177
Regulation of transcription	54	19	5	14	35	12	23	72	11	417
Information processing	35	9	3	6	25	4	21	32	3	1146
Other processes	54	17	8	9	37	13	24	52	8	865
Carbohydrate metabolism	36	14	6	8	22	5	17	83	6	293
Secondary metabolism	102	99	79	20	3	0	3	180	65	321
Other metabolism	189	73	30	43	117	46	71	251	29	1902
Unknown	360	142	64	78	218	73	145	665	81	3922
Total	943	415	216	199	528	184	344	1509	229	9626

Total number of LaeA-regulated genes in each category is shown (LaeA). Genes are indicated as highly down-regulated in the mutant (SAM d-score > 5), moderately down-regulated (3 < d-score < 5), moderately up-regulated (−5 < d-score < −3), or highly up-regulated (d-score < −5). The functional categories correspond to standard Gene Ontology category definitions (<http://www.geneontology.org>) with supercategories defined as follows: Information processing: transcription, translation, ribosomal structure and biogenesis, DNA replication, recombination and repair. Cellular processes: cell cycle, cell division, development, energy pathways, homeostasis, intracellular transport, organelle organization and biogenesis, response to stress; Cellular metabolism: amino acid and derivative metabolism, cofactor metabolism, lipid metabolism, protein catabolism, protein modification. Only genes present in secondary metabolism gene clusters were included in Secondary Metabolism. The number of LaeA-regulated genes that are within 300 kb of telomere ends is indicated for each category (Telomere-Proximal LaeA). The number of genes in each functional category for the whole genome is shown in the last column. doi:10.1371/journal.ppat.0030050.t001

To confirm these microarray results, quantitative real-time reverse-transcription (RT)-PCR (QRT-PCR) was performed on one major class of secondary metabolite genes, those encoding NRPSs [28]. As indicated in Table 3, relative expression levels for NRPSs that showed differential expression between mutant and wild-type in the microarray study also were dramatically reduced upon QRT-PCR analysis. In all cases, complementation of the *laeA* defect restored NRPS gene expression to wild-type levels (Table 3).

Notably, because the microarray analysis determines only relative expression and not absolute levels of transcript, we could not conclude whether secondary metabolite clusters not showing differential expression in mutant versus wild-type were not affected by LaeA or were simply not induced under the growth condition used. To further examine these possibilities, we assessed the expression of a subset of NRPSs thought to encode siderophore-biosynthesizing enzymes. Although siderophores do not fit neatly into a definition of secondary metabolites, which are dispensable in laboratory growth conditions [29], these molecules are produced from clustered genes and are critical for pathogen growth in blood serum [30]. Because iron was included in the media used for the microarray study, we investigated whether the *Alae* mutant was deficient in expression of siderophore gene cluster NRPSs under iron-limiting conditions.

As previously reported [29], low iron conditions induced transcriptional upregulation of several NRPS genes known or predicted to be involved in siderophore biosynthesis (Table 4). Normalized expression levels of the siderophore NRPSs in the low iron condition relative to high iron conditions were *NRPS2/sidC*, 2.245 ± 0.449 ; *NRPS3/sidE*, 68.595 ± 13.725 ; and *NRPS4/mps6/sidD*, 28.509 ± 4.704 . *NRPS7* transcripts were not detectable in these experiments. In contrast to Reiber et al. [29], *NRPS3/sidE* showed the highest induction to the low iron conditions in our experiments. This discrepancy might be explained by differential sensitivity of the semiquantitative RT-PCR method used by Reiber et al. compared to our QRT-PCR methodology or subtle differences in culture conditions. We also noted that the complemented control strain with an ectopic copy of *laeA* showed increased expression of *NRPS3/sidE* in both low and high iron conditions.

Comparison of the *AlaeA* mutant and controls by QRT-PCR analysis indicated differential expression of *NRPS3/sidE* in low (inducing) iron conditions. In high iron conditions, *NRPS4/mps6/sidD*, *NRPS3/sidE*, and possibly *NRPS2/sidC* showed decreased expression in the *AlaeA* mutant (Table 4). Interestingly, *NRPS7* was not detectable in these experiments. In the low iron condition, expression of actin did decrease in the *AlaeA* mutant. However, the dramatic decrease in expression of *NRPS3/sidE* (1,000-fold less) seen in the *AlaeA* background strongly suggests that LaeA regulates the expression of at least this NRPS. Little is known about the function of SidE, although it has been speculated to be involved in siderophore biosynthesis on the basis of homology to SidC [29]. It remains to be determined whether *NRPS3/sidE* is involved in siderophore production, a process known to be critical to virulence [31,32], or whether it synthesizes an iron-responsive compound with a distinct function. Regardless of the function of SidE, these experiments show that LaeA is also involved in controlling expression of other secondary metabolite clusters not induced by the environmental conditions used in the microarray experiments.

Table 2. Summary of Secondary Metabolite Gene Cluster Function and Regulation by LaeA

Cluster Number	Chromosome	Location	Distance to Telomere (kb)	LaeA Regulation	Product(s)	Pathogenicity	References
1	1	Afu1g10360–Afu1g10390	2,300	Yes	Unknown product of <i>Afpes1</i>	Required for virulence (<i>Galleria mellonella</i> model)	[4]
2	1	Afu1g17640–Afu1g17740	100	No	Unknown	—	—
3	2	Afu2g17510–Afu2g17600	200	Yes	DHN-melanin	Required for virulence (murine)	[5–7,24]
4	2	Afu2g17960–Afu2g18070	100	Yes	Ergot alkaloids: festuclavine, elymoclavine, fumigaclavines A, B, and C	Unknown	[22,23,25]
5	3	Afu3g01290–Afu3g01600	400	Partial	Unknown	—	—
6	3	Afu3g02520–Afu3g02720	700	No	Unknown	—	—
7	3	Afu3g03190–Afu3g03370	900	No	Probably two compounds (a siderophore and a distinct toxin) ^a	—	—
8	3	Afu3g12870–Afu3g13010	700	Yes	Putative ETP	—	—
9	3	Afu3g13580–Afu3g13750	500	No	Unknown	—	—
10	3	Afu3g14560–Afu3g14760	200	Partial	Unknown	—	—
11	3	Afu3g15200–Afu3g15340	100	No	Unknown	—	—
12	4	Afu4g00110–Afu4g00280	100	Partial	Unknown	—	—
13	4	Afu4g14380–Afu4g14850	100	Partial	Unknown	—	—
14	5	Afu5g00110–Afu5g00160	100	No	Unknown	—	—
15	5	Afu5g00340–Afu5g00400	100	No	Unknown	—	—
16	5	Afu5g12700–Afu5g12740	700	No	Unknown	—	—
17	6	Afu6g03290–Afu6g03490	800	Yes	Unknown	—	—
18	6	Afu6g09580–Afu6g09770	1,500	Yes	Gliotoxin	Immunosuppressive	[15–17,43,44]
19	6	Afu6g12040–Afu6g12080	800	Yes	Unknown	—	—
20	6	Afu6g13920–Afu6g14000	300	Yes	Unknown	—	—
21	7	Afu7g00110–Afu7g00190	100	No	Unknown	—	—
22	8	Afu8g00100–Afu8g00720	100	Yes	Fumitremorgen B; supercluster probably producing more than one product	Unknown	[26,27]

^aRevised gene cluster border merges clusters previously defined as separate [20].

PKS, polyketide synthetase; ETP, epipolythiodioxopiperazine; DHN, 1,8-dihydroxynaphthalene. doi:10.1371/journal.ppat.0030050.t002

Secondary Metabolites under Control of LaeA

Cluster 18 (Figure 1) on Chromosome 6, strongly differentially expressed in *ΔlaeA*, encodes the genes required for gliotoxin biosynthesis. Gliotoxin is arguably the most well-studied mycotoxin produced by *A. fumigatus*. First identified in 1936, this compound has immunosuppressive properties in vitro [12] and in vivo [13,14], although its direct contribution to pathogenicity is only beginning to be understood [15–17]. Like all other compounds in the epipolythiodioxopiperazine class, gliotoxin is a cyclic dipeptide with an internal disulfide bridge that can undergo redox cycling (for a recent review, see [33]). Immunosuppressive activity of gliotoxin is due at least in part to negative regulation of the transcription factor nuclear factor-κB, which occurs by inhibition of proteasome-mediated degradation of the nuclear factor-κB inhibitor IκBα [34,35]. Gliotoxin is also known to be cytotoxic and can evoke both apoptotic [36–39] and necrotic [40,41] cell death. Recently, gliotoxin was shown to trigger the release of apoptogenic factors by the host mitochondrial protein Bak [42]. The secondary metabolism gene cluster responsible for gliotoxin production was recently identified by bioinformatic

analysis [43] and has been experimentally confirmed [15,17]. Despite the known immunosuppressive activities of the molecule and its detection in blood serum of patients with invasive aspergillosis [44], three recent studies using genetic mutants of the gliotoxin gene cluster demonstrated that gliotoxin is not a virulence factor in murine models of invasive aspergillosis [15–17]. However, these same studies presented evidence that gliotoxin could adversely affect T cells, neutrophils, and mast cells and, we offer, likely acts synergistically with other LaeA-regulated toxins. The *ΔlaeA* mutant is impaired in gliotoxin production during growth in culture as well as growth in vivo in murine models of invasive aspergillosis [10,15], and the microarray results presented here confirm that LaeA strongly influences expression of genes in this cluster under the condition investigated.

Secondary metabolite cluster 1 on Chromosome 1, which is differentially expressed in *ΔlaeA*, contains an atypical NRPS called *Afpes1* that is required for virulence in an insect model of invasive aspergillosis [4]. *Afpes1* shows greatest homology to NRPSs that produce siderophores or destruxins, including one paralog required for virulence of the plant pathogen




Figure 1. Positional Bias of LaeA-Regulated Gene Expression

Left, chromosomes of *A. fumigatus* with secondary metabolite gene clusters indicated as bars and clusters numbered as indicated in Table 2. Arrows indicate clusters that are completely or partially regulated by LaeA. Graphs to the right of each chromosome show normalized expression ratio values of $\Delta laeA$ versus wild-type plotted by chromosomal position. Each point represents the average value of ten contiguous loci. The gap on Chromosome 4 indicates a region of the genome not included in the microarray that contains the rRNA gene cluster. Positive values correspond to reduced expression in $\Delta laeA$ and negative values correspond to increased expression in $\Delta laeA$.
doi:10.1371/journal.ppat.0030050.g001

Alternaria brassicae [45]. However, the *Afpes1* cluster is thought to be unlikely to produce either of these compounds, because destruxin toxin has not been detected in *A. fumigatus* [4] and expression of *Afpes1* was not responsive to iron levels [4,21]. Deletion of *Afpes1* alters conidial morphology and hydrophobicity as well as melanin synthesis and results in increased susceptibility to reactive oxygen species, implying altered conidial melanin and/or rodlet composition [4]. Most of these characteristics are common to the $\Delta laeA$ phenotype [11], possibly implicating a role of the *Afpes1* metabolite in the attenuated virulence of $\Delta laeA$.

A. fumigatus synthesizes several clavine ergot alkaloids, compounds that can be partial agonists or antagonists of serotonin, dopamine, and α -adrenalin receptors, thus affecting nervous, circulatory, reproductive, and immune system function [46]. The role of these compounds in invasive aspergillosis has not been determined. In addition to having the receptor-modulating activities mentioned, the festuclavine ergot alkaloid produced by *A. fumigatus* is cytostatic and is directly mutagenic in the Ames assay [47,48]. Recently, Coyle and Panaccione [25] showed that deletion of an *A. fumigatus* dimethylallyltryptophan synthase (DMAT synthase) homologous to *dmaW* of the ergot-producing species *Claviceps purpurea* eliminated all known ergot alkaloids, confirming its predicted function in the first committed step of ergot alkaloid production (i.e., addition of dimethylallyl diphosphate to L-tryptophan to result in 4-methylallyl-tryptophan). The biochemical activity of the *A. fumigatus* DmaW enzyme was also confirmed by Unsöld and Li [22], who subsequently

characterized a reverse prenyltransferase in the same gene cluster that converts fumigaclavine A to fumigaclavine C [23]. These genes are located in secondary metabolite gene cluster 4 on Chromosome 2, which is strongly differentially expressed in $\Delta laeA$.

Melanins found in conidia are one of the few described virulence factors in *A. fumigatus* [5,6,24]. Lack of melanins leads to increased susceptibility to reactive oxygen species produced by the host innate immune response during infection as well as altered (smooth) conidial morphology [5,7]. However, the scarcity of nonpigmented *A. fumigatus* spores in nature has drawn into question the clinical relevance of melanins as virulence factors [1]. Conidia of $\Delta laeA$ are pigmented, but altered expression of *alb1* in the mutant has been reported previously and at least one unidentified spore metabolite is missing in $\Delta laeA$ [11]. There is significant differential expression of the 1,8-dihydroxynaphthalene-melanin gene cluster in $\Delta laeA$ under the condition investigated in this study. Expression of this gene cluster is also regulated by cAMP/protein kinase A signaling [49] as is LaeA itself [10], perhaps a suggestion that in this case LaeA control of this cluster may be both directly and indirectly mediated by protein kinase A.

Additionally, a LaeA-regulated supercluster on Chromosome 8 is likely to produce multiple compounds. Recently, two genes in this cluster have been reported to encode biosynthetic enzymes for the tremorgenic mycotoxin fumitremorgin B and related compounds [26,27]. The cyclo-L-Trp-L-Pro derivative fumitremorgin B is cytotoxic, inhibiting cell

Table 3. Real-Time RT-PCR Analysis of Select *A. fumigatus* NRPSs in Wild-Type, $\Delta laeA$, and Complemented Control Strains

NRPS ^a (Gene Locus ID)	Secondary Metabolism Gene Cluster Number	Wild-Type ^b	$\Delta laeA$ ^b	Complemented Control ^b
NRPS1 (Afu1g10380)	1	25.17 ± 0.611, 1.0 ± 0.425	27.10 ± 0.400, 0.32 ± 0.090	23.20 ± 0.265, 2.83 ± 1.109
NRPS5 (Afu3g12920)	8	20.93 ± 0.058, 1.0 ± 0.057	30.20 ± 0.87, 0.002 ± 0.001	21.30 ± 0.265, 0.561 ± 0.220
NRPS9 (Afu6g09610)	18	22.43 ± 0.153, 1.0 ± 0.113	34.63 ± 1.343, 0.000 ± 0.000	21.63 ± 0.058, 1.260 ± 0.440
NRPS10 (Afu6g09660)	18	18.90 ± 0.173, 1.0 ± 0.127	31.03 ± 0.802, 0.000 ± 0.000	18.20 ± 0.100, 1.176 ± 0.415
NRPS11 (Afu6g12050)	19	19.20 ± 0.100, 1.0 ± 0.080	31.17 ± 0.723, 0.000 ± 0.000	20.37 ± 0.115, 0.322 ± 0.115
NRPS13 (Afu8g00170)	22	21.87 ± 0.115, 1.0 ± 0.089	32.23 ± 0.321, 0.001 ± 0.000	21.97 ± 0.153, 0.675 ± 0.245
NRPS14 (Afu8g00540)	22	17.87 ± 0.058, 1.0 ± 0.057	28.63 ± 0.153, 0.001 ± 0.000	17.80 ± 0.100, 0.758 ± 0.268
Actin	—	17.80 ± 0.100, 0.758 ± 0.268	17.80 ± 0.100, 0.758 ± 0.268	16.60 ± 0.500

^aNomenclature as in Cramer et al. [28].

^bValues (given as mean cycle threshold) ± SD, three replicates, and relative expression level (arbitrary units) ± SD, three replicates. Reference gene is actin, control (relative) condition wild-type strain Af293.

QRT-PCR was used to assess absolute levels (upper numbers in each cell) and relative levels (lower numbers in each cell) of NRPS transcripts. Primers and methods were as described [28].
doi:10.1371/journal.ppat.0030050.t003

Table 4. Real-Time RT-PCR Analysis of Select *A. fumigatus* Siderophore Biosynthesis NRPSs

Gene ^a + Iron Levels	Secondary Metabolism Gene Cluster Number	Wild-Type ^b	<i>ΔlaeA</i> ^b	Complemented Control ^b
NRPS2 (<i>sidC</i>), low iron	2	24.70 ± 0.173, 1.0 ± 0.200	28.90 ± 0.458, 1.149 ± 0.368	24.77 ± 0.404, 1.587 ± 0.458
NRPS3 (<i>sidE</i>), low iron	7	18.90 ± 0.173, 1.0 ± 0.200	32.80 ± 0.854, 0.001 ± 0.001	17.80 ± 0.100, 3.564 ± 0.349
NRPS4 (<i>nps6</i> , <i>sidD</i>), low iron	7	20.83 ± 0.058, 1.0 ± 0.165	25.07 ± 0.153, 1.122 ± 0.127	21.37 ± 0.208, 1.149 ± 0.184
NRPS7, low iron	11	Not detectable	Not detectable	Not detectable
Actin, low iron	—	17.57 ± 0.231	21.97 ± 0.058	18.30 ± 0.100
NRPS2 (<i>sidC</i>), high iron	2	27.03 ± 0.153, 1.0 ± 0.113	26.73 ± 0.115, 0.675 ± 0.153	26.53 ± 0.058, 0.776 ± 0.128
NRPS3 (<i>sidE</i>), high iron	7	26.17 ± 0.058, 1.0 ± 0.057	27.70 ± 0.100, 0.189 ± 0.042	18.43 ± 0.115, 116.7 ± 20.886
NRPS4 (<i>nps6</i> , <i>sidD</i>), high iron	7	26.83 ± 0.058, 1.0 ± 0.057	29.07 ± 0.306, 0.117 ± 0.035	25.63 ± 0.404, 1.260 ± 0.407
NRPS7, high iron	11	Not detectable	Not detectable	Not detectable
Actin, high iron	—	18.73 ± 0.058	17.87 ± 0.306	17.87 ± 0.231

^aNomenclature as in Cramer et al. [28].

^bValues (given as mean cycle threshold) ± SD, three replicates, and relative expression level (arbitrary units) ± SD, three replicates. Reference gene is actin, control (relative) condition wild-type strain Af293.

QRT-PCR was used to assess absolute levels (upper numbers in each cell) and relative levels (lower numbers in each cell) of NRPS transcripts. Wild-type, *ΔlaeA*, and complemented control strains were grown in media containing high levels of iron [300 μM Fe(III)Cl₃] or low levels of iron [no Fe(III)Cl₃]. Primers and methods were as described [28].

doi:10.1371/journal.ppat.0030050.t004

cycle progression at G₂/M, and thus has been of interest as a potential anticancer agent. The pathway involves generation of the cyclic dipeptide brevianamide F by the NRPS brevianamide synthetase [27], prenylation of brevianamide F by the prenyltransferase FtmPT1 to tryptrostatin B [26], and subsequent conversion in several steps to fumitremorgen B. Thus, LaeA-mediated influence on expression of *ftmPT1* and *ftmPT2* would govern the production of this entire class of diketopiperazine compounds. Once again, however, the specific effects of these compounds on pathogenicity during invasive aspergillosis are unknown.

The fact that LaeA promotes expression not only of these secondary metabolite gene clusters but an additional eight others confirms its role as a master controller of secondary metabolism. The importance of several of these compounds in toxicity studies also underscores relevance of LaeA during infection [11]. We suggest the possibility that virulence attributes are not influenced as much by individual metabolites as by the blend of LaeA-regulated toxins, which, in combination, may confer an advantage to the pathogen.

LaeA Regulation of Evolutionarily Diverse Regions of the Genome

Comparative genomic analysis between *A. fumigatus* and related species indicates overlap between *A. fumigatus*-specific genes and genes differentially expressed in *ΔlaeA* (N. D. Fedorova and W. C. Nierman, unpublished data). In total, 68% of *A. fumigatus* secondary metabolite genes do not have orthologs in the closely related species *A. clavatus* (N. D. Fedorova and W. C. Nierman, unpublished data). Additional secondary metabolite genes do not have orthologs in more distantly related *Aspergilli* such as *A. oryzae* and *A. nidulans* [20]. The variability of secondary metabolite clusters may be explained by the fact that many of them are located in highly divergent telomere-proximal regions characterized by fre-

quent chromosomal rearrangements [20,50]. For example, 54% of the clusters showing differential expression in *ΔlaeA* in the conditions described here were found within 300 kb of telomeres. It should be noted that, in addition to the secondary metabolite clusters, other genes with significantly lower expression in *ΔlaeA* also show some positional specificity within the genome but to a much lesser extent (unpublished data). Further analysis also showed that *A. fumigatus* telomere-proximal clusters tend to have larger numbers of genes than clusters located closer to the centromeres, suggesting that the former may accumulate additional genes more easily (N. D. Fedorova, J. R. Wortman, and W. C. Nierman, unpublished data).

Initial comparative genome analyses indicate that the telomere-proximal regions (and to a lesser extent, synteny breakpoints and intrasyntenic regions) appear to be a hotbed of diversity, not only between *Aspergillus* species but even between different strains of the same species [51,52]. The genomes of two *A. fumigatus* strains have been sequenced: the clinical isolate Af293 (by The Institute for Genomic Research, Rockville, Maryland, United States) and isolate CEA10 (under contract from Elitra Pharmaceutical by Celera Genomics and made available by Merck; B. Jiang and W. C. Nierman, personal communication). These strains show an overall divergence of 2%, and the majority of this variation is in telomere-proximal and synteny breakpoint regions. Similarly, microarray experiments also supported high divergence in these regions when Af293 was compared to the unsequenced *A. fumigatus* strains Af294 and Af71 [20].

Many secondary metabolite clusters appear to be associated with transposons (known to be active in *A. fumigatus* [51]) and transposase-like sequences (Table S3). The finding that these transposable elements often flank or are embedded in many of the clusters may represent one mechanism for

generating the diversity of secondary metabolites in *Aspergillus*. Whether or not there is a connection between LaeA function and transposon activity has yet to be established. In total, these analyses suggest that secondary metabolite clusters are located in the regions that undergo extensive rearrangements, which may result in subsequent alterations in secondary metabolite production and, therefore, have major impacts on niche adaptation between different species of fungi or between strains of the same species. Other examples include a non-aflatoxin-producing clade of *A. flavus*, better known as the food-fermenting *A. oryzae* used in the production of traditional Asian products such as miso and soy sauce, which may have arisen as a result of telomere-proximal rearrangements [53]. Similarly, genotypic variability between strains of *Fusarium compactum* also proved to be a major determinant of metabolite production and geographic distribution [54]. In *Fusarium graminearum*, the major cause of wheat and barley head blight, intraspecific polymorphic variations in a trichothecene mycotoxin gene cluster were correlated with chemotype differences, host range, and fitness [55].

In light of such examples, it is interesting to speculate about the role of LaeA in chemotype evolution and niche adaptation. It is possible that variation at any particular secondary metabolite gene cluster could result in less efficient control by LaeA. This potential has been demonstrated in *A. nidulans* [18]. Conversely, LaeA itself is a major target for comprehensive changes in the entire complement of secondary metabolites. The clustering of secondary metabolite biosynthetic genes has been suggested to reflect their evolutionary history [8,9,20,51,56,57]. Several models have been proposed to explain the establishment and maintenance of secondary metabolic gene clusters in filamentous fungi. The “selfish cluster” hypothesis proposes that selection occurs at the level of the cluster and promotes maintenance of the cluster as a unit, possibly through horizontal transfer events [56]. However, there is only limited evidence for widespread horizontal transfer of secondary metabolism gene clusters, with penicillin being a notable exception [58]. Alternative models suggest that clusters are maintained due to coregulation mechanisms, likely at the level of chromatin regulation [8,9]. LaeA may provide a mechanistic means of secondary metabolism gene cluster coregulation and maintenance. Certainly LaeA demonstrates a positional bias for local gene regulation, as transfer of genes into or out of a secondary metabolite cluster leads to respective gain or loss of transcriptional regulation by LaeA [18]. This has been speculated to occur through regulation of nucleosome positioning and heterochromatin formation [9].

Conclusions

Our results confirm that LaeA plays a central role in regulation of chemical diversity in *A. fumigatus*. Furthermore, genomic regions that are transcriptionally controlled by LaeA are species and even strain specific, suggesting that they may serve as niche adaptation factors. The loss of *laeA* results in a great decrease in repertoire of secondary metabolites, which appears to impact the infection process. Therefore, LaeA constitutes a novel target for the production of an array of factors critical to success during pathogenesis. Furthermore, LaeA is a tool to identify metabolite gene clusters that may impact virulence, allowing the correlation of specific

secondary metabolite clusters with virulence even in absence of knowledge about the mycotoxin itself.

Materials and Methods

Strains and growth conditions. Three prototrophic *A. fumigatus* fungal strains were used in this study. Af293 (the wild-type clinical isolate used in the *A. fumigatus* genome sequencing project [20]), TJW54.2 (*ΔlaeA*) [11], and a complemented control strain TJW68.6 (*ΔlaeA* + *laeA*) [11] were grown in triplicate at 25 °C in liquid minimal media [59] with shaking (280 rpm) for 60 h. Profiles of secondary metabolites extracted from the media with chloroform were compared by thin-layer chromatography, and the results confirmed that the *ΔlaeA* strain showed reduced levels of multiple secondary metabolites under this condition ([10] and unpublished data). Total RNA was isolated from fungal mats, labeled, and hybridized with a DNA whole-genome amplicon microarray [20,60] in three independent biological replicates.

To analyze siderophore NRPS gene expression under low- or high-iron conditions, 50-ml liquid cultures were grown as described [29], with low-iron media containing 25 g/L glucose, 3.5 g/L (NH₄)₂SO₄, 2.0 g/L KH₂PO₄, 0.5 g/L MgSO₄ (heptahydrate), and 8 mg/L ZnSO₄ (heptahydrate) (pH 6.3). High-iron media was identical except for the addition of Fe(III)Cl₃ to a final concentration of 300 μM. Cultures were grown at 37 °C, 280 rpm, and samples were collected at 24 h postinoculation. All glassware was subjected to sequential treatment with 1 mM and 5% HCl as described [29].

RNA isolation and microarray hybridization. Total RNA was extracted from *Aspergillus* strains by use of TriZOL reagent (Invitrogen, <http://www.invitrogen.com>) according to the manufacturer's instructions. RNA was further purified by two extractions with phenol:chloroform:isoamyl alcohol (25:24:1) and then labeled with Cy-3 or Cy-5 dye and hybridized as previously described [20]. The generation of the whole genome array has been described [20].

Relative expression levels of select NRPSs. QRT-PCR was used to (1) confirm the expression level trends observed in the microarray experiment and (2) investigate NRPS gene expression under iron-limiting conditions. Expression of select NRPSs putatively regulated by LaeA was examined. Total RNA from two or three biological replicates was pooled in equal amounts (2 μg per sample) for each *Aspergillus* strain, wild-type AF293, TW54.2, and TW68.6, and treated with Ambion Turbo DNA-free DNase I (Ambion, <http://www.ambion.com>) to remove contaminating genomic DNA. A total of 500 ng of DNase I-treated total RNA from each sample was reverse transcribed with Superscript III reverse transcriptase (Invitrogen). Real-time RT-PCR was conducted with 20-μl reaction volumes with the iQ SYBR green supermix (Bio-Rad, <http://www.bio-rad.com>), 2 μl of a 1:6 dilution of first-strand cDNA, and 0.4 μl of each 10 μM primer stock. Primer sequences were previously reported [28]. No reverse transcriptase controls (NRT) were used to confirm elimination of contaminating genomic DNA. Real-time RT-PCR was performed using an iQ Cycler Real-Time PCR detection system (Bio-Rad). PCRs for each NRPS were done in triplicate and melt curve analysis was performed immediately following the PCR to confirm the absence of nonspecific amplification products and primer dimers. The relative expression levels of NRPS genes between *A. fumigatus* wild-type strain AF293, the *ΔlaeA* mutant, and the complemented control strain were calculated using 2^{-ΔΔC_t} method with iQ cycler system software. All values were normalized to expression of the *A. fumigatus* actin gene and relative to the wild-type strain for each condition analyzed.

Data analysis. Gene expression ratios were determined for triplicate comparisons of (1) wild-type and *ΔlaeA* and (2) *ΔlaeA* and the complemented control strain. Prior to statistical analysis, the LOWESS normalization method was used to remove any systematic bias from the raw expression ratios [61]. Loci showing significantly different expression were identified using the SAM method for one-class designs that has been previously described in detail [19], implemented in the TM4 suite's MultiExperiment Viewer (<http://www.tm4.org>) [62,63]. This allowed identification of genes whose mean expression across experiments is significantly different from a user-specified mean (log₂ = 0, corresponding to identical mRNA levels in the mutant and wild-type strains). Genes with scores above the significance threshold and exceeding the cutoff value of zero for the false discovery rate (the most conservative setting) were designated as significantly differentially expressed between mutant and wild-type. The delta value cutoff in SAM was chosen to capture the maximum number of significant genes while maintaining the reported estimated false discovery rate at zero. Genes down-regulated in *ΔlaeA* were further analyzed by the Expression Analysis Systematic Explorer

(EASE) [64] within TM4 to identify overrepresented Gene Ontology terms and Pfam domains. Fisher's exact test probabilities and step-down Bonferroni corrected probabilities are reported from the EASE analysis to indicate which terms are overrepresented in the down-regulated gene set.

Supporting Information

Figure S1. Expression Ratio Heat Map

A subset of *LaeA*-regulated genes (total of 409), including 174 genes up-regulated in *ΔlaeA* and 235 genes down-regulated in *ΔlaeA*. First three columns show raw expression ratio values of *ΔlaeA* versus wild-type in three independent biological replicates. Last three columns show expression ratios of wild-type versus a complemented *ΔlaeA* strain. Color-coded expression ratio scale is shown at top.

Found at doi:10.1371/journal.ppat.0030050.sg001 (1.9 MB DOC).

Figure S2. Functional Categories of Genes Showing Significantly Different Gene Expression in *ΔlaeA* versus Wild-Type, as Percentages of Total

Categories are as indicated and described in detail in Table 1.

Found at doi:10.1371/journal.ppat.0030050.sg002 (1.9 MB DOC).

Table S1. Statistical Analysis of Overrepresented Gene Ontology Categories among Genes Down-regulated in *ΔlaeA*

Numbers of genes in the indicated Gene Ontology categories were subjected to statistical analysis by EASE [64] to identify categories overrepresented compared with the whole genome data set. Fisher's exact test probabilities and Bonferroni corrected (step-down) probabilities (Bonferroni SD Corr.) are reported from the EASE analysis. Only categories with Fisher's exact test probabilities below 5.00E-02 were included.

Found at doi:10.1371/journal.ppat.0030050.st001 (41 KB DOC).

Table S2. Statistical Analysis of Overrepresented Pfam Protein Domains among Genes Significantly Down-regulated in *ΔlaeA*

Numbers of genes in the indicated Pfam protein domain categories [65] were subjected to statistical analysis by EASE [64] to identify

categories overrepresented compared with the whole genome data set. Fisher's exact test probabilities and Bonferroni corrected (step-down) probabilities (Bonferroni SD Corr.) are reported from the EASE analysis. Only categories with Fisher's exact test probabilities below 5.00E-02 were included.

Found at doi:10.1371/journal.ppat.0030050.st002 (35 KB DOC).

Table S3. Expression Ratio Values for All Secondary Metabolite Gene Clusters

Normalized expression ratio values are shown for *ΔlaeA* versus wild-type (column labeled "*ΔlaeA*") or complemented control strain versus wild-type (column labeled "comp"). Expression ratios are shown for three independent biological replicates, followed by average values (last two columns). Bold text in "Gene Name" category indicates critical genes in the cluster such as polyketide synthases or NRPSs.

Found at doi:10.1371/journal.ppat.0030050.st003 (846 KB DOC).

Acknowledgments

We would like to thank Dr. Marion Brodhagen for insightful discussion and John Braisted and Alexander Saeed for their suggestions regarding the appropriate statistical analysis techniques from The Institute for Genomic Research TM4 suite.

Author contributions. RMP, JWB, WCN, and NPK conceived and designed the experiments. RMP, JWB, RAC, JRW, and HSK performed the experiments. All authors analyzed the data and participating in the writing and editorial process. NDF, RAC, JRW, HSK, WCN, and NPK contributed reagents/materials/analysis tools.

Funding. Funding in the authors' laboratories is provided by US National Institutes of Health (NIH) grant numbers U01AI48830 (WCN), R21 AI052236 (WCN), R01 AI065728-01A1 (NPK), and National Science Foundation MCB-0236393 (NPK). RAC is funded by the NIH/National Institute of Allergy and Infectious Diseases Molecular Mycology and Pathogenesis Training Program contract number 5 T32 AI052080 at Duke University Medical Center.

Competing interests. The authors have declared that no competing interests exist.

References

- Latge JP (2001) The pathobiology of *Aspergillus fumigatus*. Trends Microbiol 9: 382–389.
- Latge JP (1999) *Aspergillus fumigatus* and aspergillosis. Clin Microbiol Rev 12: 310–350.
- Tomee JF, Kauffman HF (2000) Putative virulence factors of *Aspergillus fumigatus*. Clin Exp Allergy 30: 476–484.
- Reeves EP, Reiber K, Neville C, Scheibner O, Kavanagh K, et al. (2006) A nonribosomal peptide synthetase (Pes1) confers protection against oxidative stress in *Aspergillus fumigatus*. FEBS J 273: 3038–3053.
- Tsai HF, Chang YC, Washburn RG, Wheeler MH, Kwon-Chung KJ (1998) The developmentally regulated *abl1* gene of *Aspergillus fumigatus*: Its role in modulation of conidial morphology and virulence. J Bacteriol 180: 3031–3038.
- Langfelder K, Streibel M, Jahn B, Haase G, Brakhage AA (2003) Biosynthesis of fungal melanins and their importance for human pathogenic fungi. Fungal Genet Biol 38: 143–158.
- Jahn B, Boukhallouk F, Lotz J, Langfelder K, Wanner G, et al. (2000) Interaction of human phagocytes with pigmentless *Aspergillus nidulans*. Infect Immun 68: 3736–3739.
- Zhang Y, Wilkinson H, Keller N, Tsitsigiannis D (2005) Secondary metabolite gene clusters. In: An Z, editor. Handbook of industrial microbiology. New York: Marcel Dekker. pp. 355–386.
- Keller NP, Turner G, Bennett JW (2005) Fungal secondary metabolism: From biochemistry to genomics. Nat Rev Microbiol 3: 937–947.
- Bok JW, Keller NP (2004) *LaeA*, a regulator of secondary metabolism in *Aspergillus* spp. Eukaryot Cell 3: 527–535.
- Bok JW, Balajee SA, Marr KA, Andes D, Nielsen KF, et al. (2005) *LaeA*, a regulator of morphogenetic fungal virulence factors. Eukaryot Cell 4: 1574–1582.
- Mullbacher A, Eichner RD (1984) Immunosuppression in vitro by a metabolite of a human pathogenic fungus. Proc Natl Acad Sci U S A 81: 3835–3837.
- Sutton P, Newcombe NR, Waring P, Mullbacher A (1994) In vivo immunosuppressive activity of gliotoxin, a metabolite produced by human pathogenic fungi. Infect Immun 62: 1192–1198.
- Sutton P, Waring P, Mullbacher A (1996) Exacerbation of invasive aspergillosis by the immunosuppressive fungal metabolite, gliotoxin. Immunol Cell Biol 74: 318–322.
- Bok JW, Chung D, Balajee SA, Marr KA, Andes D, et al. (2006) GliZ, a transcriptional regulator of gliotoxin biosynthesis, contributes to *Aspergillus fumigatus* virulence. Infect Immun 74: 6761–6768.
- Kupfahl C, Heinekamp T, Geginat G, Ruppert T, Hartl A, et al. (2006) Deletion of the *gliP* gene of *Aspergillus fumigatus* results in loss of gliotoxin production but has no effect on virulence of the fungus in a low-dose mouse infection model. Mol Microbiol 62: 292–302.
- Cramer RA Jr, Gamcsik MP, Brooking RM, Najvar LK, Kirkpatrick WR, et al. (2006) Disruption of a nonribosomal peptide synthetase in *Aspergillus fumigatus* eliminates gliotoxin production. Eukaryot Cell 5: 972–980.
- Bok JW, Noordermeer D, Kale S, Keller NP (2006) Metabolic gene cluster silencing in *Aspergillus nidulans*. Mol Microbiol 61: 1636–1645.
- Tusher VG, Tibshirani R, Chu G (2001) Significance analysis of microarrays applied to the ionizing radiation response. Proc Natl Acad Sci U S A 98: 5116–5121.
- Nierman WC, Pain A, Anderson MJ, Wortman JR, Kim HS, et al. (2005) Genomic sequence of the pathogenic and allergenic filamentous fungus *Aspergillus fumigatus*. Nature 438: 1151–1156.
- Neville C, Murphy A, Kavanagh K, Doyle S (2005) A 4'-phosphopantetheinyl transferase mediates non-ribosomal peptide synthetase activation in *Aspergillus fumigatus*. ChemBiochem 6: 679–685.
- Unsöld IA, Li SM (2005) Overproduction, purification and characterization of FgaPT2, a dimethylallyltryptophan synthase from *Aspergillus fumigatus*. Microbiology 151: 1499–1505.
- Unsöld IA, Li SM (2006) Reverse prenyltransferase in the biosynthesis of fumigaclavine C in *Aspergillus fumigatus*: Gene expression, purification, and characterization of fumigaclavine C synthase FGAPT1. ChemBiochem 7: 158–164.
- Tsai HF, Wheeler MH, Chang YC, Kwon-Chung KJ (1999) A developmentally regulated gene cluster involved in conidial pigment biosynthesis in *Aspergillus fumigatus*. J Bacteriol 181: 6469–6477.
- Coyle CM, Panaccione DG (2005) An ergot alkaloid biosynthesis gene and clustered hypothetical genes from *Aspergillus fumigatus*. Appl Environ Microbiol 71: 3112–3118.
- Grundmann A, Li SM (2005) Overproduction, purification and characterization of FtmPT1, a brevianamide F prenyltransferase from *Aspergillus fumigatus*. Microbiology 151: 2199–2207.
- Maiya S, Grundmann A, Li SM, Turner G (2006) The fumitremorgin gene cluster of *Aspergillus fumigatus*: Identification of a gene encoding Brevianamide F synthetase. ChemBiochem 7: 1062–1069.

28. Cramer RA Jr, Stajich JE, Yamanaka Y, Dietrich FS, Steinbach WJ, et al. (2006) Phylogenomic analysis of non-ribosomal peptide synthetases in the genus *Aspergillus*. *Gene* 383: 24–32.
29. Reiber K, Reeves EP, Neville CM, Winkler R, Gebhardt P, et al. (2005) The expression of selected non-ribosomal peptide synthetases in *Aspergillus fumigatus* is controlled by the availability of free iron. *FEMS Microbiol Lett* 248: 83–91.
30. Hissen AH, Chow JM, Pinto LJ, Moore MM (2004) Survival of *Aspergillus fumigatus* in serum involves removal of iron from transferrin: The role of siderophores. *Infect Immun* 72: 1402–1408.
31. Haas H (2003) Molecular genetics of fungal siderophore biosynthesis and uptake: The role of siderophores in iron uptake and storage. *Appl Microbiol Biotechnol* 62: 316–330.
32. Schrettl M, Bignell E, Kragl C, Joehel C, Rogers T, et al. (2004) Siderophore biosynthesis but not reductive iron assimilation is essential for *Aspergillus fumigatus* virulence. *J Exp Med* 200: 1213–1219.
33. Gardiner DM, Waring P, Howlett BJ (2005) The epipolythiodioxopiperazine (ETP) class of fungal toxins: Distribution, mode of action, functions and biosynthesis. *Microbiology* 151: 1021–1032.
34. Kroll M, Arenzana-Seisdedos F, Bachelerie F, Thomas D, Friguet B, et al. (1999) The secondary fungal metabolite gliotoxin targets proteolytic activities of the proteasome. *Chem Biol* 6: 689–698.
35. Pahl HL, Krauss B, Schulze-Osthoff K, Decker T, Traenckner EB, et al. (1996) The immunosuppressive fungal metabolite gliotoxin specifically inhibits transcription factor NF-kappaB. *J Exp Med* 183: 1829–1840.
36. Suen YK, Fung KP, Lee CY, Kong SK (2001) Gliotoxin induces apoptosis in cultured macrophages via production of reactive oxygen species and cytochrome c release without mitochondrial depolarization. *Free Radic Res* 35: 1–10.
37. Kweon YO, Paik YH, Schnabl B, Qian T, Lemasters JJ, et al. (2003) Gliotoxin-mediated apoptosis of activated human hepatic stellate cells. *J Hepatol* 39: 38–46.
38. Piva TJ (1994) Gliotoxin induces apoptosis in mouse L929 fibroblast cells. *Biochem Mol Biol Int* 33: 411–419.
39. Waring P, Eichner RD, Mullbacher A, Sjaarda A (1988) Gliotoxin induces apoptosis in macrophages unrelated to its antiphagocytic properties. *J Biol Chem* 263: 18493–18499.
40. Beaver JP, Waring P (1994) Lack of correlation between early intracellular calcium ion rises and the onset of apoptosis in thymocytes. *Immunol Cell Biol* 72: 489–499.
41. DeWitte-Orr SJ, Bols NC (2005) Gliotoxin-induced cytotoxicity in three salmonid cell lines: Cell death by apoptosis and necrosis. *Comp Biochem Physiol C Toxicol Pharmacol* 141: 157–167.
42. Pardo J, Urban C, Galvez EM, Ekert PG, Muller U, et al. (2006) The mitochondrial protein Bak is pivotal for gliotoxin-induced apoptosis and a critical host factor of *Aspergillus fumigatus* virulence in mice. *J Cell Biol* 174: 509–519.
43. Gardiner DM, Howlett BJ (2005) Bioinformatic and expression analysis of the putative gliotoxin biosynthetic gene cluster of *Aspergillus fumigatus*. *FEMS Microbiol Lett* 248: 241–248.
44. Lewis RE, Wiederhold NP, Chi J, Han XY, Komanduri KV, et al. (2005) Detection of gliotoxin in experimental and human aspergillosis. *Infect Immun* 73: 635–637.
45. Guillemette T, Sellam A, Simoneau P (2004) Analysis of a nonribosomal peptide synthetase gene from *Alternaria brassicae* and flanking genomic sequences. *Curr Genet* 45: 214–224.
46. Panaccione DG (2005) Origins and significance of ergot alkaloid diversity in fungi. *FEMS Microbiol Lett* 251: 9–17.
47. Panaccione DG, Coyle CM (2005) Abundant respirable ergot alkaloids from the common airborne fungus *Aspergillus fumigatus*. *Appl Environ Microbiol* 71: 3106–3111.
48. Glatt H, Pertz H, Kasper R, Eich E (1992) Clavine alkaloids and derivatives as mutagens detected in the Ames test. *Anticancer Drugs* 3: 609–614.
49. Brakhage AA, Liebmann B (2005) *Aspergillus fumigatus* conidial pigment and cAMP signal transduction: Significance for virulence. *Med Mycol* 43 (Suppl 1): S75–S82.
50. Galagan JE, Calvo SE, Cuomo C, Ma LJ, Wortman JR, et al. (2005) Sequencing of *Aspergillus nidulans* and comparative analysis with *A. fumigatus* and *A. oryzae*. *Nature* 438: 1105–1115.
51. Nierman WC, May G, Kim HS, Anderson MJ, Chen D, et al. (2005) What the *Aspergillus* genomes have told us. *Med Mycol* 43 (Suppl 1): S3–S5.
52. Ronning CM, Fedorova ND, Bowyer P, Coulson R, Goldman G, et al. (2005) Genomics of *Aspergillus fumigatus*. *Rev Iberoam Micol* 22: 223–228.
53. Chang PK, Ehrlich KC, Hua SS (2006) Cladal relatedness among *Aspergillus oryzae* isolates and *Aspergillus flavus* S and L morphotype isolates. *Int J Food Microbiol* 108: 172–177.
54. Talbot NJ, Vincent P, Wildman HG (1996) The influence of genotype and environment on the physiological and metabolic diversity of *Fusarium compactum*. *Fungal Genet Biol* 20: 254–267.
55. Ward TJ, Bielawski JP, Kistler HC, Sullivan E, O'Donnell K (2002) Ancestral polymorphism and adaptive evolution in the trichothecene mycotoxin gene cluster of phytopathogenic *Fusarium*. *Proc Natl Acad Sci U S A* 99: 9278–9283.
56. Walton JD (2000) Horizontal gene transfer and the evolution of secondary metabolite gene clusters in fungi: An hypothesis. *Fungal Genet Biol* 30: 167–171.
57. Keller NP, Hohn TM (1997) Metabolic pathway gene clusters in filamentous fungi. *Fungal Genet Biol* 21: 17–29.
58. Buades C, Moya A (1996) Phylogenetic analysis of the isopenicillin-N-synthetase horizontal gene transfer. *J Mol Evol* 42: 537–542.
59. Shimizu K, Keller NP (2001) Genetic involvement of a cAMP-dependent protein kinase in a G protein signaling pathway regulating morphological and chemical transitions in *Aspergillus nidulans*. *Genetics* 157: 591–600.
60. Ferreira ME, Malavazi I, Savoldi M, Brakhage AA, Goldman MH, et al. (2006) Transcriptome analysis of *Aspergillus fumigatus* exposed to voriconazole. *Curr Genet* 50: 32–44.
61. Cleveland WS (1979) Robust locally weighted regression and smoothing scatterplots. *J Am Stat Assoc* 74: 829–836.
62. Saeed AI, Sharov V, White J, Li J, Liang W, et al. (2003) TM4: A free, open-source system for microarray data management and analysis. *Biotechniques* 34: 374–378.
63. Saeed AI, Bhagabati NK, Braisted JC, Liang W, Sharov V, et al. (2006) TM4 microarray software suite. *Methods Enzymol* 411: 134–193.
64. Hosack DA, Dennis G Jr, Sherman BT, Lane HC, Lempicki RA (2003) Identifying biological themes within lists of genes with EASE. *Genome Biol* 4: R70.
65. Finn RD, Mistry J, Schuster-Bockler B, Griffiths-Jones S, Hollich V, et al. (2006) Pfam: Clans, Web tools and services. *Nucleic Acids Res* 34: D247–D251.