

# Virtual Reality as a Surgical Care Package for Patients Undergoing Weight Loss Surgery: A Narrative Review of the Impact of an Emerging Technology

Review began 09/14/2022

Review ended 09/20/2022

Published 09/26/2022

© Copyright 2022

Gendia et al. This is an open access article distributed under the terms of the Creative Commons Attribution License CC-BY 4.0., which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Ahmed Gendia <sup>1</sup>, Ayman Zyada <sup>2</sup>, Muhammad Talal Nasir <sup>3</sup>, Mohannad Elfar <sup>3</sup>, Mohamed Sakr <sup>3</sup>, Masood U. Rehman <sup>3</sup>, Alwyn Cota <sup>1</sup>, James Clark <sup>1</sup>

1. Bariatric Surgery, Royal Cornwall Hospitals NHS Trust, Truro, GBR 2. Vascular Surgery, Northampton General Hospital, Northampton, GBR 3. General Surgery, Northampton General Hospital, Northampton, GBR

Corresponding author: Ahmed Gendia, agandia@live.com

---

---

## Abstract

While bariatric surgery is regarded as the most effective treatment for people with severe and morbid obesity, its pathway is regarded as a complex one due to the multidisciplinary approaches required from pre-surgery education until long-term management. This is essential to maintain weight loss and improve the quality of life after bariatric surgery. Although these approaches are broadened, patient education, pre-operative preparation, behavioural therapy, rehabilitation, and dietary changes are regarded as the main domains in such complex care. With the increase in technological adaptation in medical services, virtual reality (VR) has shown many benefits that can be utilized in the care of bariatric patients undergoing surgery. However, VR has not been innovated to be a multidomain care package in which bariatric patients could benefit throughout their journey from the pre-operative optimization, recovery, and long-term follow-up. This review aims to give a brief description of some of the applications of VR technology and question whether it has the potential to be considered as a virtual ecosystem to improve the bariatric patients' experience and pathway throughout surgery and follow-up.

---

**Categories:** General Surgery, Healthcare Technology, Other

**Keywords:** weight loss surgery, weight loss, virtual reality, bariatric surgery, vr games, virtual reality simulation, virtual reality exposure therapy, virtual reality-based rehabilitation

## Introduction And Background

Obesity constitutes an important threat to national and global public health in terms of its prevalence and rising incidence, quality of life, life expectancy, and economic burden [1,2]. In severe obesity, bariatric surgery is the most effective therapeutic option to achieve long-term weight loss and improve the associated comorbidities [3]. This has made Roux-en-Y gastric bypass (RYGB), sleeve gastrectomy (SG), and adjustable gastric banding the most popular and commonly performed bariatric surgeries [4]. However, a small proportion of patients have also been reported to not reach their optimum goal for weight loss two years after the procedure and very few can fail or regain the weight. While anatomical factors can play a part, behavioural and psychosocial optimizations are regarded as equally important. This includes eating patterns, depression, nutritional factors, and exercise [5,6].

Virtual reality (VR) development and applications have gained wide recognition in medical services by providing solutions to improve patients' outcomes. This is through patients' education, improving mental health, and post-operative care, including pain management, physical therapy, and rehabilitation [7,8]. VR is a computer-generated simulation of a real or imagined environment. It can be immersive or non-immersive according to its ability to involve the users [9]. The former has been the focus of many medical applications due to its ability to give the user control of the reproduced environment. Immersive virtual reality (IVR) is usually delivered in a variety of ways and the most popular being head-mounted displays or simply a headset [8].

We aim to provide insight on some of these immersive applications and how they can be included to enhance the patient pathway to optimize outcomes both in the pre- and post-operative period for patients undergoing bariatric surgery.

## Review

### Methods


A systematic search following the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) research criteria was conducted from January 2015 to December 2021. PubMed was searched using the following keywords: virtual reality, patient education, anxiety and pain, physical rehabilitation,

#### How to cite this article

Gendia A, Zyada A, Nasir M, et al. (September 26, 2022) Virtual Reality as a Surgical Care Package for Patients Undergoing Weight Loss Surgery: A Narrative Review of the Impact of an Emerging Technology. Cureus 14(9): e29608. DOI 10.7759/cureus.29608


behavioural support, obesity, eating disorders, body image, and substance cessation.

Thirty-four studies were identified and included in the final manuscript (Figure 1) supporting VR technology across applications that can be applied to bariatric patients' surgical pathways. The applications were subcategorized into eight different areas of interest, which can help to shape the concept of the virtual ecosystem of bariatric patients (Figure 2).



**FIGURE 1: PRISMA flow chart for literature search and included studies**

PRISMA: Preferred Reporting Items for Systematic Reviews and Meta-Analyses; VR: virtual reality.



**FIGURE 2: Elements of ERAS and long-term recommendations for bariatric patients and established virtual reality applications with projection into the virtual ecosystem of bariatric patients**

ERAS: Enhanced Recovery After Surgery.

## Results

VR applications have been described in the eight domains mentioned below, which can be applied in relation to patients undergoing bariatric surgery.

### *Virtual Reality Patient Education (VR PE)*

VR education has been introduced to make the information more meaningful and patient-centred by enabling its users to be fully immersed in an interactive simulated and self-controllable visual and auditory experience [10]. In a study by Pandrangi et al. [11], VR was found to be a useful informative tool in educating patients about their aneurysmal disease through interactive reconstructed three-dimensional (3D) images of their aortic anatomy. The majority of the patients in this study agreed that VR 3D anatomy helped to improve their understanding and therefore felt more engaged in their healthcare decisions [11].

VR PE has also played a role in improving the stress levels of patients undergoing radiotherapy (RT) by improving clarity and levels of education about their treatment. A randomized study on 60 patients with chest malignancy showed that patients who received VR PE showed significant improvement in comprehension and reduction in stress and anxiety levels when compared to standard education [12]. Another study on 43 patients utilized VR PE by creating 3D images of patients in RT sessions and what to expect during the treatment. After the VR PE, 95% of patients agreed that they had a clear understanding of how they would feel when lying on the treatment table. Also, patients' understanding of the location and the size of their cancer had significantly improved from around 50% to 95% with an increase in the orientation of side effects of the treatment by 30% post-VR PE [13].

In bariatric surgery, there is no currently reported data on the applications of VR education. However, the potential impact of VR PE can be numerous across the weight management pathway. Preoperatively, bariatric patients could potentially utilize VR to be virtually educated about different surgical options versus conservative treatment through enhanced 3D interactive images. This could be seen to help in better understanding of their options including surgery and thereby enhancing informed consent and overall education.

Post-operatively, VR-enhanced education could provide an option for daily or weekly updates on lifestyle changes, which could help in improving compliance. Importantly, this can be done from the comfort of the patient's home with the added advantage of reducing costs and time for travelling to attend appointments.

### *Anxiety Related to Surgery*

A significant amount of anxiety related to surgery is due to the fear and uncertainty of the outcomes. Its

psychological and physical effects are associated with longer recovery, an increase in the need for analgesia, anaesthetic requirement, and unfavourable behavioural and emotional outcomes [14]. Conventional methods of mitigation of preoperative anxiety are pharmacological and non-pharmacological strategies [15].

Recently, with promising results in the management of anxiety and other psychiatric disorders, VR has been successfully applied to reduce anxiety related to surgery in different surgical settings [16]. Chan et al. [17] tested the effect of VR relaxing meditation and breathing exercises on 108 women undergoing hysteroscopy. This showed that anxiety scores were significantly reduced after the 10 minutes of VR content, which helped in reducing pain and stress related to surgery. Also, around 85% of patients reported the VR experience as good or excellent [17].

In minimally invasive abdominal surgery, Haisley et al. [18] used VR meditation as a perioperative tool with favourable results in reducing pain, anxiety, and nausea and around 75% of patients stated that they would use the VR again [18]. Similarly, VR meditation showed favourable results in reducing pain and anxiety in burns and complex pain [19,20].

The rationale for using VR to improve anxiety preoperatively is by immersing patients in a fully simulated relaxing environment with the objective of placing them in a more empowered state to deal with the triggers of their anxiety [21]. This could be applied to the bariatric population before surgery. It is to be seen from future studies whether these expected results can be validated in bariatric patients. There is therefore the potential for obtaining better evidence for patient satisfaction and reducing stress related to bariatric surgery.

#### *Pain Management*

Successful pain management is a key element of the post-operative course as it shortens recovery and reduces risks of cardiovascular and pulmonary complications. In bariatric surgery, pain management is essential to enhance recovery and prompt early mobilization, which helps to decrease venous thromboembolism, prevent other events, and reduce hospitalization [22]. Therefore, a multimodal approach through regional and systematic analgesia is considered the most effective method as it minimizes opiate use, which can induce obstructive sleep apnoea, which is more liable due to the co-morbidities of obesity [23].

Applications of VR in pain management in other surgical patients have been reported to have numerous benefits. This includes a reduction in pain scores after cardiac, knee, abdominal, and spinal surgery with overall patients reporting the use of VR as a pleasant experience and stating that they would use it again on further occasions [18,24,25]. VR pain management follows a similar concept to VR and anxiety meditation by immersing patients in a simulated relaxing environment, which can help to divert the patient's feelings from their pain. This could be playing a major role in bariatric patients' management of pain and anxiety related to surgery with proper application integration in their peri-operative pathway.

#### *Optimizing Pulmonary Function for Surgery*

Respiratory function in morbidly obese patients follows a restrictive pattern with up to 77% suffering from obstructive sleep apnoea [26]. This increases the risk of impaired post-operative oxygenation and other respiratory complications in the form of atelectasis. Optimization of pulmonary function for surgery includes smoking cessation, breathing exercises, including inspiratory muscle training, incentive spirometry, and optimization of chronic disease, for example, asthma and chronic obstructive pulmonary disease (COPD) [27].

With the increase of applications of VR in different rehabilitation programmes, VR has been aiding in pulmonary exercises in both healthy individuals and COPD patients [28,29]. VR pulmonary rehabilitation is designed to enable home-based exercises in the form of a 3D avatar instructor in an immersive relaxing environment to guide patients through breathing exercises based on traditional rehabilitation programmes [30]. In COPD patients, VR-based respiratory rehabilitation has shown to have similar outcomes when compared to a conventional programme with the additional benefit of performing the exercises from home. Moreover, VR showed enrichment of experience by also decreasing the levels of anxiety during exercise and therefore optimizing cardiorespiratory function [31].

#### *Physical Fitness Applications*

Pre- and post-operative physical activity (PA) is regarded as an important element in enhancing recovery after surgery as it improves physical state, responses to stress from surgery, and improvement of cardiovascular function, thereby reducing complications [32].

In the bariatric population, a structured exercise regime is considered a feasible and effective adjunct therapy that benefits cardiometabolic parameters when compared to those with bariatric surgery alone [33].

Exercise before surgery has shown to be beneficial in reducing body weight, improving blood pressure, general fitness, quality of life satisfaction, and decreasing fasting plasma insulin and blood lipid. Exercise after bariatric surgery has been shown to preserve dynamic muscle strength and contribute to maintaining weight loss after calorie restriction [34].

Although PA promotion is recognized as an important component of weight loss programmes, there are no current evidence-based or standardized bariatric surgery-specific PA guidelines [35]. Reported exercise regimes ranged from walking, aquatic, resistance, and supervised exercises. Also, adherence to exercise before and after surgery plays a big role in physical rehabilitation. As in the bariatric population, many can face barriers in the form of low confidence levels in their abilities and not feeling comfortable going to the gym due to real and perceived discrimination. Therefore, many come up with the belief of not having time to participate in sports [36].

VR rehabilitation has gained much recognition from dedicated platforms like treadmills, diving, cycling simulators, and medically oriented VR rehabilitation. These studies have demonstrated increased participation of users utilizing VR exercise programmes [37]. VR rehabilitation and exercise have shown to be effective in healthy individuals and different medical rehabilitations. It was reported to be equivalent and sometimes more superior to standard physiotherapy in cerebral palsy, spinal injury, and stroke [38]. In healthy individuals, VR exercise was demonstrated to increase adherence and enjoyment with positive physiological effects during exercise [39]. It was also reported that obese children performed better on treadmills while using VR than traditional walking, as VR allowed more distraction and less discomfort [40].

VR exercises during rehabilitation can therefore potentially play a major role in pre- and post-operative PA improvement in bariatric patients. Given the feasibility and the safety of these home-based devices, it can decrease the load on healthcare services, as most of the standard pre-operative programmes are resource intensive.

#### *Virtual Reality and Enhanced Cognitive Behavioural Therapy*

Eating and depressive disorders significantly affect the bariatric population with a prevalence of 24% and 17%, respectively. Both can lead to less post-operative weight loss, weight regains, impaired general psychology, and quality of life [41]. Cognitive behavioural therapy (CBT) is recommended for patients undergoing weight loss surgery (WLS). It has been shown to improve self-monitoring and control eating behaviours with significant improvement in depression and anxiety and therefore better results [42].

Over the last decades, VR-enhanced cognitive therapy (VRCBT) has been embraced for being a novel way to deliver CBT. The technique creates an interactive 3D environment to simulate successful goal achievement. This helps patients to overcome memories of previous real-life experiences through emotionally guided virtual exposure [43]. VRCBT has shown favourable results in anxiety, phobias, social anxiety disorders, and depression [21]. Moreover, randomized trials have shown VRCBT to be superior to conventional CBT in managing eating disorders and binge eating [44,45]. This helped in weight reduction therapy and adding adherence to programmes [46].

There is a paucity of evidence of the use of VR in the overweight and morbidly obese population. Phelan et al. [47] tested the use of a VR environment on 15 overweight adults for four weeks with the main hypothesis to evaluate the effect of the simulated scenes on behavioural skills related to eating habits. Although they showed no difference in weight loss among participants, VR intervention was more preferred by patients over traditional weight loss programmes [47]. Manzoni et al. [45] tested the efficacy of an enhanced VRCBT module aimed to unlock the negative memory of the body and modify its behavioural and emotional behaviour. A total of 163 female morbidly obese inpatients were randomly assigned to three CBT-based treatments: a standard behavioural inpatient programme (SBP), SBP plus standard CBT, and SBP plus VR-enhanced CBT. The study showed that patients in the VR group had a greater probability of maintaining or improving weight loss at one-year follow-up than SBP patients and, to a lesser extent, CBT patients. On the contrary, participants who received only a behavioural programme regained on average most of the weight they had lost [45].

VRCBT can therefore be a valuable tool in managing behavioural disorders related to obesity in patients undergoing WLS. This can help in maintaining weight loss and improving well-being and quality of life.

#### *Virtual Reality and Body Image (VRBI)*

Body image disorders (BIDs) are linked to various psychological and physical sequelae of impaired functions, for instance, depression, anxiety, eating disorders, and poor quality of life [48]. Among the bariatric population, body image dissatisfaction is associated with binge eating, depression, and lower self-esteem, with one in five bariatric patients identifying appearance as their main motive for surgery [49]. Improvement in body image perception after successful surgery has been linked to a decrease in compulsive eating syndromes, reduction in body mass index (BMI), and improvement in self-esteem and intimate relationships [50].

A contrary aspect of body image after surgery includes the issue of excess skin with massive weight reduction. This has been linked to poor body satisfaction, dermatitis and skin fold irritations, and impairment in daily activities and exercise. In turn, this leads 85% of bariatric patients to seek body-contouring surgery (BCS) to alleviate this problem [51].

The application of VR has been used to improve BID. This is by creating a 3D simulation of their bodies in the form of avatars through an immersive environment that reproduces situations related to their body image concerns. Through multisensory simulations, it produces an empowered feeling of ownership of one's body, which consequently promotes a healthier body image and behaviour [52]. A recent systematic review of six studies utilizing avatars and VR in weight loss programmes showed that avatar-based interventions were effective in both short- and mid-term weight loss. Also, the technology helped to improve exercise adherence in the long term [53]. VR was also used to assess the BID of 78 women with different BMIs by exposing the participants to different versions of avatars: slimmer, same weight, and overweight. The study showed that women with higher BMI reported more BID on their replicated avatar and showed satisfaction with their slimmer version. This finding indicated that VR may serve as a novel tool for measuring BID [54].

Potentially, VR avatars can also play a role in body image perception in bariatric patients. It can be integrated to improve BIDs by recreating slimmer avatars, which could promote adherence to weight loss and exercise programmes.

#### *Smoking and Alcohol*

While the increase in BMI is a risk factor for adverse outcomes related to surgical procedures, smoking's hazardous effects range from increased risks of pulmonary complications, wound infection, venous thromboembolism, and slower recovery. Similarly, alcohol consumption before surgery can lead to increased unfavourable outcomes [55]. Smoking and other substance abuse are recommended to be stopped four to six weeks pre-operatively [56]. VR has been tested as a potential solution to stop smoking and alcohol usage by inducing an advanced cue exposure therapy (CET), which was superior to static images or videos used in conventional settings [57]. Also, VR exposure therapy (VRET) has been reported to be more effective if combined with conventional cognitive behaviour therapy in relation to stopping smoking [58].

Although its applications are still under development and validation, VRET in smoking and alcohol cessation could play an important role in optimizing patients undergoing bariatric surgery as a part of a virtual reality surgical care package (VRSCP).

## Discussion

Patients who are candidates for WLS usually undergo variable preparatory phase and post-operative optimization to improve both short- and long-term results. Standard care models usually involve education and follow-up through multidisciplinary teams with reflection on the patient's progress through educational sessions and follow-up plans.

While VR applications are being investigated in many surgical and medical specialties, their application to patients undergoing WLS is limited and not yet explored. The favourable applications of VR in patient education, anxiety and pain management, preoperative optimization, and behavioural and physiological treatment can be packaged as a surgical care bundle making bariatric patients' journey more satisfactory with the potential for improved outcomes.

Despite its promising applications, VR is still an emerging technology and has its own initial drawbacks to gaining traction in the healthcare system. There are several reasons for this. Firstly, the obvious cost of the systems and the absence of adequate clinical validation could play a major role in limiting widespread adoption. Further delays in adoption would likely be seen within the education of both healthcare providers and their patients, particularly on the application and utilization of the systems. The technology is still seen to be clumsy to wear and will need educational support to use [59].

With the increased investments and advancement in VR technology, education of healthcare professionals and further studies demonstrating evidence of improved outcomes, VR will play a major role in surgical patients and more specifically bariatric patients. This could be even refined as a personalized surgical care package. This will contribute to a fully virtual ecosystem in health care.

## Conclusions

VR as a new technology in healthcare services will play an increasingly important role in improving patients' journey through surgery. VR has shown primary favourable results in managing obesity-related problems, anxiety, and pain related to surgery, improving physical fitness, physiotherapy, and other services; however, it has not yet been directed toward a patient's package in the bariatric population. It can be utilized as a virtual ecosystem in the pre-operative stage, patient education, physical preparation, and mental health support. VR applications can be also evident in admission and peri-operative period, anxiety, and pain



management. Moreover, VR applications can be ultimately useful in the post-operative phase and long-term management of bariatric patients through integration with cognitive therapy, body image disorders, and ongoing motivational fitness programmes.

Although VR is still in the early development phases, the primes of its integration into bariatric patient pathways can be limitless. Therefore, further studies into the application of VR in these patients as a surgical package (VRSCP) should contribute to a fully patient-centred virtual ecosystem for bariatric patients in the future.

## Additional Information

### Disclosures

**Conflicts of interest:** In compliance with the ICMJE uniform disclosure form, all authors declare the following: **Payment/services info:** All authors have declared that no financial support was received from any organization for the submitted work. **Financial relationships:** All authors have declared that they have no financial relationships at present or within the previous three years with any organizations that might have an interest in the submitted work. **Other relationships:** All authors have declared that there are no other relationships or activities that could appear to have influenced the submitted work.

### Acknowledgements

Contributions: A. Gendia provided the concept of the virtual reality surgical care bundle (VRSCB), wrote the manuscript, and supervised the literature search. A. Zyada and M.T. Nasir performed the literature search and extracted articles to be included in the review. M. Elfar provided the primary manuscript related to anxiety, pain management, and optimizing pulmonary function sections. M.U. Rehman provided the primary manuscript related to physical fitness, VR-enhanced cognitive behavioural therapy, and body image. A. Cota provided supervision and revision of the manuscript. J. Clark provided final revision and supervision of the manuscript.

## References

1. Tremmel M, Gerdtham UG, Nilsson PM, Saha S: Economic burden of obesity: a systematic literature review. *Int J Environ Res Public Health*. 2017, 14:435. [10.3390/ijerph14040435](https://doi.org/10.3390/ijerph14040435)
2. Fruh SM: Obesity: risk factors, complications, and strategies for sustainable long-term weight management. *J Am Assoc Nurse Pract*. 2017, 29:S3-14. [10.1002/2327-6924.12510](https://doi.org/10.1002/2327-6924.12510)
3. Maggard MA, Shugarman LR, Suttrop M, et al.: Meta-analysis: surgical treatment of obesity. *Ann Intern Med*. 2005, 142:547-59. [10.7326/0003-4819-142-7-200504050-00013](https://doi.org/10.7326/0003-4819-142-7-200504050-00013)
4. Mulita F, Lampropoulos C, Kehagias D, et al.: Long-term nutritional deficiencies following sleeve gastrectomy: a 6-year single-centre retrospective study. *Prz Menopauzalny*. 2021, 20:170-6. [10.5114/pm.2021.110954](https://doi.org/10.5114/pm.2021.110954)
5. Sheets CS, Peat CM, Berg KC, White EK, Bocchieri-Ricciardi L, Chen EY, Mitchell JE: Post-operative psychosocial predictors of outcome in bariatric surgery. *Obes Surg*. 2015, 25:330-45. [10.1007/s11695-014-1490-9](https://doi.org/10.1007/s11695-014-1490-9)
6. Neff KJ, Olbers T, le Roux CW: Bariatric surgery: the challenges with candidate selection, individualizing treatment and clinical outcomes. *BMC Med*. 2013, 11:8. [10.1186/1741-7015-11-8](https://doi.org/10.1186/1741-7015-11-8)
7. Mantovani F, Castelnuovo G, Gaggioli A, Riva G: Virtual reality training for health-care professionals. *Cyberpsychol Behav*. 2003, 6:389-95. [10.1089/10949310332278772](https://doi.org/10.1089/10949310332278772)
8. The growing value of XR in healthcare in the United Kingdom. (2021). <https://www.xrhealthuk.org/the-growing-value-of-xr-in-healthcare>.
9. Cipresso P, Giglioli IA, Raya MA, Riva G: The past, present, and future of virtual and augmented reality research: a network and cluster analysis of the literature. *Front Psychol*. 2018, 9:2086. [10.3389/fpsyg.2018.02086](https://doi.org/10.3389/fpsyg.2018.02086)
10. Adapa K, Jain S, Kanwar R, Zaman T, Taneja T, Walker J, Mazur L: Augmented reality in patient education and health literacy: a scoping review protocol. *BMJ Open*. 2020, 10:e038416. [10.1136/bmjopen-2020-038416](https://doi.org/10.1136/bmjopen-2020-038416)
11. Pandrangi VC, Gaston B, Appelbaum NP, Albuquerque FC Jr, Levy MM, Larson RA: The application of virtual reality in patient education. *Ann Vasc Surg*. 2019, 59:184-9. [10.1016/j.avsg.2019.01.015](https://doi.org/10.1016/j.avsg.2019.01.015)
12. Gao J, Liu S, Zhang S, et al.: Pilot study of a virtual reality educational intervention for radiotherapy patients prior to initiating treatment. *J Cancer Educ*. 2022, 37:578-85. [10.1007/s13187-020-01848-5](https://doi.org/10.1007/s13187-020-01848-5)
13. Wang LJ, Casto B, Luh JY, Wang SJ: Virtual reality-based education for patients undergoing radiation therapy. *J Cancer Educ*. 2022, 37:694-700. [10.1007/s13187-020-01870-7](https://doi.org/10.1007/s13187-020-01870-7)
14. Gerbershagen HJ, Dagtekin O, Rothe T, et al.: Risk factors for acute and chronic postoperative pain in patients with benign and malignant renal disease after nephrectomy. *Eur J Pain*. 2009, 13:853-60. [10.1016/j.ejpain.2008.10.001](https://doi.org/10.1016/j.ejpain.2008.10.001)
15. Ayyadhah Alanazi A: Reducing anxiety in preoperative patients: a systematic review. *Br J Nurs*. 2014, 23:387-93. [10.12968/bjon.2014.23.7.387](https://doi.org/10.12968/bjon.2014.23.7.387)
16. Koo CH, Park JW, Ryu JH, Han SH: The effect of virtual reality on preoperative anxiety: a meta-analysis of randomized controlled trials. *J Clin Med*. 2020, 9:3151. [10.3390/jcm9103151](https://doi.org/10.3390/jcm9103151)
17. Chan JJ, Yeam CT, Kee HM, Tan CW, Sultana R, Sia AT, Sng BL: The use of pre-operative virtual reality to reduce anxiety in women undergoing gynecological surgeries: a prospective cohort study. *BMC Anesthesiol*. 2020, 20:261. [10.1186/s12871-020-01177-6](https://doi.org/10.1186/s12871-020-01177-6)
18. Haisley KR, Straw OJ, Müller DT, et al.: Feasibility of implementing a virtual reality program as an adjuvant

- tool for peri-operative pain control; results of a randomized controlled trial in minimally invasive foregut surgery. *Complement Ther Med*. 2020, 49:102356. [10.1016/j.ctim.2020.102356](https://doi.org/10.1016/j.ctim.2020.102356)
19. Jeffs D, Dorman D, Brown S, et al.: Effect of virtual reality on adolescent pain during burn wound care. *J Burn Care Res*. 2014, 35:395-408. [10.1097/BCR.0000000000000019](https://doi.org/10.1097/BCR.0000000000000019)
  20. Sato K, Fukumori S, Matsusaki T, et al.: Nonimmersive virtual reality mirror visual feedback therapy and its application for the treatment of complex regional pain syndrome: an open-label pilot study. *Pain Med*. 2010, 11:622-9. [10.1111/j.1526-4637.2010.00819.x](https://doi.org/10.1111/j.1526-4637.2010.00819.x)
  21. Maples-Keller JL, Bunnell BE, Kim SJ, Rothbaum BO: The use of virtual reality technology in the treatment of anxiety and other psychiatric disorders. *Harv Rev Psychiatry*. 2017, 25:103-13. [10.1097/HRP.0000000000000158](https://doi.org/10.1097/HRP.0000000000000158)
  22. Mulita F, Karpetas G, Liolis E, Vilas M, Tchabashvili L, Maroulis I: Comparison of analgesic efficacy of acetaminophen monotherapy versus acetaminophen combinations with either pethidine or parecoxib in patients undergoing laparoscopic cholecystectomy: a randomized prospective study. *Med Glas (Zenica)*. 2021, 18:27-32. [10.17392/1245-21](https://doi.org/10.17392/1245-21)
  23. Schug SA, Raymann A: Postoperative pain management of the obese patient. *Best Pract Res Clin Anaesthesiol*. 2011, 25:73-81. [10.1016/j.bpa.2010.12.001](https://doi.org/10.1016/j.bpa.2010.12.001)
  24. Sharif F, Rahman A, Tonner E, et al.: Can technology optimise the pre-operative pathway for elective hip and knee replacement surgery: a qualitative study. *Perioper Med (Lond)*. 2020, 9:33. [10.1186/s13741-020-00166-0](https://doi.org/10.1186/s13741-020-00166-0)
  25. Mosso-Vázquez JL, Gao K, Wiederhold BK, Wiederhold MD: Virtual reality for pain management in cardiac surgery. *Cyberpsychol Behav Soc Netw*. 2014, 17:371-8. [10.1089/cyber.2014.0198](https://doi.org/10.1089/cyber.2014.0198)
  26. Jehan S, Zizi F, Pandi-Perumal SR, et al.: Obstructive sleep apnea and obesity: implications for public health. *Sleep Med Disord*. 2017, 1:00019.
  27. Azhar N: Pre-operative optimisation of lung function. *Indian J Anaesth*. 2015, 59:550-6. [10.4103/0019-5049.165858](https://doi.org/10.4103/0019-5049.165858)
  28. Rockstroh C, Blum J, Göritz AS: A mobile VR-based respiratory biofeedback game to foster diaphragmatic breathing. *Virtual Reality*. 2021, 25:539-52. [10.1007/s10055-020-00471-5](https://doi.org/10.1007/s10055-020-00471-5)
  29. Jung T, Moorhouse N, Shi X, Amin MF: A virtual reality-supported intervention for pulmonary rehabilitation of patients with chronic obstructive pulmonary disease: mixed methods study. *J Med Internet Res*. 2020, 22:e14178. [10.2196/14178](https://doi.org/10.2196/14178)
  30. Condon C, Lam WT, Mosley C, Gough S: A systematic review and meta-analysis of the effectiveness of virtual reality as an exercise intervention for individuals with a respiratory condition. *Adv Simul (Lond)*. 2020, 5:33. [10.1186/s41077-020-00151-z](https://doi.org/10.1186/s41077-020-00151-z)
  31. Rutkowski S, Szczegielniak J, Szczepańska-Gieracha J: Evaluation of the efficacy of immersive virtual reality therapy as a method supporting pulmonary rehabilitation: a randomized controlled trial. *J Clin Med*. 2021, 10:352. [10.3390/jcm10020352](https://doi.org/10.3390/jcm10020352)
  32. Małczak P, Pisarska M, Piotr M, Wysocki M, Budzyński A, Pędziwiatr M: Enhanced recovery after bariatric surgery: systematic review and meta-analysis. *Obes Surg*. 2017, 27:226-35. [10.1007/s11695-016-2438-z](https://doi.org/10.1007/s11695-016-2438-z)
  33. Coen PM, Tanner CJ, Helbling NL, et al.: Clinical trial demonstrates exercise following bariatric surgery improves insulin sensitivity. *J Clin Invest*. 2015, 125:248-57. [10.1172/JCI78016](https://doi.org/10.1172/JCI78016)
  34. Baillot A, Mampuya WM, Comeau E, Méziat-Burdin A, Langlois MF: Feasibility and impacts of supervised exercise training in subjects with obesity awaiting bariatric surgery: a pilot study. *Obes Surg*. 2013, 23:882-91. [10.1007/s11695-013-0875-5](https://doi.org/10.1007/s11695-013-0875-5)
  35. King WC, Bond DS: The importance of preoperative and postoperative physical activity counseling in bariatric surgery. *Exerc Sport Sci Rev*. 2013, 41:26-35. [10.1097/ES.0b013e31826444e0](https://doi.org/10.1097/ES.0b013e31826444e0)
  36. Hunt HR, Gross AM: Prediction of exercise in patients across various stages of bariatric surgery: a comparison of the merits of the theory of reasoned action versus the theory of planned behavior. *Behav Modif*. 2009, 33:795-817. [10.1177/0145445509348055](https://doi.org/10.1177/0145445509348055)
  37. Dębska M, Polechoński J, Mynarski A, Polechoński P: Enjoyment and intensity of physical activity in immersive virtual reality performed on innovative training devices in compliance with recommendations for health. *Int J Environ Res Public Health*. 2019, 16:3673. [10.3390/ijerph16193673](https://doi.org/10.3390/ijerph16193673)
  38. Qian J, McDonough DJ, Gao Z: The effectiveness of virtual reality exercise on individual's physiological, psychological and rehabilitative outcomes: a systematic review. *Int J Environ Res Public Health*. 2020, 17:4133. [10.3390/ijerph17114133](https://doi.org/10.3390/ijerph17114133)
  39. Liu W, Zeng N, Pope ZC, McDonough DJ, Gao Z: Acute effects of immersive virtual reality exercise on young adults' situational motivation. *J Clin Med*. 2019, 8:1947. [10.3390/jcm8111947](https://doi.org/10.3390/jcm8111947)
  40. Baños RM, Escobar P, Cebolla A, Guixeres J, Alvarez Pitti J, Lisón JF, Botella C: Using virtual reality to distract overweight children from bodily sensations during exercise. *Cyberpsychol Behav Soc Netw*. 2016, 19:115-9. [10.1089/cyber.2015.0283](https://doi.org/10.1089/cyber.2015.0283)
  41. Rudolph A, Hilbert A: Cognitive-behavioral therapy for postbariatric surgery patients with mental disorders: a pilot study. *Front Psychiatry*. 2020, 11:14. [10.3389/fpsy.2020.00014](https://doi.org/10.3389/fpsy.2020.00014)
  42. Paul L, van der Heiden C, Hoek HW: Cognitive behavioral therapy and predictors of weight loss in bariatric surgery patients. *Curr Opin Psychiatry*. 2017, 30:474-9. [10.1097/YCO.0000000000000359](https://doi.org/10.1097/YCO.0000000000000359)
  43. Vincelli F: From imagination to virtual reality: the future of clinical psychology. *Cyberpsychol Behav*. 1999, 2:241-8. [10.1089/cpb.1999.2.241](https://doi.org/10.1089/cpb.1999.2.241)
  44. Cesa GL, Manzoni GM, Bacchetta M, et al.: Virtual reality for enhancing the cognitive behavioral treatment of obesity with binge eating disorder: randomized controlled study with one-year follow-up. *J Med Internet Res*. 2013, 15:e113. [10.2196/jmir.2441](https://doi.org/10.2196/jmir.2441)
  45. Manzoni GM, Cesa GL, Bacchetta M, et al.: Virtual reality-enhanced cognitive-behavioral therapy for morbid obesity: a randomized controlled study with 1 year follow-up. *Cyberpsychol Behav Soc Netw*. 2016, 19:134-40. [10.1089/cyber.2015.0208](https://doi.org/10.1089/cyber.2015.0208)
  46. Clus D, Larsen ME, Lemey C, Berruiguet S: The use of virtual reality in patients with eating disorders: systematic review. *J Med Internet Res*. 2018, 20:e157. [10.2196/jmir.7898](https://doi.org/10.2196/jmir.7898)
  47. Phelan S, Peruvemba S, Levinson D, Stulberg N, Lacy A, Legato M, Werner JP: Feasibility of a virtual reality-



- based approach to improve behavioral weight management outcomes. *Pilot Feasibility Stud.* 2021, 7:129. [10.1186/s40814-021-00865-5](https://doi.org/10.1186/s40814-021-00865-5)
48. Lydecker JA, White MA, Grilo CM: Form and formulation: examining the distinctiveness of body image constructs in treatment-seeking patients with binge-eating disorder. *J Consult Clin Psychol.* 2017, 85:1095-105. [10.1037/ccp0000258](https://doi.org/10.1037/ccp0000258)
  49. Libeton M, Dixon JB, Laurie C, O'Brien PE: Patient motivation for bariatric surgery: characteristics and impact on outcomes. *Obes Surg.* 2004, 14:392-8. [10.1381/096089204322917936](https://doi.org/10.1381/096089204322917936)
  50. Bertoletti J, Galvis Aparicio MJ, Bordignon S, Trentini CM: Body image and bariatric surgery: a systematic review of literature. *Bariatr Surg Pract Patient Care.* 2018, 14:81-92. [10.1089/bari.2018.0036](https://doi.org/10.1089/bari.2018.0036)
  51. Kitzinger HB, Abayev S, Pittermann A, et al.: After massive weight loss: patients' expectations of body contouring surgery. *Obes Surg.* 2012, 22:544-8. [10.1007/s11695-011-0551-6](https://doi.org/10.1007/s11695-011-0551-6)
  52. Porras Garcia B, Ferrer Garcia M, Olszewska A, et al.: Is this my own body? Changing the perceptual and affective body image experience among college students using a new virtual reality embodiment-based technique. *J Clin Med.* 2019, 8:925. [10.3390/jcm8070925](https://doi.org/10.3390/jcm8070925)
  53. Horne M, Hill A, Murells T, Ugail H, Irving, Chinnadorai R, Hardy M: Using avatars in weight management settings: a systematic review. *Internet Interv.* 2020, 19:100295. [10.1016/j.invent.2019.100295](https://doi.org/10.1016/j.invent.2019.100295)
  54. Purvis CK, Jones M, Bailey JO, Bailenson J, Taylor CB: Developing a novel measure of body satisfaction using virtual reality. *PLoS One.* 2015, 10:e0140158. [10.1371/journal.pone.0140158](https://doi.org/10.1371/journal.pone.0140158)
  55. Gupta PK, Franck C, Miller WJ, Gupta H, Forse RA: Development and validation of a bariatric surgery morbidity risk calculator using the prospective, multicenter NSQIP dataset. *J Am Coll Surg.* 2011, 212:301-9. [10.1016/j.jamcollsurg.2010.11.003](https://doi.org/10.1016/j.jamcollsurg.2010.11.003)
  56. Lemanu DP, Srinivasa S, Singh PP, Johannsen S, MacCormick AD, Hill AG: Optimizing perioperative care in bariatric surgery patients. *Obes Surg.* 2012, 22:979-90. [10.1007/s11695-012-0648-6](https://doi.org/10.1007/s11695-012-0648-6)
  57. Trahan MH, Maynard BR, Smith KS, Farina ASJ, Khoo YM: Virtual reality exposure therapy on alcohol and nicotine: a systematic review. *Res Soc Work Pract.* 2019, 29:876-91. [10.1177/1049731518823073](https://doi.org/10.1177/1049731518823073)
  58. Kaganoff E, Bordnick PS, Carter BL: Feasibility of using virtual reality to assess nicotine cue reactivity during treatment. *Res Soc Work Pract.* 2012, 22:159-65. [10.1177/1049731511428617](https://doi.org/10.1177/1049731511428617)
  59. Nichols S, Patel H: Health and safety implications of virtual reality: a review of empirical evidence. *Appl Ergon.* 2002, 33:251-71. [10.1016/s0003-6870\(02\)00020-0](https://doi.org/10.1016/s0003-6870(02)00020-0)