

RESEARCH ARTICLE

Open Access

# A dose-ranging study of the bronchodilator effects of abediterol (LAS100977), a long-acting $\beta_2$ -adrenergic agonist, in asthma; a Phase II, randomized study

Dave Singh<sup>1\*</sup>, Helena Pujol<sup>2</sup>, Anna Ribera<sup>2</sup>, Beatriz Seoane<sup>2</sup>, Eric Massana<sup>2</sup>, Carol Astbury<sup>2</sup>, Sandrine Ruiz<sup>2</sup> and Gonzalo de Miguel<sup>2</sup>

## Abstract

**Background:** Long-acting  $\beta_2$ -adrenergic agonists (LABAs) are recommended in combination with inhaled corticosteroids (ICSs) for asthma management. Abediterol is a novel, selective, potent, once-daily LABA in development for treatment of asthma and chronic obstructive pulmonary disease. This study aimed to determine abediterol doses with similar peak bronchodilatory effect to salbutamol 400  $\mu\text{g}$ , and duration of action compatible with once-daily dosing in patients with persistent, stable asthma.

**Methods:** This was a Phase II, randomized, double-blind, double-dummy, crossover, placebo-controlled, dose-ranging study (ClinicalTrials.gov NCT01425801) in 62 patients with mild-to-moderate asthma who were also receiving an ICS. Patients received single doses of abediterol 0.313, 0.625, 1.25, or 2.5  $\mu\text{g}$ , salbutamol 400  $\mu\text{g}$ , or placebo in the morning. Spirometry was performed up to 36 h post-dose; safety and tolerability were assessed throughout the study. The primary endpoint was change from baseline in peak forced expiratory volume in 1 s ( $\text{FEV}_1$ ). Additional endpoints included trough  $\text{FEV}_1$ , normalized area under the  $\text{FEV}_1$  curve ( $\text{FEV}_1$  AUC) up to 24 h post-dose, and peak and trough forced vital capacity (FVC).

**Results:** Abediterol produced dose-dependent improvements in peak  $\text{FEV}_1$  from baseline compared with placebo, from 0.274 (95% CI 0.221, 0.327) to 0.405 L (95% CI 0.353, 0.458) for abediterol 0.313 to 2.5  $\mu\text{g}$ , respectively ( $p < 0.0001$  all doses). Abediterol 0.625, 1.25, and 2.5  $\mu\text{g}$  had similar magnitude of peak  $\text{FEV}_1$  effect to salbutamol. Dose-dependent changes from baseline in trough  $\text{FEV}_1$  versus placebo were 0.219 (95% CI 0.136, 0.302) to 0.400 L (95% CI 0.317, 0.483) for abediterol 0.313 to 2.5  $\mu\text{g}$ , respectively ( $p < 0.0001$ ). All abediterol doses achieved significant improvements versus placebo in  $\text{FEV}_1$  AUC 0–6, 0–12, and 0–24 h, and peak and trough FVC ( $p < 0.05$ ). Less than 10% of patients experienced treatment-related adverse events for each dose of abediterol; most were mild to moderate in intensity and the most common were headache and nasopharyngitis. There were no clinically relevant changes in heart rate.

**Conclusions:** Abediterol 0.625–2.5  $\mu\text{g}$  provided dose-dependent, clinically and statistically significant bronchodilation versus placebo in patients with asthma, with a peak effect similar to salbutamol and duration of action compatible with once-daily dosing. All doses of abediterol were well tolerated.

**Keywords:** LABA, Chronic respiratory disease, Asthma, Dose-finding, Bronchodilation

\* Correspondence: [dsingh@meu.org.uk](mailto:dsingh@meu.org.uk)

<sup>1</sup>University of Manchester, Medicines Evaluation Unit Ltd, The Langley Building, Southmoor Road, Wythenshawe, Manchester M23 9QZ, UK  
Full list of author information is available at the end of the article

## Background

The use of  $\beta_2$ -adrenergic agonist bronchodilators in combination with inhaled corticosteroids (ICSs) to manage the symptoms of asthma is recommended by Global Initiative for Asthma (GINA) guidelines [1]. Currently licensed bronchodilators include salbutamol, a short-acting  $\beta_2$ -adrenergic agonist (SABA) administered as required [2], salmeterol and formoterol, two long-acting  $\beta_2$ -adrenergic agonists (LABAs) administered twice daily in combination with an ICS [3,4], and the recently licensed vilanterol, a LABA administered once daily in fixed-dose combination with the ICS fluticasone furoate [5]. There is ongoing interest to develop once-daily medications to further simplify treatment regimens and improve patient compliance [6].

Abediterol is a novel, once-daily LABA in clinical development as a fixed-dose combination with an ICS for the treatment of asthma and chronic obstructive pulmonary disease (COPD) [7]. Abediterol displays high affinity for  $\beta_2$ -adrenoceptors, plus a higher functional selectivity for  $\beta_2$ -adrenoceptors over  $\beta_1$ -adrenoceptors than formoterol, indacaterol, vilanterol and olodaterol in cellular models [7-9]. The potency of abediterol is also higher than salmeterol and indacaterol in isolated human bronchi and, in addition to these two LABAs, vilanterol and olodaterol in animal models [7,10]. Abediterol has a rapid onset and prolonged duration of action [7,10].

A Phase II clinical trial in patients with asthma provided results that were consistent with the preclinical data; abediterol resulted in significant improvements in lung function at 5 min post-dose that were greater than improvements seen with salmeterol, and significant bronchodilation was sustained up to 24 h post-dose [11]. These characteristics make abediterol an ideal candidate for once-daily dosing. Moreover, abediterol has a favorable cardiovascular safety and tolerability profile: in anesthetized dogs, abediterol had a lesser effect on heart rate compared with salmeterol, formoterol, and indacaterol at concentrations that produced comparable bronchodilation [7,12]. In early phase clinical trials, doses of abediterol  $\leq 10$   $\mu\text{g}$  were safe and well tolerated and did not demonstrate any clinically relevant effects on heart rate in healthy subjects [13], patients with asthma [14], or patients with COPD [15].

Drug regulatory authorities have concerns that excessively high LABA doses may cause unacceptable side effects [16]. The trough forced expiratory volume in 1 s ( $\text{FEV}_1$ ), measured 24 h after dosing, is an important endpoint to assess once-daily LABA efficacy [16]. The maximum effect achieved soon after dosing, otherwise known as the 'peak' effect, is also of interest for efficacy and safety reasons. A very high peak effect may indicate high levels of acute drug exposure that could predispose patients to  $\beta$ -agonist-mediated side effects, such as

cardiovascular adverse events, when receiving chronic treatment [17]. Moreover, the ideal pharmacologic characteristic of a once-daily LABA is a low 'peak to trough ratio', ensuring relatively stable lung function throughout the day [18].

The peak effect of salbutamol is well characterized, and causes acute bronchodilation without safety concerns. The bronchodilator response following single-dose administration of salbutamol (4 x 100  $\mu\text{g}$ ) is commonly used as one of the diagnostic tests for asthma, as recommended by a joint European Respiratory Society (ERS)/American Thoracic Society (ATS) taskforce [19]. This effect is, therefore, a reasonable benchmark for a novel LABA to achieve and this degree of acute bronchodilation is considered clinically meaningful. A major regulatory agency recommended that the peak effects of abediterol should be compared to those of salbutamol to guide dose selection for further clinical trials in order to avoid a dose that induces a much greater degree of bronchodilation than is necessary for clinical effectiveness, which could increase the potential for safety concerns during chronic treatment.

We report a placebo-controlled, dose-ranging, crossover study comparing the acute bronchodilator effects of abediterol to salbutamol in patients with persistent stable asthma. The primary aim of the study was to investigate the peak bronchodilatory effect of abediterol versus placebo and to confirm it was in the same range as that of salbutamol. The full profile of lung function over 24 h, including the trough  $\text{FEV}_1$ , was also measured. The goal of this study was to enable the selection of abediterol doses for future long-term clinical trials, based on a peak effect similar to that of salbutamol 400  $\mu\text{g}$  metered dose inhaler (MDI), coupled with a trough effect compatible with a 24-h duration of action. This trial design was a specific recommendation by a major regulatory agency.

## Methods

### Study design

This was a Phase IIa, randomized, double-blind, double-dummy, 6-way crossover, single-dose, multicenter, dose-ranging study conducted between August 2011 and January 2012. The study was conducted in two sites in the UK and seven in Germany. The study protocol was approved by an independent ethics committee in each country (see Additional file 1) and complied with the Declaration of Helsinki and the International Conference on Harmonisation and Good Clinical Practice guidelines. Patients provided written, informed consent.

Following a screening visit and a 12–16 day run-in period to assess clinical stability, participants were randomized to 1 of 6 treatment sequences according to a balanced Williams design for crossover studies (1:1:1:1:1:1).

Block randomization was performed by the Sponsor, using a computer-generated schedule to assign a treatment sequence to each patient randomization number. The block size and randomization schedule were not communicated to investigators. Patients received abediterol 0.313, 0.625, 1.25, or 2.5 µg, salbutamol 400 µg (administered as four inhalations with 100 µg per inhalation), or placebo (at 09:00 h ±1 h) at each visit, with a washout period of 7–14 days. A follow-up telephone call to assess safety was made approximately 2 weeks following the final treatment visit or premature discontinuation.

Abediterol and placebo to abediterol were administered via a multidose dry powder inhaler (Genuair®). Salbutamol and placebo to salbutamol were administered via a pressurized MDI (Ventolin™ Evohaler™). In order to preserve study blinding, independent, trained personnel prepared the medication, delivered the device to patients, and collected the device following inhalation.

### Patients

Eligible participants were men or women aged 18–70 years with a clinical diagnosis of persistent asthma (according to GINA guidelines) for ≥6 months prior to screening. Eligible patients had an FEV<sub>1</sub> >60% and ≤85% of the predicted normal value, with an FEV<sub>1</sub> reversibility ≥12%, and an absolute increase in FEV<sub>1</sub> of ≥200 mL over their baseline value following inhalation of salbutamol 400 µg at screening. Comparable FEV<sub>1</sub> measurements were required at screening and pre-dose at each visit in order to ensure there were no carry-over effects from previous treatment periods (limit of variability ±20%). At enrollment, all patients were using only an ICS to control their asthma, and had been on a stable dose and regimen for at least 4 weeks prior to screening, up to the equivalent of 1600 µg/day of beclometasone dipropionate, and had not previously been exposed to abediterol.

Exclusion criteria included: a smoking history within the past 6 months or >10 pack-years; presence or history of relevant pulmonary disease or any other clinically relevant disease or abnormality as judged by the investigator; difficult-to-treat asthma; any respiratory tract infection ≤6 weeks prior to screening; hospitalization or emergency room treatment for asthma ≤3 months prior to screening; treatment with β<sub>2</sub>-antagonists; and positive laboratory tests for illicit drugs.

The use of anticholinergics, SABAs (other than salbutamol), LABAs, continuous oral or parenteral corticosteroids, methyl-xanthines, cromolyn sodium, nedocromil, leukotriene modifiers, anti-immunoglobulin E, β<sub>1</sub>-blocking agents, and other investigational drugs was prohibited during the study. Patients using prohibited medications underwent up to 6 weeks' washout before the screening visit. Salbutamol pressurized MDI (100 µg/puff) was the only reliever medication permitted

and was used on an as-needed basis and avoided 6 h prior to and during a treatment visit unless absolutely necessary.

### Efficacy assessments

Pulmonary function tests were conducted at screening, pre-dose, and at 0.25, 0.5, 1, 2, 3, 4, 6, 8, 12, 14, 23, 24, and 36 h post-dose at each treatment visit using standardized spirometers at each investigational site (eResearchTechnology, Philadelphia, PA, USA). Prior to the first spirometry measure, the technician demonstrated the procedure and allowed the patient two practice attempts. At each study time point, the maneuvers were repeated until three technically adequate measurements were made according to ERS/ATS guidelines for acceptability and repeatability [20], with a maximum of eight attempts allowed.

### Safety and tolerability assessments

Treatment-emergent adverse events (TEAEs), including serious adverse events, were recorded throughout the study and follow-up period. A physical examination was performed at screening and at 36 h post-dose following the last treatment period (Visit 6), and blood pressure and 12-lead electrocardiogram measures were made at screening, pre-dose, and at 0.75, 2, 6, 24 h post-dose for each treatment period, and 36 h post-dose at Visit 6. Clinical laboratory tests (blood chemistry, hematology, urinalysis, and a serum pregnancy test) were performed at screening and at 36 h post-dose at Visit 6. Blood glucose

**Table 1 Patient demographics at baseline: safety population**

	Patients (N = 62)
Age, mean no. years (SD)	40.9 (11.2)
Male gender, n (%)	40 (64.5)
Race, n (%)	
White	58 (93.5)
Other	4 (6.5)
Cigarette smoking status, n (%)	
Former smoker	21 (33.9)
Never smoked	41 (66.1)
Smoking history, mean no. pack-years (SD)	4.6 (3.3)
Asthma duration, mean no. years (SD)	25.5 (13.2)
Asthma severity, n (%)	
60% ≤ FEV <sub>1</sub> < 80% of predicted	46 (74.2)
FEV <sub>1</sub> ≥80% of predicted	16 (25.8)
Bronchial reversibility <sup>a</sup> , mean % (SD)	21.3 (8.2)
FEV <sub>1</sub> absolute reversibility <sup>b</sup> , mean L (SD)	0.568 (0.261)

<sup>a</sup>calculated as 100 × (FEV<sub>1</sub> [post-bronchodilator] – FEV<sub>1</sub> [pre-bronchodilator])/FEV<sub>1</sub> [pre-bronchodilator]; <sup>b</sup>calculated as FEV<sub>1</sub> (post-bronchodilator) – FEV<sub>1</sub> (pre-bronchodilator).

FEV<sub>1</sub>, forced expiratory volume in 1 s; SD, standard deviation.

and serum potassium levels were assessed at screening, pre-dose, and at 4, 24 h post-dose for each treatment period, and 36 h post-dose at Visit 6.

### Statistics

The primary efficacy endpoint was change from baseline in peak FEV<sub>1</sub> (defined as the highest FEV<sub>1</sub> value observed within 4 h of the administration of the investigational medicinal product [IMP]) on Day 1. Secondary endpoints included change from baseline in peak forced vital capacity (FVC) on Day 1, trough FEV<sub>1</sub> (defined as the highest average of the maneuvers performed at 23–24 h post IMP administration) and FVC on Day 2, change from baseline in FEV<sub>1</sub> and FVC at each time point up to 36 h post-dose, and change from baseline in normalized FEV<sub>1</sub> area under the curve (AUC; defined as the area between baseline FEV<sub>1</sub> and the FEV<sub>1</sub> curve, from 0 to  $\times$  hours divided by  $\times$  hours) at 0–6, 0–12, and 0–24 h post-dose. In accordance with the GINA guidelines, a change in FEV<sub>1</sub> of 200 mL was considered clinically relevant [1].

A population of 48 patients was necessary to achieve 80% power to detect a 150 mL change from baseline in peak FEV<sub>1</sub> for any dose of abediterol compared with placebo. Assuming a 15% withdrawal rate, the protocol required at least 54 patients to be randomized.

All reported efficacy assessments were analyzed in the intention-to-treat population, defined as all randomized patients who took at least one dose of the IMP (placebo, abediterol, or salbutamol) and had at least one baseline and one post-dose FEV<sub>1</sub> value for at least one treatment visit. Safety analyses are reported in the safety population, which consisted of all randomized patients who took at least one dose of the IMP. All statistical analyses were performed using Statistical Analysis System (SAS Institute Inc., Cary, NC, USA) version 9.1.3 software.

Efficacy variables were analyzed using an analysis of covariance model for crossover designs, with fixed-effect factors of sequence, treatment, and period, patient within sequence as a random effect, and baseline FEV<sub>1</sub> or FVC as a covariate, as appropriate. Between-treatments comparisons were performed by means of contrasts on the treatment factor. The difference between treatments was estimated by the difference between the least squares means. Statistical comparisons between active treatments and placebo were two-sided hypothesis tests with a significance level of 0.05.

## Results

### Study population

A total of 115 patients were screened, with 62 patients randomized (40/62 [64.5%] male, mean age 40.9 ± 11.2 years). Patient demographics are presented in Table 1. A diagram of patient flow is presented in Figure 1. All 62





Figure 1 Diagram of patient flow.

patients were included in the intention-to-treat and safety populations. One protocol violation occurred when a patient was inadvertently given the treatment scheduled for Visit 6 at Visit 5, and vice versa. Data for this patient were analyzed according to the treatments received. Four patients withdrew or were discontinued from the study: two due to inability to comply with the study schedule, one who did not fulfill stability criteria, and one for unknown reasons (patient did not return for the premature discontinuation visit).

### Efficacy

#### Peak FEV<sub>1</sub>

Mean baseline FEV<sub>1</sub> values were similar across all treatment periods, ranging from 2.63 L to 2.69 L. The full profile of lung function after each treatment is shown in Figure 2. The onset of action of abediterol was rapid, with significant bronchodilation compared with placebo observed at the first study time point (15 min) post-dose ( $p < 0.0001$  for all doses of abediterol versus placebo, Figure 2a). All doses of abediterol produced significantly greater improvements in peak FEV<sub>1</sub> from baseline compared with placebo ( $p < 0.0001$  for all doses, Table 2), and the magnitude of effect of abediterol 0.625, 1.25, and 2.5  $\mu\text{g}$  was not significantly different from



**Figure 2 Change from baseline in FEV<sub>1</sub> over time.** Data are presented as least squares means with 95% confidence intervals: **a)**  $p < 0.0001$  versus placebo for salbutamol and all doses of abediterol at all time points; **b)**  $p < 0.0001$  for all doses of abediterol 0–24 h post-dose and abediterol 1.25 and 2.5 µg 36 h post-dose versus placebo,  $p = 0.002$  and  $p = 0.01$  for abediterol 0.313 and 0.625 µg, respectively, at 36 h post-dose versus placebo. FEV<sub>1</sub>, forced expiratory volume in 1 s.

salbutamol 400 µg (Figure 3, Table 3). Least squares means differences in change from baseline in peak FEV<sub>1</sub> versus placebo were 0.274, 0.322, 0.371, and 0.405 L for

abediterol 0.313, 0.625, 1.25, and 2.5 µg, respectively, and 0.353 L for salbutamol 400 µg (Table 3). Median time to peak FEV<sub>1</sub> was 3 h post-dose for abediterol (all



**Table 2 Change from baseline in pulmonary function parameters (L): intention-to-treat population**

	Placebo N = 59	Salbutamol 400 µg N = 58	Abediterol 0.313 µg N = 60	Abediterol 0.625 µg N = 60	Abediterol 1.25 µg N = 60	Abediterol 2.5 µg N = 61
Peak FEV <sub>1</sub>	0.202 (0.045)	0.555**** (0.045)	0.477****†† (0.045)	0.524**** (0.045)	0.573**** (0.045)	0.608**** (0.045)
Trough FEV <sub>1</sub> <sup>a</sup>	-0.054 (0.039)	-0.076 (0.039)	0.166****†††† (0.039)	0.205****†††† (0.039)	0.278****†††† (0.039)	0.346****†††† (0.039)
Normalized FEV <sub>1</sub> AUC <sub>0-6</sub>	0.088 (0.039)	0.347**** (0.039)	0.354**** (0.038)	0.403****† (0.038)	0.461****†††† (0.038)	0.496****†††† (0.038)
Normalized FEV <sub>1</sub> AUC <sub>0-12</sub>	0.063 (0.040)	0.219**** (0.040)	0.348****†††† (0.039)	0.393****†††† (0.039)	0.455****†††† (0.039)	0.496****†††† (0.040)
Normalized FEV <sub>1</sub> AUC <sub>0-24</sub>	0.007 (0.039)	0.100** (0.039)	0.282****†††† (0.039)	0.321****†††† (0.038)	0.390****†††† (0.038)	0.446****†††† (0.039)
Peak FVC	0.222 (0.034)	0.361**** (0.034)	0.317** (0.034)	0.329*** (0.034)	0.361**** (0.034)	0.380**** (0.034)
Trough FVC <sup>a</sup>	-0.028 (0.035)	-0.065 (0.035)	0.094**†††† (0.035)	0.050*†† (0.035)	0.111****†††† (0.035)	0.120****†††† (0.035)

Data are least squares means (standard error).

<sup>a</sup>trough values are the mean of the 23 and 24 h post-dose measurements.

\*p < 0.05, \*\*p < 0.01, \*\*\*p < 0.001, \*\*\*\*p < 0.0001 versus placebo; †p < 0.05, ††p < 0.01, †††p < 0.0001 versus salbutamol 400 µg (analysis of covariance).

AUC, area under the FEV<sub>1</sub> curve; FEV<sub>1</sub>, forced expiratory volume in 1 s; FVC, forced vital capacity.

doses), 2 h post-dose for placebo, and 1 h post-dose for salbutamol 400 µg.

#### Trough FEV<sub>1</sub>

Statistically significant increases in FEV<sub>1</sub> were sustained at 24 h post-dose compared with placebo (p < 0.0001 for all doses of abediterol 0–24 h post-dose, Figure 2b). Abediterol significantly improved trough FEV<sub>1</sub> from baseline compared with placebo (p < 0.0001 for all doses; Figure 4). Least squares means differences in change from baseline in trough FEV<sub>1</sub> versus placebo were 0.219 L (95% CI 0.136, 0.302), 0.259 L (95% CI 0.176, 0.342), 0.332 L (95% CI 0.249, 0.415), and 0.400 L (95% CI 0.317, 0.483) for abediterol 0.313, 0.625, 1.25, and 2.5 µg, respectively (Table 2). The effects of salbutamol on FEV<sub>1</sub> were not sustained after 6 h post-dose, as expected for a SABA (Figure 2b).

#### FEV<sub>1</sub> AUC

Change from baseline in normalized FEV<sub>1</sub> AUC<sub>0-6</sub>, FEV<sub>1</sub> AUC<sub>0-12</sub>, and FEV<sub>1</sub> AUC<sub>0-24</sub> was significantly higher for all doses of abediterol versus placebo (p < 0.0001 for all, Table 2), with dose-related effects observed. The change from baseline in normalized FEV<sub>1</sub> AUC<sub>0-6</sub> was significantly greater for abediterol 0.625–2.5 µg versus salbutamol 400 µg (p < 0.05, Table 2).

#### FVC

Mean baseline FVC values were comparable across treatment periods, ranging from 4.23 L to 4.30 L. Abediterol produced significantly greater change from baseline in


peak FVC compared with placebo (p < 0.01 for all doses, Table 2). The change from baseline in peak FVC was similar to that observed with salbutamol. Additionally, abediterol significantly improved trough FVC from baseline compared with placebo (p < 0.05 for all doses, Table 2). All doses of abediterol significantly (p < 0.05) increased FVC versus placebo up to 24 h post-dose except for the 0.625 µg dose at 12 and 23 h.

#### Safety and tolerability

The percentage of patients experiencing TEAEs was similar across all treatment groups (Table 4). Most TEAEs were mild to moderate in intensity, no patient discontinued the study due to a TEAE, and there were no serious adverse events or deaths. The most common TEAEs were headache and nasopharyngitis, reported by 41.9 and 27.4% of patients, respectively. The percentage of patients that experienced at least one treatment-related TEAE was <10% for all doses of abediterol, and was similar to that of salbutamol and placebo. No clinically significant changes in serum glucose or potassium concentrations were observed over time in any of the treatments groups (Table 5). Minor variations in heart rate over time were observed for all treatment groups, including placebo; none were considered to be dose related or clinically relevant (Table 5). There were no clinically relevant changes in clinical laboratory tests, blood pressure, or electrocardiogram parameters.

#### Discussion

In this study, abediterol 0.313, 0.625, 1.25, and 2.5 µg provided clinically and statistically significant improvements



**Figure 3 Change from baseline in peak FEV<sub>1</sub> at Day 1.** Data are presented as least squares means ± standard error; p < 0.0001 versus placebo for all doses of abediterol and salbutamol 400 µg; \*\*p < 0.01, \*\*\*p < 0.001, \*\*\*\*p < 0.0001 (analysis of covariance). FEV<sub>1</sub>, forced expiratory volume in 1 s.

in peak bronchodilation versus placebo in patients with persistent, stable asthma. Abediterol doses  $\geq 0.625$  µg also caused bronchodilation compatible with once a day administration, as improvements in trough FEV<sub>1</sub> >250 mL compared with placebo were observed. The peak bronchodilatory effect of abediterol was similar to that of salbutamol 400 µg for all doses  $\geq 0.625$  µg.

A mean increase in peak FEV<sub>1</sub> >200 mL is considered clinically relevant in asthma [1]. This magnitude of change was exceeded for all doses of abediterol, including the lowest dose, 0.313 µg. The primary objective of this study was to define abediterol doses that caused peak bronchodilation similar to salbutamol, the most widely prescribed SABA [21]. High doses of  $\beta_2$ -agonists


may increase the degree of bronchodilation but also can cause excessive systemic side effects through systemic absorption [22]. However, the current study demonstrates that the doses of abediterol from 0.625 to 2.5 µg induce acute bronchodilation similar to salbutamol, and therefore treatment with higher doses may not be warranted.

Significant bronchodilation compared with placebo was achieved with a range of abediterol doses from 0.313 to 2.5 µg, consistent with the high potency of abediterol demonstrated in preclinical studies that suggested a potential for significant bronchodilation *in vivo* [7]. Abediterol has higher potency than formoterol, salmeterol, indacaterol, olodaterol, and vilanterol in

**Table 3 Comparison of peak FEV<sub>1</sub> for abediterol versus placebo and abediterol versus salbutamol**

	Least-squares mean difference (standard error)	95% confidence interval
Abediterol 2.5 µg vs. placebo	0.405 (0.027)	0.353, 0.458
Abediterol 1.25 µg vs. placebo	0.371 (0.027)	0.318, 0.424
Abediterol 0.625 µg vs. placebo	0.322 (0.027)	0.269, 0.375
Abediterol 0.313 µg vs. placebo	0.274 (0.027)	0.221, 0.327
Salbutamol 400 µg vs. placebo	0.353 (0.027)	0.299, 0.406
Abediterol 2.5 µg vs. salbutamol 400 µg	0.0529 (0.0270)	-0.0002, 0.1061
Abediterol 1.25 µg vs. salbutamol 400 µg	0.0181 (0.0269)	-0.0350, 0.0711
Abediterol 0.625 µg vs. salbutamol 400 µg	-0.0308 (0.0269)	-0.839, 0.0223
Abediterol 0.313 µg vs. salbutamol 400 µg	-0.0783 (0.0269)	-0.1313, -0.0253

Data included for comparisons of all doses of abediterol with placebo and with salbutamol.



**Figure 4 Change from baseline in trough FEV<sub>1</sub> at Day 2.** Data are presented as least squares means ± standard error: p < 0.0001 versus placebo for all doses of abediterol; \*\*p < 0.01, \*\*\*p < 0.001, \*\*\*\*p < 0.0001 (analysis of covariance). FEV<sub>1</sub>, forced expiratory volume in 1 s.

anesthetized guinea pigs [7,10] as well as higher potency than salmeterol and indacaterol with comparable potency to formoterol in isolated human bronchi [7]. Based on the results of this preliminary dose-ranging study and the guidelines published by Chowdhury et al. [16], the clinically effective dose in asthma is considered most likely to be in the range of 0.625–2.5 µg.

Significant bronchodilation compared with placebo was observed for salbutamol and all doses of abediterol at the first-assessed time point (15 min) post-dose; this indicates a rapid onset of action for both salbutamol and abediterol. Salmeterol and formoterol, both twice-daily LABAs, are widely used as maintenance therapy in COPD and asthma due to their longer durations of

action compared with salbutamol [6,18]. In clinical studies in patients with asthma, the onset of action of formoterol 24 µg is consistently faster than that of salmeterol 50 µg, with a median onset of action of 3.6 min compared with 31.0 min in one study [23,24]. *In vitro* experiments in isolated human bronchi suggest that the onset of action of abediterol is faster than salmeterol and is not significantly different from formoterol and indacaterol [7].

The prolonged duration of action of abediterol, with significant improvements in trough FEV<sub>1</sub>, normalized FEV<sub>1</sub> AUC<sub>0–24h</sub>, and trough FVC relative to placebo, and significant bronchodilation maintained up to 36 h post-dose is consistent with the results of previous clinical studies in healthy subjects and patients with COPD or

**Table 4 TEAEs occurring in ≥2% of patients in any treatment group: safety population**

	Number (%) of patients reporting TEAEs						Total N = 62
	Placebo N = 59	Salbutamol 400 µg N = 58	Abediterol 0.313 µg N = 60	Abediterol 0.625 µg N = 60	Abediterol 1.25 µg N = 60	Abediterol 2.5 µg N = 61	
Any TEAE	16 (27.1)	12 (20.7)	18 (30.0)	16 (26.7)	18 (30.0)	23 (37.7)	51 (82.3)
Headache	7 (11.9)	6 (10.3)	4 (6.7)	7 (11.7)	8 (13.3)	7 (11.5)	26 (41.9)
Nasopharyngitis	1 (1.7)	0	8 (13.3)	3 (5.0)	4 (6.7)	2 (3.3)	17 (27.4)
Chest discomfort	2 (3.4)	0	1 (1.7)	0	0	2 (3.3)	5 (8.1)
Wheezing	1 (1.7)	2 (3.4)	1 (1.7)	0	1 (1.7)	0	4 (6.5)
Oropharyngeal pain	0	1 (1.7)	0	1 (1.7)	0	3 (4.9)	4 (6.5)
Cough	2 (3.4)	0	0	1 (1.7)	0	0	3 (4.8)

TEAE, treatment-emergent adverse event.



**Table 5 Change in clinical safety parameters from pre-dose values following treatment: safety population**

	Placebo N = 59	Salbutamol 400 µg N = 58	Abediterol 0.313 µg N = 60	Abediterol 0.625 µg N = 60	Abediterol 1.25 µg N = 60	Abediterol 2.5 µg N = 61
Serum glucose (mmol/L)						
4 hours post-dose	-0.37 (0.56)	-0.24 (0.55)	-0.19 (0.71)	-0.27 (0.70)	-0.16 (0.94)	-0.23 (0.74)
24 hours post-dose	0.01 (0.38)	0.00 (0.33)	0.13 (0.51)	0.06 (0.62)	0.11 (0.45)	0.15 (0.59)
Serum potassium (mmol/L)						
4 hours post-dose	-0.01 (0.36)	-0.09 (0.38)	-0.01 (0.35)	0.10 (0.37)	0.02 (0.36)	-0.04 (0.35)
24 hours post-dose	-0.02 (0.37)	0.04 (0.34)	0.05 (0.33)	0.15 (0.32)	0.04 (0.28)	0.05 (0.38)
Heart rate (bpm)						
0.75 h post-dose	-0.6 (9.1)	2.0 (10.1)	-0.7 (6.1)	-0.8 (7.5)	-1.5 (9.0)	-0.8 (6.2)
2 h post-dose	-3.4 (8.7)	-1.0 (8.0)	-2.0 (6.6)	-3.4 (8.1)	-4.3 (8.4)	-3.2 (6.7)
6 h post-dose	5.1 (10.2)	7.2 (7.7)	6.2 (7.1)	6.2 (8.0)	6.4 (9.8)	5.3 (7.2)
24 h post-dose	-2.1 (9.8)	-1.9 (8.0)	-1.6 (6.7)	-0.8 (8.4)	-0.9 (8.7)	-0.6 (7.6)

Data are mean (standard deviation).  
 bpm, beats per minute.

asthma [11,13,25]. For example, a recent Phase II clinical study in patients with COPD demonstrated that abediterol 2.5, 5, and 10 µg achieved >200 mL increases in FEV<sub>1</sub> at 24 h, and that this bronchodilatory effect was greater than that of indacaterol 150 µg, and the duration of action was similar [25]. Taken together, these results are compatible with a once-daily dosing regimen in asthma and COPD.

The convenience of once-daily dosing is likely to improve compliance in patients with asthma, leading to better patient outcomes [26,27], as shown in various studies, including that of Wells et al. whereby once-daily ICS led to improved compliance in patients with asthma compared with ICS administered multiple times daily [28]. LABA monotherapy treatment is contra-indicated in asthma; co-administration with an ICS is mandated, preferably as a fixed-dose combination to reduce the risk of monotherapy administration. Development of once-daily LABAs for use in fixed-dose combinations with once-daily ICS is important for simplifying treatment regimens and, therefore, helping improve patient compliance and general asthma management.

The safety and tolerability profile of abediterol in this study is comparable to that reported in other studies in the abediterol development program [11-13,25]. Single doses of abediterol 0.313–2.5 µg were safe and well tolerated, and the TEAEs were consistent with the known safety profile of β<sub>2</sub>-agonists.

## Conclusion

Abediterol 0.625–2.5 µg demonstrated a peak bronchodilatory response similar to that of a high dose of salbutamol (400 µg) at very low doses, and showed a clear

dose–response relationship. Abediterol is a promising new, highly potent, LABA with a sustained duration of action and a favorable safety and tolerability profile. These results suggest that further studies are warranted to further evaluate the efficacy and safety of abediterol in combination with ICS in both asthma and COPD.

## Additional file

**Additional file 1: Independent Ethics Committees (IEC).**

### Abbreviations

ATS: American Thoracic Society; COPD: Chronic obstructive pulmonary disease; ERS: European Respiratory Society; FEV<sub>1</sub> AUC: Area under the FEV<sub>1</sub> curve; FEV<sub>1</sub>: Forced expiratory volume in 1 s; FVC: Forced vital capacity; GINA: Global Initiative for Asthma; ICS: Inhaled corticosteroid; IMP: Investigational medicinal product; LABA: Long-acting β<sub>2</sub>-adrenergic agonist; MDI: Metered dose inhaler; SABA: Short-acting β<sub>2</sub>-adrenergic agonist; TEAE: Treatment-emergent adverse event.

### Competing interests

DS has received sponsorship to attend international meetings, honoraria for lecturing or attending advisory boards and research grants from various pharmaceutical companies including Almirall, AstraZeneca, Boehringer Ingelheim, Chiesi, CIPLA, Forest, Genetech, GlaxoSmithKline, Merck, Novartis, Pfizer and Takeda. HP, AR, BS, EM, CA, SR and GdM are all employees of Almirall. BS, SR and GdM have a patent pending for abediterol (Novel dosage form and formulation of abediterol, WO2013178742 A1). HP, AR, BS and GdM hold Almirall shares or stocks.

### Authors' contributions

All authors contributed to the conception and design of the study, data analysis/interpretation and revision of the manuscript for important intellectual content and provided final approval of the manuscript. DS was the coordinating investigator of the study and drafted the article. There are no other contributors who fulfill the authorship criteria.

### Acknowledgments

The authors wish to thank the study investigators and PAREXEL International Ltd. Suzanne McAllister, PhD, from Complete Medical Communications provided medical writing assistance funded by Almirall S.A., Barcelona, Spain.

This study is registered with clinicaltrials.gov, trial registration number NCT01425801, and was funded by Almirall S.A., Barcelona, Spain.

#### Author details

<sup>1</sup>University of Manchester, Medicines Evaluation Unit Ltd, The Langley Building, Southmoor Road, Wythenshawe, Manchester M23 9QZ, UK. <sup>2</sup>Almirall Research and Development Centre, Barcelona, Spain.

Received: 16 April 2014 Accepted: 23 October 2014

Published: 14 November 2014

#### References

1. Global Strategy for Asthma Management and Prevention. In [www.ginasthma.org]
2. Ventolin Evohaler Summary of Product Characteristics. In [http://www.medicines.org.uk/emc/medicine/97/SPC]
3. Serevent Accuhaler Summary of Product Characteristics. In [http://www.medicines.org.uk/emc/medicine/91]
4. Formoterol Easyhaler Summary of Product Characteristics. In [http://www.medicines.org.uk/emc/medicine/19503/SPC]
5. Relvar Ellipta Summary of Product Characteristics. In [www.medicines.org.uk/emc/medicine/28496/SPC]
6. Cazzola M, Matera MG: Novel long-acting bronchodilators for COPD and asthma. *Br J Pharmacol* 2008, **155**:291–299.
7. Aparici M, Gómez-Angelats M, Vilella D, Otal R, Carcasona C, Viñals M, Ramos I, Gavalda A, De Alba J, Gras J, Cortijo J, Morcillo E, Puig C, Ryder H, Beleta J, Miralpeix M: Pharmacological characterization of abediterol, a novel inhaled  $\beta_2$ -adrenoceptor agonist with long duration of action and a favorable safety profile in preclinical models. *J Pharmacol Exp Ther* 2012, **342**:497–509.
8. Aparici M, Gómez-Angelats M, Vilella D, Cortijo J, Morcillo EJ, Carcasona C, Gavalda A, Beleta J, Puig C, Ryder H, Miralpeix M: The *in vitro* pharmacological profile of LAS100977 – a potent, selective and long-acting beta-2 receptor agonist [abstract A5675]. *Am J Respir Crit Care Med* 2010, **181**:A5675.
9. Miralpeix M, Aparici M, Carcasona C, Ramos I, Puig C, Vilella D, Gavalda A: Abediterol (LAS100977) is A Potent and Selective  $\beta_2$ -Agonist with Sustained Duration of Action and Favourable Selectivity *in Vitro* [poster]. In San Diego, CA, USA: Presented at the American Thoracic Society Annual Congress; 2014.
10. De Alba J, Otal R, Fernández-Blanco JA, Aparici M, Puig C, Gavalda A, Miralpeix M: Abediterol (LAS100977) Demonstrates a Higher Potency and a Longer Duration of Action in Comparison with Other Long-Acting  $\beta_2$ -Adrenoceptor Agonists in a Guinea Pig Bronchoconstriction Model [poster]. In San Diego, CA, USA: Presented at the American Thoracic Society Annual Congress; 2014.
11. Beier J, Fuhr R, Massana E, Jiménez E, Seoane B, de Miquel G, Ruiz S: Efficacy and safety of single inhaled doses of LAS100977, a novel long acting  $\beta_2$ -agonist, in patients with persistent asthma [abstract A5414]. *Am J Respir Crit Care Med* 2010, **181**:A5414.
12. Miralpeix M, Gomez-Angelats M, Aparici M, Vinals M, Beleta J, Gavalda A: LAS100977, a novel beta-2 receptor agonist, with a longer duration of action and more favourable safety margin than salmeterol in anesthetised dogs [abstract A5676]. *Am J Respir Crit Care Med* 2010, **181**:A5676.
13. Timmer W, Massana E, Jiménez E, Seoane B, de Miquel G, Ruiz S: Single doses of LAS100977, a novel long acting  $\beta_2$ -agonist, show high activity and long duration in healthy subjects [abstract A5663]. *Am J Respir Crit Care Med* 2010, **181**:A5663.
14. Beier J, Fuhr R, Massana E, Jiménez E, Seoane B, de Miquel G, Ruiz S: Efficacy and Safety of Single Inhaled Doses of LAS100977, A Novel Long Acting  $\beta_2$ -Agonist, in Patients with Persistent Asthma [poster]. In New Orleans, Louisiana, USA: Presented at the American Thoracic Society International Conference; 2010.
15. Beier J, Pujol H, Seoane B, Jiménez E, Astbury C, Massana E, Ruiz S, de Miquel G: Efficacy and Safety of Abediterol (LAS100977) in Patients with COPD: Phase III, Randomized, Crossover Study [poster]. In Barcelona, Spain: Presented at the European Respiratory Society Annual Congress; 2013.
16. Chowdhury B, Seymour SM, Michele TM, Durmowicz AG, Liu D, Rosebraugh CJ: The risks and benefits of indacaterol—the FDA's review. *N Engl J Med* 2011, **365**:2247–2249.
17. Whale CI, Sovani MP, Mortimer KJ, Harrison TW, Tattersfield AE: Systemic and bronchodilator effects of inhaled rac-formoterol in subjects with chronic obstructive pulmonary disease: a dose–response study. *Br J Clin Pharmacol* 2008, **65**:841–847.
18. Beeh KM, Beier J: The short, the long and the “ultra-long”: why duration of bronchodilator action matters in chronic obstructive pulmonary disease. *Adv Ther* 2010, **27**:150–159.
19. Pellegrino R, Viegi G, Brusasco V, Crapo RO, Burgos F, Casaburi R, Coates A, van der Grinten CP, Gustafsson P, Hankinson J, Jensen R, Johnson DC, MacIntyre N, McKay R, Miller MR, Navajas D, Pedersen OF, Wanger J: Interpretative strategies for lung function tests. *Eur Respir J* 2005, **26**:948–968.
20. Quanjer PH, Tammeling GJ, Cotes JE, Pedersen OF, Peslin R, Yernault JC: Lung volumes and forced ventilatory flows. Report Working Party Standardization of Lung Function Tests, European Community for Steel and Coal. Official Statement of the European Respiratory Society. *Eur Respir J Suppl* 1993, **16**:5–40.
21. Crompton G: A brief history of inhaled asthma therapy over the last fifty years. *Prim Care Respir J* 2006, **15**:326–331.
22. Cazzola M, Page CP, Calzetta L, Matera MG: Pharmacology and therapeutics of bronchodilators. *Pharmacol Rev* 2012, **64**:450–504.
23. Palmqvist M, Persson G, Lazer L, Rosenborg J, Larsson P, Lötvall J: Inhaled dry-powder formoterol and salmeterol in asthmatic patients: onset of action, duration of effect and potency. *Eur Respir J* 1997, **10**:2484–2489.
24. Bodzenta-Lukaszyk A, Dymek A, McAulay K, Mansikka H: Fluticasone/formoterol combination therapy is as effective as fluticasone/salmeterol in the treatment of asthma, but has a more rapid onset of action: an open-label, randomized study. *BMC Pulm Med* 2011, **11**:28.
25. Beier J, Pujol H, Seoane B, Jiménez E, Astbury C, Massana E, Ruiz S, de Miquel G: Efficacy and safety of abediterol (LAS100977) in patients with COPD: Phase II, randomized, crossover study [abstract 83]. *Eur Respir J* 2013, **42**(Suppl 57):183.
26. Chapman KR, Walker L, Cluley S, Fabbri L: Improving patient compliance with asthma therapy. *Respir Med* 2000, **94**:2–9.
27. Makela MJ, Backer V, Hedegaard M, Larsson K: Adherence to inhaled therapies, health outcomes and costs in patients with asthma and COPD. *Respir Med* 2013, **107**:1481–1490.
28. Wells KE, Peterson EL, Ahmedani BK, Williams LK: Real-world effects of once vs greater daily inhaled corticosteroid dosing on medication adherence. *Ann Allergy Asthma Immunol* 2013, **111**:216–220.

doi:10.1186/1471-2466-14-176

Cite this article as: Singh et al.: A dose-ranging study of the bronchodilator effects of abediterol (LAS100977), a long-acting  $\beta_2$ -adrenergic agonist, in asthma; a Phase II, randomized study. *BMC Pulmonary Medicine* 2014 **14**:176.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
www.biomedcentral.com/submit

