

RESEARCH ARTICLE

Open Access

An agonist sensitive, quick and simple cell-based signaling assay for determination of ligands mimicking prostaglandin E₂ or E₁ activity through subtype EP₁ receptor: Suitable for high throughput screening

Annirudha J Chillar¹, Parastoo Karimi¹, Kathy Tang², Ke-He Ruan^{1*}

Abstract

Background: Conventionally the active ingredients in herbal extracts are separated into individual components, by fractionation, desalting, and followed by high-performance liquid chromatography (HPLC). In this study we have tried to directly screen water-soluble fractions of herbs with potential active ingredients before purification or extraction. We propose that the herbal extracts mimicking prostaglandin E₁ (PGE₁) and E₂ (PGE₂) can be identified in the water-soluble non-purified fraction. PGE₁ is a potent anti-inflammatory molecule used for treating peripheral vascular diseases while PGE₂ is an inflammatory molecule.

Methods: We used cell-based assays (CytoFluor multi-well plate reader and fluorescence microscopy) in which a calcium signal was generated by the recombinant EP₁ receptor stably expressed in HEK293 cells (human embryonic kidney). PGE₁ and PGE₂ were tested for their ability to generate a calcium signal. Ninety-six water soluble fractions of Treasures of the east (single Chinese herb dietary supplements) were screened.

Results: After screening, the top ten stimulators were identified. The identified herbs were then desalted and the calcium fluorescent signal reconfirmed using fluorescence microscopy. Among these top ten agonists identified, seven stimulated the calcium signaling (1-40 μM concentration) using fluorescence microscopy.

Conclusions: Fluorescence microscopy and multi-well plate readers can be used as a target specific method for screening water soluble fractions with active ingredients at a very early stage, before purification. Our future work consists of purifying and separating the active ingredients and repeating fluorescence microscopy. Under ordinary circumstances we would have to purify the compounds first and then test all the extracts from 96 herbs. Conventionally, for screening natural product libraries, the procedure followed is the automated separation of all constituents into individual components using fractionation and high performance liquid chromatography. We, however, demonstrated that the active ingredients of the herbal extracts can be tested before purification using an agonist sensitive, quick and simple cell-based signaling assay for ligands mimicking the agonists, PGE₁ and PGE₂.

* Correspondence: khruan@uh.edu

¹Centre for Experimental Therapeutics and Pharmacoinformatics, Department of Pharmacological and Pharmaceutical Sciences, College of Pharmacy, University of Houston, Houston, Texas 77004 USA

Full list of author information is available at the end of the article

Background

The subtype receptors of prostaglandin E (PGE) isomer-1 (PGE₁) and isomer-2 (PGE₂) (termed EP₁, EP₂, EP₃ and EP₄) are widely distributed and have been extensively studied for their involvement in a variety of cancers and stem cell differentiation. Expression of EP₁ is frequently seen in human breast cancers and colon tumor cells. Nuclear expression of EP₁ in human breast cancers correlates with good prognosis [1,2]. EP (1-4) subtype receptor mRNAs are generally positively correlated to both COX-1 and COX-2 in tumor tissue, but not in normal colon [3]. Various studies have shown the involvement of PGE₂ via its EP receptors in growth, differentiation and metastasis of cancer; however, there are no therapeutic ligands available for these receptors [4-7]. Nevertheless, PGE₁ has been shown to have anti-inflammatory properties as compared to PGE₂, which is a pro-inflammatory mediator. PGE₁ has been used therapeutically in peripheral vascular diseases [8-10], and its importance as a potential ligand in cancer cannot be overlooked. We chose to test our herbal extracts on the EP₁ subtype receptor since it couples to calcium, which can be used for detecting the stimulation and inhibition of the receptor.

Conventionally, for screening natural product libraries, the procedure followed is the automated separation of all constituents into individual components. This is achieved by fractionation of crude extracts from natural materials using desalting followed by high-performance liquid chromatography (HPLC). Subsequently, full spectroscopic identification is carried out prior to high throughput screening (HTS) [11]. This process generates molecules or lead compounds responsible for the biological activity found in analyzed extracts. These identified relevant molecules are further employed in pre-clinical studies [12]. The structural elucidation is achieved by mass spectrometry and multi-dimensional nuclear magnetic resonance (NMR) spectroscopy, and followed by the generation of analogues [13].

We, however, propose that the water-soluble and desalted-soluble fractions of single plant extracts (Treasures of East Single Herbs, Blue Light Inc.) can be used for tentatively screening potentially active compounds. These water-soluble fractions can be microscopically evaluated on specific targets using cell-based assays. The crude soluble fractions were manually evaluated on the EP₁ subtype receptor for PGE₁- and PGE₂- mimicking extracts which could act as agonists. Specifically, this was achieved by detecting an increase or inhibition of a calcium signal. This receptor (EP₁) is involved in various cancers and stem cell differentiation [1,2,14]. The receptor activation signal was microscopically detected as fluorescence using Fluo8-AM dye in HEK293 cells stably

expressing the recombinant EP₁. This obviates the need for separation and purification of extracts before further testing, and can be accomplished with amazing reliability in a much shorter time. In fact, this separation and purification technique would be a very target-specific, easy alternative to identify the herbal extract of interest, which can be later purified by conventional processes and evaluated again. Potential ligands derived from these extracts could also be used in cancer therapeutics and stem cell proliferation and differentiation in the future.

Methods

The herbal extract granules, taken from Traditional Chinese Medicine (TCM) herbs, were purchased from Tianjiang Pharmaceutical Co., LTD, China (distributed by Blue Light Inc. Ithaca New York). PGE₁ and PGE₂ were purchased from Cayman chemicals (Ann Arbor, MI), and the Fluo8-AM was from ABD-Bioquest (Sunnyvale, CA).

Preparation of Drug Library

The crude drug (2.5 g) was dissolved in hot distilled water (10 mL). The contents were dissolved by repeated vortexing. After centrifugation, the soluble supernatants (5 μ L) from 96 individual fractions (water-soluble drug library) were tested in 96-well plates containing HEK293 cells stably expressing the recombinant EP₁ and cell assays were performed using the CytoFluor Multi-well plate reader. The stimulators of a calcium signal were identified as 'hits'. For confirming the hit compounds, the remaining supernatants (from original 10 mL) were filtered through a sephadex C18 column to desalt the samples. The acetone-eluted compounds were dried, weighed and dissolved in distilled water (0.2 mL) and distributed into 96-well plates, in which serial one-to-one dilutions were performed (using distilled water) until obtaining nearly colorless solutions (desalted drug library) for further fluorescence microscopy testing using these varied concentrations (1-40 μ M) of crude extract. LC/MS analysis for the herbal extracts revealed that the majority of molecular masses of the compounds in the herbs were approximately within the 400-500 Dalton range. Thus the molecular mass of the individual ingredients in the soluble fraction was assumed to be 450 Daltons, which is similar to that of the prostaglandin family.

EP Receptor pcDNA

A pAcSG-EP cDNA cloned by our laboratory was first subcloned into EcoRI/XhoI sites of pcDNA3.1 (+) expression vector to generate the plasmid of pcDNA: human EP₁. The pcDNA vector has a Cytomegalovirus (CMV) promoter and geneticin (G418) as the selection antibiotic.

Stable expression of Recombinant EP₁ in HEK293 cells

HEK293 cells, placed on 100-mm dishes at a density of 2.0×10^6 , were cultured overnight, at 37°C in a humidified 5% CO₂ atmosphere in Dulbecco's Modified Eagle's Medium (DMEM) containing fetal bovine serum (FBS, 10%), antibiotics, and antimycotics. The cells were transfected with the purified cDNA of the recombinant protein (EP₁ receptor) by the Lipofectamine 2000 method [15]. Approximately 48 hours after transfection, the cells were subcultured and incubated with G418 (selection antibiotic) for four weeks to generate the EP₁ receptor stable cell line.

Western Blot analysis

The HEK293 cells stably expressing the EP₁ receptor (EP₁ receptor stable cells) were collected by centrifugation using PBS buffer, pH 7.4. After washing three times, the pellet was re-suspended in a small volume of the same buffer. Protein estimation was performed using fluorescence spectroscopy. Each protein sample (15 µg) was separated by 10% polyacrylamide gel electrophoresis under denaturing conditions and then transferred to a nitrocellulose membrane. Bands for the EP₁ receptor were recognized by the EP₁ receptor polyclonal antibody (dilution 1:1000) and visualized with horseradish peroxidase-tagged goat anti-rabbit secondary antibody (dilution 1:5000). The EP₁ receptor band was detected at 42 kDa for the stable cell line with the appropriate HEK293 control.

Confocal microscopy

Confocal microscopy was also performed to confirm the surface receptor expression in the EP₁ receptor stable cell line. The EP₁ receptor stable cell line or untransfected HEK293 control cells were grown on cover-slides that were fixed with 3.7% formaldehyde and blocked with 10% goat serum and glycine. The cells were generally permeabilized by 1% saponin and then incubated with anti-EP₁ receptor polyclonal antibody. The bound antibodies were stained by FITC-labeled (fluorescein isothiocyanate) goat anti-rabbit IgG. The stained cells were examined by fluorescence confocal microscopy.

Calcium assay

The calcium assay was performed on the HEK293 cells stably expressing the EP₁ receptor using Fluo8-AM dye. The cells were cultured in 12-well glass-bottom plates and incubated with Fluo8-AM dye (excitation 490 nm, emission 525 nm), dissolved in Modified Hank's buffered salt solution (HBSS, without Calcium and Magnesium) containing 10 mM HEPES, pH 7.6, and 0.1% bovine serum albumin (HBSSHB buffer) for 20 minutes at 22°C. The cells were then washed 3 times for perturbation with wash buffer (containing HBSSHB with Probenecid Acid (2.5 mM) and Pluronic F-68 (0.1%)), and incubated for an additional 10 minutes. Afterward, the

ligands PGE₁ (1 nM) and PGE₂ (100 pM) were individually tested for a calcium signal in a reaction volume of 1 mL wash buffer using the Nikon Ti-S eclipse microscope (40× objective) with n = 3 experiments [16]. The cell viability following the experiment was confirmed by the trypan blue dye exclusion method.

Using the CytoFluor multi-well plate reader for Calcium assay

The calcium assay was performed in 96-well plates containing the HEK293 cells stably expressing the EP₁ receptor and incubated with Fluo8-AM dye as mentioned above. Keeping in mind that the calcium signal duration is approximately 30-40 seconds, the experiment was performed four wells at a time. The water-soluble supernatants (5 µL) of the herbal extracts were applied to the HEK293 cells and the calcium signals were determined with n = 3 experiments [16]. The untransfected HEK293 control cells were also tested (data not shown). The cell viability following the experiment was confirmed by the trypan blue dye exclusion method.

Using fluorescence microscopy for Calcium assay

The calcium signal stimulators identified by the above assay were confirmed by fluorescence microscopy using the Nikon Ti-S eclipse microscope (40× objective). The HEK293 cells stably expressing EP₁ receptor were cultured in 12-well glass-bottom plates. The calcium assay (with the Fluo8-AM loading dye and wash buffer) was performed as described above. A library of desalted single herbs was obtained by serially diluting the soluble fractions in the 96-well plates until nearly colorless fractions were obtained. The calcium signal increase with potential agonist extracts were identified and confirmed [16] with n = 3 experiments. The normalized calcium signal was calculated by subtracting the inherent fluorescence of the extract in the background from the calcium signal of the EP₁ receptor stable cell line.

Results

Establishing the stable cell line expressing the EP₁ receptor

The four subtype receptors which mediate the PGE₂-pathological functions that cause pain, inflammation, and cancer are the most attractive molecules that can be used as targets for developing the next generation of NSAIDs. This is a key step toward obtaining reliable, simple, and easy biological assays for the receptor signaling mediated by its ligand. High Throughput Screening (HTS), which requires fewer procedures for the assays, is the key for successful results. We are currently using cyclic AMP (cAMP) assays for detection of the signaling mediated by the prostaglandin I₂ receptor (IP) and prostaglandin E (PGE) isomer-2 (PGE₂) subtype

receptor (EP₂) in a manner similar to that described [17], which required ligand assays and immunoassays consisting of 5-10 steps. Such procedures are not suitable for cell-based HTS because this system requires many washes during which the number of cells can be greatly depleted. However, EP₁ signaling that occurs through increasing intracellular calcium levels is a fast and reliable measurement. Yet, a simple and quick measurement for the EP₁ calcium signaling that is also suitable for cell-based screening has not been established thus far. To address this concern, we have begun by generating a cell line that consistently and stably expresses the EP₁ receptor and could be used as a sensitive drug target. This has been accomplished by the addition of G418 following transfection of the EP₁ cDNA into the HEK293 cells, which ultimately yielded the EP₁ receptor stable cell line. High expression levels of the human EP₁ receptor were observed by Western blot analysis (Figure 1A). The surface expression of the EP₁ on the HEK293 cells was also identified by Confocal microscopy (Figure 1B). On the other hand, the untransfected HEK293 cells showed no expression.



Establishing a calcium signaling assay for the stable cell line expressing the human EP₁ receptor

After generating the EP₁ receptor stable cell line and confirming the receptor expression, the cells grown in glass-bottom plates were tested for their ability to

generate a calcium signal. The calcium signal procedure (using Fluo8-AM) was described above in the methods section. In comparison to the vector-transfected and control (untransfected) HEK293 cells which lacked signaling, the stable cell line was very sensitive to dilute concentrations of PGE₁ and PGE₂ (Figure 2II.&2IV.) using Fluo8-AM. This is an extremely important observation because the non-purified fractions of the desalted herbs contained multiple ingredients cumulatively amounting to μ M concentrations.

Selection of the Traditional Chinese Medicine (TCM) herbs for the EP₁ screening

The TCM herbs selected for the screening were based on the TCM principle which favors herbs that are likely to have clinical implications in the prevention and treatment of inflammation and cancers. The herbal extracts are manufactured by pharmaceutical companies in mainland China. The facility follows the GMP (Good Manufacturing Practice) standards in China. The single herb granules (5-folds concentrated) contain less than 1,000 bacteria and less than 100 fungi per gram (no *E. coli* or live mites are present) based on the manufacturer's



instructions. The water content of each herb is less than 9% (<http://www.treasureofeast.com/>).

Comparison of calcium signal of 96 herbal extracts on recombinant EP₁ receptor expressed in HEK293 cells

Many of the herbal extracts showed an increase and decrease in calcium signal using the CytoFluor Multi-well plate reader. The colored compounds possessed high intrinsic fluorescence, therefore showed high signals. The signal from compound alone plus baseline was subtracted from the total signal. Therefore, some compounds showed high signal, some no signal or negative signal after subtraction from total signal (Figure 3I. & 3II.). The HEK293 controls did not show significant increase in fluorescence (data not shown). From the original ninety-six compounds, the top ten extracts (Figure 3I.) that stimulated the calcium signal were chosen as a potential source for agonists.

Comparison of calcium signal of the top 10 herbal extracts on recombinant EP₁ receptor expressed in HEK293 cells

The ten identified water-soluble extracts (after desalting) were serially diluted in 96-well plates (desalted drug library) until they became nearly colorless solutions. The colored extracts were producing overwhelming fluorescence signals which made microscopy difficult; therefore, we continued the testing using nearly colorless fractions. These nearly colorless dilutions of crude extract (1-40 μM) were tested for their ability to stimulate a calcium signal. Untransfected HEK293 cells and wash buffer were taken as the controls, since they showed little to no activity. The background fluorescence of the extract was subtracted from the cellular signal to give the specific fluorescence. Seven out of the ten agonists were able to stimulate the calcium signal at 1-40 μM concentrations of crude extract (Figure 4). The EC₅₀ calculated for the compounds/extracts (using 450 Dalton as an avg. mass for the herbal ingredients) were as follows: for Compound 7 (2.479 μM), 1 (4.284 nM), 3 (7.941 μM), 2 (8.55 μM), 5 (3.751 μM), 4 (4.14 μM) and 6 (7.937 μM).

Discussion

Morphine was the first pharmacologically active compound isolated from a plant [18] which was followed by a quest for drugs purified from plant sources. Plant Extracts or herbal dietary supplements have been used for ages for their biological properties, such as anti-inflammation, anti-cancer, etc., relevant for Human Diseases. The first step in screening is the identification of plant extracts and essential oils which have relevant biomedical effects [19-26].

Demonstration of relevant, potential biological activity on specific molecular targets is followed by the

identification of putative lead compounds. This is done by various procedures including direct chemical analysis of plant extracts, gas chromatography/mass spectrometry (GC-MS) and high performance liquid chromatography/MS (HPLC-MS) [12].

We have, however, generated a convenient agonist-sensitive cell-based assay to perform the primary preliminary screening for potential ligands in herbal extracts. Receptors, such as the EP₁ receptor, which couple to calcium as a second messenger can be used with Fluo8-AM dye. Since a very powerful calcium signal is detected with high sensitivity, the validity of the approach becomes more significant with high target specificity. It is advantageous as the supernatants of the extracts from the soluble fraction can be conveniently tested for potential hit compounds which can be later purified.

We suggest that the multi-plate reader can be used to initially identify the compounds (Figure 3) which can be diluted and confirmed visually using fluorescence microscopy (Figure 4). The only limitation is that natural sources are likely to have complex structures with numerous oxygen-containing substituents and an abundance of centers of stereochemistry [27], which is not ideal for rapid high-throughput screening (HTS) of desirable activity of drugs [28]. Nevertheless, natural products remain an important source of potential drugs.

Among the un-purified extracts we found seven agonists for the EP₁ receptors. Decreased sensitivity due to color interference was avoided by diluting the herbal extract until producing nearly colorless concentrations. The diluted samples for microscopy contained a sufficient amount of active ingredients to stimulate the calcium signal. The diluted sample concentrations were approximately in the range of 1-40 μM of crude extract. The treasures of the east are sold as dietary supplements and consist of well-known herbs from the Chinese material medica. These herbs are used for a multitude of problems and consumed orally directly (<http://www.treasureofeast.com/>). The EP₁ receptor target that we chose is involved in cancer, and stem cell proliferation and differentiation [1,2,14]. Antagonists to EP₁ receptor have also been studied for their ability to reduce human colonic longitudinal muscle contractility in vitro [29]. The ligand, PGE₁, is already being used in diseases such as peripheral vascular diseases with cutaneous ulcers. Though it is similar to PGE₂ in structure, PGE₁ possesses good wound healing properties, which is opposite to that of PGE₂. It is short-acting and is available as liposomal delivery for cutaneous ulcers [30-32]. Molecules mimicking PGE₁ and PGE₂ will be further differentiated and evaluated after purification. Also, antagonists to the inflammatory PGE₂ molecule can be




Figure 3 Normalized calcium signals for the 96 fractions of herbal extracts using the cytofluor multi-well plate reader. The calcium assay was performed in 96-well plates containing HEK293 cells stably expressing the EP₁ receptor and incubated with Fluo8-AM dye. The water soluble supernatants (5 μ L) of the herbal extracts were added to the stable EP₁ HEK293 cells and the calcium signals were determined. To minimize loss of signal, the experiment was performed four wells at a time. The normalized calcium signal was calculated by subtracting the inherent fluorescence of the extract in the background, where n = 3. The results were presented as four individual graphs showing the extracts, as indicated. The top 17 extracts with significantly increased calcium signal (moderate ≥ 5000 , high ≥ 20000) are listed in panel I. The normalized calcium signals for the remaining herbal extracts (18-96) with little or no activity (≤ 5000) are shown in panel II.




Figure 4 Calcium signal for active compounds. The concentration-dependent increase in calcium signal of EP₁ stable cell line for the desalted, diluted potential agonists using fluorescence microscopy are shown. The top 10 water-soluble fractions of herbal extracts were serially diluted (1:1 using distilled water) in 96-well plates until obtaining nearly colorless solutions ranging from 1-40 µM. The calcium assays were performed in 12-well glass-bottom plates following the procedures described in Figure 3. Here, the specific fluorescence signal (A.-G., line a) is shown for the extracts: Extract 7 (A.), Extract 4 (B.), Extract 3 (C.), Extract 5 (D.), Extract 2 (E.), Extract 6 (F.), and Extract 1 (G.). Their respective controls using HEK293 cells alone (A.-G., line b) and wash buffer (A.-G., line c) are also shown for all seven compounds.

evaluated for their anti-inflammatory properties. The dearth of ligands for EP receptors, especially EP₁ receptor, as a therapeutic target is what we want to explore. Similarly, all receptors (such as the EP₁), which signal through calcium as the second messenger can be evaluated for cell-based drug screening. Since the recombinant EP₁ receptor stable cell line becomes a very target-specific approach, its reliability is unquestionable. The use of the multi-well plate reader followed by fluorescence microscopy becomes a very rapid screening technique. Also, the stable cell lines are able to give strong calcium signals with as low as picomolar concentrations of PGE₁. This is beneficial as we expect very low concentrations of active ingredients in soluble fractions of herbal extracts. The current difficulty, however, remains chiefly in modifying our HTS approach for calcium signal detection using cell-based assays.

Conclusion

With this technique we could screen 96 extracts and conveniently identify 7 extracts with potential target specificity. In a multi-well plate reader it becomes difficult to identify the exact normalized calcium signal for highly colored compounds as their own fluorescence is very high giving a higher error margin. The fluorescence microscope also gradually shows overwhelming fluorescence with increasing concentrations of extracts. Our future work consists of purifying and separating the active ingredients and repeating fluorescence microscopy. Under ordinary circumstances we would have to purify the compounds first and then test all the extracts from 96 herbs. Historically, the screening of natural materials for biological activity has proven to be better [33]. Due to the increase in target sites, the pharmaceutical discovery efforts currently favor HTS of pure synthetic compounds [34]. We propose that these diluted extracts in 96-well plates can be used in HTS using calcium signal detection in cell-based assays. However, the short duration of the calcium signal is a major limitation for its applicability in HTS.

Acknowledgements

We thank the graduate students Anita Mohite, Renu Singh, Fady Makram, Mohamed Nounou, Odella Ngala, Prema Latha, Ting Zhu, Ayuk Roland, and Ashley Guillory for helping in the multi-well plate reader screening. The authors of this report would like to thank Mrs. Vanessa C. Herrera for the assistance of the manuscript preparation. This work was supported by the National Institutes of Health [Grants HL56712 and HL79389 (to KHR), HL100807 (to RD and KHR)] and the American Heart Association [Grant 10GRNT4470042 (to KHR)].

Author details

¹Centre for Experimental Therapeutics and Pharmacoinformatics, Department of Pharmacological and Pharmaceutical Sciences, College of Pharmacy, University of Houston, Houston, Texas 77004 USA. ²TR Acupuncture and Herbal Clinic, Houston, TX 77005 USA.

Authors' contributions

AJC carried out the calcium signaling assays, participated in analysis and the design of the experiments and drafted the manuscript. PK prepared the herbal drug library, helped in the calcium signaling assays and participated in analysis of the data. KT and KHR designed and guided the study and helped to correct the manuscript. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 13 October 2010 Accepted: 7 February 2011

Published: 7 February 2011

References

- O'Callaghan G, Kelly J, Shanahan F, Houston A: Prostaglandin E₂ stimulates Fas ligand expression via the EP₁ receptor in colon cancer cells. *Br J Cancer* 2008, **99**:502-512.
- Thorat MA, Morimiya A, Mehrotra S, Konger R, Badve SS: Prostanoid receptor EP₁ expression in breast cancer. *Mod Pathol* 2008, **21**:15-21.
- Gustafsson A, Hansson E, Kressner U, Nordgren S, Andersson M, Wang W, Lönnroth C, Lundholm K: EP₁₋₄ subtype, COX and PPAR gamma receptor expression in colorectal cancer in prediction of disease-specific mortality. *Int J Cancer* 2007, **121**:232-240.
- Subbaramaiah K, Hudis C, Chang SH, Hla T, Dannenberg AJ: EP₂ and EP₄ receptors regulate aromatase expression in human adipocytes and breast cancer cells. Evidence of a BRCA1 and p300 exchange. *J Biol Chem* 2008, **283**:3433-3444.
- Chang SH, Ai Y, Breyer RM, Lane TF, Hla T: The prostaglandin E₂ receptor EP₂ is required for cyclooxygenase 2-mediated mammary hyperplasia. *Cancer Res* 2005, **65**:4496-4499.
- Shoji Y, Takahashi M, Kitamura T, Watanabe K, Kawamori T, Maruyama T, Sugimoto Y, Negishi M, Narumiya S, Sugimura T, Wakabayashi K: Down regulation of prostaglandin E receptor subtype EP₃ during colon cancer development. *Gut* 2004, **53**:1151-1158.
- Chell SD, Witherden IR, Dobson RR, Moorghen M, Herman AA, Qualtrough D, Williams AC, Paraskeva C: Increased EP₄ receptor expression in colorectal cancer progression promotes cell growth and anchorage independence. *Cancer Res* 2006, **66**:3106-3113.
- Miyauchi Y: Treatment of the peripheral vascular diseases with prostaglandin. *Nippon Rinsho* 1994, **52**:2182-2186.
- Murota H, Kotobuki Y, Umegaki N, Tani M, Katayama I: New aspect of anti-inflammatory action of lipo-prostaglandin E₁ in the management of collagen diseases-related skin ulcer. *Rheumatol Int* 2008, **28**:1127-1135.
- Sobota RM, Müller PJ, Heinrich PC, Schaper F: Prostaglandin E1 inhibits IL-6-induced MCP-1 expression by interfering specifically in IL-6-dependent ERK1/2, but not STAT3, activation. *Biochem J* 2008, **412**(1):65-72.
- Bugni TS, Richards B, Bhoite L, Cimbora D, Harper MK, Ireland CM: Marine Natural Product Libraries for High-Throughput Screening and Rapid Drug Discovery. *J Nat Prod* 2008, **71**:1095-1098.
- Nicolis E, Lampronti I, Dececchi MC, Borgatti M, Tamanini A, Bianchi N, Bezzeri V, Mancini I, Giri MG, Rizzotti P, Gambari R, Cabrini G: Pyrogallol, an active compound from the medicinal plant *Emblica officinalis*, regulates expression of pro-inflammatory genes in bronchial epithelial cells. *Int Immunopharmacol* 2008, **8**:1672-1680.
- Lang G, Mayhudin NA, Mitova MI, Sun L, van der Sar S, Blunt JW, Cole AL, Ellis G, Laatsch H, Munro MH: Evolving Trends in the Dereplication of Natural Product Extracts: New Methodology for Rapid, Small-Scale Investigation of Natural Product Extracts. *J Nat Prod* 2008, **71**:1595-1599.
- Chen L, Zhang W, Yue H, Han Q, Chen B, Shi M, Li J, Li B, You S, Shi Y, Zhao RC: Effects of Human Mesenchymal Stem Cells on the Differentiation of Dendritic Cells from CD34 Cells. *Stem Cells And Development* 2007, **16**:719-731.
- Hatae N, Yamaoka K, Sugimoto Y, Negishi M, Ichikawa A: Augmentation of receptor-mediated adenylyl cyclase activity by Gi-coupled prostaglandin receptor subtype EP₃ in a Gbetagamma subunit-independent manner. *Biochem Biophys Res Commun* 2002, **290**:162-168.
- Gee KR, Brown KA, Chen WU, Bishop-Stewart J, Gray D, Johnson I: Chemical and physiological characterization of fluo-4 Ca(2+)-indicator dyes. *Cell Calcium* 2000, **27**:97-106.

17. Golla R, Seethala RA: Homogeneous Enzyme Fragment Complementation Cyclic AMP Screen for GPCR Agonists. *J Biomol Screen* 2002, **7**:515-525.
18. Hamilton GR, Baskett TF: In the arms of morpheus: the development of morphine for postoperative pain relief. *Can J Anaesth* 2000, **47**:367-374.
19. Cooper R, Morre' DJ, Morre' DM: Medicinal benefits of green tea, part II: review of anticancer properties. *J Altern Complement Med* 2005, **11**:639-652.
20. Vieira JRC, de Souza IA, do Nascimento SC, Leite SP: Indigoferasuffruticosa: an alternative anticancer therapy. *eCAM* 2007, **4**:355-359.
21. Lambertini E, Piva R, Khan MT, Lampronti I, Bianchi N, Borgatti M, Gambari R: Effects of extracts from Bangladeshi medicinal plants on in vitro proliferation of human breast cancer cell lines and expression of estrogen receptor alpha gene. *Int J Oncol* 2004, **24**:419-423.
22. Khan MT, Ather A, Thompson KD, Gambari R: Extracts and molecules from medicinal plants against herpes simplex viruses. *Antiviral Res* 2005, **67**:107-119.
23. Tohda C, Nakayama N, Hatanaka F, Komatsu K: Comparison of anti-inflammatory activities of six Curcuma rhizomes: a possible curcuminoid-independent pathway mediated by Curcuma phaeocaulis extract. *eCAM* 2006, **3**:255-260.
24. Chun S, Jee S, Lee S, Park SJ, Lee J, Kim S: Anti-inflammatory activity of the methanol extract of moutan cortex in LPS-activated Raw264.7 cells. *eCAM* 2007, **4**:327-333.
25. Lee KH, Kim AJ, Choi EM: Antioxidant and anti-inflammatory activity of pine pollen extract in vitro. *Phytother Res* 2009, **23**:41-48.
26. Wu M, Gu Z: Screening of bioactive compounds from moutan cortex and their anti-inflammatory activities in rat synoviocytes. *eCAM* 2009, **6**:57-63.
27. Koehn FE, Carter GT: The evolving role of natural products in drug discovery. *Nat Rev Drug Discov* 2005, **4**:206-220.
28. Butler MS: The Role of Natural Product Chemistry in Drug Discovery. *J Nat Prod* 2004, **67**:2141-2153.
29. Smid SD, Svensson KM: Inhibition of cyclooxygenase-2 and EP₁ receptor antagonism reduces human colonic longitudinal muscle contractility in vitro. *Prostaglandins Other Lipid Mediat* 2009, **88**:117-121.
30. Mizushima Y, Yanagawa A, Hoshi A: Prostaglandin E₁ is more effective, when incorporated in lipid microspheres, for treatment of peripheral vascular diseases in man. *J Pharm Pharmacol* 1983, **35**:666-667.
31. Toyota T, Hirata Y, Ikeda Y, Matsuoka K, Sakuma A, Mizushima Y: Lipo-PGE₁ A New Lipid-Encapsulated Preparation Of Prostaglandin E₁: Placebo-And Prostaglandin E₁-Controlled Multicentric Trials In Patients With Diabetic Neuropathy And Leg Ulcers. *Prostaglandins* 1993, **46**:453-468.
32. Rie I, Mitsuko T, Jun T, Aki K, Kayo M, Yutaka M: Marked hypotensive and blood flow-increasing effects of a new lipo-PGE (lipo-AS013) due to vascular wall targeting. *Journal of Controlled Release* 2001, **71**:157-164.
33. Weissman KJ, Leadlay PF: Combinatorial biosynthesis of reduced polyketides. *Nat Rev Microbiol* 2005, **3**:925-936.
34. Newman DJ: Natural Products as Leads to Potential Drugs: An Old Process or the New Hope for Drug Discovery? *J Med Chem* 2008, **51**:2589-2599.

Pre-publication history

The pre-publication history for this paper can be accessed here:
<http://www.biomedcentral.com/1472-6882/11/11/prepub>

doi:10.1186/1472-6882-11-11

Cite this article as: Chillar et al.: An agonist sensitive, quick and simple cell-based signaling assay for determination of ligands mimicking prostaglandin E₂ or E₁ activity through subtype EP₁ receptor: Suitable for high throughput screening. *BMC Complementary and Alternative Medicine* 2011 **11**:11.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

