

Changes of Left Ventricular Systolic Function in Patients Undergoing Coronary Artery Bypass Grafting

Vasil Papestiev^{1*}, Sasko Jovev¹, Marjan Sokarovski¹, Petar Risteski^{1,2}, Valentina Andova³, Vangel Zdraveski¹, Kujtim Dzeljilji¹, Sonja Grazhdani¹, Ljubica Georgievska-Ismail³

¹University Clinic for Cardiac Surgery, Medical Faculty, Ss. Cyril and Methodius University of Skopje, Skopje, Republic of Macedonia; ²Department of Thoracic and Cardiovascular Surgery, Johann Wolfgang Goethe University, Frankfurt, Germany; ³University Clinic of Cardiology, Medical Faculty, Ss. Cyril and Methodius University of Skopje, Skopje, Republic of Macedonia

Citation: Papestiev V, Jovev S, Sokarovski M, Risteski P, Andova V, Zdraveski V, Dzeljilji K, Grazhdani S, Georgievska-Ismail Lj. Changes of Left Ventricular Systolic Function in Patients Undergoing Coronary Artery Bypass Grafting. Open Access Maced J Med Sci. 2019 Nov 15; 7(21):3574-3578. <https://doi.org/10.3889/oamjms.2019.849>

Keywords: Coronary artery bypass graft surgery; Echocardiography; Left ventricular function

***Correspondence:** Vasil Papestiev. University Clinic for Cardiac Surgery, Medical Faculty, University Ss. Cyril and Methodius University of Skopje, Skopje, Republic of Macedonia. E-mail: papestiev@gmail.com

Received: 01-Sep-2019; **Revised:** 24-Sep-2019; **Accepted:** 25-Sep-2019; **Online first:** 12-Oct-2019

Copyright: © 2019 Vasil Papestiev, Sasko Jovev, Marjan Sokarovski, Petar Risteski, Valentina Andova, Vangel Zdraveski, Kujtim Dzeljilji, Sonja Grazhdani, Ljubica Georgievska-Ismail. This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0)

Funding: This research did not receive any financial support

Competing Interests: The authors have declared that no competing interests exist

Abstract

AIM: This prospective study was designed to evaluate the changes in left ventricular (LV) systolic function after coronary artery bypass grafting (CABG) in patients with both normal and abnormal pre-operative systolic function.

METHODS: During the period from October 2017 to October 2018, forty-seven consecutive patients undergoing CABG were enrolled in this prospective study. Transthoracic echocardiography was performed within 1 week before CABG as well as 4 to 6 months after surgery. All measurements were made by a single experienced investigator.

RESULTS: While the mean LV ejection fraction (LVEF) showed neither improvement nor significant reduction in the whole group of patients following CABG (from 54.21 ± 15.36 to $53.66 \pm 11.56\%$, $p = 0.677$), significant improvement in LVEF was detected in the subgroup of patients with pre-operative LV dysfunction (from 40.05 ± 8.65 to $45.85 \pm 9.04\%$, $p = 0.008$). On the other hand, there was a significant decline in LEFT in the subgroup of patients with normal pre-operative LEFT (from 64.70 ± 9.72 to $59.44 \pm 9.75\%$, $p = 0.008$). As for the other parameters of systolic function, significant decrease in LV end-diastolic volume index (LVEDVI) ($p = 0.001$), LV end-systolic volume index (LVESVI) ($p = 0.0001$), wall motion score index (WMSI) ($p = 0.013$) and LVmass index in male patients ($p = 0.011$) was shown only in patients with decreased LVEF after CABG. Patients with improved postoperative LVEF (53.2% of all patients) had significantly lower baseline LVEF ($p = 0.0001$), higher LVESVI (0.009) and higher WMSI ($p = 0.006$) vs patients with worsened postoperative LVEF (38.3% of all patients). Postoperative improvement of LVEF was correlated with stable angina, lack of preoperative myocardial infarction and smoking, higher baseline WMSI, higher LV internal diameters and indexed volumes in diastole and systole and lower baseline LVEF. In stepwise linear regression analysis the value of baseline LVEF appeared as independent predictor of improved LVEF after CABG ($B = 0,836\%$; 95% CI 0.655-1.017; $p = 0.0001$).

CONCLUSION: Our study showed that LVEF, internal baseline diameters and indexed volumes of LV in diastole and systole are important determinants of postoperative change in LVEF. In patients with preoperative depressed myocardial function, there is an improvement in systolic function, whereas in patients with preserved preoperative myocardial function, the decline in postoperative LVEF was detected.

Introduction

Patients with multivessel coronary artery disease (CAD), especially those with stenosis of the left main (LM) coronary artery and suitable coronary anatomy benefit from coronary artery bypass grafting (CABG) [1], [2], [3]. The goal of CABG is not only to allay symptoms and improve survival [4] but also to optimise cardiovascular function and prevent progressive remodelling. Coronary artery disease (CAD) lead's to left ventricular (LV) dysfunction as a result of myocardial scarring, stunning, or hibernation [5]. The impact of CABG on regional and global LV

systolic function has been studied but with conflicting results, most probably because of patient selection. Most of the studies that evaluated the effect of CABG in patients with severe LV dysfunction showed significant improvement in LV ejection fraction (LVEF) and LV systolic parameters after revascularisation [3], [6], [7].

Moreover, those patients with ischemic symptoms and the most severe LV dysfunction appear to benefit most from surgical revascularisation. On the other hand, only a few retrospective studies evaluate the changes in LV systolic function after CABG in patients with preserved baseline LVEF. In these patients despite the apparent improvement in

cardiac function conventional echocardiography did not change significantly even showed a decline in baseline LVEF [8], [9]. Despite advances in cardiac imaging, we believe that 2-dimensional (2D) echocardiography is still most suitable for detection of myocardial function recovery after CABG to highlight the controversies. This prospective study was designed to evaluate the changes in LV systolic parameters after CABG in patients with both normal and abnormal pre-operative systolic function.

Methods

Study patients

During the period from October 2017 to October 2018, forty-seven consecutive patients undergoing CABG were enrolled in this prospective study. All procedures were done on-pump with cardiopulmonary bypass (CPB). In every case, the left internal thoracic artery (LITA) was used to bypass the left anterior descending artery (LAD). None of the patients had associated surgical procedures such as valve replacement or surgery of the ascending aorta. This study was approved by the Medical Ethics Committee of Medical School, University Ss. "Cyril and Methodius", Skopje, and all patients provided informed consent.

2-D Echocardiography parameters

Transthoracic echocardiography was performed within 1 week before CABG as well as 4 to 6 months after surgery. All measurements were made by a single experienced investigator. Standard assessments of LV dimensions, wall thickness, LV mass, LV volumes as well as LV systolic function were performed in standard views using 2D echocardiography and Tissue Doppler imaging (TDI) on commercially available equipment (Vivid 7; GE, USA) according to the professional association recommendations [10].

Statistical analysis

Categorical parameters were summarised as percentages and continuous parameters as mean \pm SD. Comparisons of preoperative vs postoperative data were performed using a Wilcoxon Signed Rank test for related samples. Continuous variables were compared using the nonparametric Mann-Whitney test for independent samples and categorical parameters were compared using Pearson's chi-square test. Assessment of correlations was done using Pearson's correlation analysis. Stepwise regression analysis was performed to define the independent significant predictive variable of postoperative LV ejection

fraction. All data analysis was performed using SPSS version 25.0 (IBM SPSS, Inc., Chicago, Illinois), and p-value \leq 0.05 was considered significant.

Results

Patient characteristics

The patients were divided into two subgroups: those with normal preoperative LVEF (LVEF $>$ 50% n = 27) and those with abnormal LVEF (LVEF \leq 50% n = 20). The baseline demographic and clinical characteristics of the patients as a whole and divided in subgroups, including coronary anatomy, and the postoperative course are shown in Table 1 and were similar in both subgroups.

Forty-three percent of patients had decreased LVEF (\leq 50%) at baseline. These patients had significantly lower body mass index, higher Euro SCORE 2 and more likely to have chronic kidney disease. There was no statistically significant difference in age, gender, prior myocardial infarction, SINTAX score and other comorbidities between the groups. Distribution of 3-vessel CAD and significant LM stenosis was also similar between the two groups. The number of bypassed vessels was not significantly different between the subgroup of patients with normal, and abnormal LVEF. The majority of patients received three bypass grafts (median 3, range 2–5), and in all patient, a left internal thoracic artery to the left anterior descending coronary artery bypass graft was used (Table 1).

Table 1: Baseline characteristics in the study population as a whole and comparison of demographic, clinical and operative characteristics of 47 patients divided according to the preoperative LVEF

Parameter	All patients N = 47	LEFT > 50% N = 27	LVEF \leq 50% N = 20	P
Age (years)	65.55 \pm 8.25	64.93 \pm 7.74	66.40 \pm 9.02	0.628
Gender (n/%)				
Male	35 / 74.5	18 / 66.7	17 / 85.0	
Female	12 / 25.5	9 / 33.3	3 / 15	0.154
BMI (kg/m ²)	27.40 \pm 4.38	28.99 \pm 4.68	25.25 \pm 2.86	0.004
Euro SCORE	2.17 \pm 0.60	1.66 \pm 0.91	2.61 \pm 1.97	0.058
Angina, stable (n/%)	26 / 55.3	18 / 66.7	8 / 40	0.064
Previous MI (n/%)	25 / 53.2	12 / 44.4	13 / 65.0	0.135
Previous PCI (n/%)	15 / 31.9	7 / 25.0	8 / 40.0	0.306
Urgent CABG (n/%)	14 / 29.8	9 / 33.3	5 / 25.0	0.748
Preoperative AF (n/%)	2 / 4.3	1 / 3.7	1 / 5.0	0.828
COPD (n/%)	8 / 17.0	4 / 14.8	4 / 20.0	0.640
PVD (n/%)	6 / 12.8	2 / 7.4	4 / 20.0	0.201
CKD (n/%)	9 / 19.1	2 / 7.4	7 / 35	0.017
Smoking (n/%)	17 / 36.2	7 / 25.9	10 / 50.0	0.089
Hypertension (n/%)	47 / 100	27 / 100	20 / 100	-
Dyslipidemia (n/%)	46 / 97.9	26 / 96.3	20 / 100	0.384
Diabetes mellitus (n/%)	23 / 48.9	13 / 48.1	10 / 50.0	0.900
SYNTAX score	31.53 \pm 6.58	31.48 \pm 5.99	31.60 \pm 7.46	0.612
Left main disease	19 / 40.4	11 / 40.7	8 / 40.0	0.599
LAD proximal disease	38 / 80.9	20 / 74.1	18 / 90.0	0.170
1 vessel disease	-	-	-	-
2 vessel disease	9 / 19.1	6 / 22.2	3 / 15.0	-
3 vessel disease	38 / 80.9	21 / 77.8	17 / 85.0	0.407
Number of grafts	2.77 \pm 0.72	2.85 \pm 0.77	2.65 \pm 0.67	0.523
Number of grafts per patient (n/%)				
1	1 / 1.2	-	1 / 5.0	
2	15 / 31.9	9 / 33.3	6 / 30.0	
3	26 / 55.3	14 / 51.9	12 / 60.0	0.597
4	4 / 8.5	3 / 11.1	1 / 5.0	
5	1 / 1.2	1 / 3.7	-	
CPB time (min)	108.91 \pm 29.73	108.37 \pm 28.37	109.65 \pm 32.20	0.763
Ischemic time (min)	66.09 \pm 20.03	65.52 \pm 18.92	66.85 \pm 21.92	0.698

CABG = coronary artery bypass graft surgery; MI = body mass index; ACS = acute coronary syndrome; MI = myocardial infarction; PCI = percutaneous coronary intervention; AF = atrial fibrillation; COPD = chronic obstructive pulmonary disease; PAD = peripheral vascular disease; CKD = chronic kidney disease; SYNTAX = SYNERGY between percutaneous intervention with TAXUS drug-eluting stents and cardiac surgery; CPB = Cardio Pulmonary Bypass; LAD = Left Anterior Descending.

Left ventricular myocardial function before and after CABG

Echocardiographic systolic parameters in the study group as a whole and in the subgroups of patients with normal and decreased LVEF before and after CABG are shown in Table 2.

In the study group as a whole there was statistically significant reduction in LVEDVI (p=0.001), LVESVI (p = 0.003), IVSd (p = 0.037) and WMSI (p = 0.016). There was a significant improvement in MAPSE (p = 0.001). Mean LVEF showed neither improvement nor significant reduction in the whole group of patients (from 54.21 ± 15.36 to 53.66 ± 11.56%, p = 0,677). There were no postoperative changes in other LV measurements including LVIDd, LVIDs, posterior and septal wall thickness, and LVmass index (Table 2).

When we divided our cohort according to the LVEF, significant improvement in LVEF was detected in the subgroup of patients with pre-operative LV dysfunction (from 40.05 ± 8.65 to 45.85 ± 9.04%, p = 0.008), resulting in a mean change in LVEF of 5.80%. On the other hand, there was a statistically significant decline in LVEF in the subgroup of patients with normal pre-operative LVEF (from 64.70 ± 9.72 to 59.44 ± 9.75%, p = 0.008), resulting in a meaningful change in LVEF of -5.26%.

As for the other parameters of systolic function, statistically significant decrease in LVEDVI (p = 0.001), LVESVI (p = 0.0001), WMSI (p = 0.013) and LVmass index in male patients (p = 0.011) was shown only in patients with decreased LVEF after CABG (Table 2).

Table 2: Comparison of echocardiographic parameters of LV systolic function before and after CABG in patients divided according to the preoperative LEFT

Parameters	All patients	p	LVEF > 50% N = 27	p	LVEF ≤ 50% N = 20	p
LVEF (%)						
Before CABG	54.21 ± 15.36	0,677	64.70 ± 9.72	0,008	40.05 ± 8.65	0,008
After CABG	53.66 ± 11.56		59.44 ± 9.75		45.85 ± 9.04	
LVIDd (mm)						
Before CABG	51.79 ± 8.76	0,981	47.52 ± 7.07	0,666	57.55 ± 7.51	0,678
After CABG	51.83 ± 8.34		48.07 ± 6.26		56.90 ± 8.23	
LVIDs (mm)						
Before CABG	33.96 ± 10.52	0,052	28.44 ± 7.50	0,241	41.40 ± 9.45	0,106
After CABG	32.26 ± 9.72		27.48 ± 7.24		38.70 ± 8.99	
IVSd (mm)						
Before CABG	13.06 ± 2.25	0,037	13.41 ± 2.27	0,111	12.60 ± 2.18	0,175
After CABG	12.34 ± 2.66		12.74 ± 2.55		11.80 ± 2.78	
PWd (mm)						
Before CABG	11.21 ± 2.04	0,345	11.56 ± 1.98	0,418	10.75 ± 2.07	0,681
After CABG	10.79 ± 1.98		11.07 ± 1.79		10.40 ± 2.21	
LVEDVI (ml/m2)						
Before CABG	64.54 ± 31.73	0,001	48.44 ± 19.88	0,107	86.29 ± 32.10	0,001
After CABG	54.41 ± 22.22		43.65 ± 14.62		68.94 ± 22.71	
LVESVI (ml/m2)						
Before CABG	33.89 ± 27.19	0,003	17.52 ± 10.60	0,614	55.99 ± 27.24	0,0001
After CABG	26.81 ± 17.24		18.12 ± 8.73		38.54 ± 19.05	
SVI (ml/m2)						
Before CABG	38.08 ± 9.99	0,804	38.97 ± 10.03	0,313	36.89 ± 10.07	0,455
After CABG	39.18 ± 10.56		38.30 ± 10.46		40.37 ± 10.85	
MAPSEaverage (mm)						
Before CABG	12.94 ± 2.26	0,001	13.81 ± 2.32	0,035	11.78 ± 1.58	0,008
After CABG	14.08 ± 2.08		14.98 ± 1.98		12.88 ± 1.57	
sTDI (cm/s)						
Before CABG	6.04 ± 1.44	0,608	6.34 ± 1.67	0,695	5.65 ± 0.96	0,281
After CABG	6.12 ± 1.21		6.20 ± 1.32		6.02 ± 1.11	
WMSI						
Before CABG	1.34 ± 0.35	0,016	1.17 ± 0.25	0,615	1.57 ± 0.33	0,013
After CABG	1.23 ± 0.23		1.12 ± 0.14		1.37 ± 0.25	
LVmass index (g/m2)						
Male						
Before CABG	143.43 ± 39.26	0,017	129.04 ± 33.39	0,420	158.67 ± 40.14	0,011
After CABG	122.31 ± 24.14		122.87 ± 26.23		143.89 ± 32.41	
Female						
Before CABG	133.08 ± 30.85	0,347	113.84 ± 20.52	0,767	147.72 ± 15.15	0,109
After CABG	115.29 ± 26.82		107.95 ± 25.80		137.31 ± 17.97	

CABG = coronary artery bypass graft surgery; CI = cardiac index; IVSd = septal wall thickness; LVEDVI = left ventricular end-diastolic volume indexed to body surface area; LEFT = left ventricular ejection fraction; LVESVI = left ventricular end-systolic volume indexed to body surface area; LVIDd = left ventricular end-diastolic dimension; LVIDs = left ventricular end-systolic dimension; MAPSE = mitral annular plane systolic excursion; PW = posterior wall thickness; sTDI = peak systolic mitral annular velocity by TDI; SVI = systolic volume indexed to body surface area; WMSI = wall motion score index; *p < 0,05 for comparison between groups.

The only parameter that significantly improved in both groups after CABG was MAPSE (p = 0,035, and p = 008 in patients with preserved and reduced LVEF respectively). Except for MAPSE, none of the systolic echocardiographic parameters improved in the subgroup of patients with preserved LVEF (Table 2).

Parameters related to LVEF change post CABG surgery

In our study, out of 47 patients, 4 patients (8.5%) had unchanged LVEF (+ / -5%) after successful CABG operation, 25 patients (53.2%) had increased LVEF (> 5%) and 18 patients (38.3%) had decreased in the postoperative LVEF (> 5%). Comparison of the three groups (Table 3) showed an only significant difference between patients with improved and decreased postoperative LVEF. Thus, patients with improved postoperative LVEF had significantly lower baseline LVEF (p = 0.0001), higher LVESVI (0.009) and higher WMSI (p = 0.006) vs patients with worsened postoperative LEFT.

Table 3: Baseline echocardiographic parameters of all patients about perioperative change in left ventricular ejection fraction

Parameter	Unchanged EF N = 4	Improved EF N = 25	Worsened EF N = 18	p
LVIDd (mm)	48.0 ± 3.9	53.7 ± 9.3	49.8 ± 8.3	0,243
LVIDs (mm)	29.0 ± 4.2	37.1 ± 11.6	30.7 ± 8.6	0,089
IVSd (mm)	13.2 ± 2.9	12.9 ± 2.4	13.1 ± 1.8	0,945
PWd (mm)	10.2 ± 1.5	10.8 ± 2.3	11.9 ± 1.5	0,173
LVEDVI (ml/m2)	64.1 ± 34.1	74.0 ± 33.6	51.4 ± 24.7	0,069
LVESVI (ml/m2)	25.7 ± 11.3	44.9 ± 30.9	20.3 ± 15.5	0,009
LVEF (%)	56.7 ± 5.1	46.6 ± 14.9	64.1 ± 11.2	0,0001
SVI (ml/m2)	36.5 ± 6.0	37.9 ± 10.2	38.6 ± 10.7	0,930
CI (L/min/m2)	2.7 ± 0.8	2.5 ± 0.7	2.6 ± 0.7	0,888
MAPSEaverage (mm)	14.1 ± 2.6	12.4 ± 1.9	13.2 ± 2.5	0,277
sTDI (cm/s)	5.4 ± 0.9	5.9 ± 1.5	6.2 ± 1.4	0,544
WMSI	1.3 ± 0.2	1.4 ± 0.4	1.3 ± 0.3	0,006

CABG = coronary artery bypass graft surgery; CI = cardiac index; IVSd = septal wall thickness; LVEDVI = left ventricular end-diastolic volume indexed to body surface area; LEFT = left ventricular ejection fraction; LVESVI = left ventricular end-systolic volume indexed to body surface area; LVIDd = left ventricular end-diastolic dimension; LVIDs = left ventricular end-systolic dimension; MAPSE = mitral annular plane systolic excursion; PW = posterior wall thickness; sTDI = peak systolic mitral annular velocity by TDI; SVI = systolic volume indexed to body surface area; WMSI = wall motion score index; *p < 0,05 for comparison between groups.

Postoperative improvement of LVEF was correlated with stable angina, lack of preoperative myocardial infarction and smoking, higher baseline WMSI, higher LV internal diameters and indexed volumes in diastole and systole and lower baseline LVEF (Table 4).

Table 4: Correlation between the change of LVEF and preoperative parameters

Parameters	ΔLVEF
Angina (%)	R = 0.386; p = 0.007
Previous MI (%)	R = -0.288; p = 0.049
Smoking (%)	R = -0.319; p = 0.029
LVIDd (mm)	R = -0.294; p = 0.045
LVIDs (mm)	R = -0.404; p = 0.005
LVEDVI (ml/m2)	R = -0.467; p = 0.001
LVESVI (ml/m2)	R = -0.557; p = 0.0001
LVEF	R = 0.652; p = 0.0001
WMSI	R = -0.480; p = 0.001

To determine the independent predictors of improvement of LVEF after CABG, we performed multiple stepwise linear regression analysis with covariates that showed a significant relation to it. The results demonstrated that the value of baseline LVEF appeared as an independent predictor of improved LVEF after CABG (Table 5, Figure 1).

Table 5: Stepwise regression analysis of LVEF after CABG as the dependent variable and clinical and echocardiographic parameters as independent variables in cases for which LVEF improved

Model	Coefficients ^{a,b}				t	Sig.	95.0% Confidence Interval for B	
	Unstandardized Coefficients	Std. Error	Standardized Coefficients	Beta			Lower Bound	Upper Bound
1	(Constant)	14,624	4,281		3,416	,002	5,769	23,479
	LVEF before CABG	,836	,088	,894	9,545	,000	,655	1,017

a. Dependent Variable: LVEF post CABG
 b. Selecting only cases for which LVEF improved = improved

Thus, for every 1% absolute decrease in pre-operative LVEF, there is postoperative improvement of LVEF of 0,836% (95% CI 0.655-1.017; p = 0.0001).




Figure 1: Graphical presentation of regression standardised predicted value for LVEF after CABG as the dependent variable in cases for which LVEF improved

Discussion

CABG surgery can improve the myocardial blood supply in the hibernating regions of the heart. This results in increased contractility and better performance of the myocardium [11], [12]. The 2D biplane echocardiography is a widely used method to obtain pre and postoperative systolic parameters of the right and left ventricle in patients undergoing CABG surgery.

In this study, the parameters of LV ejection fraction, LV internal dimensions and LV indexed volumes in systole and diastole, cardiac index, mitral annular plane systolic excursion, wall motion score index and LV mass index obtained with conventional 2D echocardiography were used to assess the global systolic function in patients with CAD undergoing CABG.

We showed deterioration in LVEF after CABG in patients with normal baseline LVEF. Except for MAPSE, other systolic parameters did not change significantly even after successful CABG treatment in patients with preserved baseline LVEF. On the other hand, an improvement in LV systolic function was observed in patients with decreased pre-operative LVEF. There were significant improvement in LVEF, LV systolic and diastolic indexed volumes, WMSI and LV mass index in this subgroup of patients.

Prior studies have similarly found an improvement in LVEF and other systolic parameters in patients with pre-operative LV systolic dysfunction. In the largest prospective, randomised, controlled trial, the STICH trial, Michler et al., [13] in a post hoc subgroup analysis showed a significant improvement in LV size and function in the subgroup of patients with higher baseline LV end-systolic dimensions. Our study also showed improvement in LVEF in the subgroup of patients with preoperative LV systolic dysfunction and higher baseline internal diameters as well as indexed volumes of LV in diastole and systole.

While many studies evaluate changes of perioperative systolic parameters in patients with reduced LVEF, only a few studies examine changes in LV systolic function in patients with preserved baseline LVEF. In a small prospective study, Diller et al. demonstrated improvement in LV diastolic function and did not find a significant reduction in LVEF immediately after CABG [14]. In the largest study to assess pre and post-operative echocardiograms in a population including both normal and reduced pre-operative LV function, Koene et al., [8] showed a decrease in LVEF with CABG in patients with normal baseline LV systolic function. In this study, the magnitude of decrease in LVEF was 3% mean and ranged from -33% to 15%. Our study is in agreement with these findings demonstrating a decrease in LVEF in patients with preserved baseline LVEF resulting in a mean decrease in LVEF of 5.26%. This postoperative decrease in LVEF might result from myocardial stunning [15], reperfusion injury [16] and early postoperative graft failure [17].

In our study, a total of 18 patients (38.3%) had decreased in the postoperative LVEF (> 5%). This suggests that CABG itself contributes to postoperative myocardial dysfunction. Although these patients were angina free 4-6 months after CABG, the relative decline in LVEF suggests that myocardial recovery might take longer time. We strongly believe that these results are worthy of further investigation to understand the effect of CABG on myocardial function. Another issue that should be investigated is whether the lack of improvement of LVEF post-CABG portends a worse outcome.

The major limitations in our study are that we used only conventional 2-D echocardiography imaging to assess pre and postoperative systolic LV function. Other technologies such as magnetic resonance

imaging, positron emission tomography and speckle tracking imaging might have yielded other results, but 2-D echocardiography is a widely used method for quantifying perioperative LV function. This study has the advantage of being prospective and all consecutive patients that met inclusion criteria were enrolled in the study but we believe that their number is too small and is thus hypothesis-generating rather than definitive. Another disadvantage is that paired echocardiograms were done a maximum of 6 months after surgery, time that might be too short for complete myocardial recovery after surgery. In our study, all patients were done on the pump with crystalloid cardioplegia and this might affect postoperative LV function in a certain percentage of patients.

In conclusion, our study showed that LVEF, internal baseline diameters and indexed volumes of LV in diastole and systole are important determinants of postoperative change in LVEF. In patients with the preoperative depressed myocardial function, we should expect improvement in systolic function, whereas in patients with preserved myocardial function, decline in postoperative LVEF should be anticipated, despite successful CABG. The present study suggests further investigations in order to understand the effect of CABG on myocardial function.

References

1. Win Decker S, Strobeck S, Stefani GG, et al. Revascularization versus medical treatment in patients with stable coronary artery disease: Network meta-analysis. *BMJ*. 2014; 348:g3859.
2. Yusuf S, Zucker D, Peduzzi P, et al. Effect of coronary artery bypass graft surgery on survival: Overview of 10-year results from randomized trials by the Coronary Artery Bypass Graft Surgery Trialists Collaboration. *Lancet*. 1994; 344:563-70. [https://doi.org/10.1016/S0140-6736\(94\)91963-1](https://doi.org/10.1016/S0140-6736(94)91963-1)
3. Velazquez EJ, Lee KL, Jones RH, et al. STICHES Investigators. Coronary-artery bypass surgery in patients with ischemic cardiomyopathy. *N Engl J Med*. 2016; 374:1511-20. <https://doi.org/10.1056/NEJMoa1602001> PMID:27040723 PMID:PMC4938005
4. Neumann FJ, Sousa-Uva M, Ahlsson A, et al. ESC Scientific Document Group. 2018 ESC/EACTS Guidelines on myocardial revascularization. *Eur Heart J*. 2018; 00:1-96. <https://doi.org/10.5603/KP.2018.0228> PMID:30566213
5. Anavekar NS, Chareonthitawee P, Narula J, Gersh BJ. Revascularization in Patients With Severe Left Ventricular Dysfunction: Is the Assessment of Viability Still Viable? *J Am Coll Cardiol*. 2016; 67:2874-87. <https://doi.org/10.1016/j.jacc.2016.03.571> PMID:27311527
6. Rubenson DS, Tucker CR, London E, et al. Two-dimensional echocardiographic analysis of segmental left ventricular wall motion before and after coronary bypass surgery. *Circulation*. 1982; 66:1025-33. <https://doi.org/10.1161/01.CIR.66.5.1025> PMID:6982113
7. Elefteriades JA, Tolls G, Levi E, Mills LK, Zaret BL. Coronary artery bypass grafting in severe left ventricular dysfunction: excellent survival with improved ejection fraction and functional state. *J Am Coll Cardiol*. 1993; 22:1411-7. [https://doi.org/10.1016/0735-1097\(93\)90551-B](https://doi.org/10.1016/0735-1097(93)90551-B)
8. Koene RJ, Kealhofer JV, Adabag S, et al. Effect of coronary artery bypass graft surgery on left ventricular systolic function. *J Thorac Dis*. 2017; 9:262-70. <https://doi.org/10.21037/jtd.2017.02.09> PMID:28275473 PMID:PMC5334093
9. Yin ZY1, Li XF, Tu YF, et al. Speckle-tracking imaging to monitor myocardial function after coronary artery bypass graft surgery. *J Ultrasound Med*. 2013; 32:1951-6. <https://doi.org/10.7863/ultra.32.11.1951> PMID:24154899
10. Lang RM, Badano LP, Mor-Avi V, et al. Recommendations for cardiac chamber quantification by echocardiography in adults: An update from the American Society of Echocardiography and the European Association of Cardiovascular Imaging. *Eur Heart J Cardiovasc Imaging*. 2015; 16:233-71. <https://doi.org/10.1093/ehjci/jev014> PMID:25712077
11. Bax JJ, Van der Wall EE. Assessment of myocardial viability: guide to prognosis and clinical management. *Eur Heart J*. 2000; 21:961-963. <https://doi.org/10.1053/euhj.2000.2110> PMID:10901505
12. Bair TL, Muhlestein JB, May HT, et al. Surgical revascularization is associated with improved long-term outcomes compared with percutaneous stenting in most subgroups of patients with multivessel coronary artery disease: results from the Intermountain Heart Registry. *Circulation*. 2007; 116(11):1226-1231. <https://doi.org/10.1161/CIRCULATIONAHA.106.681346> PMID:17846308
13. Michler RE, Rouleau JL, Al-Khalidi HR, et al. Insights from the STICH trial: change in left ventricular size after coronary artery bypass grafting with and without surgical ventricular reconstruction. *J Thorac Cardiovasc Surg*. 2013; 146:1139-45.e6. <https://doi.org/10.1016/j.jtcvs.2012.09.007> PMID:23111018 PMID:PMC3810307
14. Diller GP, Wasan BS, Kyriacou A, et al. Effect of coronary artery bypass surgery on myocardial function as assessed by tissue Doppler echocardiography. *Eur J Cardiothorac Surg*. 2008; 34:995-9. <https://doi.org/10.1016/j.ejcts.2008.08.008> PMID:18829341
15. Leung JM. Clinical evidence of myocardial stunning in patients undergoing CABG surgery. *J Card Surg*. 1993; 8:220-3. <https://doi.org/10.1111/j.1540-8191.1993.tb01310.x> PMID:8461506
16. Lin PJ, Chang CH, Lee YS, et al. Acute endothelial reperfusion injury after coronary artery bypass grafting. *Ann Thorac Surg*. 1994; 58:782-8. [https://doi.org/10.1016/0003-4975\(94\)90749-8](https://doi.org/10.1016/0003-4975(94)90749-8)
17. Oshima H, Tokuda Y, Araki Y, et al. Predictors of early graft failure after coronary artery bypass grafting for chronic total occlusion. *Interact Cardiovasc Thorac Surg*. 2016; 23:142-9. <https://doi.org/10.1093/icvts/ivw084> PMID:27030683 PMID:PMC4986749