

Research

Open Access

Molecular characterisation of drug-resistant *Plasmodium falciparum* from Thailand

Dinora Lopes¹, Kanchana Rungsihirunrat², Fátima Nogueira¹, Aree Seugorn², José Pedro Gil^{1,3}, Virgilio E do Rosário¹ and Pedro Cravo*¹

Address: ¹Centro de Malária e Outras Doenças Tropicais/IHMT/UNL, Rua da Junqueira, 96, 1349-008, Lisbon, Portugal, ²Malaria Research Unit, Institute of Health Research, Chulalongkorn University, Bangkok 10330, Thailand and ³Present address: Malaria Research Laboratory, Division of Infectious Diseases, Institutionen för Medicin, Karolinska Institutet, Karolinska Sjukhuset, Malaria Research Laboratory L7, Plan 3, 17176 Stockholm, Sweden

E-mail: Dinora Lopes - dmlopes@ihmt.unl.pt; Kanchana Rungsihirunrat - Kanchana.R@Chula.ac.th; Fátima Nogueira - fnogueira@ihmt.unl.pt; Aree Seugorn - Aree.S@Chula.ac.th; José Gil - pedro.gil@ks.se; Virgilio E do Rosário - cmdt@ihmt.unl.pt; Pedro Cravo* - pcravo@ihmt.unl.pt

*Corresponding author

Published: 14 October 2002

Received: 28 June 2002

Malaria Journal 2002, 1:12

Accepted: 14 October 2002

This article is available from: <http://www.malariajournal.com/content/1/1/12>

© 2002 Lopes et al; licensee BioMed Central Ltd. This article is published in Open Access: verbatim copying and redistribution of this article are permitted in all media for any purpose, provided this notice is preserved along with the article's original URL.

Abstract

Background: The increasing levels of *Plasmodium falciparum* resistance to chloroquine (CQ) in Thailand have led to the use of alternative antimalarials, which are at present also becoming ineffective. In this context, any strategies that help improve the surveillance of drug resistance, become crucial in overcoming the problem.

Methods: In the present study, we have established the *in vitro* sensitivity to CQ, mefloquine (MF), quinine (QUIN) and amodiaquine (AMQ) of 52 *P. falciparum* isolates collected in Thailand, and assessed the prevalence of four putative genetic polymorphisms of drug resistance, *pfcr* K76T, *pfmdr1* N86Y, *pfmdr1* D1042N and *pfmdr1* Y1246D, by PCR-RFLP.

Results: The percentage of isolates resistant to CQ, MF, and AMQ was 96% (50/52), 62% (32/52), and 58% (18/31), respectively, while all parasites were found to be sensitive to QUIN. In addition, 41 (79%) of the isolates assayed were resistant simultaneously to more than one drug; 25 to CQ and MF, 9 to CQ and AMQ, and 7 to all three drugs, CQ, MF and AMQ. There were two significant associations between drug sensitivity and presence of particular molecular markers, i) CQ resistance / *pfcr* 76T ($P = 0.001$), and ii) MF resistance / *pfmdr1* 86N ($P < 0.001$)

Conclusions: i) In Thailand, the high levels of CQ pressure have led to strong selection of the *pfcr* 76T polymorphism and ii) *pfmdr1* 86N appears to be a good predictor of *in vitro* MF resistance.

Background

Malaria due to *Plasmodium falciparum* affects 300 million people and claims an estimated 1.5 million lives every year. Our present inability to synthesise a fully protective vaccine means that chemotherapy stands as the only effective measure in the control of the disease. However, in

many parts of the world the parasite *P. falciparum* has become resistant to most drugs presently used [1], seriously undermining efforts for controlling malaria.

Chloroquine (CQ) has long been the drug of choice for the treatment of malaria; however, CQ-resistant parasites

are now present in most areas where malaria is endemic [2]. Chloroquine resistance is especially well established in Thailand, after having been first described in that country in the late 1950's [3]. The decline in the efficacy of chloroquine has led to the use of alternative antimalarials, such as antifolates, mefloquine and artemisinin derivatives, but parasite resistance to these drugs is also becoming a real problem [2]. In this context, understanding the genetic basis of drug resistance is essential for implementing rational measures to overcome the problem.

Although significant progress has been made in trying to understand how resistance to CQ may occur, many aspects of it remain unclear, and the genetic mechanisms responsible for mefloquine and quinine resistance are largely unknown. Nevertheless, two main genes have been implicated in quinoline resistance; the *pfmdr1* (*P. falciparum* multi-drug resistance1) and the *Pfcr1* (*P. falciparum* chloroquine resistance transporter). There is evidence from the analysis of a genetic cross which indicates that point polymorphisms in the *pfmdr1* gene may modulate sensitivity to both mefloquine (MF) and artemisinin in *P. falciparum*[4]. Furthermore, recent genetic transfection work has suggested that single nucleotide polymorphisms in the *pfmdr1* gene encoding changes in aminoacids 1034, 1042 and 1246 can influence parasite responses to mefloquine, quinine and halofantrine as well as to the structurally unrelated drug artemisinin, and modulate sensitivity to chloroquine depending on the genetic background of the parasites strains [5]. However, chloroquine resistance was shown to segregate independently of the *pfmdr1* gene, following a genetic cross between a CQ-sensitive parasite, *P. falciparum* HB3, and a CQ-resistant one, Dd2 [6], and the absence of a clear association between *pfmdr1* and chloroquine responses in natural parasite populations [7–15], strongly suggests the involvement of other gene(s). Recently, detailed linkage analysis and fine chromosome mapping of progeny clones of the HB3 × Dd2 cross has allowed the identification of another gene, *pfcr1*, in which a mutation at aminoacid 76 (*pfcr1* K76T) is highly correlated with increased CQ tolerance among field parasite isolates of *P. falciparum*[16–24]. In addition, a causal relationship between *pfcr1* 76T and chloroquine resistance has been confirmed by genetic transfection experiments [16].

The study of the correlation between drug resistance in natural parasite populations and genetic polymorphisms may allow the development of molecular tools to help predict responses to drugs and, as mentioned above, the *pfcr1* and *pfmdr1* genes have been identified as putative markers of quinoline resistance. In the present work we have investigated possible associations between four molecular markers in these genes and sensitivity to chloroquine, mefloquine, quinine (QUIN) and amodiaquine (AMQ) of *P. falciparum* parasites collected in Thailand.

Methods

Study site and method of parasite collection

After confirmation of *P. falciparum* infection by microscopical observation of thin and thick Giemsa-stained blood films, approximately 20 µl of blood were collected by finger-prick from consenting subjects living in 4 different areas of Thailand where malaria is highly endemic; the Tak province (North-western Thailand), Kanchanaburi (Western Thailand: Thai-Myanmar border), Chonburi (Eastern Thailand), Chantaburi and Trat (Eastern Thailand: Thai-Cambodia border). Samples were placed in sterile 1.5 ml micro-centrifuge tubes, together with 0.5 ml of transport medium (10 ml RPMI 1640 complete medium without serum, plus 20 µl of heparin 5000 i.u./ml), carried to Chulalongkorn University of Bangkok on the day of collection, at ambient temperature and placed in *in vitro* culture, following established procedures [25].

Parasite phenotyping (micro-tests)

Assessment of *P. falciparum* susceptibility to CQ, MF, QUIN and AMQ was performed according to the M.I.C. test [26,27], whereby samples were exposed to a range of concentrations of the four drugs, in microtitre culture plates, for 72 hours, with daily changes of medium (in the presence or absence of drugs). After this period, thin film Giemsa-stained preparations from each culture plate well were observed under light microscopy for the presence of parasites. The results were expressed in "minimum inhibiting concentration" (M.I.C.) units, i.e., the lowest drug concentration required to kill all or nearly all parasites (I.C.99) after 72 hours [26,27].

Parasite reference strains of known sensitivity to the drugs concerned were tested in parallel to all field-collected isolates. In this manner, the drug response of each isolate was determined by comparison of the M.I.C. between each sample and the relevant reference strains. Accordingly, samples tested for CQ, MF, QUIN and AMQ responses were clustered into two groups; sensitive (S) or resistant (R) based on sensitivity thresholds established in previous studies [26–29].

Genotyping by PCR-RFLP

Genomic DNA was extracted prior to drug testing, by a standard phenol:chloroform DNA extraction method, precipitated with absolute ethanol and stored in TE Buffer (Tris-EDTA), following established protocols [30]. The resulting DNA was used as template in 50 µl PCR reactions, containing 1 µM of each oligonucleotide primer, 1 × PCR buffer (Promega™), 2.5 mM MgCl₂, 0.2 mM dNTP's and 0.025 U/µl of Promega™ *Taq* DNA polymerase. Accordingly, a fragment of the *pfcr1* gene containing codon 76 was amplified by PCR using a Nested-PCR approach. For amplification of DNA fragments containing *pfmdr1* polymorphisms we used oligonucleotide primers published

Table 1: Polymerase Chain Reaction for amplification of fragments containing *pfcr* and *pfmdr1* gene polymorphisms

| Primer | Sequence (5' → 3') | PCR |
|---------------------------|------------------------------|--|
| <i>Pfcr</i> 76 | | |
| Ist round sense | CAAGAAGGAAGTAAGTATCCAAAAATGG | 94°C, 30"; 56°C, 30"; 60°C, 60"; 45 cycles |
| Antisense | GTAGTTCTTGTAGACCTATGAAGGC | |
| Nested sense | GCAAAAATGACGAGCGTTATAGAG | 94°C, 30"; 59°C, 30"; 60°C, 60"; 45 cycles |
| Antisense | CTGAACAGGCATCTAACATGGATATAGC | |
| <i>Pfmdr1</i> 86 | | |
| Sense | ATGGGTAAAGAGCAGAAAGAG | 94°C, 30"; 53°C, 30"; 68°C, 60"; 10 cycles, followed by 94°C, 30"; 50°C, 30"; 68°C, 60", 35 cycles |
| Antisense | CGTACCAATTCCTGAACTCAC | |
| <i>Pfmdr1</i> 1042 | | |
| Ist round sense | TATGTCAAGCGGAGTTTTTTCG | 94°C, 30"; 50°C, 30"; 68°C, 60"; 45 cycles |
| Antisense | TCTGAATCTCCTTTTAAGGAC | |
| Semi-nested sense | GTAAATGCAGCTTTATGGG | 94°C, 30"; 50°C, 30"; 68°C, 60"; 45 cycles |
| Antisense | TCTGAATCTCCTTTTAAGGAC | |
| <i>Pfmdr1</i> 1246 | | |
| Sense | CTACAGCAATCGTTGGAGAAA | 94°C, 30"; 53°C, 30"; 68°C, 60"; 10 cycles, followed by 94°C, 30"; 50°C, 30"; 68°C, 60", 35 cycles |
| Antisense | GCTCTAGCTATAGCTATTCTC | |

elsewhere [31], as well as newly designed ones. In this manner, the fragments of the *pfmdr1* gene containing codons 86 and 1246 were amplified in a single-step PCR, whereas the sequence of codon *pfmdr1* 1042 was determined following amplification by semi-nested PCR. All primer sequences and respective PCR conditions are presented in Table 1.

Restriction enzymes generating RFLPs

Following amplification of the fragments concerned, polymorphisms in the *pfcr* and *pfmdr1* genes were assessed as follows: *pfcr* 76K and *pfmdr1* 86N were detected by incubation of the corresponding PCR fragments with *ApoI* (r/aaty), *pfmdr1* 1042N was detected using *AsnI* (at/taat), and *pfmdr1* 1246Y was determined by incubation with *EcoRV* (gat/atc). Endonucleases *ApoI*, *AsnI* and *EcoRV* had been obtained from New England BioLabs™, Roche Molecular Biochemicals™ and Stratagene™ respectively, and incubations were setup following the manufacturers instructions. Appropriate control DNA of samples with known *pfcr* and *pfmdr1* sequences was used in parallel with field-collected parasite isolates in every PCR-RFLP protocol; these were 3D7 (genotype *pfcr* 76K, *pfmdr1* 86N, *pfmdr1* 1042N, *pfmdr1* 1246D), HB3 (genotype *pfmdr1* 1042D), Dd2 (genotype *pfcr* 76T, *pfmdr1* 86Y) and 180/92 (genotype *pfmdr1* 1246Y). The products resulting from restrictions of *pfmdr1* 1042 were resolved in 8% acrylamide gels, whereas *pfmdr1* 86, *pfmdr1* 1246 and *pfcr* 76 digests were run on 2% agarose gels, with both types of gels made in 1 × TBE buffer. All gels were stained

with ethidium bromide and visualised under UV (ultraviolet) transillumination.

Statistical analysis of the association between *pfcr* and *pfmdr1* markers and drug responses

We searched for statistically significant associations between sensitivity to each of the drugs among all isolates and the presence of each of the particular markers included in our study by using Fisher's Exact Test (2-tailed) after having arranged the data in 2 × 2 contingency tables (mixed infections were excluded from this analysis). An association between a particular marker and resistance to a given drug was considered to be significant if the *P* value was found to be lower than 0.05 (*P* < 0.05).

Results

Responses to drugs

In total, 52 samples were analysed *in vitro* for their susceptibility to chloroquine, mefloquine and quinine, and 31 for amodiaquine. The results of these tests are presented in Table 2.

Pfcr and *pfmdr1* genotypes

The *pfcr* 76 PCR primers amplified a sub-fragment of the gene comprising 479 nucleotides, containing both a monomorphic and a polymorphic *ApoI* restriction site. In this manner, incubation of this fragment with endonuclease *ApoI* generated either 3 fragments of 122, 124 and 233 base-pairs (b.p.), or 2 fragments of 233 and 246 b.p., if K or T were present respectively (Figure 1). The PCR of *pfmdr1* 86 amplified a fragment of 504 b.p., within which

Table 2: Results of *in vitro* drug tests (CQ-chloroquine; MF-mefloquine; QUIN-quinine; AMQ-amodiaquine; S-sensitive; R-resistant)

| | CQ | MF | QUIN | AMQ | CQ+MF only | CQ+AMQ only | MF+AMQ only | CQ+MF+AMQ |
|---|-------------|-------------|--------------|-------------|-------------|-------------|-------------|------------|
| S | 4% (2/52) | 38% (20/52) | 100% (52/52) | 42% (13/31) | - | - | - | - |
| R | 96% (50/52) | 62% (32/52) | - | 58% (18/31) | 48% (25/52) | 29% (9/31) | -(0/31) | 23% (7/31) |

Table 3: Frequencies of allelic polymorphisms in the *pfcr* and *pfmdr1* genes

| | <i>Pfcr</i> | | | | | <i>Pfmdr1</i> | | | | | | |
|-----------|-------------|-----|-------|-----|-----|---------------|-------|-------|-------|-------|-------|-------|
| Allele | 76K | 76T | Mixed | 86N | 86Y | Mixed | 1042N | 1042D | Mixed | 1246D | 1246Y | Mixed |
| Frequency | .04 | .83 | .13 | .60 | .23 | .17 | .92 | .08 | 0 | .92 | .02 | .06 |

the presence of asparagine (N) was determined by incubation with *ApoI*, originating segments of 255 and 249 b.p., which resolved as a single band in 2% agarose gels (Figure 1). For detection of 1042 polymorphisms (N or D) the 188 b.p. fragment obtained by PCR could be restricted with endonuclease *AsnI*, after which 3 segments were obtained of 116, 46 and 26 b.p. if 1042N were present in the sample, and two fragments of 162 and 26 b.p. in the presence of the alternative aminoacid (as with *pfcr* 76, one of the restriction sites for *AsnI* within this segment is also monomorphic, therefore always resulting in at least one cut) (Figure 1). Codon 1246 polymorphisms (D or Y) were assayed following incubation of the corresponding 508 b.p. PCR product with *EcoRV*, which produced 2 fragments of 268 and 240 b.p. if the target DNA contained tyrosine (Y) (Figure 1). All data resulting from this analysis is compiled in Table 3, where allele frequencies for each of the markers are shown.

Association between *pfcr* and *pfmdr1* markers and responses to drugs

Two significant correlations were detected between the presence of a particular marker and *in vitro* outcomes (Tables 4 and 5); one between CQ resistance and the presence of *pfcr* 76T ($P = 0.001$) and the other, between MF resistance and the presence of *pfmdr1* 86N ($P < 0.001$). There was an evident lack of an association between AMQ responses and all of the markers studied (Table 6), a fact that was corroborated after statistical analysis (data not shown). In addition, since all isolates proved to be quinine sensitive, any correlations between response to this drug and corresponding genotypes could not be established. The data of the *in vitro* tests for the 4 drugs, and corresponding genotypes is compiled in Table 7.

Discussion

The increasing failure rates of several antimalarial drugs in the majority of malaria-affected areas means that close monitoring of the epidemiology and dynamics of drug resistance are necessary if we are to implement measures to circumvent the problem. The identification and validation of easy, rapid molecular markers of drug resistance would greatly facilitate this process, and would allow us to overcome difficulties in the use of traditional methods for assaying drug sensitivity.

In Thailand, CQ resistance was first reported more than forty years ago [3], and after ten years, resistance to chloroquine had become so widespread that use of the drug against *P. falciparum* was abandoned. At present, even though the drug is used only against *P. vivax*, it is perhaps not surprising to find that most *P. falciparum* are largely unaffected by CQ. Our *in vitro* observations show a near total prevalence of CQR (96%) in the present study area, confirming what has been widely reported [1]. Among all CQR isolates tested in our study, the presence of *pfcr* 76T was universal ($P = 0.001$), indicating complete selection of this polymorphism by the drug, a result that is in agreement with recently published work not only from Thai-originated parasites [22], but also from parasites of different areas of the globe [32].

The correlation between *pfmdr1* genotypes and quinoline resistance has often generated conflicting results; although it has been suggested that *pfmdr1* 86Y can be correlated with increased CQ resistance in parasites which originated from different areas of the world [33–36], other field studies have not corroborated these findings [7–15], and the results of a *P. falciparum* genetic cross indicat-




Figure 1

Agarose (*pfmdr1* 86, *pfmdr1* 1246, *pfcr76*), and acrylamide (*pfmdr1* 1042) gels showing PCR products and corresponding restriction digests, of control and field-collected samples of *P. falciparum*. (PCR-RFLP controls – 3D7 (genotype *pfcr76* K, *pfmdr1* 86N, *pfmdr1* 1042N, *pfmdr1* 1246D), HB3 (genotype *pfmdr1* 1042D), Dd2 (genotype *pfcr76* T, *pfmdr1* 86Y) and I80/92 (genotype *pfmdr1* 1246Y); IV, V and VIII – DNA Molecular Weight Markers IV, Bioline; V and VIII, Roche)

ed that CQR did not depend on the *pfmdr1* gene [6]. Our present findings do not implicate *pfmdr1* 86 in CQ resistance in Thailand, since the presence of both N and Y in our samples was largely independent of their CQ response, indicating that chloroquine does not appear to exert selective pressure on this area of the gene. Interestingly though, *pfmdr1* 1042D and *pfmdr1* 1246N, previously associated with increased CQ sensitivity following genetic transfection experiments [5], were largely absent in our samples, possibly suggesting a mechanism of chloroquine resistance that may in part depend on the presence of the alternative polymorphic alleles, *pfmdr1* 1042N and *pfmdr1* 1246D, respectively.

Mefloquine was introduced in Thailand in the form of Fansimef (mefloquine-sulfadoxine-pyrimethamine) during 1984 with an initial cure rate of 95%, but MF-resistant *P. falciparum* parasites have arisen and present a real threat to the control of malaria, especially in the Thai/Cambodia and Thai/Myanmar border areas [37]. In the present work the prevalence of MF resistance was 62%, indicating a worrying trend. A correlation between *pfcr76* T and mefloquine sensitivity would always be difficult to establish, since the near complete presence of this polymorphism is likely to have been selected by chloroquine pressure, whose mechanisms of action and resistance are probably distinct from those of MF. Earlier work from Thailand and other areas of the world has indicated that an increase in the level of mefloquine sensitivity among field isolates of

Table 4: *Plasmodium falciparum* isolates listed according to sensitivity to chloroquine (CQ), with corresponding *pfcr* and *pfmdr1* markers

| Isolate | CQ | <i>pfcr</i> 76 | 86 | <i>pfmdr1</i> 1042 | 1246 |
|---------|----|----------------|----|--------------------|------|
| T9/94b3 | S | K | Y | N | D |
| TM408 | S | K | N | N | D |
| T101 | R | T | N | N | D |
| T108 | R | T | N | N | D |
| T113 | R | T | N | N | D |
| T115 | R | T | N | N | D |
| T116 | R | T | N | N | D |
| T120 | R | T | Y | N | D |
| T123 | R | T | N | D | D |
| T130 | R | T | N | N | D |
| T131 | R | T | N | N | D |
| T132 | R | T | N | N | D |
| T133 | R | T | N | N | D |
| S3 | R | T | Y | N | D |
| S64 | R | T | Y | N | D |
| S71 | R | T | Y | N | D |
| S90 | R | T | Y | D | D |
| S118 | R | T | Y | N | D |
| S149 | R | T | NY | N | DY |
| S151 | R | KT | NY | N | D |
| S152 | R | T | Y | N | D |
| S153 | R | T | Y | N | D |
| S157 | R | KT | NY | N | D |
| S160 | R | KT | NY | N | DY |
| CH1 | R | T | Y | D | D |
| CH3 | R | T | Y | D | D |
| CH7 | R | T | N | N | D |
| TD2 | R | T | N | N | D |
| TD3 | R | T | N | N | D |
| TD8 | R | T | N | N | D |
| TD14 | R | T | N | N | D |
| TD21 | R | T | N | N | D |
| TD27 | R | T | N | N | D |
| TD49 | R | T | N | N | D |
| TD56 | R | T | N | N | D |
| TD61 | R | KT | N | N | D |
| TD62 | R | T | N | N | D |
| TD64 | R | T | N | N | DY |
| TD79 | R | KT | N | N | D |
| TD134 | R | T | N | N | D |
| TD328 | R | KT | N | N | D |
| TP4 | R | T | Y | N | D |
| TP7 | R | T | NY | N | D |
| TP13 | R | T | NY | nd | D |
| TP17 | R | T | NY | N | D |
| TP18 | R | T | N | N | D |
| TP20 | R | T | NY | N | D |
| TP21 | R | KT | N | nd | D |
| TP26 | R | T | N | N | D |
| TP34 | R | T | N | N | D |
| TP40 | R | T | N | nd | Y |
| RC17 | R | T | NY | N | D |

(S – sensitive; R – resistant; KT, NY and DY – mixed infections; nd – not determined; $P = 0.001$ for the association between CQR and presence of *pfcr* 76T, mixed alleles excluded)

Table 5: *Plasmodium falciparum* isolates listed according to sensitivity to mefloquine (MF), with corresponding *pfcr* and *pfmdr1* markers

| Isolate | MF | <i>pfcr</i> 76 | 86 | <i>pfmdr1</i> 1042 | 1246 |
|---------|----|----------------|----|--------------------|------|
| T9/94b3 | S | K | Y | N | D |
| S3 | S | T | Y | N | D |
| S64 | S | T | Y | N | D |
| S71 | S | T | Y | N | D |
| S90 | S | T | Y | D | D |
| S118 | S | T | Y | N | D |
| S149 | S | T | NY | N | DY |
| S151 | S | KT | NY | N | D |
| S152 | S | T | Y | N | D |
| S153 | S | T | Y | N | D |
| S157 | S | KT | NY | N | D |
| S160 | S | KT | NY | N | DY |
| CH1 | S | T | Y | D | D |
| CH3 | S | T | Y | D | D |
| TP4 | S | T | Y | N | D |
| TP7 | S | T | NY | N | D |
| TP13 | S | T | NY | nd | D |
| TP17 | S | T | NY | N | D |
| TP18 | S | T | N | N | D |
| TP40 | S | T | N | nd | Y |
| TM408 | R | K | N | N | D |
| TP20 | R | T | NY | N | D |
| TP21 | R | KT | N | nd | D |
| RC17 | R | T | NY | N | D |
| T108 | R | T | N | N | D |
| T113 | R | T | N | N | D |
| T115 | R | T | N | N | D |
| T130 | R | T | N | N | D |
| T131 | R | T | N | N | D |
| T132 | R | T | N | N | D |
| TD61 | R | KT | N | N | D |
| TD14 | R | T | N | N | D |
| T101 | R | T | N | N | D |
| T116 | R | T | N | N | D |
| T120 | R | T | Y | N | D |
| T123 | R | T | N | D | D |
| T133 | R | T | N | N | D |
| CH7 | R | T | N | N | D |
| TD2 | R | T | N | N | D |
| TD3 | R | T | N | N | D |
| TD8 | R | T | N | N | D |
| TD21 | R | T | N | N | D |
| TD27 | R | T | N | N | D |
| TD49 | R | T | N | N | D |

Table 5: *Plasmodium falciparum* isolates listed according to sensitivity to mefloquine (MF), with corresponding *pfcr* and *pfmdr1* markers

| | | | | | |
|-------|---|----|---|---|----|
| TD62 | R | T | N | N | D |
| TD134 | R | T | N | N | D |
| TD328 | R | KT | N | N | D |
| TD56 | R | T | N | N | D |
| TD64 | R | T | N | N | DY |
| TD79 | R | KT | N | N | D |
| TP26 | R | T | N | N | D |
| TP34 | R | T | N | N | D |

(S – sensitive; R – resistant; KT, NY and DY – mixed infections; nd – not determined; $P < 0.001$ for the association between MFR and presence of *pfmdr1* 86N, mixed alleles excluded)

Table 6: *Plasmodium falciparum* isolates listed according to sensitivity to amodiaquine (AQ), with corresponding *pfcr* and *pfmdr1* markers

| Isolate | AQ | <i>pfcr</i> 76 | 86 | <i>Pfmdr1</i> 1042 | 1246 |
|---------|----|----------------|----|--------------------|------|
| SI53 | S | T | Y | N | D |
| SI57 | S | KT | NY | N | D |
| SI60 | S | KT | NY | N | DY |
| TD3 | S | T | N | N | D |
| TD49 | S | T | N | N | D |
| TD56 | S | T | N | N | D |
| TD62 | S | T | N | N | D |
| TD79 | S | KT | N | N | D |
| TD134 | S | T | N | N | D |
| TD328 | S | KT | N | N | D |
| TP4 | S | T | Y | N | D |
| TP18 | S | T | N | N | D |
| TP26 | S | T | N | N | D |
| T9/94b3 | R | K | Y | N | D |
| TM408 | R | K | N | N | D |
| S3 | R | T | Y | N | D |
| SI18 | R | T | Y | N | D |
| SI49 | R | T | NY | N | DY |
| SI51 | R | KT | NY | N | D |
| SI52 | R | T | Y | N | D |
| TD14 | R | T | N | N | D |
| TD61 | R | KT | N | N | D |
| TD64 | R | T | N | N | DY |
| TP7 | R | T | NY | N | D |
| TP13 | R | T | NY | nd | D |
| TP17 | R | T | NY | N | D |
| TP20 | R | T | NY | N | D |
| TP21 | R | KT | N | nd | D |
| TP34 | R | T | N | N | D |
| TP40 | R | T | N | nd | Y |
| RC17 | R | T | NY | N | D |

(S – sensitive; R – resistant; KT, NY and DY – mixed infections; nd – not determined)

Table 7: Number of isolates clustered according to sensitivity to each drug and corresponding *pfprt* and *pfmdr1* genotypes

| Drug | | <i>Pfprt</i> | | | | | | <i>Pfmdr1</i> | | | | | |
|-------------|--------------|--------------|-----|------|-----|-----|------|---------------|-------|--------|-------|-------|--------|
| | | 76K | 76T | 76KT | 86N | 86Y | 86NY | 1042N | 1042D | 1042ND | 1246D | 1246Y | 1246DY |
| CQ | Total | | | | | | | | | | | | |
| S | 2 | 2 | 0 | 0 | 1 | 1 | 0 | 2 | 0 | 0 | 2 | 0 | 0 |
| R | 50 | 0 | 43 | 7 | 30 | 11 | 9 | 43 | 4 | 0 | 46 | 1 | 3 |
| MF | | | | | | | | | | | | | |
| S | 35 | 2 | 27 | 6 | 2 | 11 | 7 | 29 | 3 | 0 | 32 | 1 | 2 |
| R | 17 | 0 | 16 | 1 | 29 | 1 | 2 | 16 | 1 | 0 | 16 | 0 | 1 |
| QUIN | | | | | | | | | | | | | |
| S | 52 | 2 | 43 | 7 | 31 | 12 | 9 | 45 | 4 | 0 | 48 | 1 | 3 |
| R | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| AMQ | | | | | | | | | | | | | |
| S | 13 | 0 | 9 | 4 | 9 | 2 | 2 | 13 | 0 | 0 | 12 | 0 | 1 |
| R | 18 | 2 | 13 | 3 | 7 | 4 | 7 | 15 | 0 | 0 | 15 | 1 | 2 |

(CQ – chloroquine; MF – mefloquine; QUIN – quinine; AMQ – amodiaquine; S – sensitive; R – resistant; 76KT, 86NY, 1042ND and 1246DY – mixed infections; shaded boxes indicate the two cases where there was a significant association between the presence of a particular marker and drug sensitivity: *pfprt* 76T/CQR, $P = 0.001$ and *pfmdr1* 86N/MFR, $P < 0.001$)

P. falciparum may be correlated with a mutation in codon 86 of the *pfmdr1* gene (N86Y) [38–40]. In the present work, the occurrence of *pfmdr1* 86N was significantly associated with MF resistance as 31/32 resistant isolates carried this polymorphism ($P < 0.001$), strongly suggesting that 86N is an important event in the generation of MF resistance and may be a useful marker to monitor MF resistance in this country.

Interestingly, although the prevalence of AMQ resistance was high (58%), we did not detect a significant correlation between AMQ responses and any of the markers studied, contrary to what could be expected considering that AMQ is chemically very similar to CQ. These observations indicate that the mechanism of action and/or resistance differ between the two drugs, which may raise interesting questions about the design of new CQ-derivative compounds.

Quinine is one of the most effective drugs for the treatment of malaria in Thailand. Although the efficacy of this drug has been reduced, it has not yet become a serious problem, since the combination with tetracycline increases its cure rate. In fact, *P. falciparum* quinine sensitivity was total in all regions covered by our study, showing that despite its adverse side effects, quinine can still be used as a reliable resource of malaria therapy in Thailand. The reasons for the long-lasting efficacy of quinine may be manifold, but its explanations lie outside the scope of this article. Most importantly, quinine may prove especially valuable in the treatment of multi-drug resistant *Falci-parum* malaria, which we found to be largely present in

our study as 41 out of 52 isolates were found to be resistant to more than one compound and 7 were unaffected by all drugs (CQ, MF and AMQ) except quinine.

Conclusions

Taken together, our results seem to be suggesting that CQ and MF are the major selective forces on the *pfprt* and *pfmdr1* genes, whereby the presence of *pfprt* 76T, and possibly *pfmdr1* 1042N and *pfmdr1* 1246D in Thai-originated parasites has been selected by chloroquine pressure. The *pfmdr1* 86N mutation seems to be important only for mefloquine resistance, and may represent a useful marker for monitoring resistance in this country, although its validation may require *in vivo* correlates and the analysis of a larger number of samples.

Authors' contributions

DL carried out a proportion of the parasite phenotyping, performed the molecular analysis of the majority of the samples and contributed for the elaboration of the manuscript. KR performed the molecular typing of a number of the isolates. FN participated in the parasite's phenotyping and genotyping. AS performed parasite collection and did a proportion of the micro-tests. JPG, VR and PC, conceived the study, participated in its design and co-ordination and were involved in phases of the experimental work.

Competing interests

None declared.

Acknowledgements

Dinora Lopes, Fátima Nogueira, José Pedro Gil, Virgílio do Rosário and Pedro Cravo were funded by the Ministério da Ciência e Tecnologia, Portugal. This work was partially supported by Fundação para a Ciência e Tecnologia (FCT), project PRAXIS XXI P/SAU/14070/98. Dinora Lopes and Pedro Cravo are grateful to David Walliker, Paul Hunt and Hamza Babiker from the Institute of Cell, Animal and Population Biology, University of Edinburgh, for excellent scientific advice and support. The work in Thailand was supported by malaria grants from Institute of Health Research, Faculty of Science, Chulalongkorn University. We thank the Malaria Division, Ministry of Health (Bangkok) and especially the staff of the Malaria Clinics for their co-operation.

References

- White NJ: **Drug resistance in malaria.** *Br Med Bull* 1998, **54**:703-715
- Peters W: **Drug resistance in malaria parasites of animals and man** *Adv Parasitol* 1998, **41**:1-62
- Harinasuta T, Migasen S, Boonag D: **Chloroquine resistance in *Plasmodium falciparum* in Thailand.** *UNESCO First Regional Symposium on Scientific Knowledge of Tropical Parasites, Singapore University, Singapore* 1962
- Duraisingh MT, Jones P, Sambou I, Von Seidlein L, Pinder M, Warhurst DC: **The tyrosine-86 of the *pfmdr1* gene of *Plasmodium falciparum* is associated with increased sensitivity to the anti-malarials mefloquine and artemisinin.** *Mol Biochem Parasitol* 2000, **108**:13-23
- Reed MB, Saliba KJ, Caruana SR, Kirk K, Cowman AF: **PghI modulates sensitivity and resistance to multiple antimalarials in *Plasmodium falciparum*.** *Nature* 2000, **403**:906-909
- Wellemans TE, Panton LJ, Gluzman IY, Rosário VE, Gwadz RW, Walker-Jonah A, Krogstad DJ: **Chloroquine resistance not linked to *mdr*-like genes in a *Plasmodium falciparum* cross.** *Nature* 1990, **345**:253-255
- Awad-el-Kariem FM, Miles MA, Warhurst DC: **Chloroquine-resistant *Plasmodium falciparum* isolates from the Sudan lack two mutations in the *pfmdr1* gene thought to be associated with chloroquine resistance.** *Trans R Soc Trop Med Hyg* 1992, **86**:587-589
- Wilson C, Volkman S, Thaithong S, Martin R, Kyle D, Milhous W, Wirth D: **Amplification of *Pfmdr1* associated with mefloquine and halofantrine resistance in *Plasmodium falciparum* from Thailand.** *Mol Biochem Parasitol* 1993, **57**:151-160
- Cremer G, Basco LK, Le Bras J, Camus D, Slomianny C: ***Plasmodium falciparum*: detection of P-glycoprotein in chloroquine-susceptible and chloroquine-resistant clones and isolates.** *Exp Parasitol* 1995, **81**:1-8
- Basco L, de Pecoulas PE, Le Bras J, Wilson CM: ***Plasmodium falciparum*: molecular characterisation of multi-drug resistant Cambodian isolates.** *Exp Parasitol* 1996, **82**:97-103
- Bhattacharya PR, Biswas S, Kabilan L: **Alleles of the *Plasmodium falciparum* *pfmdr1* gene appear not to be associated with chloroquine resistance in India.** *Trans R Soc Trop Med Hyg* 1997, **91**:454-455
- von Seidlein L, Duraisingh MT, Drakeley CJ, Bailey R, Greenwood BM, Pinder M: **Polymorphisms in the *pfmdr1* gene and chloroquine resistance in *Plasmodium falciparum* in the Gambia.** *Trans R Soc Trop Med Hyg* 1997, **91**:450-453
- Povoa MM, Adagu IS, Oliveira SG, Machado RL, Miles MA, Warhurst DC: ***Pfmdr1* Asn1042Asp and Asp1246Tyr polymorphisms, thought to be associated with chloroquine resistance, are present in chloroquine-resistant and sensitive Brazilian field isolates of *Plasmodium falciparum*.** *Exp Parasitol* 1998, **88**:64-68
- Bhattacharya PR, Pillai CR: **Strong association, but incomplete correlation between chloroquine resistance and allelic variation in the *pfmdr1* gene of *Plasmodium falciparum* isolates from India** *Ann Trop Med Parasitol* 1999, **93**:679-684
- Chaiyaroj SC, Buranakiti A, Angkasekwinai P, Looressuwan S, Cowman AF: **Analysis of mefloquine resistance and amplification of *pfmdr1* in multidrug-resistant *Plasmodium falciparum* isolates from Thailand.** *Am J Trop Med Hyg* 1999, **61**:780-783
- Fidock DA, Nomura T, Talley AT, Cooper RA, Dzekunov S., Ferdig MT, Ursos LM, Sidhu AS, Naude B, Deitsch K., Su XZ, Wootton JC, Roepe PD, Wellemans TE: **Mutations in the *Plasmodium falciparum*. Digestive vacuole transmembrane protein PfCRT and evidence for their role in chloroquine resistance.** *Mol Cell* 2000, **6**:861-871
- Babiker A, Pringle SJ, Abdel-Muhsin A, Mackinnon M, Hunt P, Walliker D: **High level of chloroquine resistance in Sudanese isolates of *Plasmodium falciparum* is associated with mutations in the chloroquine resistance transporter gene *pfCRT* and the multidrug resistance gene *pfmdr1*** *J Inf Dis* 2001, **183**:1535-1538
- Djimdé A, Doumbo OK, Steketee RW, Plowe CV: **Application of a molecular marker for surveillance of chloroquine-resistant malaria.** *Lancet* 2001, **358**:890-891
- Djimdé A, Doumbo OK, Cortese JF, Kayentao K, Doumbo S, Diourte Y, Dicko , Su XZ, Nomura T, Fidock DA, Wellemans TE, Plowe CV, Coulibaly D.: **A molecular marker for chloroquine-resistant falciparum malaria.** *New Engl J Med* 2001, **344**:257-263
- Dorsey G, Kanya MR, Singh A, Rosenthal PJ: **Polymorphisms in the *Plasmodium falciparum* *pfCRT* and *pfmdr1* genes and clinical response to chloroquine in Kampala, Uganda.** *J Inf Dis* 2001, **183**:1417-1420
- Durand R, Jafari R, Vauzelle J, Delabre JF, Jesic Z, Le Bras J: **Analysis of *pfCRT* point mutations and chloroquine susceptibility in isolates of *Plasmodium falciparum*.** *Mol Biochem Parasitol* 2001, **114**:95-102
- Labbé AC, Bualombai P, Pillai DR, Zhong KJY, Vanisaveth V, Hongvathong B, Looareesuwan S, Kain KC: **Molecular markers for chloroquine-resistant *Plasmodium falciparum* malaria in Thailand and Laos.** *Ann Trop Med Parasitol* 2001, **95**:781-788
- Maguire JD, Susanti AI, Krisin S, Sismadi P, Fryauff DJ, Baird JK: **The T76 mutation in the *pfCRT* gene of *Plasmodium falciparum* and clinical chloroquine resistance phenotypes in Papua, Indonesia.** *Ann Trop Med Parasitol* 2001, **95**:559-572
- Pillai DR, Labbé AC, Vanisaveth V, Hongvathong B, Pomphida S, Inkathone S, Zhong KJY, Kain KC: ***Plasmodium falciparum* malaria in Laos: chloroquine treatment outcome and predictive value of molecular markers.** *J Inf Dis* 2001, **183**:789-795
- Thaithong S, Seugorn A, Beale GH: **Culturing *Plasmodium falciparum* from finger-prick samples of infected blood.** *Trans R Soc Trop Med Hyg* 1994, **88**:490
- Thaithong S, Beale GH: **"Malaria Parasites".** *Chulalongkorn University research report Series, No. 1* 1992
- Thaithong S, Beale GH, Chutmongkonkul M: **Susceptibility of *Plasmodium falciparum* to five drugs: an *in vitro* study of isolates mainly from Thailand.** *Trans R Soc Trop Med Hyg* 1983, **77**:228-231
- Congpuong K, Sirtichaisinthop J, Tippawangkosol P, Suprakrob K, Nabangchang K, Tan-ariya P, Karbwang J: **Incidence of antimalarial pretreatment and drug sensitivity *in vitro* in multidrug-resistant *Plasmodium falciparum* infection in Thailand.** *Trans R Soc Trop Med Hyg* 1998, **92**:84-6
- Creasey A, Fenton B, Walker A, Thaithong S, Oliveira S, Mutambu S, Walliker D: **Genetic diversity of *Plasmodium falciparum* shows geographical variation.** *Am J Trop Med Hyg* 1990, **42**:403-413
- Sambrook J, Fritsch EF, Maniatis T: **"Molecular Cloning: A Laboratory Manual".** *Cold Spring Harbour Laboratory Cold Spring Harbour, New York* 1989
- Cox-Singh J, Singh B, Alias A, Abdullah MS: **Assessment of the association between three *pfmdr1* point mutations and chloroquine resistance *in vitro* of Malaysian *Plasmodium falciparum* isolates.** *Trans R Soc Trop Med Hyg* 1995, **89**:436-437
- Wellemans TE, Plowe CV: **Chloroquine-resistant malaria.** *J Inf Dis* 2001, **184**:770-776
- Footo SJ, Kyle DE, Martin RK, Oduola AM, Forsyth K, Kemp DJ, Cowman AF: **Several alleles of the multidrug-resistance gene are closely linked to chloroquine resistance in *Plasmodium falciparum*.** *Nature* 1990, **345**:255-258
- Basco L, Le Bras J, Rhoades Z, Wilson C: **Analysis of *Pfmdr1* and drug susceptibility in fresh isolates of *Plasmodium falciparum* from Sub-Saharan Africa.** *Mol Biochem Parasitol* 1995, **74**:157-166
- Duraisingh M, Drakeley C, Muller O, Bailey R, Snounou G, Targett G, Greenwood B, Warhurst D: **Evidence for selection for the tyrosine-86 allele of the *Pfmdr1* gene of *Plasmodium falciparum* by chloroquine and amodiaquine.** *Parasitology* 1997, **114**:205-211
- Adagu IS, Warhurst DC: **Association of *cg2* and *pfmdr1* genotypes with chloroquine resistance in field samples of *Plasmodium falciparum* from Nigeria.** *Parasitology* 1999, **119**:343-348
- Wernsdorfer W: **Epidemiology of drug resistance in malaria.** *Acta Trop* 1994, **56**:143-156

38. Mungthin M, Bray PG, Ward AS: **Phenotypic and genotypic characteristics of recently adapted isolates of *Plasmodium falciparum* from Thailand.** *Am J Trop Med Hyg* 1999, **60**:469-74
39. Price RN, Cassar C, Brockman A, Duraisingh M, Van Vugt M, White NJ, Nosten F, Krishna S: **The *pfmdr1* gene is associated with a multidrug-resistant phenotype in *Plasmodium falciparum* from the western border of Thailand.** *Antimicrob Agents Chemother* 1999, **43**:2943-2949
40. Duraisingh MT, Roper C, Walliker D, Warhurst DC: **Increased sensitivity to the antimalarials mefloquine and artemisinin is conferred by mutations in the *pfmdr1* gene of *Plasmodium falciparum*** *Mol Microbiol* 2000, **36**:955-961

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMedcentral will be the most significant development for disseminating the results of biomedical research in our lifetime."

Paul Nurse, Director-General, Imperial Cancer Research Fund

Publish with **BMC** and your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours - you keep the copyright

Submit your manuscript here:

<http://www.biomedcentral.com/manuscript/>



BioMedcentral.com

editorial@biomedcentral.com