

CCL21/CCR7 Promotes G₂/M Phase Progression via the ERK Pathway in Human Non-Small Cell Lung Cancer Cells

Ying Xu^{1,2}, Lifeng Liu³, Xueshan Qiu^{1,2*}, Lili Jiang^{1,2}, Bo Huang^{1,2}, Haiying Li^{1,2}, Zixuan Li^{1,2}, Wenting Luo^{1,2}, Enhua Wang^{1,2}

1 Department of Pathology, First Affiliated Hospital and College of Basic Medical Sciences, China Medical University, Shenyang, Liaoning, China, **2** Institute of Pathology and Pathophysiology, China Medical University, Shenyang, Liaoning, China, **3** Department of Orthopaedics, First Affiliated Hospital of China Medical University, Shenyang, Liaoning, China

Abstract

C-C chemokine receptor 7 (CCR7) contributes to the survival of certain cancer cell lines, but its role in the proliferation of human non-small cell lung cancer (NSCLC) cells remains vague. Proliferation assays performed on A549 and H460 NSCLC cells using Cell Counting Kit-8 indicated that activation of CCR7 by its specific ligand, exogenous chemokine ligand 21 (CCL21), was associated with a significant linear increase in cell proliferation with duration of exposure to CCL21. The CCL21/CCR7 interaction significantly increased the fraction of cells in the G₂/M phase of the cell cycle as measured by flow cytometry. In contrast, CCL21/CCR7 had no significant influence on the G₀/G₁ and S phases. Western blot and real-time PCR indicated that CCL21/CCR7 significantly upregulated expression of cyclin A, cyclin B1, and cyclin-dependent kinase 1 (CDK1), which are related to the G₂/M phase transition. The expression of cyclin D1 and cyclin E, which are related to the G₀/G₁ and G₁/S transitions, was not altered. The CCL21/CCR7 interaction significantly enhanced phosphorylation of extracellular signal-regulated kinase (P-ERK) but not Akt, as measured by Western blot. LY294002, a selective inhibitor of PI3K that prevents activation of the downstream Akt, did not weaken the effect of CCL21/CCR7 on P-ERK. Coimmunoprecipitation further confirmed that there was an interaction between P-ERK and cyclin A, cyclin B1, or CDK1, particularly in the presence of CCL21. CCR7 small interfering RNA or PD98059, a selective inhibitor of MEK that disrupts the activation of downstream ERK, significantly abolished the effects of exogenous CCL21. These results suggest that CCL21/CCR7 contributes to the time-dependent proliferation of human NSCLC cells by upregulating cyclin A, cyclin B1, and CDK1 potentially via the ERK pathway.

Citation: Xu Y, Liu L, Qiu X, Jiang L, Huang B, et al. (2011) CCL21/CCR7 Promotes G₂/M Phase Progression via the ERK Pathway in Human Non-Small Cell Lung Cancer Cells. *PLoS ONE* 6(6): e21119. doi:10.1371/journal.pone.0021119

Editor: Lin Zhang, University of Pennsylvania, United States of America

Received: May 7, 2011; **Accepted:** May 19, 2011; **Published:** June 16, 2011

Copyright: © 2011 Xu et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported by grants from National Natural Science Foundation of China (No. 30972967) and Research Fund for the Doctoral Program of the Higher Education of China (No. 20092104110018). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: xueshan-qiu@hotmail.com

Introduction

C-C chemokine receptor 7 (CCR7) is expressed on all naive T-cells and on some memory T-cells, B-cells, and mature dendritic cells [1]. Upon interaction with its ligands, chemokine ligand 19 (CCL19) or chemokine ligand 21 (CCL21) [2], CCR7 contributes to lymphocyte trafficking and homing to lymph nodes during immune and inflammatory reactions [3–5]. CCR7 is highly expressed in non-small cell lung cancer (NSCLC), breast cancer, and squamous cell carcinoma of the head and neck and is responsible for mediating metastasis in certain cancer cell lines [6–15]. To date, the role of CCR7 in the proliferation of human NSCLC cells has not been elucidated.

Activation of CCR7 can increase phosphorylation of extracellular signal-regulated kinase (P-ERK) or Akt (P-Akt) via G_i proteins to enhance cell proliferation or survival [16–20]. ERK belongs to the mitogen-activated protein kinase (MAPK) family, which also includes c-Jun N-terminal kinase (JNK) and p38. The ERK cascade, activated by mitogenic stimuli, is critical for proliferation and survival [21,22] and is required for normal progression into mitosis [23,24]. The JNK and p38 pathways are

activated in response to chemicals and environmental stress [25–27]. Akt (also known as Akt1), a mediator of growth factor-induced cell survival [28–30], may promote cell proliferation via phosphorylation [31].

The purpose of this study was to examine the effect and regulatory mechanism of the CCL21/CCR7 interaction on the proliferation of A549 and NCI-H460 (H460) human NSCLC cells. Here, we demonstrated that CCL21/CCR7 contributed to the time-dependent proliferation of human NSCLC cells by upregulating the expression of cyclin A, cyclin B1, and CDK1 via the ERK pathway. Information garnered from this study lends insight to the mechanisms of survival of CCR7-mediated cancer cells and has implications for treatment targets in NSCLC.


Results

CCL21/CCR7 promotes proliferation of A549 and H460 cells. In a previous study, we identified a higher CCR7 expression level in A549 and H460 human NSCLC cell lines compared with other cell lines [32]. To investigate the role of CCR7 in the functioning of A549 and H460 cells, CCR7 activation and inhibition were

induced with exogenous CCL21 and with CCR7 small interfering RNA (siRNA), respectively. After transfection with CCR7 siRNA (siCCR7) or control siRNA, the expression of CCR7 was evaluated using Western blot and reverse transcriptase (RT)-PCR. We found that siCCR7 significantly downregulated the protein and mRNA levels of CCR7, compared with control siRNA (Figure 1).

To determine the effect of CCL21/CCR7 on cell proliferation, the CCK-8 assay was performed on A549 and H460 cells. According to the published data [33,34] and the results of our preliminary experiment, at 100 ng/mL concentrations CCL21 significantly promoted cell proliferation, compared with 50 ng/mL concentrations, while there were no significant difference between 100 ng/mL and 200 ng/mL concentrations (Figure 2).

A



B




Figure 1. Efficiency of CCR7 siRNA in A549 or H460 cells. A549 (A) and H460 (B) cells were transfected with control siRNA or CCR7 siRNA (siCCR7). After transfection, the expression of CCR7 protein (a) and mRNA (b) was evaluated using Western blot (a) and RT-PCR (b) and compared to untransfected A549 or H460 cells. Each bar represents the mean \pm SD of three independent experiments. * $p < 0.05$, compared with control cells. doi:10.1371/journal.pone.0021119.g001

Therefore, at 100 ng/mL concentrations CCL21 was used in following experiment. The CCL21/CCR7 interaction significantly promoted cell proliferation, whereas siCCR7 significantly abrogated the action of CCL21 (Figure 3). siCCR7 alone had no significant effect on cell proliferation, compared with control cells. Significant differences were observed between all time point examined (all $p < 0.01$), indicating a linear increase in proliferation with increasing exposure times to CCL21 (all $p < 0.01$).

CCL21/CCR7 augments the proportion of cells in G₂/M. To verify whether the action of CCL21/CCR7 on the proliferation of A549 and H460 cells is associated with a change in cell cycle distribution, cell cycle analysis was performed using flow cytometry. The CCL21/CCR7 interaction significantly enhanced the proportion of cells in the G₂/M phase, whereas there was no significant effect of this interaction on the proportion of cells in G₀/G₁ or the S phase, compared with control cells (Table 1). siCCR7 significantly abolished this effect of CCL21, whereas siCCR7 alone had no significant effect on cell cycle distribution.

CCL21/CCR7 upregulates the expression of cyclin A, cyclin B1, and CDK1. To determine the possible mechanism by which CCL21/CCR7 influences the G₂/M phase distribution in A549 and H460 cells, the expression of cyclins and cyclin-dependent kinase 1 (CDK1) was assessed using Western blot and real-time PCR. Compared with control cells, the CCL21/CCR7 interaction significantly upregulated the protein and mRNA levels of cyclin A, cyclin B1, and CDK1, which are related to G₂/M phase progression (Figure 4). siCCR7 significantly abrogated the effects of CCL21, whereas siCCR7 alone had no significant effect on cyclin or CDK1 expression. CCL21 had no significant effect on the levels of cyclin D1 or cyclin E, which are related to G₀/G₁ and the G₁/S transition.

CCL21/CCR7 upregulates the expression of P-ERK but not P-Akt. Others have reported that CCR7 may increase the phosphorylation of ERK, JNK, or Akt, which are related to cell survival [16–20,33,35–42]. To verify whether the CCL21/CCR7 interaction may also enhance the expression of MAPK family members and Akt in A549 and H460 cells, the expression of these components was assessed using Western blot. The CCL21/CCR7 interaction significantly upregulated the expression of P-ERK at 24 h and 48 h, whereas there was no significant impact on the

expression of ERK (Figure 5). CCL21/CCR7 had no significant influence on the expression or phosphorylation of JNK, p38, or Akt. Because Akt is possibly upstream of ERK [43], A549 and H460 cells were treated with CCL21 for 24 h after a 1-h exposure to LY294002, a selective inhibitor of PI3K that inhibits activation of the downstream Akt pathway. Following this treatment, CCL21/CCR7 still significantly upregulated the expression of P-ERK (Figure 6).

Inhibition of P-ERK abolishes the impellent effects of CCL21/CCR7 on proliferation and G₂/M phase progression. To verify whether PD98059, a selective inhibitor of MEK that disrupts activation of downstream ERK, can abolish the effects of CCL21/CCR7 on the proliferation and G₂/M phase progression of A549 and H460 cells, cell viability and cell cycle distribution assays were performed using CCK-8 and flow cytometry, respectively. PD98059 significantly abrogated the effects of CCL21/CCR7 on cell proliferation and the G₂/M phase progression (Figure 7 and Table 2, respectively). PD98059 also abolished the influence of CCL21/CCR7 on the expression of P-ERK, cyclin A, cyclin B1, and CDK1 (Figure 8). In addition, PD98059 alone had a significant inhibitory effect on the cell proliferation, the G₂/M phase progression and the expression of P-ERK, cyclin A, and cyclin B1 (Figure 7, Table 2, and Figure 8, respectively).

P-ERK, induced by CCL21/CCR7, interacts with cyclin A, cyclin B1, and CDK1. To further identify whether there is an interaction between P-ERK and cyclin A, cyclin B1, or CDK1, coimmunoprecipitation was performed. A549 and H460 cells in the absence or presence of CCL21 for 24 h were subjected to immunoprecipitation with antibodies against P-ERK or IgG, followed by Western blotting for cyclin A, cyclin B1, and CDK1. A pronounced, specific interaction between P-ERK and cyclin A, cyclin B1, or CDK1 was observed, especially when the cells were treated with CCL21 for 24 h (Figure 9a). Reciprocal immunoprecipitation with antibodies against cyclin A, cyclin B1, CDK1, or IgG was assessed by Western blotting for P-ERK, and again, the interaction between P-ERK and cyclin A, cyclin B1, or CDK1 was salient, especially in the presence of CCL21 (Figure 9b). A549 and H460 cells in the absence or presence of PD98059 for 1 h were subjected to immunoprecipitation with antibodies against P-ERK or IgG, followed by Western blotting for cyclin A, cyclin B1,




Figure 2. Effect of CCL21 at various concentrations on proliferation of A549 and H460 cells. A549 (A) and H460 (B) cells were treated with CCL21 (50, 100 or 200 ng/mL) for 24, 48, or 72 h, and cell vitality was estimated using the CCK-8 assay. Each bar represents the mean \pm SD of three independent experiments. ** $p < 0.01$, compared with cells treated with CCL21 (50 ng/mL). doi:10.1371/journal.pone.0021119.g002




Figure 3. Effect of CCL21/CCR7 on proliferation of A549 and H460 cells. A549 (A) and H460 (B) cells were treated with CCL21 (100 ng/mL) for 24, 48, or 72 h, and cell vitality was estimated using the CCK-8 assay. Each bar represents the mean \pm SD of three independent experiments. ** $p < 0.01$, compared with control cells. doi:10.1371/journal.pone.0021119.g003

and CDK1. The interaction between P-ERK and cyclin A, cyclin B1, or CDK1 was weakened in response to PD98059 exposure (Figure 9c).

Discussion

Several studies have documented that the activation of CCR7 is responsible for mediating survival of certain cancer cell lines by promoting migration and proliferation or by inhibiting apoptosis [33,35–42]. However, the role of CCR7 in the proliferation of human NSCLC cells has not been well documented. In the present study, we confirmed that the CCL21/CCR7 interaction can significantly enhance human NSCLC cell proliferation in a time-dependent manner, involving the upregulation of cyclin A, cyclin B1, and CDK1, possibly via the ERK, but not the Akt, pathway.

Consistent with previous studies using different cell models [16,37,40], the activation of CCR7 with CCL21 promoted cell proliferation. In contrast to our current findings, a prior study suggested that murine CCL21 had no significant action on the proliferation of A549 cells [44]. A possible explanation for the discrepancy would be that mouse CCL21 can interact with CXCR3 but not CCR7, whereas human CCL21 can bind CCR7 but not CXCR3 [45]. This suggests that activation of CXCR3 does not affect proliferation of A549 cells.

Traditionally, the cell cycle is segregated into four phases: DNA replication occurs during S phase, and chromosome segregation occurs during M phase. The S and M phases are separated by the so-called gap phases, G₁ (before DNA replication) and G₂ (before mitosis). We demonstrated that CCL21/CCR7 significantly altered cell cycle distribution such that more cells populated the G₂/M phase. No significant differences were observed regarding the distribution of cells in the G₀/G₁ and S phases.

The cell cycle is regulated by cyclins and CDKs. D-type cyclins are regarded as key regulators of G₁ progression [46]. Cyclin E is required for the G₁/S transition [47,48] and cyclin A is essential for progression through the S phase [49,50]. Both cyclin A and the B-type cyclins associate with CDK1 to promote entry into mitosis [51,52]. In this study, both protein and mRNA levels of cyclin A, cyclin B1, and CDK1 were significantly upregulated when cells were treated with CCL21 for 24 h. This indicates that CCL21/CCR7 accelerates the G₂/M phase progression to promote cell proliferation. No significant differences in cyclin D1 or cyclin E expression levels were measured, indicating that CCL21/CCR7 has no significant effect on the G₀/G₁ phase or the G₁/S transition. This finding demonstrates for the first time that CCL21/CCR7 has an impellent effect on cell cycle progression involving the G₂/M phase.

Several studies using other cell types have indicated that activation of CCR7 is associated with enhanced cell survival via increased phosphorylation of ERK, JNK, or Akt [16–20,33,35–42]. We examined whether CCL21/CCR7 may enhance the expression of MAPK components and Akt in A549 and H460 cells. Our results suggested that CCL21/CCR7 enhanced the phosphorylation of ERK but not JNK, p38, or Akt. These findings are inconsistent with previously published reports that suggested a

Table 1. Effect of CCL21/CCR7 on cell cycle distribution in A549 and H460 cells.

Group	G ₀ /G ₁ phase	S phase	G ₂ /M phase
	Mean \pm SD (%)	Mean \pm SD (%)	Mean \pm SD (%)
(A549) Control	74.35 \pm 9.67	23.42 \pm 12.29	2.23 \pm 2.82
(A549) CCL21	66.92 \pm 1.23	21.60 \pm 2.91	11.48 \pm 2.64*
(A549) siCCR7-CCL21	69.08 \pm 10.08	23.90 \pm 9.28	6.60 \pm 2.46
(A549) siCCR7	75.73 \pm 2.32	18.09 \pm 3.67	6.19 \pm 2.45
(H460) Control	61.50 \pm 2.70	31.33 \pm 3.41	7.17 \pm 1.63
(H460) CCL21	60.38 \pm 6.85	27.08 \pm 4.13	12.54 \pm 2.75*
(H460) siCCR7-CCL21	69.25 \pm 4.99	21.40 \pm 3.23	6.02 \pm 2.53
(H460) siCCR7	67.57 \pm 2.40	28.00 \pm 4.10	4.63 \pm 1.20

A549 and H460 cells were treated with CCL21 (100 ng/mL) for 24 h, and cell cycle distribution was estimated using flow cytometry. Data were present as mean \pm SD of three independent experiments.

* $p < 0.05$, compared with control cells.

doi:10.1371/journal.pone.0021119.t001




Figure 4. Effect of CCL21/CCR7 on the expression of cyclins and cyclin-dependent kinases in A549 and H460 cells. A549 (A) and H460 (B) cells were treated with CCL21 (100 ng/mL) for 24 h, and the protein (a) and mRNA (b) levels of cyclins A, B1, D1, and E and of CDK1 were estimated using Western blot (a) and real-time PCR (b). Each bar represents the mean ± SD of three independent experiments. *p<0.05 or **p<0.01, compared with control cells. doi:10.1371/journal.pone.0021119.g004

facilitative effect of CCR7 on the Akt, but not ERK, pathway [16,19]. A separate study indicated that Akt is possibly upstream of ERK [43]. We found that CCL21/CCR7 still could significantly

upregulate the expression of P-ERK after the cells were treated with LY294002 for 1 h. Because LY294002 selectively inhibits PI3K to prevent downstream activation of Akt, our results strongly




Figure 5. Effect of CCL21/CCR7 on the expression of ERK, JNK, p38, and Akt in A549 and H460 cells. A549 (A) and H460 (B) cells were treated with CCL21 (100 ng/mL) for 12, 24, or 48 h, and normal and phosphorylated (P-) expression levels were estimated by Western blot. Each bar represents the mean ± SD of three independent experiments. *p<0.05 or **p<0.01, compared with control cells. doi:10.1371/journal.pone.0021119.g005




Figure 6. Effects of CCL21/CCR7 on the expression of P-ERK after inhibiting activation of Akt in A549 and H460 cells. A549 (A) and H460 (B) cells were treated with CCL21 (100 ng/mL) for 24 h after LY294002, a selective inhibitor of PI3K that consequently prevents activation of the downstream Akt, was applied for 1 h. The expression of P-ERK was estimated using Western blot. Each bar represents the mean \pm SD of three independent experiments. ** $p < 0.01$, compared with control cells. doi:10.1371/journal.pone.0021119.g006




Figure 7. Effect of CCL21/CCR7 on proliferation of A549 and H460 cells after inhibiting ERK activation. A549 (A) and H460 (B) cells were treated with CCL21 (100 ng/mL) for 24, 48, or 72 h after PD98059, a selective inhibitor of MEK that disrupts activation of downstream ERK, was applied for 1 h. Cell vitality was estimated using the CCK-8 assay. Each bar represents the mean \pm SD of three independent experiments. * $p < 0.05$ or ** $p < 0.01$, compared with control cells. doi:10.1371/journal.pone.0021119.g007

Table 2. Effect of CCL21/CCR7 on cell cycle distribution after inhibiting ERK in A549 and H460 cells.

Group	G ₀ /G ₁ phase	S phase	G ₂ /M phase
	Mean ± SD (%)	Mean ± SD (%)	Mean ± SD (%)
(A549) Control	80.49±0.67	14.57±1.48	4.94±0.91
(A549) CCL21	68.58±3.17	20.21±0.40	11.48±2.64**
(A549) PD98059-CCL21	73.06±6.94	22.11±7.42	4.84±0.58
(A549) PD98059	76.65±11.28	24.64±5.83	2.05±0.31*
(H460) Control	65.73±1.72	27.95±0.94	6.33±1.76
(H460) CCL21	58.58±8.68	29.41±6.16	12.01±2.70**
(H460) PD98059-CCL21	67.80±3.00	26.91±3.17	5.29±1.29
(H460) PD98059	67.07±2.17	30.50±3.02	2.43±1.18*

A549 and H460 cells were treated with CCL21 (100 ng/mL) for 24 h after PD98059, a selective inhibitor of MEK that disrupts activation of downstream ERK, was used for 1 h. After treatment, cell cycle distribution was estimated using flow cytometry. Data were present as mean ± SD of three independent experiments.

*p<0.05 or

**p<0.01, compared with control cells.

doi:10.1371/journal.pone.0021119.t002

suggest that Akt does not play a critical role in the interaction between CCL21/CCR7 and ERK. This is in concordance with a previous report [53]. The reason for these discrepancies remains unclear.

Since CCL21/CCR7 was capable of increasing the expression of P-ERK, we sought to determine whether there is an interaction between P-ERK and cyclin A, cyclin B1, or CDK1. Coimmunoprecipitation and reciprocal immunoprecipitation results strongly suggested an interaction between P-ERK and cyclin A, cyclin B1, or CDK1, especially in the presence of CCL21. This interaction could be weakened by inhibiting ERK with PD98059. In addition, PD98059 abolished the effect of CCL21/CCR7 on A549 and H460 cell proliferation and G₂/M phase progression, as well as downregulating the expression of P-ERK, cyclin A, cyclin B1, and CDK1. These results demonstrate that the effect of CCL21/CCR7 on cell proliferation and upregulation of cyclin A, cyclin B1, and CDK1 may occur via the ERK pathway in human NSCLC cells.

This study suggests that activation of CCR7 with CCL21 can significantly promote proliferation of NSCLC cells in a time-dependent manner involving cyclin A, cyclin B1, and CDK1, possibly via the ERK, but not the Akt, pathway. This information may help clarify the mechanisms of cancer cell survival and identify potential targets for treatment of NSCLC.

Materials and Methods

Cell culture and reagents

A549 and H460 cell lines from our previous published paper [32] were cultured in RPMI-1640 or DMEM-F12 supplemented with 10% HyClone fetal bovine serum (FBS) (ThermoFisher Scientific, Fremont, CA, USA) in an atmosphere of 5% CO₂ at 37°C. Cells were grown in 75 cm² culture flasks and harvested in a solution of trypsin-EDTA at the logarithmic growth phase.

Cyclin A, cyclin B1, cyclin D1, cyclin E, CDK1, P-ERK, ERK, P-JNK, JNK, P-p38, p38, P-Akt, Akt, IgG, and β-actin mouse or rabbit monoclonal antibodies were purchased from Santa Cruz Biotechnology (Santa Cruz, CA, USA). Recombinant human

CCL21 was purchased from Pepro Tech (Rocky Hill, NJ, USA). siCCR7 and Lipofectamine 2000 were purchased from Invitrogen (Carlsbad, CA, USA). Cell Counting Kit-8 (CCK-8) was purchased from Dojindo Laboratories (Beijing, China). PD98059 and LY294002 were obtained from Sigma (St. Louis, MO, USA) and used at 50 μM or 100 μM (final concentrations), respectively, in accordance with previous reports [40,54]. Protein A/G beads were obtained from Beyotime (Haimen, China).

siRNA treatment of cells

A549 and H460 cells were plated onto 6 cm² cell culture dishes and grown to 30–50% confluence before transfection with Lipofectamine 2000 as previously described [55]. The transfection efficiency was assessed by flow cytometry. Efficiencies of siCCR7 and non-silencing control siRNA were tested using Western blot and RT-PCR. The sequences of siCCR7 and control siRNA were: CCR7, 5'-GCGUCAACCCUUCUUGUATT-3' and 3'-UA-CAAGAAAGGGUUGACGCAG-5'; control, 5'-UUCUCCGA-ACGUGUCACGUTT-3' and 3'-ACGUGACACGUUCGGA-GAATT-5'.

Cell proliferation assay

Cell proliferative activities were examined using CCK-8. A549 and H460 cells were seeded onto 96-well plates (1000 cells/well) and treated with CCL21 for 0, 24, 48, or 72 h. After treatment, CCK-8 was added to each well according to the manufacturer's instructions and incubated for 4 h at 37°C. The optical density (OD) value of each well was measured using a microplate reader (Spectra Thermo, Männedorf, Switzerland) with a test wavelength of 450 nm.


Cell cycle analysis

After treatment with CCL21 for 24 h, cells were harvested and washed twice with cold phosphate-buffered saline (PBS) and fixed in 75% ethanol for 2 h at 4°C. The fixed cells were washed twice with 500 μL of cold PBS. Cells then were stained with 500 μL of propidium iodide (PI) staining solution (50 μg/mL PI, 0.1% Triton X-100, 200 μg/mL DNase-free RNase in PBS) for 30 min at room temperature in the dark. Ten thousand events per sample were acquired using a FACS-scan flow cytometer (Becton-Dickinson, San Jose, CA, USA), and the percentage of cells in G₀/G₁, S, and G₂/M phases of the cell cycle were determined using Modfit LT 3.0 (Becton-Dickinson).

Western blot analysis

After treatment with CCL21 for 24 h, cells were extracted with lysis buffer (150 mM NaCl, 1% NP-40, 0.1% SDS, 2 μg/mL aprotinin and 1 mM PMSF) for 30 min at 4°C. Extracts were centrifuged at 12,000× g for 15 min at 4°C. Supernatants containing total protein then were harvested. Aliquots, each containing 50 μg protein, were separated by 10% SDS-PAGE and transferred to PVDF membranes at 55 V (cyclin A, cyclin B1, P-Akt, Akt) or 40 V (the others) for 2.5 h at low temperature. The membranes were blocked in 5% skim milk for 2 h, and proteins were detected using monoclonal antibodies at 1:1000 (P-JNK, JNK, P-p38, p38 and β-actin) or 1:200 (the others) dilution overnight at 4°C. Proteins were visualized using anti-mouse or anti-rabbit IgG conjugated with horse radish peroxidase (HRP) at 1:6000 or 1:8000 dilution for 2 h at room temperature, respectively. Bands were imaged with an EC3 Imaging System (UVP LLC, Upland, CA, USA), and the OD was measured using ImageJ (NIH, Bethesda, MD, USA). The OD difference between tested proteins and β-actin of the same sample was calculated as relative content and expressed graphically.

A



B




Figure 8. Effect of CCL21/CCR7 on the expression of P-ERK, cyclin A, cyclin B1, and CDK1 after inhibiting the activation of ERK. A549 (A) and H460 (B) cells were treated with CCL21 (100 ng/mL) for 24 h after PD98059, a selective inhibitor of MEK that disrupts activation of downstream ERK, was applied for 1 h. The expression levels of these components were estimated using Western blot. Each bar represents the mean \pm SD of three independent experiments. * $p < 0.05$ or ** $p < 0.01$, compared with control cells. doi:10.1371/journal.pone.0021119.g008

RT-PCR and real-time PCR

Total RNA was isolated from cells using TRIzol (Invitrogen) according to the manufacturer's instructions. β -Actin was used as an internal control. To determine the efficiencies of siCCR7 and control siRNA, semi-quantitative RT-PCR was performed on a G-STORM thermal cycler (GRI Ltd, Byfleet, UK) using the TaKaRa RNA PCR Kit (AMV) Ver.3.0 (TaKaRa, Dalian, China). PCR primers were as follows: CCR7 F: 5'-GAGGCTATTGTCCCCTAAACC-3', R: 5'-TGGAGGACAGTGAAGAAAACG-3'. The CCR7 amplicon length was 305 bp. β -actin F: 5'-AAATCGTGCGTGACATTAA-3', R: 5'-CTCGTCATACTCCTGCTTG-3'. The β -actin amplicon length was 513 bp. The heat cycling conditions were as follows: 30 cycles of 40 s each, including denaturation at 95°C, annealing at 53°C, and extension at 72°C.

Real-time PCR was performed on an ABI Prism 7900HT Fast System (Applied Biosystems, Foster, CA, USA) using SYBR

Premix Ex Taq II (TaKaRa). Amplifications were carried out in a total volume of 20 μ L and cycled 40 times after initial denaturation (95°C for 30 s) with the following parameters: 95°C for 5 s and 60°C for 30 s. β -actin F: 5'-AGCACAGAGCCTCGCCTTTG-3', R: 5'-ACATGCCGAGCCGTTGT-3'. The β -actin amplicon length was 107 bp. Other primer sequences have been published in another study [56]. The reliability of PCR results was supported by analyzing the dissociation curve. Real-time PCR data were calculated using the $2^{-\Delta\Delta CT}$ method on the SDS 2.4 software package (Applied Biosystems) [57].

Coimmunoprecipitation

After treatment with CCL21 for 24 h, cells were extracted with lysis buffer (10 mM KCl, 1.5 mM MgCl₂, 10 mM HEPES [pH 7.9], 1 mM PMSF, 1 mM DTT) and homogenized for




Figure 9. The interaction between P-ERK and cyclin A, cyclin B1, or CDK1 in the absence or presence of CCL21 or PD98059. A549 (A) and H460 (B) cells in the absence or presence of CCL21 (100 ng/mL) for 24 h were subjected to immunoprecipitation with antibodies against P-ERK or IgG, followed by Western blotting for cyclin A, cyclin B1, and CDK1 (a). Reciprocal immunoprecipitation with antibodies against cyclin A, cyclin B1, CDK1, or IgG were analyzed by Western blotting for P-ERK (b). A549 and H460 cells in the absence or presence of PD98059 for 1 h were subjected to immunoprecipitation with P-ERK or IgG antibodies followed by Western blotting for cyclin A, cyclin B1, and CDK1 (c). doi:10.1371/journal.pone.0021119.g009

30 min at 4°C. The extracts were centrifuged at 12,000× *g* for 15 min at 4°C, and the supernatants containing total protein were harvested. Equal amounts of protein were exposed to antibodies against P-ERK, IgG, cyclin A, cyclin B1, or CDK1, which were immobilized on protein A/G beads. Following 3 h incubation at 4°C with gentle rotation, beads were washed extensively five times with lysis buffer, boiled, and microcentrifuged. Proteins were detected with antibodies against P-ERK, cyclin A, cyclin B1, or CDK1 by Western blot.

Statistical analysis

Data were analyzed using SPSS 16.0 software. One-way analysis of variance (ANOVA) was used to evaluate the differences between groups with various treatments, and the least significant

difference (LSD) test or Dunnett T3 test was used for post hoc subgroup analysis. Polynomial contrast was used for trend analysis. All data are presented as the mean ± SD of three independent experiments. Results were considered statistically significant for $p < 0.05$. N-fold values for gene expression change up to 0.5 and below 2 were taken as nonsignificant in accordance with values obtained from negative control genes [57,58].

Author Contributions

Conceived and designed the experiments: YX XQ. Performed the experiments: YX LL IJ BH HL ZL WL EW. Analyzed the data: YX LL. Contributed reagents/materials/analysis tools: XQ. Wrote the paper: YX.

References

1. Baggiolini M (1998) Chemokines and leukocyte traffic. *Nature* 392: 565–8.
2. Zlotnik A, Yoshie O (2000) Chemokines: a new classification system and their role in immunity. *Immunity* 12: 121–7.
3. Dieu MC, Vanbervliet B, Vicari A, Bridon JM, Oldham E, et al. (1998) Selective recruitment of immature and mature dendritic cells by distinct chemokines expressed in different anatomic sites. *J Exp Med* 188: 373–86.
4. Hirao M, Onai N, Hiroishi K, Watkins SC, Matsushima K, et al. (2000) CC chemokine receptor-7 on dendritic cells is induced after interaction with apoptotic tumor cells: critical role in migration from the tumor site to draining lymph nodes. *Cancer Res* 60: 2209–17.
5. Sozzani S, Allavena P, D'Amico G, Luini W, Bianchi G, et al. (1998) Differential regulation of chemokine receptors during dendritic cell maturation: a model for their trafficking properties. *J Immunol* 161: 1083–6.
6. Muller A, Homey B, Soto N, Ge N, Catron D, et al. (2001) Involvement of chemokine receptors in breast cancer metastasis. *Nature* 410: 50–6.
7. Takanami I (2003) Overexpression of CCR7 mRNA in nonsmall cell lung cancer: correlation with lymph node metastasis. *Int J Cancer* 105: 186–9.
8. Ding Y, Shimada Y, Maeda M, Kawabe A, Kaganoi J, et al. (2003) Association of CC chemokine receptor 7 with lymph node metastasis of esophageal squamous cell carcinoma. *Clin Cancer Res* 9: 3406–12.
9. Wang J, Xi L, Hunt JL, Gooding W, Whiteside TL, et al. (2004) Expression pattern of chemokine receptor 6 (CCR6) and CCR7 in squamous cell carcinoma of the head and neck identifies a novel metastatic phenotype. *Cancer Res* 64: 1861–6.
10. Takeuchi H, Fujimoto A, Tanaka M, Yamano T, Hsueh E, et al. (2004) CCL21 chemokine regulates chemokine receptor CCR7 bearing malignant melanoma cells. *Clin Cancer Res* 10: 2351–8.
11. Mori T, Kim J, Yamano T, Takeuchi H, Huang S, et al. (2005) Epigenetic up-regulation of C-C chemokine receptor 7 and C-X-C chemokine receptor 4 expression in melanoma cells. *Cancer Res* 65: 1800–7.
12. Yu S, Duan J, Zhou Z, Pang Q, Wuyang J, et al. (2008) A Critical Role of CCR7 in Invasiveness and metastasis of SW620 Colon Cancer Cell in Vitro and in Vivo. *Cancer Biol Ther* 7: 1037–43.
13. Ishigami S, Natsugoe S, Nakajo A, Tokuda K, Uenosono Y, et al. (2007) Prognostic value of CCR7 expression in gastric cancer. *Hepatogastroenterology* 54: 1025–8.
14. Kodama J, Hasengaowa, Kusumoto T, Seki N, Matsuo T, et al. (2007) Association of CXCR4 and CCR7 chemokine receptor expression and lymph node metastasis in human cervical cancer. *Ann Oncol* 18: 70–6.
15. Wiley HE, Gonzalez EB, Maki W, Wu MT, Hwang ST (2001) Expression of CC chemokine receptor-7 and regional lymph node metastasis of B16 murine melanoma. *J Natl Cancer Inst* 93: 1638–43.
16. Wang J, Zhang X, Thomas SM, Grandis JR, Wells A, et al. (2005) Chemokine receptor 7 activates phosphoinositide-3 kinase-mediated invasive and pro-survival

- pathways in head and neck cancer cells independent of EGFR. *Oncogene* 24: 5897–904.
17. Li Z, Li J, Mo B, Hu C, Liu H, et al. (2008) Genistein induces G₂/M cell cycle arrest via stable activation of ERK1/2 pathway in MDA-MB-231 breast cancer cells. *Cell Biol Toxicol* 24: 401–9.
 18. Wu Y, Peng H, Cui M, Whitney NP, Huang Y, et al. (2009) CXCL12 increases human neural progenitor cell proliferation through Akt-1/FOXO3a signaling pathway. *J Neurochem* 109: 1157–67.
 19. Sánchez-Sánchez N, Riol-Blanco L, de la Rosa G, Puig-Kröger A, García-Bordas J, et al. (2004) Chemokine receptor CCR7 induces intracellular signaling that inhibits apoptosis of mature dendritic cells. *Blood* 104: 619–25.
 20. Jackson ML, Neuzil KM, Thompson WW, Shay DK, Yu O, et al. (2004) The burden of community-acquired pneumonia in seniors: results of a population-based study. *Clin Infect Dis* 39: 1642–50.
 21. Chang L, Karin M (2001) Mammalian MAP kinase signalling cascades. *Nature* 410: 37–40.
 22. Johnson GL, Lapadat R (2002) Mitogen-activated protein kinase pathways mediated by ERK, JNK, and p38 protein kinases. *Science* 298: 1911–2.
 23. Wright JH, Munar E, Jameson DR, Andreassen PR, Margolis RL, et al. (1999) Mitogen-activated protein kinase activity is required for the G₂/M transition of the cell cycle in mammalian fibroblasts. *Proc Natl Acad Sci U S A* 96: 11335–40.
 24. Hayne C, Tzivion G, Luo Z (2000) Raf-1/MEK/MAPK pathway is necessary for the G₂/M transition induced by nocodazole. *J Biol Chem* 275: 31876–82.
 25. Davis RJ (2000) Signal transduction by the JNK group of MAP kinases. *Cell* 103: 239–52.
 26. Minden A, Karin M (1997) Regulation and function of the JNK subgroup of MAP kinases. *Biochim Biophys Acta* 1333: F85–104.
 27. Roulston A, Reinhard C, Amiri P, Williams LT (1998) Early activation of c-Jun N-terminal kinase and p38 kinase regulate cell survival in response to tumor necrosis factor alpha. *J Biol Chem* 273: 10232–9.
 28. Kulik G, Klippel A, Weber MJ (1997) Antiapoptotic signalling by the insulin-like growth factor I receptor, phosphatidylinositol 3-kinase, and Akt. *Mol Cell Biol* 17: 1595–606.
 29. Kauffmann-Zeh A, Rodriguez-Viciana P, Ulrich E, Gilbert C, Coffer P, et al. (1997) Suppression of c-Myc-induced apoptosis by Ras signalling through PI(3)K and PKB. *Nature* 385: 544–8.
 30. Dudek H, Datta SR, Franke TF, Birnbaum MJ, Yao R, et al. (1997) Regulation of neuronal survival by the serine-threonine protein kinase Akt. *Science* 275: 661–5.
 31. Brunet A, Bonni A, Zigmond MJ, Lin MZ, Juo P, et al. (1999) Akt promotes cell survival by phosphorylating and inhibiting a Forkhead transcription factor. *Cell* 96: 857–68.
 32. Li Y, Qiu X, Zhang S, Zhang Q, Wang E (2009) Hypoxia induced CCR7 expression via HIF-1alpha and HIF-2alpha correlates with migration and invasion in lung cancer cells. *Cancer Biol Ther* 8: 322–30.
 33. Banas B, Wörnle M, Merkle M, Gonzalez-Rubio M, Schmid H, et al. (2004) Binding of the chemokine SLC/CCL21 to its receptor CCR7 increases adhesive properties of human mesangial cells. *Kidney Int* 66: 2256–63.
 34. Wörnle M, Schmid H, Merkle M, Banas B (2004) Effects of chemokines on proliferation and apoptosis of human mesangial cells. *BMC Nephrol* 5: 8.
 35. Kim JW, Ferris RL, Whiteside TL (2005) Chemokine C receptor 7 expression and protection of circulating CD8⁺ T lymphocytes from apoptosis. *Clin Cancer Res* 11: 7901–10.
 36. Escribano C, Delgado-Martín C, Rodríguez-Fernández JL. CCR7-dependent stimulation of survival in dendritic cells involves inhibition of GSK3beta. *J Immunol* 2009; 183: 6282–95.
 37. Gollmer K, Asperti-Boursin F, Tanaka Y, Okkenhaug K, Vanhaesebroeck B, et al. (2009) CCL21 mediates CD4⁺ T-cell costimulation via a DOCK2/Rac-dependent pathway. *Blood* 114: 580–8.
 38. Shannon LA, Calloway PA, Welch TP, Vines CM (2010) CCR7/CCL21 migration on fibronectin is mediated by PLC{gamma}1 and ERK1/2 in primary T lymphocytes. *J Biol Chem* 285: 38781–7.
 39. Iijima N, Yanagawa Y, Clingan JM, Onoć K (2005) CCR7-mediated c-Jun N-terminal kinase activation regulates cell migration in mature dendritic cells. *Int Immunol* 17: 1201–12.
 40. Liu FY, Zhao ZJ, Li P, Ding X, Zong ZH, et al. (2010) Mammalian target of rapamycin (mTOR) is involved in the survival of cells mediated by chemokine receptor 7 through PI3K/Akt in metastatic squamous cell carcinoma of the head and neck. *Br J Oral Maxillofac Surg* 48: 291–6.
 41. Lee SJ, Park K, Ha SD, Kim WJ, Moon SK (2010) Gleditsia sinensis thorn extract inhibits human colon cancer cells: the role of ERK1/2, G₂/M-phase cell cycle arrest and p53 expression. *Phytother Res* 24: 1870–6.
 42. Badr G, Borhis G, Treton D, Richard Y (2005) IFN{alpha} enhances human B-cell chemotaxis by modulating ligand-induced chemokine receptor signaling and internalization. *Int Immunol* 17: 459–67.
 43. Curmook AP, Logan MK, Ward SG (2002) Chemokine signalling: pivoting around multiple phosphoinositide 3-kinases. *Immunology* 105: 125–36.
 44. Arenberg DA, Zlotnick A, Strom SR, Burdick MD, Strieter RM (2001) The murine CC chemokine, 6C-kine, inhibits tumor growth and angiogenesis in a human lung cancer SCID mouse model. *Cancer Immunol Immunother* 249: 587–92.
 45. Jenh CH, Cox MA, Kaminski H, Zhang M, Byrnes H, et al. (1999) Cutting edge: species specificity of the CC chemokine 6Ckine signaling through the CXCR3 chemokine receptor CXCR3: human 6Ckine is not a ligand for the human or mouse CXCR3 receptors. *J Immunol* 162: 3765–9.
 46. Nigg EA (1995) Cyclin-dependent protein kinases: key regulators of the eukaryotic cell cycle. *Bioessays* 17: 471–80.
 47. Knoblich JA, Sauer K, Jones L, Richardson H, Saint R, et al. (1994) Cyclin E controls S phase progression and its down-regulation during Drosophila embryogenesis is required for the arrest of cell proliferation. *Cell* 77: 107–20.
 48. Ohtsubo M, Roberts JM (1993) Cyclin-dependent regulation of G1 in mammalian fibroblasts. *Science* 259: 1908–12.
 49. Girard F, Strausfeld U, Fernandez A, Lamb NJ (1991) Cyclin A is required for the onset of DNA replication in mammalian fibroblasts. *Cell* 67: 1169–79.
 50. Pagano M, Pepperkok R, Verde F, Ansorge W, Draetta G (1992) Cyclin A is required at two points in the human cell cycle. *EMBO J* 11: 961–71.
 51. King RW, Jackson PK, Kirschner MW (1994) Mitosis in transition. *Cell* 79: 563–71.
 52. Knoblich JA, Lehner CF (1993) Synergistic action of Drosophila cyclins A and B during the G₂-M transition. *EMBO J* 12: 65–74.
 53. Riol-Blanco L, Sánchez-Sánchez N, Torres A, Tejedor A, Narumiya S, et al. (2005) The chemokine receptor CCR7 activates in dendritic cells two signaling modules that independently regulate chemotaxis and migratory speed. *J Immunol* 174: 4070–80.
 54. Wegiel B, Bjartell A, Culig Z, Persson JL (2008) Interleukin-6 activates PI3K/Akt pathway and regulates cyclin A1 to promote prostate cancer cell survival. *Int J Cancer* 122: 1521–9.
 55. Rudd BD, Burstein E, Duckett CS, Li X, Lukacs NW (2005) Differential role for TLR3 in respiratory syncytial virus-induced chemokine expression. *J Virol* 79: 3350–7.
 56. Liu L, Zhu Y, Xu Y, Reiter RJ (2011) Melatonin delays cell proliferation by inducing G(1) and G(2)/M phase arrest in a human osteoblastic cell line hFOB 1.19. *J Pineal Res* 50: 222–31.
 57. Livak KJ, Schmittgen TD (2001) Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods* 25: 402–8.
 58. Pizarro JG, Yeste-Velasco M, Esparza JL, Verdaguier E, Pallás M, et al. (2008) The antiproliferative activity of melatonin in B6 rat dopaminergic neuroblastoma cells is related to the downregulation of cell cycle-related genes. *J Pineal Res* 45: 8–16.