

# Postpartum Hemorrhage and its Associated Factors Among Women who Gave Birth at Yirgalem General Hospital, Sidama Regional State, Ethiopia

Health Services Research and Managerial Epidemiology  
 Volume 8: 1-7  
 © The Author(s) 2021  
 Article reuse guidelines:  
[sagepub.com/journals-permissions](https://sagepub.com/journals-permissions)  
 DOI: 10.1177/23333928211062777  
[journals.sagepub.com/home/hme](https://journals.sagepub.com/home/hme)



Tedla Amanuel<sup>1</sup>, Azmach Dache<sup>2</sup> , and Aregahegn Dona<sup>2</sup>

## Abstract

**Background:** Globally, postpartum hemorrhage (PPH) is one of the leading causes of maternal mortality. In developing countries, it accounts for more than 30% of all maternal deaths. So, understanding its burden in the health care setting is significant. Thus, this study aimed to assess the magnitude of PPH and its associated factors among women who gave birth at Yirgalem General Hospital, Sidama Region, Ethiopia.

**Methods:** A cross-sectional study was conducted from March 12 to 26, 2020 among randomly selected 298 women. Data were collected using an interviewer-administered, structured, and pretested questionnaire. EpiData version 3.1 and SPSS version 20 were used to enter and analyze the data, respectively. Descriptive statistics, bivariable, and multivariable logistic regression analysis were done. Adjusted odds ratio with 95% confidence interval (CI) was used to measure the presence and strength of association between the independent and the outcome variables. A *P*-value  $\leq .05$  was considered to declare statistical significance.

**Result:** The magnitude of PPH was 9.4% [95% CI: 6.0, 12.8]. Prolonged labor ( $\geq 24$  h) [AOR = 3.4, 95% CI: 1.1, 9.9], giving birth by cesarean section [AOR = 5.8, 95% CI: 1.1, 22.0], and instrumental vaginal delivery [AOR = 3.7, 95% CI: 1.1, 12.7], and having a history of the uterine atony [AOR = 4.8, 95% CI: 1.4, 16.6] during their last delivery were factors significantly associated with PPH.

**Conclusion:** The magnitude of PPH was high. Healthcare professionals should manage the progress of labor and take all necessary measures at right time. Also, giving attention to the safety of delivery-related procedures and early related potential risks is crucial.

## Keywords

postpartum hemorrhage, Yirgalem, Ethiopia

## Background

According to the World health organization (WHO), postpartum hemorrhage (PPH) is defined as the blood loss of more than 500 mm following a vaginal delivery or more than 1000 mm following cesarean section. Also, it is defined as any amount of vaginal bleeding following delivery that causes vital sign derangement or loss of 10% hemoglobin from the baseline.<sup>1</sup>

PPH is termed as primary when it occurs within 24 h of delivery, whereas termed late or secondary PPH when it occurs after 24 h to 6 weeks of delivery.<sup>2,3</sup>

PPH is the major cause of maternal mortality and morbidity across the world, responsible for more than 25% of deaths

annually. WHO statistics suggested that 60% of maternal deaths in developing countries were due to PPH, accounting for more than 100,000 maternal deaths per year worldwide.<sup>4</sup>

<sup>1</sup> Loka Abaya District Health Office, Sidama Regional State, Hawassa, Ethiopia

<sup>2</sup> Yirgalem Hospital Medical College, Yirgalem, Sidama, Ethiopia

Submitted September 24, 2021. Revised October 26, 2021. Accepted November 4, 2021.

## Corresponding Author:

Aregahegn Dona, Department of Social and Population Health, Yirgalem Hospital Medical College, Yirgalem 184, Sidama, Ethiopia.  
 Email: aregahegndona@gmail.com



Creative Commons Non Commercial CC BY-NC: This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits non-commercial use, reproduction and distribution of the work without further permission provided the original work is attributed as specified on the SAGE and Open Access page (<https://us.sagepub.com/en-us/nam/open-access-at-sage>).

PPH remains the leading cause of maternal mortality, exacting huge tolls on economic and human capital, particularly in those countries with inadequate resources. Also, it causes serious consequences like severe anemia, shock, and death particularly in poor countries where women do not have access to quality health care services due to financial constraints.<sup>5</sup>

The major risk factors that contribute to PPH are uterine atony (failure of the uterus to contract adequately after childbirth) and a prolonged third phase of labor, retained placenta, maternal age of above 35 years, gestational age below 37 weeks or above 41 weeks, pregnancy-induced hypertension, anemia, having previous history of PPH, uterine rupture, abruptio placenta, placenta previa.<sup>6,7</sup>

Similarly, the number of pregnancy, mode of delivery (cesarean section), not attending antenatal care during pregnancy, not receiving adequate iron-folate supplementation, episiotomy, and any tear or injury to genital tract are factors related to manifestation of PPH.<sup>8-10</sup>

In African and Asian countries, where most maternal deaths occurred, PPH alone accounts for more than 30% of all maternal deaths. However, the figures of maternal death attributed to PPH differ when compared between low-income and developed countries.<sup>11</sup> In Sub-Saharan Africa, the probability of dying during child birth is significantly higher, and uterine atony (failure of the uterus to contract adequately after childbirth) accounting for 60% to 80% of PPH cases.<sup>12,13</sup>

The high magnitude of PPH and its bad outcome is still a major health problem in developing countries including Sub-Saharan African countries. Ethiopia is one of the countries with a high burden of postpartum maternal morbidity.<sup>14</sup> It was revealed that the main cause of maternal death in poor settings is hemorrhage (54%) that occurs after delivery followed by pregnancy-induced hypertension (20%).<sup>15</sup>

Although reducing maternal mortality continues as the most challenging issue of the health system in Ethiopia, there are efforts to address the profound inequities in maternal and perinatal health.<sup>16</sup> The government of Ethiopia provides free maternal and pre-delivery services regardless of social and economic status of the women. However, still maternal mortality ratio remains high, and PPH is the leading cause of the morbidity and mortality in the country.<sup>17</sup>

In addition, the Ethiopia Federal Ministry of Health has applied multi-direction approaches to improve access to and strengthen facility-based maternal health care services as a strategy to reduce high burden of postpartum maternal morbidity and mortality.<sup>18,19</sup>

Even though there is a decline in maternal mortality from time to time, in Ethiopia it is still high as 412 per 100,000 live births.<sup>20</sup> As the study revealed, shortage of skill birth attendant, inadequate blood supply, lack of ambulance, and poor-quality services at facilities are some of the challenges to prevent and early treatment of PPH in low-resource settings.<sup>18</sup>

The risk of maternal death from PPH represents one of the greatest challenges in global health. Assessing the prevalence of PPH and identifying its risk factors could help to prevent and control its negative consequences. Also, it could have

paramount implications to focus target group as well as reduce the cost of its consequence. To address these problems, there is need for strategies that are based on evidence from different health care settings and situations.<sup>15,16</sup>

Additionally, giving more emphasis on potential risk factors that contribute to PPH and related complications is significant to reduce its burden, particularly in poor countries. As a result, more understanding about the burden of PPH in health care setting is important to prevent and control its consequences.<sup>11,13</sup>

Despite of this, there are inadequacy of literature regarding the extent of PPH and its contributing factors in Ethiopia including the study setting. Therefore, this study aimed to assess the magnitude of PPH and its associated factors among women who delivered at Yirgalem General Hospital, Sidama Regional State, Ethiopia in 2020.

## Methods

### *Study Setting and Period*

This study was carried out in Yirgalem General Hospital which is found in the Sidama Regional State, Ethiopia. It is located 322 km south of Addis Ababa and 47 km from Hawassa City. The hospital provides preventive, curative, and diagnostic services to the catchment population of 5.2 million. The hospital contains different departments, and is staffed with 13 specialist doctors, 49 general physicians, 36 midwives, and with other health professionals. Obstetrics ward is one of the major departments that serve around 2964 clients per year.

### *Study Design, Period, Sample Size Determination, and Population*

A facility-based cross-sectional study was conducted from March 12 to 26, 2020. The sample size was determined using single population proportion formula with the following assumption: taking prevalence of PPH 16.6% from the previous study,<sup>21</sup> 95% confidence interval (CI), and 4% marginal of error. A 1-year recorded data (from July 1, 2018 to June 30, 2019) in Yirgalem General Hospital showed that the number of mothers who delivered in the hospital was 2935. To obtain the final sample, finite population correction formula was considered. Accordingly, the final sample size calculated for this study was 298.

All women who gave birth at Yirgalem General Hospital from July 1, 2018 to June 30, 2019 were source population. All selected records of women that fulfilled eligibility criteria were considered as the study population. All maternal records found in the delivery registration logbook during the study period were included. However, maternal records with incomplete/missed information were excluded.

## **Sampling Procedures**

In Yirgalem General Hospital, there were a total of 2935 deliveries from July 1, 2018 to June 30, 2019. All unique medical

registration numbers of clients who gave birth in the last 1 year were selected from Gynecology and obstetrics ward, and sorted accordingly. Finally, a simple random sampling technique was used to select the study subjects.

### **Data Collection Tools, Procedures, and Quality Assurance**

Data were collected using an interviewer-administered structured questionnaire. The registration logbooks and individual cards were used to retrieve important information. The tool also contained sociodemographic factors, obstetrics factors, and maternal health related or medical history. Two nurses were assigned as data collectors. One Bachelor of Science midwife was assigned as a supervisor. Cases were identified from the clinical records office through reviewing all records. Finally, the data collection process was supervised by the assigned supervisor and the principal investigator was guiding the overall activities.

To assure the quality of data, training was given to the data collectors and supervisor for 1 day. A pretest was conducted on 5% of the sample in other hospitals. Based on the pretest, the necessary modification was made. The data collectors were instructed to write card numbers on the checklist during data collection. So that any identified error was traced back using the card numbers. The filled checklist was checked for its completeness by the Principal investigator on a daily basis.

### **Data Processing and Analysis**

The collected data were coded and entered into EpiData version 3.1 and exported to SPSS version 20.0 for further analysis. Descriptive statistics were used to describe the study subjects. Frequencies and cross-tabulations were used to check for missed values and variables. Bivariable analysis was used primarily to check which variables were associated with the dependent variable.

Variables with a *P*-value of <.25 in the bivariate analysis were entered into multivariable logistic regression model. Multivariable logistic regression with odds ratio and 95% CI was used to measure factors associated with the outcome variable, and control for possible confounders.

To assess the goodness of fit of the final model, Hosmer and Lemeshow goodness-of-fit test was checked to see good fitness. To check the multicollinearity effect, variance inflation factor less than 10 and tolerance test greater than 0.1 was considered. A *P*-value ≤.05 was considered to declare statistical significance. Finally, the results were presented using texts, tables, and figures.

### **Variables**

For this study, the outcome variable was PPH (yes or no) based on a clinician's diagnosis. The independent variables were sociodemographic (residence, maternal age, marital status, occupation), obstetric history (parity, gravidity, ANC utilization, status of labor, uterine atony, retained placenta, mode of delivery, and

previous PPH), and maternal health-related factors (anemia, preeclampsia, pregnancy-induced hypertension, different infections)

### **Operational Definitions**

PPH: according to this study, a PPH was considered based on a clinician diagnose who attended the delivery that was obtained from the women's records/cards.

Current history uterine atony: failure of the uterus to contract adequately after childbirth according to a clinician's diagnosis.

Prolonged labor: according to this study, it was defined as delayed labor for more than 24 h.

## **Results**

### **Sociodemographic Characteristics of the Study Participants**

A total of 298 woman charts were reviewed in this study. About 187 (62.8%) of them were aged 20 to 34 years with a mean age of 25.7 ( $\pm 5.6$ ) years. Majority of them (96.6%) were married. In terms of occupational status, 149 (50%) were housewives (Table 1).

### **Obstetrics-Related Characteristic of the Study Participants**

About 207 (69.5%) were multiparous. Majority (81.2%) had a history of ANC follow-up for their last pregnancy. Concerning the mode of delivery, 247 (82.9%) was spontaneous vaginal delivery. Fifty five (18.5%) had history of prolonged labor, 58 (19.5%) had history of delivery with episiotomy, and 21(7%) had history of

**Table 1.** Sociodemographic Characteristics of Women who Gave Birth at Yirgalem General Hospital, Sidama Regional State, Ethiopia, 2020.

Variable (n = 298)	Frequency	Percentage
Age in years		
<20	72	24.2
20 to 34	187	62.8
≥35	39	13.0
Residence		
Urban	108	36.2
Rural	190	63.8
Marital status		
Married	288	96.6
<sup>a</sup> Others	10	3.4
Occupation		
Housewife	149	50
Merchant	90	30.2
Government employee	54	18.1
<sup>b</sup> Others	5	1.7

<sup>a</sup>Others: single, divorced, and widowed.

<sup>b</sup>Others: farmer and others.

uterine atony. Regarding onset of the labor, 12 (4%) was induced, and the main reason was hypertensive disorders (preeclampsia and eclampsia) (Table 2).

### The Magnitude of PPH, and Other Medical and Obstetrics Complications

In this study, the magnitude of PPH was 9.4% (95% CI: 6.0, 12.8). About 66 (20.5%) had history of medical illness during their last pregnancy. Of which, 27 (43.3%) had anemia followed by urinary tract infections (39.3%). Twenty nine (9.7%) had history of hypertensive disorder during their last

pregnancy. Majorities (98.7%) were screened for HIV, and out of them, 0.3% were seropositive (Table 3).

### Factors Associated with Postpartum Hemorrhage

In the bivariable logistic regression analysis age of the woman, ANC follow-up, previous history of PPH, mode of delivery, duration of labor, delivery with episiotomy, and current history of uterine atony were associated with the outcome variable.

However, in multivariable logistic regression analysis, duration of labor, uterine atony, and mode of delivery (cesarean section and instrumental vaginal delivery) were factors significantly associated with PPH.

Women who had history of the prolonged labor ( $\geq 24$  h) were 3.4 [AOR = 3.4, 95% CI: 1.1, 9.9] times more likely to have PPH when compared with those who had less than 24 h duration of labor.

Those women who gave birth by cesarean section were 5.8 [AOR = 5.8, 95% CI: 1.1, 22.0] times, and instrumental vaginal delivery 3.7 [AOR = 3.7, 95% CI: 1.1, 12.7] times more likely developed PPH when compared to these who gave birth by spontaneous vaginal delivery.

Similarly, women who gave birth by instrumental vaginal delivery were 3.7 [AOR = 3.7, 95% CI: 1.1, 12.7] times more likely to develop PPH when compared to those who gave birth by spontaneous vaginal delivery.

This study also examined that the likelihood of developing PPH was 4.8-folds [AOR = 4.8, 95% CI: 1.4, 16.6] higher among women who had experienced uterine atony (Table 4).

**Table 2.** Obstetric Characteristics of Women who Gave Birth at Yirgalem General Hospital, Sidama Regional State, Ethiopia, 2020.

Variable (n = 298)	Frequency	Percentage
<b>History of antenatal care</b>		
Yes	242	81.2
No	56	18.8
<b>Parity</b>		
Primipara	91	30.5
Multipara	207	69.5
<b>History of current obstructed labor</b>		
Yes	61	20.5
No	237	79.5
<b>On set of labor</b>		
Spontaneous	286	96.0
Induced	12	4.0
<b>Duration of labor</b>		
<24 h	243	81.5
$\geq 24$ h	55	18.5
<b>Delivery with episiotomy performed</b>		
Yes	58	19.5
No	240	81.5
<b>Mode of delivery</b>		
Vaginal delivery (SVD)	247	82.9
Cesarean section	22	7.4
Instrumental delivery	29	9.7
<b>Duration of third stage labor</b>		
$\leq 30$ min	235	78.9
$> 30$ min	63	21.1
<b>Current history of retained placenta</b>		
Yes	38	12.8
No	260	87.2
<b>Current history of uterine atony</b>		
Yes	21	7
No	277	93
<b>History of previous PPH</b>		
Yes	51	17.1
No	247	82.9
<b>Current antepartum hemorrhage</b>		
Yes	26	8.7
No	272	91.3

Abbreviations: PPH: postpartum hemorrhage; SVD: spontaneous vaginal delivery.

**Table 3.** Medical Illness-Related Characteristics of Women who Gave Birth at Yirgalem General Hospital, Sidama Regional State, Ethiopia, 2020.

Variable	Frequency	Percentage
<b>History of current medical illness (n = 298)</b>		
Yes	61	20.5
No	237	79.5
<b>Type of current medical illness (n = 61)</b>		
Anemia	27	44.3
UTI	24	39.3
Malaria	9	14.8
Diabetes mellitus	1	1.6
<b>Have hypertensive disorder (n = 298)</b>		
Yes	29	9.7
No	269	90.3
<b>Type of hypertensive disorder (n = 29)</b>		
Preeclampsia	17	60.7
Eclampsia	7	25
Gestational HTN	4	14.3
<b>Screened for syphilis (n = 298)</b>		
Yes	271	90.9
No	27	9.1

Abbreviations: HTN: hypertension; UTI: urinary tract infection.

**Table 4.** Bivariable and Multivariable Logistic Regression Analysis for Factors Associated with PPH Among Women who Gave Birth at Yirgalem Hospital Sidama Regional State, Ethiopia, 2020.

Variables (n = 298)	Postpartum hemorrhage		Crude odds ratio (95% CI)	Adjusted odds ratio (95% CI)
	Yes	No		
Age group in years				
<20	6	66	1.2 (0.4,3.3)	1.6 (0.3,7.2)
20 to 34 (ref)	13	174	1	1
≥35 year	9	30	4.0 (1.5,10.2)	2.1 (0.6,7.3)
History of antenatal care				
No	12	44	3.8 (1.7,8.7)	2.1 (0.7,5.6)
Yes	16	226	1	1
Parity				
Multipara	23	184	2.1 (0.7,5.8)	1.6 (0.3,7.3)
Primipara	5	86	1	1
Previous history of PPH				
Yes	11	40	3.7 (1.6,8.5)	1.5 (0.5,4.7)
No	17	230	1	1
Mode of delivery				
Instrumental delivery	7	22	4.5 (1.7,12.7)	3.7 (1.1,12.7)*
Cesarean section	5	17	4.2 (1.3,12.9)	5.8 (1.5,22.0)*
Spontaneous vaginal delivery	16	231	1	1
Duration of labor				
≥24 h	16	40	7.6 (3.3,17.4)	3.4 (1.2,9.9)*
<24 h	12	230	1	1
Duration of third stage labor				
>30 min	10	53	2.2 (0.9,5.2)	1.5 (0.5,4.4)
≤30 min	18	217	1	1
Episiotomy performed				
Yes	12	45	3.7 (1.6,8.4)	2.3 (0.8,7.0)
No	16	225	1	1
History of current uterine atony				
Yes	7	15	5.6 (2.1,15.4)	4.8 (1.4,16.6)**
No	21	255	1	1

\* =  $P < .05$ , \*\* =  $P < .001$ , 1 = reference.

Abbreviations: CI: confidence interval; PPH: postpartum hemorrhage.

## Discussion

Globally, PPH is the major cause of maternal mortality and morbidity, accountable for more than 25% of deaths annually.<sup>4</sup> Subsequently, understanding its burden in the health care setting is important to prevent its negative outcomes.<sup>11,13</sup> Hence, this study attempted to assess the extent of PPH and its associated factors among postpartum women.

Accordingly, the magnitude of PPH was 9.4% [95% CI: 6.0, 12.8] in the study area. This result was comparable with the findings from previous study conducted in central Uganda (9%),<sup>22</sup> Bedele Hospital (9.6%),<sup>14</sup> and Debre Tabor General Hospital (7.6%).<sup>18</sup> However, this finding was lower when compared with the previous study conducted in Cameroon

(23.6%),<sup>23</sup> Pakistan (21.6%),<sup>24</sup> and Ethiopia (16.6%).<sup>25</sup> Meanwhile, our study findings were relatively higher than the previous findings from India (3.4%),<sup>26</sup> Zimbabwe (1.6%),<sup>27</sup> and Ethiopia (5.8%).<sup>28</sup>

The variation might be due to the differences in the study period, socioeconomic status, and upgrading of the health service delivering systems in terms of its availability and accessibility. Furthermore, the difference in the study setting and target population could contribute to this variation. Additionally, upgrading of data recording and documentation system could contribute to this variation.

This study showed that women who had experienced prolonged labor (≥24 h duration of labor) were 3.4 times more likely to develop PPH when compared with those who had less than 24 h duration of labor. Similarly, the study conducted in Bonassama Hospital in Cameroon supports this finding.<sup>24</sup> This finding was also in line with the previous results from Ethiopia.<sup>28,29</sup> Moreover, as our findings supported by different literatures, when the duration of labor increases, the risk of developing PPH also increased.<sup>30</sup>

The possible explanation for this could be that the pronged labor might increase the risk for a laceration to the blood vessels and soft tissue in the pelvis. This reduces uterine contraction by a significant level. As a result, the chance of losing blood after delivery could be increased by a substantial amount.

This study also revealed that delivery by cesarean section increased the probability of developing PPH by nearly sixfolds. Similarly, those women who gave birth by instrumental vaginal delivery were four times more likely to develop PPH when compared with those who gave birth by spontaneous vaginal delivery. This finding was in agreement with the previous results.<sup>21,28</sup>

A likely clarification for this might be that conducting a cesarean section could result in an organ cut, injury to the blood vessels, and tear of nearby tissue which make more bleeding. Furthermore, during instrumental vaginal delivery, there is an increased risk for cervical, vaginal and perineal lacerations which increases bleeding tendency later. Thus, those mothers who give birth by these procedures might easily get these risks that increase the chance of facing PPH and its related complications when compared with their counterparts.

The current study also indicated that woman who had history of uterine atony during her last delivery was nearly five times more likely to develop PPH than those who had normal contraction after childbirth. This finding was in line with previous reports from different settings.<sup>24,26,30</sup>

That might mainly happen because of many conditions that interfere with uterine contraction, such as retained placenta or retained small remnants of placental tissue, and blood clots could profoundly diminish contractility of the uterine muscle. That could lead to potentially heavy bleeding. In addition to this, physiological changes (cardiovascular changes) that occur during pregnancy and childbirth could also contribute to this problem.

## **Limitations of the Study**

The study might not be a real representative of the catchment area since women who received delivery services other than this hospital were not included. Since we used secondary data, we could not address cultural practices and social believes of the client. Also, factors related to the health system and the service providers were not addressed.

## **Conclusion**

The magnitude of PPH was 9.4% in the study setting. Prolonged labor, giving birth by cesarean section and instrumental vaginal delivery as well as having a history of uterine atony were factors significantly associated with PPH.

Healthcare professionals should manage the progress of labor and take all necessary measures at right time. Additionally, attention should be given to the safety of delivery-related procedures to reduce the risk of damaging other adjacent organs. Also, early recognition of potential risks for pregnancy-related complications, and following all suggested guidelines for interventions are key actions to decrease the probability of getting PPH. Further study is recommended with a consideration of factors related to the health system, service providers, and cultural aspects.

## **Authors' Contribution**

TA made considerable contributions to the conception, design, acquisition, analysis, and interpretation of the result. AD and AD contributed to analysis, interpretation of the result, drafting and revising the final document. All authors read and gave final approval of the version to be published.

## **Declaration of Conflicting Interests**

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

## **Funding**

The author(s) received no financial support for the research, authorship, and/or publication of this article.

## **Ethics Approval and Consent to Participate**

Ethical clearance was obtained from the Institutional Review Board of the Hawassa University College of Medicine and Health Science. Official permission letters were obtained from Yirgalem General Hospital. The purpose of the study also was informed to the manager, staff members, and all responsible body who are working at Yirgalem General Hospital.

## **Availability of Data and Materials**

The finding of this study is generated from the data collected and analyzed based on stated methods and materials. The original data supporting this finding are available from the corresponding author on reasonable request.

## **Consent to Publish**

Not applicable.

## **ORCID iDs**

Azmach Dache  <https://orcid.org/0000-0002-5961-6905>  
Aregahegn Dona  <https://orcid.org/0000-0001-5418-6662>

## **References**

1. World Health Organization. *WHO Recommendations Uterotonics for the Prevention of Postpartum Haemorrhage*. World Health Organization; 2018.
2. El-Refaey H, Rodeck C. Post-partum haemorrhage: definitions, medical and surgical management. A time for change. *Br Med Bull*. 2003;67(1):205-217.
3. Gürmezoglu AM. *WHO Guidelines for the Management of Postpartum Hemorrhage and Retained Placenta*. World Health Organization; 2009.
4. Ford JB, Patterson JA, Seeho SK, Roberts CL. Trends and outcomes of postpartum hemorrhage, 2003-2011. *BMC Pregnancy Childbirth*. 2015;15(1):334.
5. Abrams ET, Rutherford JN. Framing postpartum hemorrhage as a consequence of human placental biology: an evolutionary and comparative perspective. *Am Anthropol*. 2011;113(3):417-430.
6. Devi KP, Singh LR, Singh LB, Singh MR, Singh NN. Postpartum hemorrhage and maternal deaths in north east India. *Open J Obstet Gynecol*. 2015;5(11):635.
7. Habitemu D, Goshu YA, Zeleke LB. The magnitude and associated factors of postpartum hemorrhage among mothers who delivered at Debre Tabor general hospital 2018. *BMC Res Notes*. 2019;12(1):618.
8. Kodla CS. A study of prevalence, causes, risk factors and outcome of severe obstetrics hemorrhage. *J Sci Innov Res*. 2015;4(2):83-87.
9. Temesgen M. Magnitude of postpartum hemorrhage among women delivered at Dessie Referral Hospital, South Wollo, Amhara Region, Ethiopia. *J Women's Health Care*. 2017;6(391):2167-0420.1000391.
10. Sheldon W, Blum J, Vogel J, et al. Postpartum hemorrhage management, risks, and maternal outcomes: findings from the World Health Organization Multicountry Survey on Maternal and Newborn Health. *BJOG: Int J Obstet Gynaecol*. 2014;121(Supply 1):5-13.
11. Bulindi LB, Seljul RM-C, Ogundeko OT, et al. Knowledge and management of post partum haemorrhage among skilled birth attendants in Primary Health Centers of Jos North LGA, Plateau State. *World J Public Health*. 2017;2(4):124.
12. Bai J, Sun Q, Zhai H. A comparison of oxytocin and carboprost tromethamine in the prevention of postpartum hemorrhage in high-risk patients undergoing cesarean delivery. *Exp Ther Med*. 2014;7(1):46-50.
13. Aberra K. *Magnitude, Associated Factors and Maternal Outcome of Postpartum Hemorrhage at Black Lion Specialized Hospital from Jan. 1, 2009 to Dec. 30, 2013*. Addis Ababa University; 2014.
14. Gudeta T, Kebede D, Nigeria G, Dow M, Hassen S. Magnitude of post-partum hemorrhage among women who received postpartum care at Bedele Hospital South West, Ethiopia, 2018. *J Preg Child Health*. 2018;5(396):2.p.4/4.

15. Legesse T, Misra Abdulahi AD. Trends and causes of maternal mortality in Jimma University specialized hospital. Southwest Ethiopia: a matched case-control study. *Int J Womens Health.* 2017;9:307-313.
16. Miyake S, Speakman EM, Currie S, Howard N. Community midwifery initiatives in fragile and conflict-affected countries: a scoping review of approaches from recruitment to retention. *Health Policy Plan.* 2017;32(1):21-33.
17. Tesfaye G, Loxton D, Chojenta C, Assefa N, Smith R. Magnitude, trends and causes of maternal mortality among reproductive aged women in Kersa health and demographic surveillance system, eastern Ethiopia. *BMC Womens Health.* 2018;18(1):198.
18. Ethiopian Public Health Institute AMD, Disability. *Ethiopian Emergency Obstetric and Newborn Care (EmONC) Assessment 2016.* Ethiopian Public Health Institute, Federal Ministry of Health; 2017.
19. Central Statistical Agency. *Ethiopia Demographic and Health Survey 2016.* CSA and ICF; 2016.
20. Kebede BA, Abdo RA, Anshebo AA, Gebremariam BM. Prevalence and predictors of primary postpartum hemorrhage: an implication for designing effective intervention at selected hospitals, Southern Ethiopia. *PLoS One.* 2019;14(10):e0224579.
21. Ononge S, Mirembe F, Wandabwa J, Campbell OM. Incidence and risk factors for postpartum hemorrhage in Uganda. *Reprod Health.* 2016;13(1):38.
22. Gani GN, Ali AT. Prevalence and factors associated with maternal postpartum hemorrhage in Khyber agency. Pakistan. *J Ayub Med Coll Abbottabad.* 2013;25(1-2):81-85.
23. Halle-Ekane GE, Emade FK, Bechem NN, et al. Prevalence and risk factors of primary postpartum hemorrhage after vaginal deliveries in the Bonassama District Hospital, Cameroon. *Int J Trop Dis Health.* 2016;13(2):1-12.
24. Ngwenya S. Postpartum hemorrhage: incidence, risk factors, and outcomes in a low-resource setting. *Int J Womens Health.* 2016;8:647-650.
25. Abate T, Kebede B, Feleke A, Misganaw E, Rogers N. Prospective study on birth outcome and prevalence of postpartum morbidity among pregnant women who attended for antenatal Care in Gondar Town, North West Ethiopia. *Andrology.* 2014; 3(125):2167-0250.100012.
26. Nyflot LT, Stray-Pedersen B, Forsen L, Vangen S. Duration of labor and the risk of severe postpartum hemorrhage: a case-control study. *PLoS One.* 2017;12(4):e0175306.
27. Kramer MS, Berg C, Abenhaim H, et al. Incidence, risk factors, and temporal trends in severe postpartum hemorrhage. *Am J Obstet Gynecol.* 2013;209(5):449.E1-e7.
28. American College of Obstetricians & Gynecologists. ACOG practice bulletin no. 76: Postpartum hemorrhage. *Obs. Gynecol.* 2006;108:10391047. doi:10.1097/00006250-200610000-00046.
29. Smit M, Chan K-LL, Middeldorp JM, van Roosmalen J. Postpartum hemorrhage in midwifery care in the Netherlands: validation of quality indicators for midwifery guidelines. *BMC Pregnancy Childbirth.* 2014;14(1):397.
30. Montufar-Rueda C, Rodriguez L, Jarquin JD, et al. Severe postpartum hemorrhage from uterine atony: a multicentric study. *J Pregnancy.* 2013;525914. doi:10.1155/2013/525914.

## Author Biographies

**Tedla Amanuel** has a Master Degree in Public Health from Hawassa University and MCH Coordinator at Loka Abaya District Health Office.

**Azmach Dache** has a Master Degree in Public Health from Hawassa University and Lecturer at Yirgalem Hospital Medical College.

**Aregahegn Dona** has a Master Degree in Reproductive Health from Jimma University and Lecturer at Yirgalem Hospital Medical College.