

Article

Toxicity and Sublethal Effects of Lambda-Cyhalothrin Insecticide on Parent and Filial Generations of *Henosepilachna vigintioctomaculata* (Coleoptera: Coccinellidae)

Wenbo Li ¹, Muhammad Naeem ¹, Juan Cui ², Guochuan Du ^{3,*} and Huanhuan Chen ^{1,*}

¹ College of Biological Resource and Food Engineering, Qujing Normal University, Qujing 655011, China; lwbnicklwb@163.com (W.L.); naeem1633@yahoo.com (M.N.)

² College of Agriculture, Jilin Agriculture Science and Technology College, Jilin 132101, China; cuijuanjilin@163.com

³ College of Geography and Tourism, Qujing Normal University, Qujing 655011, China

* Correspondence: dgc0411@sina.cn (G.D.); chhuanhuan@163.com (H.C.)

Simple Summary: The ladybug *Henosepilachna vigintioctomaculata* is a widely distributed leaf-eating pest. Lambda-cyhalothrin, a synthetic pyrethroid insecticide, is widely used to control leaf-eating pests. However, the effects of this insecticide on biological activity, cross-generations, and detoxification enzyme activity of this ladybug are imperfectly known. We reared *H. vigintioctomaculata* for at least three generations under experimental conditions to determine the effects of lambda-cyhalothrin on biological activity. We evaluated the sublethal effects on F_0 generation adults, and cross-generational effects on F_1 adults, using age-stage bisexual life tables, and examined the detoxification enzyme activity of F_0 adults. Sublethal concentrations of lambda-cyhalothrin significantly reduced F_0 adult longevity and average fecundity, and inhibited various life table parameters in the F_1 population; there was also a cross-generational genetic effect, with population growth being inhibited. Low concentrations of lambda-cyhalothrin significantly inhibit *H. vigintioctomaculata* population growth. Multifunctional oxidase, carboxylesterase, and glutathione S-transferase play important roles in *H. vigintioctomaculata* resistance to lambda-cyhalothrin.



Academic Editors: Panagiotis J. Skouras and George J. Stathas

Received: 25 January 2025

Revised: 14 February 2025

Accepted: 21 February 2025

Published: 3 March 2025

Citation: Li, W.; Naeem, M.; Cui, J.; Du, G.; Chen, H. Toxicity and Sublethal Effects of Lambda-Cyhalothrin Insecticide on Parent and Filial Generations of *Henosepilachna vigintioctomaculata* (Coleoptera: Coccinellidae). *Insects* **2025**, *16*, 259. <https://doi.org/10.3390/insects16030259>

Copyright: © 2025 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: Lambda-cyhalothrin is a synthetic pyrethroid insecticide that is widely used to control leaf-eating pests. Because of increased insecticide resistance, an understanding of sublethal cross-generational effects of insecticides is important. We examine the effects of sublethal concentrations (SLCs) (LC_{10} , LC_{20} , and LC_{40}) of lambda-cyhalothrin on the growth, reproduction, and detoxification enzyme activities of F_0 and F_1 generation *Henosepilachna vigintioctomaculata*. Lambda-cyhalothrin is toxic to adult *H. vigintioctomaculata*, with an LC_{40} at 48 h of 0.355 mg L^{-1} . At SLCs, lambda-cyhalothrin significantly reduces the longevity and average fecundity of F_0 and F_1 adults, and prolongs the durations of the egg, larval, and pupal stages and adult preoviposition period. Additionally, an increased lambda-cyhalothrin concentration significantly decreases net reproductive rates, and both finite and intrinsic rates of increase in the F_1 generation, and significantly increases the average generation cycle. The detoxification enzyme activity of F_1 adults treated with SLCs of lambda-cyhalothrin for 48 h trends upwards. Results indicate that low concentrations of lambda-cyhalothrin induce glutathione S-transferase and carboxylesterase activities and inhibit multifunctional oxidase activity. The growth, development, and reproduction of the *H. viltioctomaculata* F_1 population remain inhibited by lambda-cyhalothrin treatment in the adult stage, and inhibitory effects increase with increased lambda-cyhalothrin concentration. The control efficacy of lambda-cyhalothrin against *H. viltioctomaculata* shows cross-generational effects.

Keywords: *Henosepilachna vigintioctomaculata*; lambda-cyhalothrin; life table; transgenerational studies; detoxifying enzyme

1. Introduction

The 28-spotted ladybug *Henosepilachna vigintioctomaculata* (Coleoptera: Coccinellidae) is a crop pest that occurs widely throughout China, and elsewhere in countries such as Korea, Japan, Russia, and Australia. This ladybug is a phytophagous pest that feeds mainly on crops such as Solanaceae, Leguminosae, Cruciferae, and Cucurbitaceae, with adults and larvae eating the tender leaves and stems [1,2]. When leaves are eaten, a semi-transparent parallel-shaped depression is formed, and in severe cases, only leaf veins and the epidermis remain, causing the leaves to wilt, yellow, and die. China is an important producer of potato—a globally important food crop—and *H. vigintioctomaculata* seriously affects potato agriculture [2–4]. Changes in microclimates and promotion of intercropping of corn, vegetables, and potatoes have also led to *H. vigintioctomaculata* becoming both an increasingly serious pest in potato plantations in the Yunnan, Guizhou, and Sichuan regions, and a main spreader of potato brown spot disease, which also poses a threat to local potato production. In severe cases, yields can be reduced by 50% [5–7].

The insecticide lambda-cyhalothrin is highly efficient at controlling leaf-eating pests [8]. After spraying, insecticide residue, in general, gradually decreases over time, and with crop growth [9]. The physiology and behavior of some pests exposed to insecticides at sublethal doses may change [10,11]. Most sublethal insecticide doses have inhibitory or delayed effects on pest populations, but some can also stimulate growth and development, and promote pest proliferation [12]. These sublethal effects on pests can lead to physiological and biochemical changes and insecticide resistance. Therefore, understanding the sublethal effects of insecticides on pests is important to evaluate their efficacy and assess the risks associated with their use [13]. Pests can reduce the effects of insecticides by regulating detoxification enzyme activities [14]. Among enzymes, glutathione S-transferase (GST), carboxylesterase (CarE), and multifunctional oxidase (MFO) are the most important. Low concentrations of insecticides or plant secondary metabolites can induce or inhibit the activity of various enzymes, thereby affecting pest metabolic processes, promoting resistance and providing continuous selection pressure, reducing the effectiveness of insecticides for pest control [15,16].

While the median lethal concentration (LC₅₀) or median lethal dose (LD₅₀) have been commonly used to evaluate the effects of insecticides on pests, they only report the response to insecticides at certain developmental stages [17]. Life tables can be used to more comprehensively analyze the effects of pesticides on pests at the population level. Compared with traditional life tables, an age–age bisexual life table increases statistics for males, accurately describes age differentiation of insects, differentiates between the pre- and total-oviposition periods of adults, and can comprehensively describe changes in the entire population [18–20]. Beta-cypermethrin, phoxim, and abamectin pesticides are mostly used to control potato ladybird adults [4]. While the sublethal effects of insecticides on target species are being increasingly explored, the effects of the lethal concentration (LC) of lambda-cyhalothrin on the growth, development, and fecundity of *H. vigintioctomaculata* are imperfectly known. We first determine the sublethal concentration (SLC) of lambda-cyhalothrin on the F₀ generation, and its effect on the F₁ generation at SLCs. We construct an age–instar bisexual life table for the F₁ population and compare changes in parameters such as developmental duration, survival rate, fecundity, and longevity at different lambda-cyhalothrin concentrations. We also report the detoxifying enzyme activities after

SLC treatment of F_0 adults. The results provide a theoretical basis for a comprehensive assessment of the potential of lambda-cyhalothrin to control *H. vigintioctomaculata*, and for more informed applications of this insecticide.

2. Materials and Methods

2.1. Insect Rearing

Henosepilachna vigintioctomaculata were collected from Luliang County, Qujing City potato fields, Yunnan Province (103°48′52.02″ E, 25°17′20.74″ N, altitude 1878 m) in September 2023. Insects were placed in an insect feeding cage (50 × 50 × 50 cm) at room temperature. Potted potatoes (Yunshu 108) grown at (25 ± 1) °C, relative humidity 70% ± 5%, and a photoperiod 16:8 h L/D were cultivated as a host; fresh leaves were cut to feed the ladybugs when the plants had grown to 15–20 cm. Before experimentation, adult ladybugs were reared for at least three generations with no exposure to insecticide. Three-day-old adults were used as the initial insect source. Insect feeding conditions were consistent with potato cultivation conditions.

2.2. Insecticide and Chemicals and Enzyme Assay Kits

The stock insecticide solution was 2.5% lambda-cyhalothrin WG (Henan Yongguan Qiaodi Agricultural Science and Technology Co., Ltd, (Zhengzhou, China). Multifunctional oxidase (MFO) activity detection kits (A162-1-1), Carboxylesterase (CarE) activity detection kits (A133-1-1), and Glutathione S-transferase (GSH-ST) activity detection kits (A004-1-1) were purchased from Nanjing Jiancheng Bioengineering Institute Co., Ltd (Nanjing, China).

2.3. Determining Biological Activity

The sensitivity of *H. vigintioctomaculata* to lambda-cyhalothrin was determined using a leaf film method [21]. Five lambda-cyhalothrin treatments (1.25, 1.00, 0.75, 0.50, and 0.25 mg L⁻¹) were initially established; water was used as a control. Each treatment contained 30 × 3 d old similarly sized adult ladybugs, and was replicated three times. Before experimentation, adult ladybugs were starved for 48 h. Potato leaves were immersed in each treatment concentration for 20 s, then removed, placed in an insect box (15 × 12 × 5 cm), dried, and their petiole wrapped with water-soaked cotton wool. Ladybugs were then added. Ladybug mortality was recorded after 48 h of treatment. Feeding conditions were the same as in Section 2.1. Ladybugs were pronounced dead if touched by a brush and no reaction occurred. A virulence regression curve, 95% confidence intervals, chi-square values (χ^2), and degrees of freedom (*df*) were calculated for LC₁₀, LC₂₀, and LC₄₀ concentrations.

2.4. Effects of Lambda-Cyhalothrin Sublethal Treatment on F_0 and F_1 *H. vigintioctomaculata*

Potato leaves were immersed in LC₁₀, LC₂₀, and LC₄₀ concentrations for 20 s, then removed and dried; water was used as a control. Sixty healthy 3 d old similarly sized male and female adults (1:1, morphological identification) were fed potato leaves and reared in an artificial climate box [4]. After 48 h of exposure to leaves immersed in different concentrations of insecticide (without changing leaves), surviving *H. vigintioctomaculata* were removed, and males and females were paired, placed in a Petri dish, and fed fresh, untreated potato leaves. The initial stage of the F_0 generation experiment involved approximately 30 individuals, with three biological replicates. At least 15 pairs of ladybugs from each insecticide treatment were paired; the fresh potato leaves in the insect box were replaced daily. The oviposition date, oviposition amount, and adult longevity of single females were observed regularly (8:00 and 18:00 h). If a male died before a female, it was replaced. We refer to adult contemporary *H. vigintioctomaculata* treated with SLCs

of lambda-cyhalothrin as the parent generation (F_0), and offspring produced by natural mating of these parents as the filial generation (F_1) [2].

Eggs (90, oviposition < 5 h) laid on the same day by an F_0 female in each lambda-cyhalothrin treatment and the control group were used as the F_1 generation. These eggs were placed into circular plastic, numbered Petri dishes (diameter 60 mm) with moist filter paper, and then moved to an artificial climate box for hatching, where the hatching rate was determined. The feeding conditions were the same as in Section 2.1. Individual newly hatched F_1 -generation larvae treated with different insecticide concentrations were inoculated separately into Petri dishes for feeding. Fresh potato leaves were placed into Petri dishes, and ladybug development was monitored every 24 h. After 24 h of pupation, pupae were weighed, then transferred into individual centrifuge tubes sealed with absorbent cotton, and returned to the climate chamber. After adult eclosion, pupal duration was determined. For each insecticide concentration treatment, female and male adults that emerged on the same day were randomly selected and paired 1:1 and placed into new culture dishes. A wet cotton ball was placed in the dish to provide moisture, and fresh leaves were replaced the next day. Test insects ate and deposited eggs. The number of eggs laid by a single female and the longevity of adult females and males were observed daily (8:00 and 18:00 h) until all adults had died. The number of eggs hatched/the total number of eggs laid by females represents the egg-hatching rate. The time from newly hatched larvae to pupation represents the larval development period, and the time from pupation to adult eclosion represents the pupal development period. The percentage of the number of pupae that could break the shell and become adults divided by the total number of pupae $\times 100$ represents the eclosion rate.

2.5. Determining Detoxifying Enzyme Activities

Detoxifying enzyme activities were determined using the method of Jiang et al. [22]. F_0 -generation adult ladybugs were fed fresh leaves treated with LC_{10} , LC_{20} , and LC_{40} concentrations of lambda-cyhalothrin, with four replicates per treatment. After 48 h, surviving adults were collected and placed into centrifuge tubes. Each replicate contained 10 individuals, frozen in liquid nitrogen and stored at -80°C for enzyme activity determination. Ten similarly sized adults were selected from each concentration treatment. PBS (pH 7.4) was added to an ice bath homogenate at a weight (g)–volume (mL) ratio of 1:9; the homogenate was centrifuged at 4°C and 12,000 r/min for 30 min. The supernatant was recovered to determine GST, CarE, and MFO enzyme activities, and protein concentration, following kit instructions.

2.6. Data Analysis

A probability unit analysis was used to analyze the toxicity regression equation of lambda-cyhalothrin using SPSS 26.0 (IBM Co., Ltd., Armonk, NY, USA); SLCs were determined. One way ANOVA was used on data related to detoxification enzymes. Tukey's multiple range test was used for multiple comparisons, and Student's *t* tests were used to identify significant differences between treatments in pairwise comparisons. Original growth and development, survival rate, and fecundity data for the F_1 generation were collated and imported into Twosex-MSChart software (v 5/7/2024) to calculate developmental duration, adult longevity, adult preoviposition period (APOP), total preoviposition period (TPOP), fecundity, net reproductive rate (R_0), finite rate of increase (λ), intrinsic rate of increase (r), and average generation cycle (T) for each stage [23–26]. All variances and standard errors were obtained by using 100,000 random sampling bootstraps, and the differences for each parameter of the insecticide treatments were evaluated by paired

bootstrap tests [27,28]. Plots were generated using SigmaPlot 14.0. The specific calculation formulas are as follows:

$$l_x = \sum_{j=1}^{\beta} s_{xj} \quad (1)$$

$$M_x = \frac{\sum_{j=1}^{\beta} s_{xj} f_{xj}}{\sum_{j=1}^{\beta} s_{xj}} \quad (2)$$

$$R_0 = \sum_{x=0}^{\infty} l_x m_x \quad (3)$$

$$\sum_{x=0}^{\infty} e^{-r(x+1)} l_x m_x = 1 \quad (4)$$

$$\lambda = e^r \quad (5)$$

$$T = \frac{\ln(R_0)}{r} \quad (6)$$

3. Results

3.1. Toxicity of Lambda-Cyhalothrin to *Henosepilachna vigintioctomaculata* Adults

After 48 h treatment, the LC₁₀, LC₂₀, and LC₄₀ values for lambda-cyhalothrin exposure to *H. vigintioctomaculata* adults were 0.193, 0.251, and 0.355 mg L⁻¹, respectively. Three concentration values overlapped with other 95% CL values for subsequent experiments (Table 1).

Table 1. Sublethal concentrations of lambda-cyhalothrin for newly hatched adult *H. vigintioctomaculata* (F₀) 48 h after treatment.

Insecticide	Number of Samples	LC ₁₀ (95% CL) (mg L ⁻¹)	LC ₂₀ (95% CL) (mg L ⁻¹)	LC ₄₀ (95% CL) (mg L ⁻¹)	χ ² (df)	Slope ± SE	p-Value
Lambda-cyhalothrin	450	0.193 (0.112–0.265)	0.251 (0.184–0.352)	0.355 (0.258–0.432)	9.53	3.91 ± 0.59	0.023

3.2. Effect of Sublethal Concentration of Lambda-Cyhalothrin on Longevity and Fecundity of F₀ *H. vigintioctomaculata*

LC₁₀, LC₂₀, and LC₄₀ concentrations of lambda-cyhalothrin significantly affected the longevity and average fecundity of F₀ adults (Table 2). Compared with the control group, the higher the concentration of lambda-cyhalothrin, the shorter the adult life span (female and male), with the life span of females being slightly longer. After exposure to lambda-cyhalothrin concentrations (LC₁₀, LC₂₀, and LC₄₀), individual egg production and F₁ egg hatching rates also significantly decreased compared with the control group.

Table 2. Impact of sublethal concentrations of lambda-cyhalothrin on F₀ generation adult *H. vigintioctomaculata* longevity and fecundity.

Parameter	Control	Lambda-Cyhalothrin (LC ₁₀)	Lambda-Cyhalothrin (LC ₂₀)	Lambda-Cyhalothrin (LC ₄₀)
	Mean ± SE	Mean ± SE	Mean ± SE	Mean ± SE
Male longevity (d)	24.39 ± 0.30 a	24.00 ± 0.23 a	21.45 ± 0.29 b	20.09 ± 0.33 c
Female longevity (d)	26.27 ± 0.34 a	26.00 ± 0.21 a	24.29 ± 0.39 b	21.79 ± 0.29 c
Fecundity (eggs laid/female)	65.66 ± 9.21 a	59.11 ± 7.09 b	42.12 ± 5.56 c	31.09 ± 4.76 d
F ₁ egg Hatching rate	84.29 ± 11.55 a	68.73 ± 9.79 b	58.26 ± 9.60 c	49.46 ± 7.88 d

Note: Different letters within a row indicate significant differences ($p < 0.05$).

3.3. Effect of Sublethal Concentration of Lambda-Cyhalothrin on Growth, Development, Fecundity, and Pupal Weight of *F₁* *H. vigintioctomaculata*

LC₁₀, LC₂₀, and LC₄₀ concentrations of lambda-cyhalothrin affected each *F₁* generation developmental stage (Table 3). Compared with the control, the duration of the egg, larval, and pupal stages in the LC₁₀ and LC₂₀ treatment groups differed significantly ($p < 0.05$); the lifespan of male and female adults in the LC₄₀ concentration treatment group was 16.57% and 19.76%, significantly shorter than control group adults. APOP and TPOP were prolonged with increased lambda-cyhalothrin concentrations; the control group's APOP and TPOP durations were the shortest (6.20 d and 23.53 d), and those in the LC₄₀ treatment were the longest (7.10 d and 30.45 d, respectively). Compared with other treatments, the numbers of eggs laid by females exposed to the LC₄₀ concentration was lowest (13.40 eggs/female); the number of eggs laid by females in the control group (55.59 eggs/female) was 1.13×, 1.70×, and 4.15× that of the LC₁₀, LC₂₀, and LC₄₀ treatments, respectively. Additionally, compared with controls, the pupal weight and adult emergence rate of the *F₁* generation were also significantly lower in the LC₁₀, LC₂₀, and LC₄₀ treatments ($p < 0.05$), with the effect stronger with increased insecticide concentration (Figure 1, Table 3).

Table 3. Fecundity and duration of various life-history parameters of progeny of lambda-cyhalothrin-treated *H. vigintioctomaculata*.

Parameter (Duration, Days)	Treatment			
	Control	Lambda-Cyhalothrin (LC ₁₀)	Lambda-Cyhalothrin (LC ₂₀)	Lambda-Cyhalothrin (LC ₄₀)
	Mean ± SE	Mean ± SE	Mean ± SE	Mean ± SE
Egg (d)	5.66 ± 0.07 c	6.48 ± 0.07 b	6.70 ± 0.07 b	7.12 ± 0.05 a
Larva (d)	15.05 ± 0.17 c	16.32 ± 0.23 b	16.60 ± 0.21 b	18.69 ± 0.30 a
Pupa (d)	4.59 ± 0.08 c	5.40 ± 0.13 b	5.71 ± 0.15 b	6.17 ± 0.19 a
Male longevity (d)	23.53 ± 0.25 a	22.52 ± 0.31 b	20.12 ± 0.17 c	18.88 ± 0.21 d
Female longevity (d)	25.40 ± 0.27 a	24.86 ± 0.18 b	23.21 ± 0.46 c	21.19 ± 0.26 d
APOP (d)	6.20 ± 0.47 b	7.00 ± 1.36 a	7.00 ± 0.59 a	7.10 ± 0.82 a
TPOP (d)	23.53 ± 0.52 d	26.75 ± 1.44 c	29.79 ± 1.24 b	30.45 ± 1.70 a
Eclosion rate %	85.98 ± 0.13 a	80.54 ± 0.76 b	71.02 ± 0.54 c	60.66 ± 0.73 d
Fecundity (eggs laid/female)	55.59 ± 8.72 a	48.79 ± 7.08 b	32.71 ± 3.23 c	13.40 ± 3.06 d

Note: APOP, adult pre-oviposition period; TPOP, total pre-oviposition period. Data are presented as means ± SE. Different letters within a row indicate significant differences based on paired bootstrap tests ($p < 0.05$). Standard errors are estimated from 100,000 bootstrap resamples.




Figure 1. Pupal weight of progeny of lambda-cyhalothrin-treated *H. vigintioctomaculata*. Note: Data are means ± SE; different lowercase letters above whiskers indicate significant differences ($p < 0.05$, Tukey's multiple range test).

3.4. Population Parameters

Compared with the controls, the LC₁₀, LC₂₀, and LC₄₀ treatments significantly affected the F₁ generation's life table parameters. The values for the r , λ , and R_0 of the F₁ population decreased with increased lambda-cyhalothrin concentration, and T increased (Table 4).

Table 4. Population parameters of F₁ *H. vigintioctomaculata* descended from the F₀ generation exposed to lambda-cyhalothrin.

Parameters	Concentration Treatments			
	Control	Lambda-Cyhalothrin (LC ₁₀)	Lambda-Cyhalothrin (LC ₂₀)	Lambda-Cyhalothrin (LC ₄₀)
	Mean ± SE	Mean ± SE	Mean ± SE	Mean ± SE
Intrinsic rate of increase, r d ^{−1}	0.087 ± 0.006 a	0.062 ± 0.007 b	0.041 ± 0.005 c	0.017 ± 0.007 d
Finite rate of increase, λ d ^{−1}	1.081 ± 0.006 a	1.064 ± 0.007 b	1.041 ± 0.005 c	1.017 ± 0.009 d
Net reproductive rate, R_0 (offspring/individual)	25.98 ± 6.75 a	13.66 ± 3.64 b	9.16 ± 1.25 c	3.68 ± 0.96 d
Mean generation time, T (d)	38.97 ± 0.46 c	41.69 ± 0.74 b	53.98 ± 1.80 a	54.52 ± 1.91 a

Note: Different letters within a row indicate significant differences ($p < 0.05$).

3.5. Age–Stage Specific Maternity

At each concentration, the l_x curves follow a similar trend: there is a period of stability, and then a gradual and then steep decline (Figure 2), indicating that the death of individuals in the *H. vigintioctomaculata* F₁ population mainly occurred in the later stages. The f_x curves are more variable, indicating greater variation in the emergence and oviposition of female adults, resulting in the curve appearing high and low. For the LC₁₀, LC₂₀, and LC₄₀ lambda-cyhalothrin concentrations, peak adult female f_x values were 13.77 (37 d), 14.11 (55 d), and 8.46 (58 d) (Figure 2). The peak number of eggs of f_x in the control group was greatest (15.62 eggs d^{−1}). Over time, the age-specific survival rate l_x of the control group trended downwards. The m_x curve indicated that the control group began reproducing at 33 d, whereas in other concentrations reproduction began 2–11 d later. Multiple peaks appear in the m_x curve, indicating changes in individual spawning periods. The age-specific oviposition rate $l_x m_x$ in the LC₄₀-treated individuals decreased sharply, from 5.00 individuals d^{−1} in the control group to 1.70 individuals d^{−1} (Figure 2).




Figure 2. l_x , f_x , m_x , and $l_x m_x$ values of F₁ *H. vigintioctomaculata* descended from F₀ ladybugs treated with lambda-cyhalothrin and control values.

3.6. Effect of Lambda-Cyhalothrin Exposure on Detoxifying Enzyme Activity

The detoxifying enzyme activities in the adult *H. vigintioctomaculata* treated with SLCs of lambda-cyhalothrin trended upwards (Figure 3). After 48 h exposure at LC₂₀ and LC₄₀ concentrations, female GST activities decreased significantly by 28.88% and 44.88% compared with control values, and male activities decreased significantly by 39.39% and 44.95%, respectively ($p < 0.05$); at LC₁₀ there was an increase in GST activity. CarE activity first increased, then decreased with increased lambda-cyhalothrin concentration; the enzyme activities in the LC₄₀ treatment were significantly lower than those in either of the other treatments ($p < 0.05$). With an increased treatment concentration, MFO activity was inhibited. Additionally, the GST activity of adults in the LC₂₀ treatment and CarE activity in the LC₁₀ and LC₄₀ treatments differed significantly ($p < 0.05$).




Figure 3. (A) GST, (B) CarE, and (C) MFO activities in *H. vigintioctomaculata* adults treated with lambda-cyhalothrin. Note: Data are means \pm SE; different uppercase and lowercase letters indicate significant differences between males and females, respectively ($p < 0.05$, Tukey's multiple range test). Asterisks indicate significant differences in detoxifying enzyme activities between females and males (* $p < 0.05$, independent samples t -test).

4. Discussion

Insecticides are used extensively in agriculture, and they contribute to the development of pest resistance [17]. Integrated pest-management programs underscore the importance and necessity of moderate insecticide use [29]. Lambda-cyhalothrin is an efficient, broad-spectrum and quick-acting pyrethroid insecticide developed by ICI, UK. It has contact and stomach poisoning effects, and has no internal absorption effect. It is mainly used to control pests with chewing or piercing and sucking mouthparts [8,9,14]. Using a leaf-dip method, we evaluated the bioactivity of lambda-cyhalothrin against adult *H. vigintioctomaculata* [4]. Our LC₄₀ value of 0.355 mg L⁻¹ indicates that this insecticide exhibits potent toxicity toward *H. vigintioctomaculata*. Chemical control is the main strategy to manage *H. vigintioctomaculata*. However, post-application in the field, insecticide toxicity generally decreases because of various abiotic environmental factors, producing SLCs [30]. This reduced toxicity can induce sublethal effects in certain pest individuals that may be passed on to subsequent generations. Sublethal effects may influence the growth, develop-

ment, behavior, and reproductive capacity of target pests and their offspring, and change population dynamics [30–32].

Many studies have reported that sublethal insecticide concentrations exert significant inhibitory effects. For instance, treatment with chlorantraniliprole at LC_{10} concentrations significantly reduces the survival rate and longevity of F_1 generations of *Sogatella furcifera* [33]. Studies on the development of *Cydia pomonella* under lambda-cyhalothrin stress have yielded similar results [34]. We report SLCs of lambda-cyhalothrin (LC_{10} , LC_{20} , and LC_{40}) to significantly reduce the lifespan and average oviposition of the F_0 generation *H. vigintioctomaculata* compared with the control group ladybugs (Table 2). Furthermore, exposure to LC_{40} concentrations significantly reduced the F_1 population size, with delays observed in egg, larval, and pupal developmental duration, further indicative of sublethal effects (Table 3). These findings are consistent with those for exposure of *Paracoccus marginatus* to SLCs of spirotetramat, with extended nymphal periods from the F_0 to F_2 generations and pre-adult delays [35]. We also report the pupal weight—an important indicator of insect stress resistance and environmental adaptability—of the F_1 generation of *H. vigintioctomaculata* to decrease with increased SLC of lambda-cyhalothrin [36]. Leaves treated with LC_{40} concentrations also showed negligible feeding damage, likely attributable to the antifeedant properties of lambda-cyhalothrin [37]. Reduced feeding reduced the growth and development of *H. vigintioctomaculata*, and reduced body weight.

The impact of insecticides on pest fecundity manifests as sublethal effects [31]. The toxic excitatory response is influenced by various factors, with exposure duration (or generational exposure) being a key determinant [38]. Low doses of imidacloprid stimulate peach aphids, leading to increased methylation levels in the F_2 generation compared with the F_1 generation. This phenomenon may be linked to genetic adaptability induced by low-dose insecticide stress [39]. We report that SLCs of lambda-cyhalothrin also induce transgenerational effects in *H. vigintioctomaculata*. Specifically, the average oviposition, r , λ , and R_0 values of both the F_0 and F_1 generations decreased with increased lambda-cyhalothrin concentration, limiting F_1 population growth (Tables 2 and 3). A similar phenomenon is reported for pests such as *Paracoccus marginatus*, *Aphis gossypii*, and *Rhizoglyphus robini* [9,14,40]. These findings suggest that insecticide treatments can effectively slow transgenerational population growth of a variety of species. This inhibitory effect may be attributed to the downregulation of gene expression associated with vitellogenin (Vg) and its receptor synthesis in females exposed to sublethal doses of insecticide. For example, the reduced expression of Vg and vitellogenin receptor (VgR) genes leads to decreases in their respective contents [40–42]. Furthermore, age-stage-specific maternity curves (Figure 2) indicate that lambda-cyhalothrin stress significantly suppresses the F_1 generation's reproductive capacity, with the effect increasing at higher concentrations. Interestingly, many insecticides (e.g., triazophos and thiamethoxam) regulate pest fecundity by modulating the expression of Vg and VgR through the TOR protein kinase and juvenile hormone signaling pathways [43,44]. Whether lambda-cyhalothrin similarly regulates the reproductive capacity of *H. vigintioctomaculata* adults via these pathways is unknown.

The reduced sensitivity of insects to insecticides typically involves multiple mechanisms, with the most prominent being the regulation of insecticide metabolism through detoxifying enzymes [15]. These enzymes play important roles in the metabolic processing of chemicals in insects. Insecticides can induce or inhibit their activity, and by exerting continuous selective pressure, drive the evolution of insecticide resistance [45]. Insecticides can affect various population parameters in insects such as developmental periods, reproductive capacity, and adult longevity. Additionally, they can influence the activity of detoxifying enzymes [46]. Different insecticides or plant secondary metabolites have distinct inhibitory or inducing effects on detoxifying enzyme activities in the same pest

species [31]. For example, after 24 h of spirotetramat exposure, low concentrations induced a significant increase in GST and CarE activity in *Bradysia odoriphaga* [47]. Similarly, treatment with lambda-cyhalothrin significantly increased GST activity in *Cydia pomonella*, but suppressed CarE activity [48]. We report that, after 48 h of exposure to lambda-cyhalothrin, LC₁₀ concentrations induced an increase in both GST and CarE activities in adult *H. vigintioctomaculata*. At LC₂₀, the CarE activity in adult males and females was significantly higher than in the control group, but lower than in the LC₁₀ treatment group, possibly because of the lower insecticide dose (where CarE was activated to participate in insecticide metabolism). Additionally, the GST activity of adults in the LC₂₀ treatment and CarE activity in the LC₁₀ and LC₄₀ treatments differed significantly ($p < 0.05$). Similar GST and CarE activity results were reported by Kinareikina [49]. At increased dosage, enzyme activity was progressively suppressed. The degree of MFO activity correlated positively with insecticide dosage. At low lambda-cyhalothrin concentrations, the activities of CarE and GST in *H. vigintioctomaculata* were induced, but MFO activity was significantly reduced. Further molecular-level studies on higher resistance strains are required to elucidate the underlying mechanisms of resistance to lambda-cyhalothrin.

After the exposure of adult *H. vigintioctomaculata* in the F₀ generation to SLCs of lambda-cyhalothrin, F₁ population growth was significantly reduced, and the degree of reduction increased with increased insecticide concentration. This suggests that field application of lambda-cyhalothrin exerts transgenerational effects on *H. vigintioctomaculata*. In agricultural practice, the rational use of lambda-cyhalothrin to mitigate damage caused by *H. vigintioctomaculata* to crops is viable. However, the dosage and frequency of application must be managed to avoid enhancing pest fitness or promoting resistance through prolonged use of a single class of insecticide. Furthermore, the implementation of insecticide rotation—using two or more insecticides with different modes of action—could delay the development of resistance and support sustainable pest management strategies.

5. Conclusions

As a new insecticide with a high efficiency, low toxicity, and low residue, lambda-cyhalothrin has high insecticidal activity against Coleoptera pests. We report the effects of SLCs of this insecticide on the growth and development, longevity, and fecundity of *H. vigintioctomaculata* using an age–age bisexual life table. Compared with the control group, male and female longevity and fecundity all decreased, and other life table parameters decreased with increased SLCs. At low concentrations, this insecticide also increases the activity of detoxifying metabolic enzymes. Field application of lambda-cyhalothrin has a cross-generational effect on *H. vigintioctomaculata* that may affect F₁ generations. This provides a reference for how to improve the evaluation of the effects of insecticides.

Author Contributions: H.C. conceived and designed the experiments. W.L. and J.C. conducted the experiments. M.N. and G.D. analyzed the data. W.L. and H.C. obtained the funding for the study. All authors contributed to writing and editing of the manuscript. Writing—original draft, W.L. and M.N.; Writing—review and editing, J.C.; Supervision, H.C. All authors have read and agreed to the published version of the manuscript.

Funding: This study was financially supported by Yunnan Fundamental Research Projects Association (grant no. 202301AU070005), the Special Basic Cooperative Research Programs of Yunnan Provincial Undergraduate Universities (202301BA070001-076), the Special Basic Cooperative Research Innovation Programs of Qujing Science and Technology Bureau & Qujing Normal University (KJLH2024ZD05), and Innovative Research Team in Qujing Normal University.

Data Availability Statement: The data presented in this study are available on request from the corresponding author.

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. Matsishina, N.V.; Ermak, M.V.; Fisenko, P.V.; Sobko, O.A. *Henosepilachna vigintioctomaculata* Motschulsky (Coleoptera: Coccinellidae): Morphotypes in an East Asian population. *J. Insect Biodivers.* **2023**, *38*, 15–23. [\[CrossRef\]](#)
2. Li, W.; Bashir, N.H.; Chen, H.; Li, X.; Wang, Z.; Du, L.; Tian, R. Effect of different host plants on the growth, development, and reproduction of *Henosepilachna vigintioctomaculata* (Coleoptera: Coccinellidae). *Chin. J. Appl. Entomol.* **2024**, *61*, 287–294.
3. Ze, L.J.; Jin, L.; Li, G.Q. The compatible effects of RNA interference targeting laccase2 with biocontrol in *Henosepilachna vigintioctopunctata*. *Entomol. Gen.* **2023**, *43*, 117–126. [\[CrossRef\]](#)
4. Ding, X.; Aerziguli, R.; Ji, L.; Liu, S.; Fu, K.; Tursun, A.; Guo, W. The occurrence and harm of *Henosepilachna vigintioctomaculata*. *Xinjiang Agric. Sci.* **2022**, *59*, 983–989. [\[CrossRef\]](#)
5. Tamang, S.; Alam, T.; Rana, L. Biointensive integrated pest management of potato. In *Biointensive Integrated Pest Management for Horticultural Crops*, 1st ed.; Kumar, A., Saha, S., Choudhary, J.S., Eds.; CRC Press: London, UK, 2021; pp. 207–240.
6. Nithish, A. Study on management of damage caused by *Henosepilachna vigintioctopunctata* in brinjal. *J. Pharmacog. Phytochem.* **2020**, *9*, 2874–2881.
7. Zhou, J.; Xu, R.; Chen, Z.; Jia, Y.; Xu, K. Phototactic behavior of *Henosepilachna vigintioctomaculata* motschulsky (Coleoptera: Coccinellidae). *Coleopt. Bull.* **2015**, *69*, 806–812. [\[CrossRef\]](#)
8. Keyhanian, A.A.; Barari, H.; Mobasheri, M.T. Comparison of the efficacy of insecticides, alphacypermethrin and lambda-cyhalothrin, against canola flea beetles. *Appl. Entomol. Phytopathol.* **2021**, *44*, 113–122.
9. Qiu, Y.; Chen, Z. Intergenerational effects of sublethal lambda-cyhalothrin exposure on *Aphis gossypii* glover (Hemiptera: Aphididae) reproduction and development. *Insects* **2024**, *15*, 173. [\[CrossRef\]](#)
10. Wang, D.; Wang, Y.M.; Liu, H.Y.; Xin, Z.; Xue, M. Lethal and sublethal effects of spinosad on *Spodoptera exigua* (Lepidoptera: Noctuidae). *J. Econ. Entomol.* **2013**, *106*, 1825–1831. [\[CrossRef\]](#)
11. Boina, D.R.; Onagbola, E.O.; Salyani, M.; Stelinski, L.L. Antifeedant and sublethal effects of imidacloprid on Asian citrus psyllid, *Diaphorina citri*. *Pest Manag. Sci.* **2009**, *65*, 870–877. [\[CrossRef\]](#)
12. Cutler, G.C.; Amichot, M.; Benelli, G.; Guedes, R.N.C.; Qu, Y.; Rix, R.R.; Ullah, F.; Desneux, N. Hormesis and insects: Effects and interactions in agroecosystems. *Sci. Total Environ.* **2022**, *825*, 153899. [\[CrossRef\]](#) [\[PubMed\]](#)
13. Gul, H.; Güncan, A.; Ullah, F.; Desneux, N.; Liu, X. Intergenerational sublethal effects of flonicamid on cotton aphid, *Aphis gossypii*: An age-stage, two-sex life table study. *Insects* **2024**, *15*, 529. [\[CrossRef\]](#) [\[PubMed\]](#)
14. Li, J.Y.; Chen, Y.T.; Chen, Y.; Zhao, J.W.; Fu, J.W.; Shi, M.Z. Sublethal effects of lambda-cyhalothrin on the biological characteristics, detoxification enzymes, and genes of the papaya mealybug, *Paracoccus marginatus*. *J. Pest Sci.* **2024**, *7*, 1–15. [\[CrossRef\]](#)
15. van Leeuwen, T.; Dermauw, W. The molecular evolution of xenobiotic metabolism and resistance in chelicerate mites. *Ann. Rev. Entomol.* **2016**, *61*, 475–498. [\[CrossRef\]](#)
16. Guo, W.; Yang, H.; Duan, M.; Xu, H.; Zhao, W.; Wang, C. Sublethal effects of tolfeprad and its impact on detoxifying enzymes to *Spodoptera exigua*. *Plant Protect.* **2024**, *50*, 168–175. [\[CrossRef\]](#)
17. Batool, N.; Abubakar, M.; Nourelddeen, A.; Naqqash, M.N.; Alghamdi, A.; Al Dhafar, Z.M.; Baakdah, F.; Mozūratīs, R. Toxicity and sublethal effect of chlorantraniliprole on multiple generations of *Aedes aegypti* L. (Diptera: Culicidae). *Insects* **2024**, *15*, 851. [\[CrossRef\]](#)
18. Li, W.; Bashir, N.H.; Naeem, M.; Tian, R.; Tian, X.; Chen, H. Age-stage, two-sex life table of *Atractomorpha lata* (Orthoptera: Pyrgomorphidae) at different temperatures. *Insects* **2024**, *15*, 493. [\[CrossRef\]](#)
19. Borges, I.; Dury, G.J.; Soares, A.O. Population growth parameters of *Scymnus nubilus* fed single-aphid diets of *Aphis fabae* or *Myzus persicae*. *Insects* **2024**, *15*, 486. [\[CrossRef\]](#)
20. Ahn, J.J.; Choi, K.S. Population parameters and growth of *Riptortus pedestris* (Fabricius) (Hemiptera: Alydidae) under fluctuating temperature. *Insects* **2022**, *13*, 113. [\[CrossRef\]](#)
21. Guo, L.; Bian, Q.; Zhang, H.; Gao, X.; Liang, P. Bioassay technique for *Plutella xylostella*: Leaf-dip method. *Chin. J. Appl. Entomol.* **2013**, *50*, 556–560.
22. Jiang, M.; Qian, X.; Zhou, Z.; Liu, Y.; Zhang, M.; Yang, Y. Impacts of sublethal doses of spinetoram on the biological traits and detoxifying enzymes of the tomato leaf miner, *Tuta absoluta* (Lepidoptera: Gelechiidae). *Insects* **2024**, *15*, 990. [\[CrossRef\]](#) [\[PubMed\]](#)
23. Chi, H.; Liu, H.S.I. Two new methods for study of insect population ecology. *Bull. Inst. Zool Acad. Sin.* **1985**, *24*, 225–240.
24. Chi, H. Life-table analysis incorporating both sexes and variable development rates among individuals. *Environ. Entomol.* **1988**, *17*, 26–34. [\[CrossRef\]](#)
25. Chi, H.; Su, H.Y. Age-stage, two-sex life tables of *Aphidius gifuensis* (Ashmead) (Hymenoptera: Braconidae) and its host *Myzus persicae* (Sulzer) (Homoptera: Aphididae) with mathematical proof of the relationship between female fecundity and the net reproductive rate. *Environ. Entomol.* **2006**, *35*, 10–21. [\[CrossRef\]](#)

26. Tuan, S.J.; Lee, C.C.; Chi, H. Population and damage projection of *Spodoptera litura* (F.) on peanuts (*Arachis hypogaea* L.) under different conditions using the age-stage, two-sex life table. *Pest Manag. Sci.* **2014**, *70*, 805–813. [\[CrossRef\]](#)
27. Chi, H. *TWOSEX-MSChart: A Computer Program for the Age-Stage, Two-Sex Life Table Analysis*; National Chung Hsing University in Taiwan: Taichung City, Taiwan, 2023.
28. Efron, B.; Tibshirani, R.J. *An Introduction to the Bootstrap*; Chapman and Hall: New York, NY, USA, 1993.
29. Dean, A.N.; Niemi, J.B.; Tyndall, J.C.; Hodgson, E.W.; O’Neal, M.E. Developing a decision-making framework for insect pest management: A case study using *Aphis glycines* (Hemiptera: Aphididae). *Pest Manag. Sci.* **2021**, *77*, 886–894. [\[CrossRef\]](#)
30. Havasi, M.; Zahedi Golpayegani, A.; Bandani, A. The sublethal concentration of Cyflumetofen adversely affect demographic parameters of *Tetranychus urticae* (Acari: Tetranychidae): Using age-stage, two-sex life tables. *Int. J. Acarol.* **2022**, *48*, 331–337. [\[CrossRef\]](#)
31. Fan, R.; Fan, Z.; Sun, Z.; Chen, Y.; Gui, F. Insecticide susceptibility and detoxification enzyme activity of frankliniella occidentalis under three habitat conditions. *Insects* **2023**, *14*, 643. [\[CrossRef\]](#)
32. Wang, D.; Gong, P.; Li, M.; Qiu, X.; Wang, K. Sublethal effects of spinosad on survival, growth and reproduction of *Helicoverpa armigera* (Lepidoptera: Noctuidae). *Pest Manag. Sci.* **2009**, *65*, 223–227. [\[CrossRef\]](#)
33. Yang, H.; Wang, Z.; Jin, D. Sublethal effects of chlorantraniliprole on the experimental populations of the white-backed planthopper, *Sogatella furcifera* (Hemiptera: Delphacidae). *Acta Entomol. Sin.* **2012**, *55*, 1161–1167. [\[CrossRef\]](#)
34. Ju, D.; Liu, Y.X.; Liu, X.; Dewar, Y.; Mota-Sanchez, D.; Yang, X.Q. Exposure to lambda-cyhalothrin and abamectin drives sublethal and transgenerational effects on the development and reproduction of *Cydia pomonella*. *Ecotoxicol. Environ. Saf.* **2023**, *252*, 114581. [\[CrossRef\]](#) [\[PubMed\]](#)
35. Li, J.Y.; Chen, Y.T.; Wang, Q.Y.; Zheng, L.Z.; Fu, J.W.; Shi, M.Z. Sublethal and transgenerational toxicities of chlorfenapyr on biological traits and enzyme activities of *Paracoccus marginatus* (Hemiptera: Pseudococcidae). *Insects* **2022**, *13*, 874. [\[CrossRef\]](#)
36. Li, D.; Peng, C.; Ma, R.; Chen, Y.; Gui, F.; Sun, Z. Effects of sublethal concentrations chlorfenapyr on growth, development and enzyme activity of *Tuta absoluta*. *J. Environ. Entomol.* **2024**, *46*, 1349–1357. [\[CrossRef\]](#)
37. Bian, D.; Ren, Y.; Ye, W.; Dai, M.; Li, F.; Wei, J.; Sun, H.; Li, B. Evaluation of tolerance to λ -cyhalothrin and response of detoxification enzymes in silkworms reared on artificial diet. *Ecotoxicol. Environ. Saf.* **2022**, *232*, 113232. [\[CrossRef\]](#) [\[PubMed\]](#)
38. Agathokleous, E.; Brown, P.H.; Calabrese, E.J. A gift from parent to offspring: Transgenerational hormesis. *Trends Plant Sci.* **2021**, *26*, 1098–1100. [\[CrossRef\]](#)
39. Ayyanath, M.-M.; Cutler, G.C.; Scott-Dupree, C.D.; Prithiviraj, B.; Kandasamy, S.; Prithiviraj, K. Gene expression during imidacloprid-induced hormesis in green peach aphid. *Dose Response* **2014**, *12*, 480–497. [\[CrossRef\]](#)
40. Alimirzaee, S.; Khajehali, J.; Van Leeuwen, T. Hormetic effects of neonicotinoid insecticides on *Rhizoglyphus robini* (Acari: Acaridae). *Pestic. Biochem. Physiol.* **2023**, *192*, 105396. [\[CrossRef\]](#)
41. Ullah, F.; Gul, H.; Tariq, K.; Desneux, N.; Gao, X.; Song, D. Thiamethoxam induces transgenerational hormesis effects and alteration of genes expression in *Aphis gossypii*. *Pestic. Biochem. Physiol.* **2020**, *165*, 104557. [\[CrossRef\]](#)
42. Wang, L.; Zhu, J.; Wang, Q.; Ji, X.; Wang, W.; Huang, W.; Rui, C.; Cui, L. Hormesis effects of sulfoxaflor on *Aphis gossypii* feeding, growth, reproduction behaviour and the related mechanisms. *Sci. Total Environ.* **2023**, *872*, 162240. [\[CrossRef\]](#)
43. Zhou, C.; Yang, X.; Yang, H.; Gong, M.; Long, G.; Jin, D. Role of SfJHMT and SfFAMeT in the reproductive regulation of *Sogatella furcifera* and its expression under insecticide stress. *Pestic. Biochem. Physiol.* **2021**, *173*, 104779. [\[CrossRef\]](#)
44. Zhou, C.; Yang, X.; Yang, H.; Gong, M.; Long, G.; Jin, D. Role of insecticide-mediated transcription of the TOR and JH signaling pathway-related genes in the regulation of reproduction in *Sogatella furcifera*. *Entomol. Gen.* **2022**, *42*, 771–779. [\[CrossRef\]](#)
45. Hafeez, M.; Liu, S.; Jan, S.; Ali, B.; Shahid, M.; Fernández-Grandón, G.M.; Nawaz, M.; Ahmad, A.; Wang, M. Gossypol-induced fitness gain and increased resistance to deltamethrin in beet armyworm, *Spodoptera exigua* (Hübner). *Pest Manag. Sci.* **2019**, *75*, 683–693. [\[CrossRef\]](#) [\[PubMed\]](#)
46. van Leeuwen, T.; van Pottelberge, S.; Tirry, L. Biochemical analysis of a chlorfenapyr-selected resistant strain of *Tetranychus urticae* Koch. *Pest Manag. Sci.* **2006**, *62*, 425–433. [\[CrossRef\]](#) [\[PubMed\]](#)
47. Zhao, Y.; Wang, Q.; Ding, J.; Wang, Y.; Zhang, Z.; Liu, F.; Mu, W. Sublethal effects of chlorfenapyr on the life table parameters, nutritional physiology and enzymatic properties of *Bradysia odoriphaga* (Diptera: Sciaridae). *Pestic. Biochem. Physiol.* **2018**, *148*, 93–102. [\[CrossRef\]](#)
48. Li, P.R.; Shi, Y.; Ju, D.; Liu, Y.X.; Wang, W.; He, Y.S.; Zhang, Y.Y.; Yang, X.Q. Metabolic functional redundancy of the CYP9A subfamily members leads to P450-mediated lambda-cyhalothrin resistance in *Cydia pomonella*. *Pest Manag. Sci.* **2023**, *79*, 1452–1466. [\[CrossRef\]](#)
49. Kinareikina, A.; Silivanova, E. Impact of insecticides at sublethal concentrations on the enzyme activities in adult *Musca domestica* L. *Toxics* **2023**, *11*, 47. [\[CrossRef\]](#)

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.