

## Expanding utility of cardiac computed tomography in infective endocarditis: A contemporary review

Diarmaid Hughes, Richard Linchangco, Reza Reyaldeem, Bo Xu

**Specialty type:** Radiology, nuclear medicine and medical imaging

**Provenance and peer review:**

Invited article; Externally peer reviewed.

**Peer-review model:** Single blind

**Peer-review report's scientific quality classification**

Grade A (Excellent): 0

Grade B (Very good): B

Grade C (Good): C

Grade D (Fair): D

Grade E (Poor): 0

**P-Reviewer:** Fan YW, China; Teragawa H, Japan

**Received:** January 5, 2022

**Peer-review started:** January 5, 2022

**First decision:** June 16, 2022

**Revised:** June 26, 2022

**Accepted:** July 6, 2022

**Article in press:** July 6, 2022

**Published online:** July 28, 2022



**Diarmaid Hughes, Reza Reyaldeem, Bo Xu**, Section of Cardiovascular Imaging, Robert and Suzanne Tomsich Department of Cardiovascular Medicine, Sydell and Arnold Miller Family Heart and Vascular Institute, Cleveland Clinic, Cleveland, OH 44195, United States

**Richard Linchangco**, Section of Cardiovascular Imaging, Imaging Institute, Cleveland Clinic, Cleveland, OH 44195, United States

**Corresponding author:** Bo Xu, MD, FACC, FASE, Multimodality Imaging Staff Cardiologist, Section of Cardiovascular Imaging, Robert and Suzanne Tomsich Department of Cardiovascular Medicine, Sydell and Arnold Miller Family Heart and Vascular Institute, Cleveland Clinic, 9500 Euclid Avenue, Cleveland, OH 44195, United States. [xub@ccf.org](mailto:xub@ccf.org)

### Abstract

There is increasing evidence on the utility of cardiac computed tomography (CCT) in infective endocarditis (IE) to investigate the valvular pathology, the extra-cardiac manifestations of IE and pre-operative planning. CCT can assist in the diagnosis of perivalvular complications, such as pseudoaneurysms and abscesses, and can help identify embolic events to the lungs or systemic vasculature. CCT has also been shown to be beneficial in the pre-operative planning of patients by delineating the coronary artery anatomy and the major cardiovascular structures in relation to the sternum. Finally, hybrid nuclear/computed tomography techniques have been shown to increase the diagnostic accuracy in prosthetic valve endocarditis. This manuscript aims to provide a contemporary update of the existing evidence base for the use of CCT in IE.

**Key Words:** Infective endocarditis; Cardiac computed tomography; Multimodality cardiac imaging; Cardiovascular structures; Hybrid nuclear

©The Author(s) 2022. Published by Baishideng Publishing Group Inc. All rights reserved.

**Core Tip:** Cardiac computed tomography (CCT) has an expanding role in the management of infective endocarditis (IE). It has been shown to be superior to echocardiography for diagnosing perivalvular complications such as pseudoaneurysms and abscesses. CCT can also diagnose extra-cardiac manifestations of IE such as septic emboli to the lungs. It can assist in pre-operative planning by delineating the coronary anatomy and assessing vascular structures. Herein, we review the role of CCT in IE including the evidence base comparing CCT to echocardiography in diagnosing the valvular complications of IE and the use of CT in IE beyond valvular assessment.

**Citation:** Hughes D, Linchangco R, Reyalde R, Xu B. Expanding utility of cardiac computed tomography in infective endocarditis: A contemporary review. *World J Radiol* 2022; 14(7): 180-193

**URL:** <https://www.wjgnet.com/1949-8470/full/v14/i7/180.htm>

**DOI:** <https://dx.doi.org/10.4329/wjr.v14.i7.180>

## INTRODUCTION

Infective endocarditis (IE) is an infection of the endocardium, heart valves or intra-cardiac devices. It remains a challenging disease to diagnose and manage with high rates of morbidity and mortality[1,2]. Echocardiography remains the main imaging modality used in IE; more recently, however, there is an increasing evidence base for a multimodality imaging approach for IE. Complementary imaging modalities including cardiac computed tomography (CCT) now play increasingly important roles in diagnosis, risk stratification and management of IE. CCT has certain advantages compared to echocardiography in being able to investigate for perivalvular extension, extra-cardiac complications of IE, including metastatic spread and planning for surgery including assessing for coronary artery disease. Advancements in CT technologies, including the use of dedicated cardiac gated four-dimensional CCT, have expanded the applications of CT in IE, demonstrating good sensitivity and specificity for diagnosing the complications of IE. This article aims to review the available evidence for the use of CCT in IE.

## CLINICAL CONSIDERATIONS

The incidence of IE in the United States is estimated to be approximately 15 per 100000 persons annually [3,4], with *Staphylococcus aureus* (SA) being the most common pathogen followed by Viridans group Streptococci[5]. A number of risk factors have been identified for acquiring IE, including the presence of a prosthetic valve, a previous episode of IE, patients with untreated cyanotic congenital heart disease, injection drug use, poor dentition and pre-existing valvular heart disease[6]. The clinical presentation of IE can vary significantly from an acute life threatening illness to a more indolent chronic disease[7]. The most common presenting symptoms are: fever, cardiac murmur, heart failure or complications from septic emboli[8].

The Duke criteria were developed in 1994 to assist in the risk stratification of patients with suspected IE into definite, possible and rejected cases of IE[9]. These criteria have been since validated by a number of retrospective analyses, and underwent further modification in 2000 to reflect changing clinical practice and the emergence of SA as the most common pathogen encountered[10-12]. Despite the updated clinical criteria for diagnosis of IE, there often remains a delay in diagnosis for many patients, commonly due to a lack of microbiological criteria from inappropriate antibiotic use, with worse outcomes seen in these patients[13,14]. Advances in CT technologies including improvements in both temporal and spatial resolution have enabled greater use of CCT for the diagnosis of IE. The European Society of Cardiology guidelines for the management of infective endocarditis reflect these advances in imaging techniques and include paravalvular lesions detected by CCT to be a major imaging criterion [15]. The 2020 American College of Cardiology/American Heart Association Guideline for the management of patients with valvular heart disease also recommend the use of CCT as an adjunctive imaging modality for IE[16].

### Echocardiography

Echocardiography, including transthoracic echocardiography (TTE) and transesophageal echocardiography (TEE) where appropriate, remain the first line imaging modality for the diagnosis and monitoring of IE[17,18]. There are three main echocardiographic findings that are considered major criteria for the diagnosis of IE: vegetations, abscesses/pseudoaneurysms and new dehiscence of a prosthetic valve[15]. In native valve endocarditis, the sensitivities for the diagnosis of a vegetation are approximately 70% for TTE and 96% for TEE, respectively[17]. For prosthetic valve endocarditis (PVE),

the sensitivities for diagnosing a vegetation are approximately 50% for TTE and 92% for TEE, respectively[17]. There are many challenges in the diagnosis of IE with echocardiography, including small vegetations or embolization of the vegetation prior to imaging, difficulty visualizing the lesion in the setting of pre-existing valvular disease or with prosthetic valves. Small abscesses may also be challenging to diagnose, especially by TTE[19]. There are also many mimics of IE that could result in a false positive diagnosis, such as Lambl's excrescence, fibroelastomas, thrombus, degenerative lesions, prosthetic material/sutures or marantic lesions[20]. CCT can therefore be helpful to assist in the diagnosis of IE, when there are equivocal findings by echocardiography or in challenging cases involving prosthetic valves[21].

### ***Dedicated cardiac CT protocol for endocarditis evaluation***

For evaluation of cardiac valves, multiphase imaging using a retrospectively ECG-gated acquisition is required to obtain an isotropic data set. Images are acquired in spiral mode utilizing a low pitch of 0.16 to 0.5 during 5-10 R-R intervals with a section thickness of 0.60 mm. Thin collimation is used for optimal visualization of the valve leaflets, typically 64 mm × 0.6 mm. The tube voltage used is adapted to the patient's weight, and can vary between 100 to 120 kV. Gantry rotation time of 0.28 s to 0.35 s is used[22-27]. The scan is typically performed in a limited field of view from the level of the carina to the cardiac apex.

Timing of the iodinated contrast bolus is important to optimize visualization of the involved cardiac valves. A monophasic contrast injection is most commonly used, with timing of the contrast bolus chosen for optimal visualization of the expected involved valve and cardiac chambers. Alternatively, biphasic contrast injection may be performed, which allows evaluation of all cardiac chambers and valves[28]. Premedication with beta blockers may be used to regulate heart rate to less than 65-70 bpm if not contraindicated. This improves image quality by reducing artefacts related to cardiac motion and valvular motion.

The isotropic data set acquired from the retrospectively gated acquisition allows for reconstruction in any desired plane. In addition to static images, imaging at multiple points during the cardiac cycle also allows for creation of 4D cine images, allowing for evaluation of valve leaflet motion and planimetry.

Because images are acquired throughout the entire cardiac cycle, this results in a significantly higher radiation dose penalty compared to prospectively gated CT as used typically in CT angiography of the coronary arteries. Radiation dose may be lowered utilizing methods such as iterative reconstruction and ECG-triggered radiation dose modulation. However, ECG-triggered dose modulation may result in suboptimal evaluation during the phase of reduced tube current, typically the systolic phase[29]. There are specific protocols used for visualizing the various cardiac and extra manifestations of IE and for pre-operative planning. Herein, we group all of these into an umbrella term of CCT, referring to ECG-gated CT of the chest with contrast. There are some situations, such as during investigation for septic emboli to the visceral organs, when abdominal imaging may also be needed.

---

## **UTILITY OF CARDIAC CT IN INFECTIVE ENDOCARDITIS**

---

CCT has the ability to assess for valvular lesions, perivalvular extension, metastatic spread/embolization, as well as aortic anatomy. CCT may also be used in appropriate cases for the assessment of coronary artery disease[30]. This is particularly relevant in patients with aortic valve IE and vegetations, whereby invasive coronary angiography may be relatively contraindicated, due to the potential risk of causing the vegetations to embolize during the procedure. CCT can detect valvular lesions, such as vegetations, prosthetic valve dehiscence in addition to perivalvular lesions such as abscesses, fistulae and pseudoaneurysms[26,31]. Table 1 compares the various definitions for IE detected on CCT *vs* TEE.

---

## **PERIVALVULAR COMPLICATIONS**

---


### ***Pseudoaneurysms and abscesses***

Perivalvular extension of IE, which includes pseudoaneurysms, abscesses and fistulae are associated with a higher rate of operative management and mortality[32-34]. A pseudoaneurysm is a perivalvular cavity that is in communication with the cardiovascular lumen which results from an abscess rupturing into a cavity[17]. On echocardiography, this appears as a pulsatile echo-free space with detectable Doppler color flow, while on CCT, it appears as an abnormal cavity close to the valve with direct communication with the heart chambers or major blood vessel[30]. An abscess is a closed cavity with necrosis and purulent material not in communication with a cardiovascular cavity[17]. On echocardiography, this appears as a thickened perivalvular area with a homogenous echo-dense or echo-lucent appearance. On CCT, abscesses appear as perivalvular collections of fluid encased in a thick layer of inflammatory tissue enhanced by the injection of contrast medium. See Figures 1-4 for examples of

**Table 1 Comparison of cardiac computed tomography vs transesophageal echocardiography findings in infective endocarditis[15,30]**

	CCT	TEE
Vegetation	An irregular mass or thickening associated with the endocardium, native valve or prosthetic valve with low to intermediate attenuation	Mobile or non-mobile intracardiac mass on valve or other endocardial structures, including on implanted intracardiac material
Pseudoaneurysm	Perivalvular collection of contrast enhanced material usually adjacent to a valve with a visible direct communication	Abnormal perivalvular echo-free space with color-Doppler flow showing connection with the cardiovascular lumen
Abscess	Usually perivalvular collection of low attenuation material. Often has a thick layer of tissue in the wall of the collection that enhances with contrast	Usually perivalvular collection that can have an echodense or echolucent appearance without a communication to a lumen
Dehiscence of a prosthetic valve	Prosthetic valve misalignment with a tissue defect between the annulus and prosthesis	Evidence of excessive motion of a prosthetic valve. Occasionally, it is possible to see a defect between annulus and prosthesis and/or evidence of paravalvular leak on Doppler assessment
Perforation	Leaflet tissue defect that can be observed in two different views	Defect in a valve leaflet that may be seen visually as an interruption of tissue or by color flow across the defect
Fistula	An abnormal communication between two cardiac chambers that is contrast filled	An abnormal connection two neighboring lumen detected by color Doppler flow

CCT: Cardiac computed tomography; TEE: Transesophageal echocardiography.



DOI: 10.4329/wjr.v14.i7.180 Copyright © The Author(s) 2022.

**Figure 1 Prosthetic aortic valve infective endocarditis with valve dehiscence and pseudoaneurysm.** Patient with extensive peri-aortic root abscess cavity/pseudo-aneurysm (stars) related to prosthetic valve endocarditis. Note the extensive peri-valvular space around the prosthetic aortic valve and evidence of valve dehiscence on transesophageal echocardiography. Comparable short-axis and long-axis images from cardiac computed tomography and short-axis and long-axis images from transesophageal echocardiography.

comparisons of TEE and CCT images in patients with perivalvular complications of IE (Table 2).


A 2009 study by Feuchter *et al*[24] to investigate the value of CCT for the assessment of valvular abnormalities included 37 consecutive patients (29 of whom went on to have surgery) with clinically suspected IE who underwent both CCT and TEE. CCT identified all pseudoaneurysms and abscesses in this study with sensitivity and specificity of 100%, which was superior to TEE (sensitivity of 89% and a specificity of 100%)[24]. CCT was also shown to be superior to TEE for perivalvular extension of the IE, identifying myocardial and pericardial extension more often than TEE[24]. In a 2009 prospective study,



**Table 2 Strengths and limitations of various imaging modalities for assessing infective endocarditis**

Modality	CCT	TTE	TEE	PET/CT
Strengths	Ability to image the entire thorax; Improved detection of perivalvular complications; CAD Assessment; Pre-Operative planning; Detection of extra-cardiac emboli	Good Spatial resolution; Availability and portable; Low cost; Lack of radiation; Lack of contrast; Chamber quantification; Assess hemodynamics	Improved spatial and temporal resolution over TTE; Availability and low cost; Lack of radiation; Lack of contrast; Better sensitivity than TTE in PVE; Assess Hemodynamics	Improved detection of perivalvular complications; Improved diagnostic accuracy in PVE detection of embolic events
Weaknesses	Higher cost; Radiation exposure; Nephrotoxicity; Lower sensitivity for small vegetations and leaflet perforation; Availability may be limited	Limited value in PVE; No tissue characterization; Low sensitivity for peri-valvular complications	No tissue characterization; May miss some peri-valvular complications; Invasive procedure requiring sedation (cannot be performed in some patients with esophageal issues)	Limited availability; Higher cost; Radiation exposure

CCT: Cardiac computed tomography; TEE: Transesophageal echocardiography; TTE: Transthoracic echocardiography; PET: Positron emission tomography; PVE: Prosthetic valve endocarditis; CAD: Coronary artery disease.




DOI: 10.4329/wjr.v14.i7.180 Copyright © The Author(s) 2022.

**Figure 2 Native mitral valve infective endocarditis with pseudoaneurysm.** A and B: Mitral annular pseudo-aneurysm (dotted box) detected on cardiac computed tomography; C: Cardiac computed tomography with evidence of native mitral valve endocarditis on transesophageal echocardiography (star); D: Cardiac computed tomography with a prominent peri-annular cavity, consistent with a mitral annular pseudo-aneurysm.

19 patients with aortic valve endocarditis requiring surgical intervention underwent CCT pre-operatively[25]. The majority of patients (approx. 90%) had native valve IE. This study showed that CCT had sensitivity and specificity for diagnosing pseudoaneurysms of 100% and 92%, respectively, and CCT correctly identified all cases where there was extension of IE into the intervalvular fibrosa[25]. This paper did not report the TEE findings for their participants [25].

A paper by Fagman *et al*[23] reported in 2012 on 27 consecutive patients who had TEE findings of aortic valve PVE and investigated the strength of agreement between the TEE and CT results. They found a strength of agreement compared to TEE was 0.68 for abscesses and 0.75 for dehiscence[23]. However, using surgery as the reference standard (16 patients went on to have surgery), CCT had sensitivity of 91% to detect pseudoaneurysms /abscesses compared to 82% for TEE[23]. A 2013 study investigated the additional value of CCT beyond the usual evaluation with TEE in PVE in 28 patients, with a final diagnosis being either determined clinically or at the time of surgery as reported by Habets *et al*[35]. They reported that usual evaluation had sensitivity of 68% for detecting periannular complica-



DOI: 10.4329/wjr.v14.i7.180 Copyright © The Author(s) 2022.


**Figure 3 Mechanical aortic valve infective endocarditis with aortic root abscess.** A and B: Patient with extensive peri-aortic root abscess, with a contrast outpouching and surrounding soft-tissue density (star) nearly encasing the left main artery; C and D: Transesophageal echocardiography demonstrated similar findings, although not as well defined due to shadowing from the aortic valve prosthesis.

ations (mycotic aneurysms and abscesses), which was increased to 100% with the use of CCT[35]. Koo *et al*[27] also compared CCT *vs* TEE using intra-operative findings as the reference standard in 2018. They enrolled 49 patients, 12 of whom had PVE[27]. The overall detection of IE by CCT was 94%, compared to 96% by TEE[27]. CCT performed better than TEE at detecting abscess/pseudoaneurysms, with sensitivities of 60% for CCT and 40% for TEE, respectively[27]. A retrospective study from 2018 by Sims *et al* [36] investigated the performance of CCT in the pre-operative evaluation of IE. In total, they had 251 patients undergoing TEE with 34 of these patients also having a CCT[36]. The sensitivity of CCT for detecting abscesses/pseudoaneurysms was 91%, which was superior to TEE at 78%[36]. CCT was reported to have a lower sensitivity for detecting fistulae at 50% *vs* 79% for TEE, and dehiscence at 57% *vs* 70% for TEE[36].

Two studies from 2018 (Ouchi *et al*[31] and Koneru *et al*[37]) retrospectively investigated the utility of CCT in IE with intra-operative findings as the reference standard. CCT performed better than TEE in detecting abscess/pseudoaneurysm in prosthetic valves with sensitivity of 81% (versus 64% for TEE) in the study by Koneru *et al*[31]. CCT had sensitivity of 100% sensitivity for detecting perivalvular complications, such as pseudoaneurysms in the paper by Ouchi *et al*[37]. A 2019 study by Hryniewicz *et al*[26] investigated 53 consecutive patients who had perivalvular complications from IE, who also underwent CCT and TEE pre-operatively. They showed the sensitivity and specificity for detecting abscesses/pseudoaneurysms for CCT were 81% and 90%, respectively, compared to 63% and 90%, respectively for TEE[26]. A 2020 study of 68 patients reported by Sifaoui *et al* with definite left-sided IE who underwent CCT and TEE reported the comparison of CCT and TEE to detect perivalvular complications[38]. They showed again that CT had a higher sensitivity for detecting pseudoaneurysms at 100%, compared to TEE at 67%[38].

Overall, the current evidence base suggests that the diagnostic performance of CCT is likely superior to that of TEE for the detection of pseudoaneurysms and abscesses in appropriately selected cases. A recent meta-analysis reported pooled sensitivity and specificity for CCT for the detection of peri-annular complications of 88% and 93%, respectively, compared to TEE at 70% and 96%, respectively[39].

The identification of perivalvular complications is important for prognostic and management considerations. These sequelae of invasive IE, which are more common with aortic valve endocarditis and PVE, have been associated with increased rates of surgical management, and may confer an increased risk of mortality[32,33]. Therefore, a multimodality imaging strategy for IE that includes CCT would have the ability to identify more of these complications, compared to using TEE alone, and therefore impact on decision making for patients.




DOI: 10.4329/wjrv.v14.i7.180 Copyright © The Author(s) 2022.

**Figure 4** Aortic repair with left ventricular outflow tract pseudoaneurysm. A and B: This patient had a complex aortic repair with a clear peri-valvular space (arrow) on transesophageal echocardiography (TEE); however, the exact origin was difficult to define by TEE imaging; C and D: Cardiac computed tomography demonstrated multiple pseudo-aneurysms arising from the left ventricular outflow tract (LVOT) with the largest, fistulous LVOT pseudo-aneurysm (arrow) best appreciated on 3-dimensional volume-rendering.

### Vegetations

A vegetation is a mass-like lesion of infected material composed of fibrin, platelets and microorganisms attached to an endocardial structure or on an implanted cardiac device (CIED)[40,41]. On echocardiography, this appears as an oscillating or non-oscillating intracardiac echodensity, which can be attached to a valve, other endocardial surface or cardiac device[15]. A vegetation tends to move with the cardiac cycle, and is more frequently found on the atrial side of the atrioventricular valves and the ventricular side of the semi-lunar valves[15]. On CT, vegetations appear as hypodense homogeneous irregular masses, which can be attached to a valve or other cardiac structures[30].

In the 2009 paper by Feuchter *et al*[24] using surgical/pathological diagnosis as the reference standard, CCT had sensitivity of 96% and specificity of 97% for the diagnosis of vegetations. 5 vegetations were missed by CCT (11%) either due to artefact or small size ( $\leq 4\text{mm}$ )[24]. The performance of TEE was similar to CCT with sensitivity of 96% and specificity of 100%[24]. CCT was found to be inferior to TEE for detecting leaflet perforations[24]. The study by Gahide *et al*[25] on aortic valve IE showed CCT had a sensitivity of 71% and a specificity of 100% for detecting vegetations, though the sensitivity was increased to 100% for large vegetations ( $> 10\text{ mm}$ ). The 2012 paper by Fagman *et al*[23] found that CCT detected vegetations in 7 out of 13 cases (54%), with a lower detection rate being potentially explained by artefact from the prosthetic valves obscuring the CCT images. The 2013 study by Habets *et al*[35] found additional benefit with CCT in addition to usual work-up with TEE in PVE, with a final diagnosis being either determined clinically or at the time of surgery. They reported that usual work-up had sensitivity of 63% for detecting vegetations, which was increased to 100% with the use of CCT[35]. The 2018 study by Koo *et al*[27] reported sensitivity for CCT to detect vegetations of 91%, compared to 100% by TEE. Missed vegetations were smaller, and the authors also listed motion artefact and beam hardening from mechanical valves as reasons for the failure of CCT to detect the vegetations[27]. Sims *et al*[36] reported the sensitivity for detecting vegetations to be 70% for CCT (34 patients) and 96% for TEE (251 patients). The study by Oucho *et al*[37] reported sensitivity of 92% for detecting vegetations for CCT, correctly identifying 12 of 13 cases who had vegetations confirmed at the time of surgery. The retrospective review on 122 patients by Koneru *et al*[31] showed TEE to have a statistically significantly higher sensitivity for detecting vegetations compared to CCT at 85% *vs* 16%, though CCT did have a higher specificity at 96% compared to 69% for TEE. The lower sensitivity in detecting vegetations by CCT in this study may be related to only reviewing single-phase images, and the fact that the slice thickness used was 3 mm which was thicker than the other studies referenced above[31]. In the 2019 paper by Hryniewicz *et al*[26] the sensitivity and specificity for detecting



DOI: 10.4329/wjrv.v14.i7.180 Copyright © The Author(s) 2022.

**Figure 5 Prosthetic aortic valve with hypoattenuating lesion.** A-C: This patient presented with elevated prosthetic aortic valve gradients, and although transesophageal echocardiography imaging was suspicious for an echodensity (white arrow), imaging quality was challenging due to prosthetic metallic valve shadowing; D-F: Multi-phase 4-dimensional cardiac computed tomography was utilized which clearly demonstrated a hypoattenuating mass (white arrows) attached to the prosthetic valve. Despite suspicion for infection, intra-operative pathology revealed thrombus.

vegetations by CT were 89% and 71%, respectively, compared to TEE at 97% and 42%, respectively. The 2020 study by Sifaoui *et al*[38] showed that TEE had a higher area under the curve (AUC) than CCT for detecting vegetations, with AUC for TEE of 0.86 *vs* AUC for CT of 0.69.


Overall, the current evidence base suggests that TEE is overall superior to CCT for the detection of vegetations, particular small vegetations, with pooled sensitivity for TEE from a recent meta-analysis of 94%[42]. CCT demonstrated a lower pooled sensitivity, at 64% for the detection of vegetations[42]. There was a wide range of results reported likely related to small sample sizes, differing patient populations and different protocols used for imaging. While CCT should not replace echocardiography as the first line imaging tool in the majority of patients primarily to detect vegetations, in a small subset of patients who could not undergo clinical indicated TEE (*e.g.*, esophageal pathology), CCT may add diagnostic value. CCT has also been shown to improve the diagnostic accuracy overall, when used in combination with TEE[35]. For example Figure 5 shows a hypoattenuating lesion on a mechanical aortic valve that was more clearly defined on CCT. Figures 6 and 7 shows an examples of an aortic graft and aortic stent infections that can be difficult to image with TEE.

## INCREMENTAL VALUE OF CCT IN INFECTIVE ENDOCARDITIS

### Pre-operative assessment


In addition to the advantages related to the management of IE as outlined above, CCT can also assist in the pre-operative planning of IE surgery. In patients with prior cardiothoracic surgery, CCT can delineate the relationship of cardiovascular structures to the sternum and the location of the coronary artery bypass grafts (CABG). See Figure 8 for CCT images of a patient with a prior CABG. For all patients, CCT can identify calcification of the ascending aorta ('porcelain aorta'), which may preclude surgery as well as give precise anatomic location and extent of the degree of calcification of the subclavian, axillary and femoral arteries. The advantage of having a pre-operative CCT was described by Merlo *et al*[43] with reported lower rates of stroke and mortality in patients undergoing pre-operative CCT followed by primary cardiac surgery, *vs* those without pre-operative CCT imaging. Figure 9 shows a CCT in a patient with vascular calcification.





DOI: 10.4329/wjrv.v14.i7.180 Copyright © The Author(s) 2022.

**Figure 6 Aortic graft infection.** A and B: Infection of aortic grafts can be difficult to visualize on transesophageal echocardiography (TEE); C and D: This case demonstrates evidence of a peri-aortic graft echolucent space with stranding, which was challenging to image on TEE, and further characterization with cardiac computed tomography clearly demonstrated peri-aortic graft thickening (star) consistent with infection.




DOI: 10.4329/wjrv.v14.i7.180 Copyright © The Author(s) 2022.

**Figure 7 Aortic endovascular stent infection.** Aortic endovascular stents are also prone to infection, and negative transesophageal echocardiography for valvular vegetations with a high suspicion should prompt consideration for other sources of infections, as in this case, which was demonstrated on cardiac computed tomography with prominent soft-tissue thickening (star) around the proximal descending thoracic aorta stent.

### Pre-operative assessment


In addition to the advantages related to the management of IE as outlined above, CCT can also assist in the pre-operative planning of IE surgery[44-50]. In patients with prior cardiothoracic surgery, CCT can delineate the relationship of cardiovascular structures to the sternum and the location of the coronary artery bypass grafts (CABG). See Figure 8 for CCT images of a patient with a prior CABG[50-60]. For all patients, CCT can identify calcification of the ascending aorta ('porcelain aorta'), which may preclude surgery as well as give precise anatomic location and extent of the degree of calcification of the subclavian, axillary and femoral arteries[61,62]. The advantage of having a pre-operative CCT was described by Merlo *et al*[43] with reported lower rates of stroke and mortality in patients undergoing pre-operative CCT followed by primary cardiac surgery, *vs* those without pre-operative CCT imaging. Figures 9 and 10 shows a CCT in a patient with vascular calcification.






DOI: 10.4329/wjr.v14.i7.180 Copyright © The Author(s) 2022.

**Figure 8 Prior coronary artery bypass graft.** A: Cardiac computed tomography has added utility for pre-operative planning by identifying areas of aortic calcification (which in this case was prominent at the aortic root graft); B and C: Relevant sternal distance for cardiovascular structures, such as the left brachiocephalic vein and right ventricle, as well as ilio-femoral anatomy in cases where peripheral vascular access may be needed.



DOI: 10.4329/wjr.v14.i7.180 Copyright © The Author(s) 2022.

**Figure 9 Aortic calcification and thoracic structures.** Cardiac computed tomography is also important to identify prior coronary artery bypass graft locations (stars) and sternal distance to avoid complications, which can be increased in redo-surgery particularly with increased adhesions and friable structures due to infection.



DOI: 10.4329/wjr.v14.i7.180 Copyright © The Author(s) 2022.

**Figure 10 Embolic phenomena of infective endocarditis.** A: It demonstrates multiple pulmonary septic emboli, including cavitory lesions (star) from tricuspid valve endocarditis. Cardiac computed tomography is also incremental in demonstrating embolic phenomena which can have implications for surgical urgency and overall prognosis; B: It demonstrates a splenic infarct (arrow) from mitral valve endocarditis).

## CONCLUSION

With improvements in the temporal and spatial resolution of CCT technology, including the use of dedicated 4D CCT, there has been an expanding role of CCT imaging in IE. CCT has been shown to be superior to TEE for the identification of pseudoaneurysms and abscesses in appropriately selected cases,

while the combination of both modalities results in the greatest sensitivity for detection. TEE is superior to CCT for small vegetations; however this advantage is less marked for larger vegetations. In addition, CCT has a number of adjunctive uses in IE beyond evaluation of valvular pathology. CCT can aid in the diagnosis of embolic events, such as pulmonary complications in RSIE. It can also be used to diagnose significant CAD in low to intermediate risk patients preoperatively, or when there is a contraindication to ICA, such as when there is a large aortic valve vegetation. CCT can also be helpful for pre-operative planning to assess the relationship of the cardiovascular structures in relation to the sternum, which is particularly helpful in re-do sternotomy cases. The addition of hybrid techniques such as positron emission computed tomography or SPECT/CT, has been shown to improve the diagnostic accuracy in challenging cases of PVE. Greater awareness of the strengths, weaknesses and appropriate applications of CCT in IE will assist in its optimal use for improved diagnosis and management of this challenging condition.

## FOOTNOTES

**Author contributions:** Hughes D, Lingchangco R, Reyalden R and Xu B were involved in the conceptualization, writing, revision and final approval of the manuscript.

**Conflict-of-interest statement:** No conflicts of interest to declare for all authors for this article.

**Open-Access:** This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

**Country/Territory of origin:** United States

**ORCID number:** Diarmaid Hughes 0000-0003-3110-6759; Richard Linchangco 0000-0002-9265-9851; Reza Reyalden 0000-0001-7501-2354; Bo Xu 0000-0002-2985-7468.

**S-Editor:** Wang LL

**L-Editor:** A

**P-Editor:** Wang LL

## REFERENCES

- 1 Sy RW, Kritharides L. Health care exposure and age in infective endocarditis: results of a contemporary population-based profile of 1536 patients in Australia. *Eur Heart J* 2010; **31**: 1890-1897 [PMID: 20453066 DOI: 10.1093/eurheartj/ehq110]
- 2 Mansur AJ, Grinberg M, da Luz PL, Bellotti G. The complications of infective endocarditis. A reappraisal in the 1980s. *Arch Intern Med* 1992; **152**: 2428-2432 [PMID: 1456853]
- 3 Pant S, Patel NJ, Deshmukh A, Golwala H, Patel N, Badheka A, Hirsch GA, Mehta JL. Trends in infective endocarditis incidence, microbiology, and valve replacement in the United States from 2000 to 2011. *J Am Coll Cardiol* 2015; **65**: 2070-2076 [PMID: 25975469 DOI: 10.1016/j.jacc.2015.03.518]
- 4 Toyoda N, Chikwe J, Itagaki S, Gelijns AC, Adams DH, Egorova NN. Trends in Infective Endocarditis in California and New York State, 1998-2013. *JAMA* 2017; **317**: 1652-1660 [PMID: 28444279 DOI: 10.1001/jama.2017.4287]
- 5 Murdoch DR, Corey GR, Hoen B, Miró JM, Fowler VG Jr, Bayer AS, Karchmer AW, Olaison L, Pappas PA, Moreillon P, Chambers ST, Chu VH, Falcó V, Holland DJ, Jones P, Klein JL, Raymond NJ, Read KM, Tripodi MF, Utili R, Wang A, Woods CW, Cabell CH; International Collaboration on Endocarditis-Prospective Cohort Study (ICE-PCS) Investigators. Clinical presentation, etiology, and outcome of infective endocarditis in the 21st century: the International Collaboration on Endocarditis-Prospective Cohort Study. *Arch Intern Med* 2009; **169**: 463-473 [PMID: 19273776 DOI: 10.1001/archinternmed.2008.603]
- 6 Strom BL, Abrutyn E, Berlin JA, Kinman JL, Feldman RS, Stolley PD, Levison ME, Korzeniowski OM, Kaye D. Risk factors for infective endocarditis: oral hygiene and nondental exposures. *Circulation* 2000; **102**: 2842-2848 [PMID: 11104742 DOI: 10.1161/01.cir.102.23.2842]
- 7 Baddour LM, Wilson WR, Bayer AS, Fowler VG Jr, Tleyjeh IM, Rybak MJ, Barsic B, Lockhart PB, Gewitz MH, Levison ME, Bolger AF, Steckelberg JM, Baltimore RS, Fink AM, O'Gara P, Taubert KA; American Heart Association Committee on Rheumatic Fever, Endocarditis, and Kawasaki Disease of the Council on Cardiovascular Disease in the Young, Council on Clinical Cardiology, Council on Cardiovascular Surgery and Anesthesia, and Stroke Council. Infective Endocarditis in Adults: Diagnosis, Antimicrobial Therapy, and Management of Complications: A Scientific Statement for Healthcare Professionals From the American Heart Association. *Circulation* 2015; **132**: 1435-1486 [PMID: 26373316 DOI: 10.1161/CIR.0000000000000296]
- 8 Bayer AS. Infective endocarditis. *Clin Infect Dis* 1993; **17**: 313-20; quiz 321 [PMID: 8218670 DOI: 10.1093/clinids/17.3.313]

- 9 **Durack DT**, Lukes AS, Bright DK. New criteria for diagnosis of infective endocarditis: utilization of specific echocardiographic findings. Duke Endocarditis Service. *Am J Med* 1994; **96**: 200-209 [PMID: [8154507](#) DOI: [10.1016/0002-9343\(94\)90143-0](#)]
- 10 **Sekeres MA**, Abrutyn E, Berlin JA, Kaye D, Kinman JL, Korzeniowski OM, Levison ME, Feldman RS, Strom BL. An assessment of the usefulness of the Duke criteria for diagnosing active infective endocarditis. *Clin Infect Dis* 1997; **24**: 1185-1190 [PMID: [9195080](#) DOI: [10.1086/513657](#)]
- 11 **Hoen B**, Béguinot I, Rabaud C, Jaussaud R, Selton-Suty C, May T, Canton P. The Duke criteria for diagnosing infective endocarditis are specific: analysis of 100 patients with acute fever or fever of unknown origin. *Clin Infect Dis* 1996; **23**: 298-302 [PMID: [8842267](#) DOI: [10.1093/clinids/23.2.298](#)]
- 12 **Li JS**, Sexton DJ, Mick N, Nettles R, Fowler VG Jr, Ryan T, Bashore T, Corey GR. Proposed modifications to the Duke criteria for the diagnosis of infective endocarditis. *Clin Infect Dis* 2000; **30**: 633-638 [PMID: [10770721](#) DOI: [10.1086/313753](#)]
- 13 **Fukuchi T**, Iwata K, Ohji G. Failure of early diagnosis of infective endocarditis in Japan--a retrospective descriptive analysis. *Medicine (Baltimore)* 2014; **93**: e237 [PMID: [25501088](#) DOI: [10.1097/MD.0000000000000237](#)]
- 14 **Naderi HR**, Sheybani F, Erfani SS. Errors in diagnosis of infective endocarditis. *Epidemiol Infect* 2018; **146**: 394-400 [PMID: [29310727](#) DOI: [10.1017/S0950268817002977](#)]
- 15 **Habib G**, Lancellotti P, Antunes MJ, Bongiorni MG, Casalta JP, Del Zotti F, Dulgheru R, El Khoury G, Erba PA, Iung B, Miro JM, Mulder BJ, Plonska-Gosciniak E, Price S, Roos-Hesselink J, Snygg-Martin U, Thuny F, Tornos Mas P, Vilacosta I, Zamorano JL; ESC Scientific Document Group. 2015 ESC Guidelines for the management of infective endocarditis: The Task Force for the Management of Infective Endocarditis of the European Society of Cardiology (ESC). Endorsed by: European Association for Cardio-Thoracic Surgery (EACTS), the European Association of Nuclear Medicine (EANM). *Eur Heart J* 2015; **36**: 3075-3128 [PMID: [26320109](#) DOI: [10.1093/eurheartj/ehv319](#)]
- 16 **Otto CM**, Nishimura RA, Bonow RO, Carabello BA, Erwin JP 3rd, Gentile F, Jneid H, Krieger EV, Mack M, McLeod C, O'Gara PT, Rigolin VH, Sundt TM 3rd, Thompson A, Toly C. 2020 ACC/AHA Guideline for the Management of Patients With Valvular Heart Disease: Executive Summary: A Report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *Circulation* 2021; **143**: e35-e71 [PMID: [33332149](#) DOI: [10.1161/CIR.0000000000000932](#)]
- 17 **Habib G**, Badano L, Tribouilloy C, Vilacosta I, Zamorano JL, Galderisi M, Voigt JU, Sicari R, Cosyns B, Fox K, Aakhus S; European Association of Echocardiography. Recommendations for the practice of echocardiography in infective endocarditis. *Eur J Echocardiogr* 2010; **11**: 202-219 [PMID: [20223755](#) DOI: [10.1093/ejehocord/jeq004](#)]
- 18 **Haq IU**, Haq I, Griffin B, Xu B. Imaging to evaluate suspected infective endocarditis. *Cleve Clin J Med* 2021; **88**: 163-172 [PMID: [33648969](#) DOI: [10.3949/ccjm.88a.19142](#)]
- 19 **Daniel WG**, Mügge A, Martin RP, Lindert O, Hausmann D, Nonnast-Daniel B, Laas J, Lichtlen PR. Improvement in the diagnosis of abscesses associated with endocarditis by transesophageal echocardiography. *N Engl J Med* 1991; **324**: 795-800 [PMID: [1997851](#) DOI: [10.1056/NEJM199103213241203](#)]
- 20 **Zmaili MA**, Alzubi JM, Kocyigit D, Bansal A, Samra GS, Grimm R, Griffin BP, Xu B. A Contemporary 20-Year Cleveland Clinic Experience of Nonbacterial Thrombotic Endocarditis: Etiology, Echocardiographic Imaging, Management, and Outcomes. *Am J Med* 2021; **134**: 361-369 [PMID: [32827467](#) DOI: [10.1016/j.amjmed.2020.06.047](#)]
- 21 **Lo Presti S**, Elajami TK, Zmaili M, Reyaldéen R, Xu B. Multimodality imaging in the diagnosis and management of prosthetic valve endocarditis: A contemporary narrative review. *World J Cardiol* 2021; **13**: 254-270 [PMID: [34589164](#) DOI: [10.4330/wjc.v13.i8.254](#)]
- 22 **Altiok E**, Koos R, Schröder J, Brehmer K, Hamada S, Becker M, Mahnken AH, Almalla M, Dohmen G, Autschbach R, Marx N, Hoffmann R. Comparison of two-dimensional and three-dimensional imaging techniques for measurement of aortic annulus diameters before transcatheter aortic valve implantation. *Heart* 2011; **97**: 1578-1584 [PMID: [21700756](#) DOI: [10.1136/hrt.2011.223974](#)]
- 23 **Fagman E**, Perrotta S, Bech-Hanssen O, Flinck A, Lamm C, Olaison L, Svensson G. ECG-gated computed tomography: a new role for patients with suspected aortic prosthetic valve endocarditis. *Eur Radiol* 2012; **22**: 2407-2414 [PMID: [22622348](#) DOI: [10.1007/s00330-012-2491-5](#)]
- 24 **Feuchtner GM**, Stolzmann P, Dichtl W, Schertler T, Bonatti J, Scheffl H, Mueller S, Plass A, Mueller L, Bartel T, Wolf F, Alkadh H. Multislice computed tomography in infective endocarditis: comparison with transesophageal echocardiography and intraoperative findings. *J Am Coll Cardiol* 2009; **53**: 436-444 [PMID: [19179202](#) DOI: [10.1016/j.jacc.2008.01.077](#)]
- 25 **Gahide G**, Bommart S, Demaria R, Sportouch C, Dambia H, Albat B, Vernhet-Kovacsik H. Preoperative evaluation in aortic endocarditis: findings on cardiac CT. *AJR Am J Roentgenol* 2010; **194**: 574-578 [PMID: [20173130](#) DOI: [10.2214/AJR.08.2120](#)]
- 26 **Hryniewicz T**, Zatorska K, Abramczuk E, Zakrzewski D, Szymański P, Kuśmierczyk M, Michałowska I. The usefulness of cardiac CT in the diagnosis of perivalvular complications in patients with infective endocarditis. *Eur Radiol* 2019; **29**: 4368-4376 [PMID: [30643945](#) DOI: [10.1007/s00330-018-5965-2](#)]
- 27 **Koo HJ**, Yang DH, Kang JW, Lee JY, Kim DH, Song JM, Kang DH, Song JK, Kim JB, Jung SH, Choo SJ, Chung CH, Lee JW, Lim TH. Demonstration of infective endocarditis by cardiac CT and transoesophageal echocardiography: comparison with intra-operative findings. *Eur Heart J Cardiovasc Imaging* 2018; **19**: 199-207 [PMID: [28329276](#) DOI: [10.1093/ehjci/jex010](#)]
- 28 **Vrachliotis TG**, Bis KG, Haidary A, Kosuri R, Balasubramaniam M, Gallagher M, Raff G, Ross M, O'neil B, O'Neill W. Atypical chest pain: coronary, aortic, and pulmonary vasculature enhancement at biphasic single-injection 64-section CT angiography. *Radiology* 2007; **243**: 368-376 [PMID: [17400761](#) DOI: [10.1148/radiol.2432060447](#)]
- 29 **Jakobs TF**, Becker CR, Ohnesorge B, Flohr T, Suess C, Schoepf UJ, Reiser MF. Multislice helical CT of the heart with retrospective ECG gating: reduction of radiation exposure by ECG-controlled tube current modulation. *Eur Radiol* 2002; **12**: 1081-1086 [PMID: [11976849](#) DOI: [10.1007/s00330-001-1278-x](#)]
- 30 **Grob A**, Thuny F, Villacampa C, Flavian A, Gaubert JY, Raoult D, Casalta JP, Habib G, Moulin G, Jacquier A. Cardiac

- multidetector computed tomography in infective endocarditis: a pictorial essay. *Insights Imaging* 2014; **5**: 559-570 [PMID: 25225108 DOI: 10.1007/s13244-014-0353-1]
- 31 **Koneru S**, Huang SS, Oldan J, Betancor J, Popovic ZB, Rodriguez LL, Shrestha NK, Gordon S, Pettersson G, Bolen MA. Role of preoperative cardiac CT in the evaluation of infective endocarditis: comparison with transesophageal echocardiography and surgical findings. *Cardiovasc Diagn Ther* 2018; **8**: 439-449 [PMID: 30214859 DOI: 10.21037/cdt.2018.07.07]
- 32 **Baumgartner FJ**, Omari BO, Robertson JM, Nelson RJ, Pandya A, Milliken JC. Annular abscesses in surgical endocarditis: anatomic, clinical, and operative features. *Ann Thorac Surg* 2000; **70**: 442-447 [PMID: 10969660 DOI: 10.1016/s0003-4975(00)01363-1]
- 33 **Graupner C**, Vilacosta I, SanRomán J, Ronderos R, Sarriá C, Fernández C, Mújica R, Sanz O, Sanmartín JV, Pinto AG. Periannular extension of infective endocarditis. *J Am Coll Cardiol* 2002; **39**: 1204-1211 [PMID: 11923047 DOI: 10.1016/s0735-1097(02)01747-3]
- 34 **Harris WM**, Sinha S, Caputo M, Angelini GD, Ahmed EM, Rajakaruna C, Benedetto U, Vohra HA. Surgical outcomes and optimal approach to treatment of aortic valve endocarditis with aortic root abscess. *J Card Surg* 2022; **37**: 1917-1925 [PMID: 35384049 DOI: 10.1111/jocs.16464]
- 35 **Habets J**, Tanis W, van Herwerden LA, van den Brink RB, Mali WP, de Mol BA, Chamuleau SA, Budde RP. Cardiac computed tomography angiography results in diagnostic and therapeutic change in prosthetic heart valve endocarditis. *Int J Cardiovasc Imaging* 2014; **30**: 377-387 [PMID: 24293045 DOI: 10.1007/s10554-013-0335-2]
- 36 **Sims JR**, Anavekar NS, Chandrasekaran K, Steckelberg JM, Wilson WR, Gersh BJ, Baddour LM, DeSimone DC. Utility of cardiac computed tomography scanning in the diagnosis and pre-operative evaluation of patients with infective endocarditis. *Int J Cardiovasc Imaging* 2018; **34**: 1155-1163 [PMID: 29450741 DOI: 10.1007/s10554-018-1318-0]
- 37 **Ouchi K**, Sakuma T, Ojiri H. Cardiac computed tomography as a viable alternative to echocardiography to detect vegetations and perivalvular complications in patients with infective endocarditis. *Jpn J Radiol* 2018; **36**: 421-428 [PMID: 29713878 DOI: 10.1007/s11604-018-0740-5]
- 38 **Sifaoui I**, Oliver L, Tacher V, Fiore A, Lepeule R, Moussafeur A, Huguet R, Teiger E, Audureau E, Derbel H, Luciani A, Kobeiter H, Lim P, Ternacle J, Deux JF. Diagnostic Performance of Transesophageal Echocardiography and Cardiac Computed Tomography in Infective Endocarditis. *J Am Soc Echocardiogr* 2020; **33**: 1442-1453 [PMID: 32981789 DOI: 10.1016/j.echo.2020.07.017]
- 39 **Jain V**, Wang TKM, Bansal A, Farwati M, Gad M, Montane B, Kaur S, Bolen MA, Grimm R, Griffin B, Xu B. Diagnostic performance of cardiac computed tomography vs transesophageal echocardiography in infective endocarditis: A contemporary comparative meta-analysis. *J Cardiovasc Comput Tomogr* 2021; **15**: 313-321 [PMID: 33281097 DOI: 10.1016/j.jcct.2020.11.008]
- 40 **Habib G**, Hoen B, Tornos P, Thuny F, Prendergast B, Vilacosta I, Moreillon P, de Jesus Antunes M, Thilen U, Lekakis J, Lengyel M, Müller L, Naber CK, Nihoyannopoulos P, Moritz A, Zamorano JL; ESC Committee for Practice Guidelines. Guidelines on the prevention, diagnosis, and treatment of infective endocarditis (new version 2009): the Task Force on the Prevention, Diagnosis, and Treatment of Infective Endocarditis of the European Society of Cardiology (ESC). Endorsed by the European Society of Clinical Microbiology and Infectious Diseases (ESCMID) and the International Society of Chemotherapy (ISC) for Infection and Cancer. *Eur Heart J* 2009; **30**: 2369-2413 [PMID: 19713420 DOI: 10.1093/eurheartj/ehp285]
- 41 **Fava AM**, Xu B. Tricuspid valve endocarditis: Cardiovascular imaging evaluation and management. *World J Clin Cases* 2021; **9**: 8974-8984 [PMID: 34786381 DOI: 10.12998/wjcc.v9.i30.8974]
- 42 **Oliveira M**, Guittet L, Hamon M. Comparative Value of Cardiac CT and Transesophageal Echocardiography in Infective Endocarditis: A Systematic Review and Meta-Analysis. *Radiol Cardiothorac Imaging* 2020; **2**: e190189 [PMID: 33778583 DOI: 10.1148/ryct.2020190189]
- 43 **Merlo A**, Chen K, Deo S, Markowitz A. Does routine preoperative computed tomography imaging provide clinical utility in patients undergoing primary cardiac surgery? *Interact Cardiovasc Thorac Surg* 2017; **25**: 659-662 [PMID: 28962500 DOI: 10.1093/icvts/ivx098]
- 44 **Gulati M**, Levy PD, Mukherjee D, Amsterdam E, Bhatt DL, Birtcher KK, Blankstein R, Boyd J, Bullock-Palmer RP, Conejo T, Diercks DB, Gentile F, Greenwood JP, Hess EP, Hollenberg SM, Jaber WA, Jneid H, Joglar JA, Morrow DA, O'Connor RE, Ross MA, Shaw LJ. 2021 AHA/ACC/AASE/CHEST/SAEM/SCCT/SCMR Guideline for the Evaluation and Diagnosis of Chest Pain: Executive Summary: A Report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *Circulation* 2021; **144**: e368-e454 [PMID: 34709928 DOI: 10.1161/CIR.0000000000001030]
- 45 **SCOT-HEART investigators**. CT coronary angiography in patients with suspected angina due to coronary heart disease (SCOT-HEART): an open-label, parallel-group, multicentre trial. *Lancet* 2015; **385**: 2383-2391 [PMID: 25788230 DOI: 10.1016/S0140-6736(15)60291-4]
- 46 **SCOT-HEART Investigators**, Newby DE, Adamson PD, Berry C, Boon NA, Dweck MR, Flather M, Forbes J, Hunter A, Lewis S, MacLean S, Mills NL, Norrie J, Roditi G, Shah ASV, Timmis AD, van Beek EJ, Williams MC. Coronary CT Angiography and 5-Year Risk of Myocardial Infarction. *N Engl J Med* 2018; **379**: 924-933 [PMID: 30145934 DOI: 10.1056/NEJMoa1805971]
- 47 **Knuuti J**, Ballo H, Juarez-Orozco LE, Saraste A, Kolh P, Rutjes AWS, Jüni P, Windecker S, Bax JJ, Wijns W. The performance of non-invasive tests to rule-in and rule-out significant coronary artery stenosis in patients with stable angina: a meta-analysis focused on post-test disease probability. *Eur Heart J* 2018; **39**: 3322-3330 [PMID: 29850808 DOI: 10.1093/eurheartj/ehy267]
- 48 **Opolski MP**, Staruch AD, Jakubczyk M, Min JK, Gransar H, Staruch M, Witkowski A, Kepka C, Kim WK, Hamm CW, Möllmann H, Achenbach S. CT Angiography for the Detection of Coronary Artery Stenoses in Patients Referred for Cardiac Valve Surgery: Systematic Review and Meta-Analysis. *JACC Cardiovasc Imaging* 2016; **9**: 1059-1070 [PMID: 27344418 DOI: 10.1016/j.jcmg.2015.09.028]
- 49 **Vahanian A**, Beyersdorf F, Praz F, Milojevic M, Baldus S, Bauersachs J, Capodanno D, Conradi L, De Bonis M, De Paulis



- R, Delgado V, Freemantle N, Gilard M, Haugaa KH, Jeppsson A, Jüni P, Pierard L, Prendergast BD, Sádaba JR, Tribouilloy C, Wojakowski W; ESC/EACTS Scientific Document Group. 2021 ESC/EACTS Guidelines for the management of valvular heart disease. *Eur Heart J* 2022; **43**: 561-632 [PMID: 34453165 DOI: 10.1093/eurheartj/ehab395]
- 50 **Hekimian G**, Kim M, Passefort S, Duval X, Wolff M, Leport C, Leplat C, Steg G, Iung B, Vahanian A, Messika-Zeitoun D. Preoperative use and safety of coronary angiography for acute aortic valve infective endocarditis. *Heart* 2010; **96**: 696-700 [PMID: 20424151 DOI: 10.1136/hrt.2009.183772]
- 51 **Saby L**, Laas O, Habib G, Cammilleri S, Mancini J, Tessonnier L, Casalta JP, Gouriet F, Riberi A, Avierinos JF, Collart F, Mundler O, Raoult D, Thuny F. Positron emission tomography/computed tomography for diagnosis of prosthetic valve endocarditis: increased valvular 18F-fluorodeoxyglucose uptake as a novel major criterion. *J Am Coll Cardiol* 2013; **61**: 2374-2382 [PMID: 23583251 DOI: 10.1016/j.jacc.2013.01.092]
- 52 **Wang TKM**, Sánchez-Nadales A, Igbinomwanhia E, Cremer P, Griffin B, Xu B. Diagnosis of Infective Endocarditis by Subtype Using <sup>18</sup>F-Fluorodeoxyglucose Positron Emission Tomography/Computed Tomography: A Contemporary Meta-Analysis. *Circ Cardiovasc Imaging* 2020; **13**: e010600 [PMID: 32507019 DOI: 10.1161/CIRCIMAGING.120.010600]
- 53 **Mahmood M**, Kendi AT, Ajmal S, Farid S, O'Horo JC, Chareonthaitawee P, Baddour LM, Sohail MR. Meta-analysis of 18F-FDG PET/CT in the diagnosis of infective endocarditis. *J Nucl Cardiol* 2019; **26**: 922-935 [PMID: 29086386 DOI: 10.1007/s12350-017-1092-8]
- 54 **Swart LE**, Gomes A, Scholtens AM, Sinha B, Tanis W, Lam MGEH, van der Vlugt MJ, Streukens SAF, Aarntzen EHJG, Bucerius J, van Assen S, Bleeker-Rovers CP, van Geel PP, Krestin GP, van Melle JP, Roos-Hesselink JW, Slart RHJA, Glaudemans AWJM, Budde RPJ. Improving the Diagnostic Performance of <sup>18</sup>F-Fluorodeoxyglucose Positron-Emission Tomography/Computed Tomography in Prosthetic Heart Valve Endocarditis. *Circulation* 2018; **138**: 1412-1427 [PMID: 30018167 DOI: 10.1161/CIRCULATIONAHA.118.035032]
- 55 **Erba PA**, Conti U, Lazzeri E, Sollini M, Doria R, De Tommasi SM, Bandera F, Tascini C, Menichetti F, Dierckx RA, Signore A, Mariani G. Added value of 99mTc-HMPAO-labeled leukocyte SPECT/CT in the characterization and management of patients with infectious endocarditis. *J Nucl Med* 2012; **53**: 1235-1243 [PMID: 22787109 DOI: 10.2967/jnumed.111.099424]
- 56 **Rouzet F**, Chequer R, Benali K, Lepage L, Ghodbane W, Duval X, Iung B, Vahanian A, Le Guludec D, Hyafil F. Respective performance of 18F-FDG PET and radiolabeled leukocyte scintigraphy for the diagnosis of prosthetic valve endocarditis. *J Nucl Med* 2014; **55**: 1980-1985 [PMID: 25453046 DOI: 10.2967/jnumed.114.141895]
- 57 **Di Salvo G**, Habib G, Pergola V, Avierinos JF, Philip E, Casalta JP, Vailloud JM, Derumeaux G, Gouvernet J, Ambrosi P, Lambert M, Ferracci A, Raoult D, Luccioni R. Echocardiography predicts embolic events in infective endocarditis. *J Am Coll Cardiol* 2001; **37**: 1069-1076 [PMID: 11263610 DOI: 10.1016/s0735-1097(00)01206-7]
- 58 **Rossi SE**, Goodman PC, Franquet T. Nonthrombotic pulmonary emboli. *AJR Am J Roentgenol* 2000; **174**: 1499-1508 [PMID: 10845470 DOI: 10.2214/ajr.174.6.1741499]
- 59 **Huang JS**, Ho AS, Ahmed A, Bhalla S, Menias CO. Borne identity: CT imaging of vascular infections. *Emerg Radiol* 2011; **18**: 335-343 [PMID: 21424803 DOI: 10.1007/s10140-011-0946-7]
- 60 **Thuny F**, Avierinos JF, Tribouilloy C, Giorgi R, Casalta JP, Milandre L, Brahim A, Nadji G, Riberi A, Collart F, Renard S, Raoult D, Habib G. Impact of cerebrovascular complications on mortality and neurologic outcome during infective endocarditis: a prospective multicentre study. *Eur Heart J* 2007; **28**: 1155-1161 [PMID: 17363448 DOI: 10.1093/eurheartj/ehm005]
- 61 **García-Cabrera E**, Fernández-Hidalgo N, Almirante B, Ivanova-Georgieva R, Noureddine M, Plata A, Lomas JM, Gálvez-Acebal J, Hidalgo-Tenorio C, Ruíz-Morales J, Martínez-Marcos FJ, Reguera JM, de la Torre-Lima J, de Alarcón González A; Group for the Study of Cardiovascular Infections of the Andalusian Society of Infectious Diseases; Spanish Network for Research in Infectious Diseases. Neurological complications of infective endocarditis: risk factors, outcome, and impact of cardiac surgery: a multicenter observational study. *Circulation* 2013; **127**: 2272-2284 [PMID: 23648777 DOI: 10.1161/CIRCULATIONAHA.112.000813]
- 62 **Heiro M**, Nikoskelainen J, Engblom E, Kotilainen E, Marttila R, Kotilainen P. Neurologic manifestations of infective endocarditis: a 17-year experience in a teaching hospital in Finland. *Arch Intern Med* 2000; **160**: 2781-2787 [PMID: 11025788 DOI: 10.1001/archinte.160.18.2781]





Published by **Baishideng Publishing Group Inc**  
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA

**Telephone:** +1-925-3991568

**E-mail:** [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)

**Help Desk:** <https://www.f6publishing.com/helpdesk>

<https://www.wjgnet.com>

