



OPEN ACCESS

A snotty nose: more than just a cold

David Lim,¹ Robert Barker,² Karyn Maltby,¹ Elizabeth George¹

¹Paediatrics, Frimley Park Hospital, Camberley, UK
²Radiology, Frimley Park Hospital, Camberley, UK

Correspondence to
 Dr David Lim,
 davidlimbn@gmail.com

Accepted 24 August 2018

DESCRIPTION

A healthy 3-year-old girl presented with a 5-day history of right-sided cheek and eye swelling and pain, associated with right-sided green nasal discharge. Examination revealed diffuse erythematous swelling around the right cheek, jawline and eye. Oral antibiotics for suspected periorbital cellulitis were ineffective. Blood inflammatory markers and ultrasound of her neck and parotid glands were normal. She was discharged as having viral parotitis. Antibiotics and nasal drops were later restarted for clinical sinusitis; however, she showed no improvement, now having 3 weeks of right-sided facial swelling with thick green and intermittently bloody nasal discharge. Further history revealed possible squint, mastication pain and breathing difficulty through the right nostril.

Cranial imaging (figure 1) demonstrated a large central but predominantly right-sided nasopharyngeal tumour, extending into the ethmoid sinuses and orbits (with medial recti distortion), infratemporal fossa and pterygoid muscles and anterior and middle cranial fossae. There was accompanying skull base destruction.



Figure 1 MRI axial T2-TSE (turbo spin echo) image showing the tumour filling the posterior nasal passages and nasopharynx which erodes the anterior border of the clivus and extends into the anterior right middle cranial fossa. There is extension into the right orbit distorting the medial rectus muscle.

Staging and biopsy confirmed the diagnosis of non-metastatic parameningeal rhabdomyosarcoma (PM-RMS). Chemoradiotherapy ensued.

RMS is the most common type of childhood soft-tissue sarcoma, mainly affecting boys at 4–6 years of age. Approximately 20% of RMS are PM, involving sites around the meninges (eg, nasopharynx, middle ear).¹ Symptoms are variable and correspond to the disease site. Pain may coexist.

Diagnosing and treating PM-RMS is challenging as it is poorly visualised and can invade critical structures. Although the PM site previously carried an unfavourable prognosis, advances in multimodality management have improved childhood survival rates.^{2,3}

Learning points

- ▶ Clinicians should consider nasopharyngeal malignancy in persistent unilateral rhinorrhea, among other symptoms.
- ▶ Para-meningeal rhabdomyosarcoma may invade critical facial and intracranial structures and hence diagnosis should not be delayed.

Contributors DL: drafted the manuscript. RB: provided the radiological images. KM, EG: edited the manuscript.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent Parental/guardian consent obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

REFERENCES

- 1 Merks JH, De Salvo GL, Bergeron C, *et al*. Parameningeal rhabdomyosarcoma in pediatric age: results of a pooled analysis from North American and European cooperative groups. *Ann Oncol* 2014;25:231–6.
- 2 Byaruhanga R, Kakande E, Buname G, *et al*. Paediatric parameningeal rhabdomyosarcoma: a case report post-multimodal treatment. *International Journal of Clinical Case Reports* 2014;4:1–5.
- 3 Rahman HA, Sedky M, Mohsen I, *et al*. Outcome of pediatric parameningeal rhabdomyosarcoma. The children cancer hospital, Egypt, experience. *J Egypt Natl Canc Inst* 2013;25:79–86.



© BMJ Publishing Group Limited 2018. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Lim D, Barker R, Maltby K, *et al*. *BMJ Case Rep* Published Online First: [please include Day Month Year]. doi:10.1136/bcr-2018-226686

Copyright 2018 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow